



## ECG Localization in ST-Elevation Myocardial Infarction: Spotting the Zebra Among the Horses

Jaskaran Singh Gujral

Abhinav Shrivastava

Follow this and additional works at: <https://www.j-saudi-heart.com/jsha>

 Part of the [Cardiology Commons](#)



This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](#).

---

# ECG Localization in ST-Elevation Myocardial Infarction: Spotting the Zebra Among the Horses

Jaskaran Singh Gujral\*, Abhinav Shrivastava

Department of Cardiology, PGIMER: Dr Ram Manohar Lohia Hospital and Post Graduate Institute of Medical Education and Research, New Delhi, India

**Keywords:** ECG Localization, ST-Elevation myocardial infarction

A 50-year-old male presented with angina pectoris at rest for four hours. On clinical examination, the patient was hemodynamically stable. A twelve-lead ECG was done (Fig. 1, Panel A) which showed a sinus rate of 72 bpm with ST-segment elevation in the precordial leads V1 to V6 and the inferior leads II, III and aVF with ST-segment depression in aVL >1 mm. In the frontal plane, the ST-segment elevation was more pronounced in lead III than lead II. The frontal ST-segment axis was 90°. The ECG post revascularization (Fig. 1, Panel B) showed resolution of ST elevation with presence of Q waves in the inferolateral leads and a distinctive absence of Q waves in the anterior precordial leads V1–V3.

The coronary angiogram (Fig. 1, Panel C, D) showed a tight stenosis in the proximal dominant right coronary artery with slow flow in the right ventricular branch and with a large posterolateral branch with no significant stenosis in the left coronary artery. Primary percutaneous coronary intervention with stenting was done, and there was restoration of flow with resolution of chest pain and ECG changes.

ST-segment elevation throughout the precordial leads is almost always interpreted as indicative of occlusion of left anterior descending artery. However, it is important to remember that when combined with ST-segment elevation in the inferior leads, a close differential is inferior wall STEMI with

predominant right ventricular infarction caused by the occlusion of the right coronary artery proximal to the right ventricular branch, as reflected in this case. The ST-elevation in leads V1–V4 may occur due to the occlusion of the right ventricular branch [1]. The ST-elevation in V5–V6 may be explained by the occlusion of a dominant right coronary artery which is also perfusing the postero-lateral wall [2]. In addition, right ventricular dilation, as was present in this case, can lead to clockwise rotation of the heart leading to ST elevation in the left precordial leads.

The clue here is to recognize that the maximum amount of ST-elevation occurs in the inferior leads. In addition, the presence of ST elevation in the right sided precordial leads, as in this case, and the absence of Q waves development in the anterior leads [3], favours the occlusion of the proximal right coronary artery with right ventricular branch involvement over the distal left anterior descending artery.

## Author's contribution

Jaskaran Singh Gujral: Conception and design of Study, Literature review, Acquisition of data, Research investigation and analysis, Data collection, Drafting of manuscript, Research coordination and management.

Abhinav Shrivastava: Conception and design of Study, Acquisition of data, Analysis and interpretation of data, Research investigation and analysis,

Received 2 December 2020; revised 2 February 2021; accepted 4 February 2021.  
Available online 15 April 2021

\* Corresponding author at: Department of Cardiology, PGIMER: Dr Ram Manohar Lohia Hospital and Post Graduate Institute of Medical Education and Research Hospital, New Delhi 110001, India.  
E-mail address: [jaskarangujral@gmail.com](mailto:jaskarangujral@gmail.com) (J.S. Gujral).






Fig. 1. Panel A: 12 Lead ECG displaying the limb leads and the left precordial leads at presentation (1 mV = 10 mm, paper speed 25 mm/s). Panel B: 12 Lead ECG displaying the limb leads and the left precordial leads post revascularization (1 mV = 10 mm, paper speed 25 mm/s). Panel C: Coronary angiogram depicting no significant stenosis in the left coronary system. Panel D: Coronary angiogram depicting tight stenosis in the proximal dominant right coronary artery with slow flow in right ventricular branch and with a large posterolateral branch.

Drafting of manuscript, Revising and editing the manuscript critically for important intellectual contents, Data preparation and presentation, Supervision of the research, Research coordination and management, Funding for the research.

#### Declaration of interest

There are no conflicts of interest.

#### References

[1] Fiol M, Carrillo A, Cygankiewicz I, Ayestarán J, Caldés O, Feral V, et al. New criteria based on ST changes in 12-lead

surface ECG to detect proximal versus distal right coronary artery occlusion in a case of acute inferoposterior myocardial infarction. *Ann Noninvasive Electrocardiol* 2004 Oct;9(4): 383–8.

- [2] Assali AR, Sclarovsky S, Herz I, Adler Y, Porter A, Solodky A, et al. Comparison of patients with inferior wall acute myocardial infarction with versus without ST-segment elevation in leads V5 and V6. *Am J Cardiol* 1998 Jan 1;81(1): 81–3.
- [3] Geft IL, Shah PK, Rodriguez L, Hulse S, Maddahi J, Berman DS, et al. ST elevations in leads V1 to V5 may be caused by right coronary artery occlusion and acute right ventricular infarction [cited 2020 Jun 27] *Am J Cardiol* [Internet] 1984 Apr 1;53(8):991–6. Available from: <https://pubmed.ncbi.nlm.nih.gov/6702712/>.