



Educational disparities in disability-free life expectancy across Europe: A focus on the East-West gaps from a gender perspective

Donata Stonkute^{a,b,*}, Angelo Lorenti^{a,c}, Jeroen J.A. Spijker^d

^a Max Planck Institute for Demographic Research, Konrad-Zuse-Strasse 1, Rostock, 18057, Germany

^b The Faculty of Social Sciences, University of Helsinki, Finland

^c Max Planck – University of Helsinki Center for Social Inequalities in Population Health, Rostock, Germany and Helsinki, Finland

^d Centre d'Estudis Demogràfics, Centres de Recerca de Catalunya (CERCA), Carrer de Ca n'Altayó, Edifici E2, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

ARTICLE INFO

Keywords:

Disability-free life expectancy
European east-west health divide
Educational inequalities

ABSTRACT

Education plays a crucial role in shaping the health outcomes of adults. This study examines the relationship between educational attainment and health across Europe. Using data from the Survey of Health, Ageing and Retirement in Europe (SHARE), we estimate educational inequalities in disability-free life expectancy (DFLE) by gender in seven Western European (2004–2019) and three Central and Eastern European (CEE) (2010–2019) countries. We exploit a novel approach that combines the Sullivan method and multivariate life tables to calculate DFLE using SHARE data. We find that educational differences in DFLE favoring the better-educated exist in both CEE and Western European countries, but also that the differences across countries are more pronounced among the low-educated. While the absolute gaps in DFLE between low- and high-educated individuals in CEE and Western European countries are similar, the educational disparities in DFLE impose a more significant burden on the CEE populations due to their overall lower life expectancy. Educational inequalities are larger among women than among men in CEE countries, while the results for Western European countries are mixed. Our findings further highlight the important role of the institutional context in mitigating or exacerbating educational inequalities in health.

1. Introduction

Health outcomes have been shown to vary widely across states in the United States and across countries in Europe, with this heterogeneity likely reflecting differences not only in demographic and epidemiological profiles, but also in contextual settings (Jagger et al., 2020; Montez et al., 2019). However, studies on health outcomes in Europe typically focus on Western European countries, while neglecting the substantial variation in health across all European regions. One of the most striking health disparities in Europe is the East-West divide, with Central and Eastern European countries having markedly lower life expectancy than the rest of the subcontinent, with a gap of almost 10 years (Eurostat, 2022). However, life expectancy alone is an insufficient measure of population health. Health expectancy estimates, which consider both mortality and morbidity information, indicate that CEE populations not only have shorter lifespans, but also spend fewer years in good health (Eikemo et al., 2008).

While CEE countries have worse health outcomes than other European countries, there are also substantial health disparities within these populations, particularly by gender and education (OECD, 2019). Women have lower mortality rates across all ages, but report poorer health and higher use of health services (Oksuzyan et al., 2008). At the same time, low education is associated with higher morbidity and mortality (Ross et al., 2012). The intersection of gender and education is also important, as educational returns to health are gendered (Ross et al., 2012).

Both gender- and education-based inequalities in life expectancy seem to be more pronounced in CEE countries than in other European regions (Mackenbach, 2017; Pinho-Gomes et al., 2022). Distinctive causal mechanisms that relate gender and education to health outcomes could be of particular importance in explaining European East-West health disparities (Geyer et al., 2006). However, studies on inequalities in health expectancies by both gender and educational attainment levels in CEE are still scarce.

* Corresponding author. Max Planck Institute for Demographic Research, Konrad-Zuse-Strasse 1, Rostock, 18057, Germany.
E-mail address: stonkute@demogr.mpg.de (D. Stonkute).

<https://doi.org/10.1016/j.ssmph.2023.101470>

Received 6 June 2023; Received in revised form 13 July 2023; Accepted 20 July 2023

Available online 22 July 2023

2352-8273/© 2023 Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Therefore, this study aims to determine the extent of gender-specific educational disparities in the duration of healthy life across European countries. More specifically, we estimate disability-free life expectancy (DFLE) at age 50 by gender and education for 10 European countries. The study also attempts to identify any emerging patterns by comparing a number of CEE countries with their Western European counterparts. Building on previous research, we hypothesize that women and populations within CEE region experience shorter DFLE and spend larger proportions of their lives with disability. Furthermore, we expect that educational disparities will exhibit a more pronounced magnitude among women and in the context of CEE countries.

Several aspects of this paper contribute to the existing literature. First, we combine two well-established estimation methods, the multivariate life table (Brown et al., 2012; Teachman & Hayward, 1993) and the Sullivan method (Sullivan, 1971), to provide an innovative approach that exploits the information available in the Survey of Health, Ageing and Retirement in Europe (SHARE) to estimate gender- and education-specific life tables and healthy life years for each country. To the best of our knowledge, this is the first paper using SHARE to estimate life tables by gender and education. In addition, estimating differential mortality and health from a single data source enables us to study countries that are traditionally overlooked in the literature: namely, the CEE countries. Thus, we also contribute to the discussion on how educational inequalities in health influence the European East-West divide.

2. Background

The differential impact of gender on mortality has been widely recognized, with women exhibiting lower mortality rates than men across all age groups (Oksuzyan et al., 2008). Despite women generally having a longer life expectancy than men, it is often assumed that they experience poorer overall health (Author 2020; 2022). This phenomenon may be due, at least in part, to the tendency of women to report poorer health, a greater burden of morbidity and disability, and more frequent use of health services (Nusselder et al., 2010). The observation that women have both a mortality advantage and a health disadvantage, which is also referred to as the female-male health-survival paradox, is often explained by citing intricate connections between gender-related social, behavioral, and biological factors (Oksuzyan et al., 2008). In addition, women's longevity advantage itself contributes to their health disadvantage, as it results in a larger proportion of life spent in poor health (Luy & Minagawa, 2014). However, the magnitude of gender disparity varies significantly, and it is important to note that female disadvantage in health does not hold true uniformly across countries or health indicators selected (Crimmins et al., 2019). Moreover, despite the fundamental importance of gender in health, prompting studying health outcomes for men and women separately, health disparities are not limited to differences by gender.

Socioeconomic factors are also considered key determinants of health differences within populations. Evidence suggests that an individual's social standing within a hierarchical structure, as determined by factors such as income, occupation, and education, can either ameliorate or exacerbate their health prospects (Siegrist & Marmot, 2004). Education, in particular, holds unique value as an indicator of socioeconomic status. It is a reliable and easily accessible indicator that tends to remain relatively stable after individuals complete their formal education. Moreover, education is fundamental in establishing both income and occupation. Notably, the impact of education on health outcomes persists even after adjusting for factors such as wealth and occupation (Cutler & Lleras-Muney, 2010). While this is an important empirical piece of evidence, theoretical support reinforces the importance of education when examining health disparities.

Higher levels of education directly contribute to better health, beyond a simple correlation. Ross and Mirowsky (2010) argued that the effects of education are embedded in the nexus between two main

pathways: commodity and human capital. Commodity theories posit that having a higher level of education provides individuals with advantages, such as access to better employment, higher income, and health insurance. Human capital theories posit that beyond the financial gains, education promotes health by equipping individuals with the knowledge and resources they need to adopt health-promoting behaviors. However, it is essential to recognize that as well as fostering favorable conditions for good health, education also plays a pivotal role as a social determinant of inequalities in health (Ross et al., 2012).

While the pathways through which gender and education act upon health are distinct, it is also crucial to consider the intersection between the two. Examining the intersection between education and gender inequalities in European countries requires a multi-dimensional understanding of the various social, economic, and cultural factors that shape health opportunities and outcomes for individuals of different genders and educational levels. Moreover, even as gender intersects with education in shaping health, there are country-specific differences in health inequalities by both gender and education (Mackenbach, 2017; Romero-Ortuno et al., 2014).

Gender differences in health vary across countries (Crimmins et al., 2011). While the differences in life expectancy between men and women are large in many Western and Southern countries, they are relatively small compared to those in Eastern European countries (Author 2020). Due to their significant mortality disadvantage, men in CEE countries are, on average, expected not only to have shorter lifespans, but also to experience fewer healthy years of life than women (Van Oyen et al., 2013).

Educational returns to health also differ across Europe (Mackenbach et al., 2018; Valverde et al., 2021), with the disparities being largest in CEE countries and smallest in Southern European countries (Mackenbach, 2017). For instance, the life expectancy difference between men with high and low education is more than 10 years in CEE countries, while it is approximately four years in Sweden, Italy, and the Netherlands (OECD, 2019). Although previous literature has primarily focused on mortality, recent studies have observed a more pronounced educational gradient in transitioning to both poor health and recovery among CEE populations (Solé-Auró & Gumà, 2022). In addition, CEE countries have been consistently shown to have the lowest health expectancies in Europe (Jagger et al., 2008, 2011), contributing to the concept of the European East-West health divide (Carlson, 2004).

Both the health of the general population and disparities by gender and education are associated with country-specific macro-level indicators (Eikemo et al., 2008; Van Oyen et al., 2010). Therefore, the persistent health divide between East and West might reflect the role of social structures, as conceptualized by the institutional theory of health inequalities (Beckfield et al., 2015). The theoretical framework outlining the effects of welfare states on social health inequalities has suggested that sociohistorical and policy contexts may configure a scenario in which health gains follow educational attainment (Ross et al., 2012). CEE is a particularly interesting region to focus on when studying this association due to its history of state socialism, early educational expansion, historically high levels of female employment, and atypical relationship between education and income (Haggard & Kaufman, 2009). However, these positive factors may be undermined by the lack of progress in the region across many health indicators, such as mortality from preventable causes (Mackenbach et al., 2015).

Despite the evidence of substantial gender and educational health disparities within CEE countries, the current research on this topic is fragmented, and shares several limitations. Existing studies primarily focused on population-level estimates and the period immediately following the fall of communism (Carlson, 1998; Shkolnikov et al., 2006), thereby failing to account for more recent developments. Furthermore, previous research predominantly focused on mortality outcomes, which offer limited insights into the overall health status of the population (Minagawa, 2013).

A growing interest has emerged in applying measures that extend

beyond mortality to gain a comprehensive understanding of population health (Robine & Jagger, 2003). One such measure is disability-free life expectancy (DFLE), which is defined as the average duration of a person's life without health limitations within a given population (Robine et al., 1999). DFLE is derived from disability indicators such as the Global Activity Limitation Indicator (GALI), and when calculated on this basis, it is known as healthy life years. Although disparities in DFLE by gender and education have been extensively documented (Mäki et al., 2013; Solé-Auró et al., 2015), limited research has been conducted on differential health expectancy for CEE countries. Thus, whether gender and educational gaps in disability-free life expectancy display a consistent pattern across European countries, as has been observed for overall life expectancy, remains an open question.

Unfortunately, however, there is not enough data available to apply traditional methods of estimating health expectancies for a wide range of countries by gender and educational attainment (Avendano et al., 2009; Jagger et al., 2020). Health expectancy measures rely on access to vital statistics, registries, censuses, and health prevalence data. National statistical offices provide data on age- and gender-specific mortality, which are then standardized and disseminated by organizations like the European Union's statistical office Eurostat and databases such as the Human Mortality Database (HMD). However, multivariate life tables by education are scarce, except for a few countries like the Nordic European countries and Spain (Klotz et al., 2017). This is because mortality records linked to population databases are not broken down by educational level, and have different time periods and stratification variables.

This study seeks to address the deficits in the literature by estimating gender- and education-specific DFLE for 10 European countries. The added value of this study is twofold. First, we combine two established estimation methods – the multivariate life table (Brown et al., 2012; Teachman & Hayward, 1993) and the Sullivan method (Sullivan, 1971) – to obtain DFLE estimates. This approach allows us to use SHARE as a single data source. To support this approach, we also generate gender-specific life tables to use for comparisons with the HMD and European Health and Life Expectancy Information System (www.eurohex.eu; henceforth EurOhex data), and to identify any discrepancies. Second, our study contributes to the literature on health disparities across Europe, specifically between CEE and Western European countries, in two ways. First, we analyze health disparities in CEE countries, which have been under-represented in prior research. Second, we shift our focus from mortality to health expectancy, estimating remaining disability-free life expectancy at age 50.

3. Data and methods

3.1. Data

This study uses data from the Harmonized Survey of Health, Ageing and Retirement in Europe (SHARE) developed by the Gateway to Global Aging Data (<https://g2aging.org/>). SHARE is a biennial survey conducted since 2004 that provides cross-national longitudinal information on demographic, socioeconomic, and health measures, as well as vital status data. The survey covers private households with members aged 50 or older. Our study employs comprehensive information on health and survival, such as the age at the time of death and the health status of participants. Depending on when each country initially participated in the survey, we combined data from either the 1st (2004) or the 4th wave (2010) to the 8th (2019) wave of SHARE. We excluded wave 3, which is called SHARELIFE, because it only focuses on people's life histories, rather than relying on a regular questionnaire. Additional information about the SHARE project and waves used (Börsch-Supan, 2022a, 2022b, 2022c, 2022d, 2022e, 2022f), as well as sampling details for each country (Bergmann et al., 2019) can be found elsewhere.

Three out of the 10 countries studied are located in Central and Eastern Europe (CEE): Estonia, Czechia and Slovenia. They therefore include one Baltic country, one Central European and one Southeast

European country that are known for having a shared sociohistorical context and welfare regime (Eikemo & Bambra, 2008). The remaining analyzed countries were classified as Western European countries.

We aimed to maximize the number of countries included in our study. However, some countries, particularly those in the CEE region, participated in fewer waves. To ensure consistency, we included only those CEE countries that have participated in the survey since at least wave 4, and that have not missed more than one consecutive round of participation. The seven Western European countries were selected on the same criteria for comparative purposes. We dropped individuals with missing data on the variables of interest or with a baseline age below 50. Additional restrictions were applied to individual parts of the statistical analysis.

3.2. Indicators

In this study, we used SHARE's single-question measure of functional limitation or disability (GALI) as our health indicator. GALI is a reliable and valid tool for measuring function and disability, and can be used for cross-country comparisons. Specifically, this indicator refers to the observed prevalence of any long-standing activity limitations, which could denote general health problems or limitations in engaging in activities people usually perform (Robine & Jagger, 2003; Van Oyen et al., 2006). In the harmonized dataset, the responses were re-coded by identifying individuals as healthy if they reported no limitations in usual activities, and as limited if they reported any level of limitation. We also use interview status and year of death variables to obtain survival information.

The level of educational attainment was determined using a harmonized measure based on the International Standard Classification of Education (Statistics, 2012). This measure categorized individuals into three groups: 1. lower secondary education or less (ISCED 0–2); 2. upper secondary or vocational training (ISCED 3–4); and 3. tertiary education (ISCED 5–6). Sample characteristics are presented in Table 1.

3.3. Analysis

In this study, we used a multivariate life table approach (Teachman & Hayward, 1993) to create gender- and education-specific life tables. Following an approach similar to that used in Brown et al. (2012), we employed a set of discrete-time regression models for each country, stratified by gender, to estimate the probability of death by age, gender, and education. To estimate this probability, survival information was needed for at least two consecutive time points for each individual. The approach assumed that individuals are alive at age t and may die at age $t+1$. This implied that age was the time scale we used in the models. The population discrete-time hazard was defined as the probability of death occurring in the current or future time period, represented by $h(t_{ij}) = \Pr(T_i = j | T_i \geq j)$, where T_i is a random variable indicating the time period j when individual i dies.

The models were as follows:

$$\ln[-\ln(1 - h(t_j|x_i))] = \alpha_j + x_i'\beta \quad (1)$$

where t_j is the time interval, α_j is the complementary log-log transformation of the baseline hazard and x_i' are a set of covariates: namely, age, age-squared, and educational level, specific to the individual i . Age was added with a second-degree polynomial; the quadratic term accounted for nonlinearities in the age pattern of deaths. The models were run separately for men and women, and for each country.

To predict death probabilities by country, gender, education, and age, we used two-year intervals to reflect the biannual panel data structure. We obtained gender- and education-specific life tables from estimated survival probabilities by simulating the death of a synthetic cohort of 500,000 individuals. In the simulation, each person in the age interval x survives to $x + n$ if the predicted probability of death is lower

Table 1
Sample characteristics and educational composition for men and women aged 50 and older.

	Years	Sample size	Proportion of men	Proportion across educational levels					
				Women			Men		
				Low	Medium	High	Low	Medium	High
Central and Eastern Europe									
Estonia	2010–2019	21 514	0.40	0.27	0.49	0.23	0.30	0.47	0.22
Czechia	2010–2019	18 141	0.41	0.44	0.46	0.11	0.39	0.44	0.17
Slovenia	2010–2019	12 111	0.43	0.42	0.43	0.15	0.25	0.58	0.17
Western Europe									
Austria	2004–2019	16 371	0.42	0.34	0.45	0.21	0.15	0.55	0.31
Belgium	2004–2019	24 572	0.45	0.44	0.27	0.29	0.39	0.26	0.35
Denmark	2004–2019	15 016	0.47	0.24	0.33	0.43	0.14	0.48	0.39
France	2004–2019	18 906	0.43	0.49	0.31	0.20	0.38	0.39	0.23
Italy	2004–2019	19 174	0.46	0.74	0.20	0.06	0.70	0.22	0.08
Spain	2004–2019	21 771	0.45	0.84	0.08	0.08	0.78	0.11	0.11
Sweden	2004–2019	16 664	0.46	0.39	0.30	0.31	0.43	0.30	0.27

Source: Own calculations, using pooled data from SHARE.

than a random draw from a uniform distribution. We repeated this procedure iteratively from ages x to $x + n$ for age intervals [50–52], [52–54], ..., 110, and obtained life table quantities, such as the number of survivors at the corresponding ages. The period life table was constructed following standard steps (Preston et al., 2001). Life tables were closed at age 90+ due to limited observations at the oldest ages.

Finally, remaining DFLE at age 50 was estimated using the Sullivan method (Sullivan, 1971), following that:

$$e_{50}^{DF} = \frac{1}{l_{50}} \sum_{x=50}^{x=90+} (1 - n\pi_x) \cdot nL_x \tag{2}$$

where n is the length of the age interval, $n\pi_x$ is a proportion of individuals with health limitations between ages x and $x + n$, l_{50} is a number of survivors at age 50 and nL_x is a number of person-years lived between ages x and $x + n$. The idea underlying the method is to apply age-specific proportions of a population in an unhealthy state to the age-specific person-years lived from the life table. Summing up only the disability-free person-years lived across all ages yielded the DFLE. The age-specific proportion of people with health limitations within subgroups was determined using a sample of individuals who were still alive. Confidence intervals for LE and DFLE were obtained using a nonparametric bootstrap approach (Cameron & Trivedi, 2005) by resampling the data 1000 times. All the analyses were conducted using R language (version 4.1.1).

We utilized a distinctive approach that capitalized on the availability of mortality data within the SHARE dataset. However, life tables calculated from survey data may differ considerably from those generated using high-quality data sources. To support our findings, we compared life expectancy estimates at the population level with HMD and EurOhex data, with the aim of identifying any substantial differences. More details on how we constructed comparative life tables using HMD data can be found in the supplementary material.

4. Results

Table 1 shows characteristics of the working sample, categorized by educational level and country for men and women. The sample included 184,240 person-wave observations of individuals aged 50+ across 10 countries. It should be noted that the data for Central and Eastern Europe countries cover the years 2010–2019, whereas the data for Western European countries cover the years 2004–2019. Estimates for both life and disability-free life expectancies are reflective of the mid-points of these intervals, namely 2015 and 2012, respectively.

Although CEE countries joined SHARE later, the number of observations from CEE countries was not substantially lower than that from Western European countries, except for Slovenia, which had the smallest number of observations.

The distribution of individuals across educational levels varied by country and gender. Most CEE countries had a large share of individuals with upper secondary or vocational training, while Western European countries were more heterogeneous, with some having a large share of individuals with low education (e.g., Spain and Italy), and others having an evenly distributed educational composition across all three levels (e.g., Denmark and Sweden). Finally, as expected, most countries had a larger proportion of women than of men in the lowest education category.

Table 2 displays data on the proportions of individuals experiencing long-term limitations in usual activities due to health problems (indicated by the GALI indicator), categorized by education and country for men and women. Women, and especially those with low education, were more likely than men to report limitations. Significant gender gaps were observed among individuals with medium education in Denmark and Sweden, with a 9–10% difference in the prevalence of health limitations. Our results showed a consistent educational pattern across all analyzed countries, with a lower proportion of individuals in the high education group reporting limitations. With the exception of Slovenia, CEE countries had the highest prevalence of health limitations. For example, up to 76% of low-educated women in Estonia reported limitations, while the Southern European countries, and particularly Spain and Italy, had the lowest prevalence of health limitations.

4.1. Life and health expectancies at the population level

Fig. 1 shows estimates of remaining life expectancy at age 50 for men and women for all the analyzed countries compared with the HMD and

Table 2
Proportions of women and men aged 50 and older experiencing long-term limitations in usual activities due to a health problem by educational level.

	Women			Men		
	Low	Medium	High	Low	Medium	High
Estonia	0.76	0.61	0.50	0.69	0.59	0.50
Czechia	0.66	0.50	0.46	0.62	0.55	0.48
Slovenia	0.63	0.46	0.37	0.63	0.47	0.37
Austria	0.62	0.48	0.43	0.63	0.47	0.43
Belgium	0.56	0.48	0.44	0.49	0.43	0.38
Denmark	0.55	0.46	0.33	0.48	0.36	0.32
France	0.55	0.40	0.36	0.52	0.40	0.31
Italy	0.51	0.32	0.25	0.44	0.30	0.26
Spain	0.50	0.24	0.25	0.41	0.27	0.24
Sweden	0.53	0.46	0.40	0.47	0.37	0.34

Note: The data used for the calculations were pooled from country-specific waves of SHARE from either 2004–2019 or 2010–2019, based on country-specific data availability.




Fig. 1. Remaining life expectancy at age 50 and their 95% confidence intervals obtained from SHARE, compared to estimates obtained from HMD and EurOhex. Note: HMD estimates are based on the calendar year that represented the midpoint population given the period of the country’s participation in the survey. For Austria, Belgium, Denmark, France, Italy, Spain, and Sweden, we used life tables from 2012. For Czechia, Estonia, and Slovenia, we used life tables from 2015. In addition, we used EurOhex estimates from 2012, as it is the latest year available in the database for all countries.

EurOhex data. While population-level estimates were not the main focus of this study, we included them because they not only provided an overview of the health status of the population, they also allowed for comparisons with data from other reliable sources. In most cases, the HMD and EurOhex estimates fell within the 95% confidence intervals of our estimates or were within the two-year difference, except for Danish and Slovenian women.

In addition, estimates at the population level revealed some important patterns. First, we did not observe greater between-country disparities in disability-free life expectancy (DFLE) than in overall life expectancy at age 50. Second, we found that individuals in CEE countries, and particularly men, spent a much smaller share of their life disability-free than their Western European counterparts. Complete information on estimates at the population level can be found in the [supplementary material Table A1](#).

4.2. Life and health expectancies by gender and education

Our findings indicated that women in all educational groups could

expect to live longer than their male counterparts across all of the countries studied. The differences were striking: in most of the studied countries, the highest-educated men could expect to live fewer than or the same number of years as the least-educated women. The full set of estimates by gender and education is provided in [supplementary materials Table A2](#).

When considering life expectancy free of disability, we found that women retained their advantage compared to men in the majority of countries, except for Belgium and Sweden. Central and Eastern European (CEE) countries generally had a wider gender gap in life expectancy without disability, especially among individuals with medium education. Women in CEE countries with medium education could expect to live 2–3 years longer without disability than their male counterparts.

Fig. 2 displays DFLE estimates by gender and education for the studied countries. Higher educational levels were associated with a longer disability-free life, but the differences between the highest and the lowest educational levels varied greatly by gender and across countries. Among women, Belgium had the smallest educational gap in




Fig. 2. Remaining disability-free life expectancy at age 50 by gender and education.

DFLE (3.7 years), while Slovenia and Italy had the largest differences (10 years). Among men, Belgium had the smallest educational gap in DFLE (3.6 years), while Slovenia and Austria had the largest differences (9.2 and 8.4 years, respectively). For Southern European and CEE countries (except for Spanish and Czech women), the health benefits of higher education were particularly noteworthy, as there was a marked increase in the differences in DFLE between individuals with medium and with high educational levels compared to the differences in DFLE between individuals with low and medium educational levels.

Although the gaps in DFLE between individuals with low and high educational levels were significant, the absolute disparities within CEE countries were not greater than those found in Western European countries. To illustrate, the difference in DFLE between men with low and high

educational levels was approximately six years in Denmark, Italy, and France, which was comparable to the gap in Czechia. Furthermore, among men in Estonia, this gap was narrower by one year. However, it is important to interpret these results in relative terms; specifically, in terms of the proportion of the overall life expectancy that an individual could expect to live disability-free.

Higher education was associated with a greater proportion of disability-free life, but the size of this advantage varied by gender and by country. The estimates indicated that individuals in Italy and Spain had the highest shares of disability-free life at age 50 across all educational groups. Notably, the estimates showed that high-educated men in Italy and Spain could expect to live 72–73% of their remaining years of life at age 50 disability-free. In contrast, the shares of disability-free life at age 50 were smallest for individuals in CEE countries (with the exception of medium- and high-educated women in Slovenia) and Austria. Across all educational groups and irrespective of gender, individuals living in CEE countries and Austria could expect to live less than 50% of their remaining years of life at age 50 free of disability. Generally, women could expect to spend a smaller share of their life without health limitations. For example, while low-educated Italian women could expect to live 60% (18.7 years) of their remaining years of life at age 50 disability-free, high-educated Estonian women could expect to live less than half (46%; 14.5 years) of their remaining years of life at age 50 free of disability.

The estimates indicated that Danish, Italian, and French men had a remaining DFLE at age 50 of 17.4–19.4 years. By contrast, Czech and Estonian men could expect to live 12.8 and 11.0 years, respectively, of their remaining life at age 50 disability-free. These findings imply that the six-year difference in DFLE between low and high educational groups mentioned earlier was proportionally smaller in Western European countries, and that the educational disparities in DFLE imposed a more significant burden on the populations in the CEE countries.

5. Discussion

Health disparities based on gender and education in Europe have been widely studied, but limited data has made it challenging to estimate these disparities in some countries. To address this issue, researchers have been compelled to adapt existing methodological approaches, and to develop new and original approaches. Our study emphasized the potential of using SHARE as a single source of data for differential life and health expectancy estimation, in particular for countries for which no information is available from other sources. We combined two established estimation techniques – discrete-time modeling and simulation – to estimate multivariate life tables, and the Sullivan method to estimate disability-free life expectancy.

The results of this paper first point to important gender differences, in line with prior research (Pinho-Gomes et al., 2022). Across all educational groups, women were shown to live longer than men, with the highest-educated men living fewer or the same number of years as the least-educated women in their respective countries. However, the differences narrowed in the DFLE estimates, which is consistent with the

female-male health-survival paradox (Oksuzyan et al., 2008). Women's health disadvantage may be due to limited access to resources such as income, education, and health care (Tawiah et al., 2021). On the other hand, women exhibit higher levels of compliance with treatments (Christensen et al., 2009), and greater resilience when living with chronic diseases (Kiely et al., 2019). Given the complexity of gender differences, it is vital to approach their interpretation with caution, recognizing that their manifestation is influenced by social, behavioral, and epidemiological contexts (Crimmins et al., 2019). Our findings, consistent with previous research, uncovered an educational gradient in health outcomes. The advantage associated with high education was first reflected in the prevalence of health limitations, as the findings indicated that individuals with higher educational levels tended to report fewer limitations than those with lower educational levels. Health disparities by education were also observed in the estimates of DFLE, but the extent of these disparities was not exclusively driven by the disadvantage associated with low education.

In certain countries, particularly in Southern European and Central and Eastern European countries, the differences in DFLE between individuals with medium and high educational levels were found to be much greater than the differences in DFLE between individuals with low and medium educational levels. The fact that the DFLE of the medium educated was barely higher in some instances than that of the lower educated could be because these social disadvantages extend to the middle class and mainly the high educated are able to obtain good employment conditions with higher income, health insurance, as well as the knowledge and resources they need to adopt health-promoting behaviors (Montez et al., 2018). Previous research has suggested that national contexts might be able to modify these patterns (Cambois et al., 2016). In Southern European countries, the proportion of high-educated individuals was still relatively small among the oldest members of the studied cohorts. This is likely because educational expansion occurred later in these countries than in the other European countries studied, resulting in a sub-population of high-educated individuals, and particularly of men, that is more selective. Meanwhile, in CEE countries, where education is more evenly distributed, institutional differences may have been more important for the more affluent. Given the socio-historical context, individuals with low educational attainment in the CEE countries tended to have consistently lower DFLE than their Western European counterparts. This could be due to the cumulative effects of exposures tied to social disadvantages, such as low education, across the life course. However, the degree to which additional levels of education were associated with health appeared to differ across countries. This finding is supported by the institutional theory of health inequalities (Beckfield et al., 2015) and empirical evidence of differential educational returns to health from other geographical contexts (Montez et al., 2019). The empirical findings of this study are also consistent with this theoretical contextualization, as the education-specific differences observed across the studied countries were more pronounced among the low-educated.

A particular interest of this study was investigating the differences between CEE and Western European countries. Additional insights into the East-West divide in Europe, consistent with prior research, emerged from our examination of gender-specific educational disparities in DFLE across European countries. First, a substantial gender gap in DFLE favoring women was observed in CEE countries, particularly among individuals with medium levels of education. Second, with a few exceptions (in particular Slovenia), the three CEE countries were shown to have lower levels of disability-free life expectancy (DFLE). Thus, the findings indicated that regardless of their educational level or gender, individuals in these CEE countries spent a smaller percentage of their remaining life disability-free than their counterparts in the seven analyzed Western European countries. However, only the lowest educated in CEE had a lower DFLE than in any of the Western European country. Third, although the differences in DFLE between high and low educational groups were not necessarily larger in CEE countries than in

Western European countries, the relative disparities by educational attainment were more significant in CEE countries due to their generally lower level of life expectancy.

Finally, an examination of the intersection of gender and education also produced a distinctive pattern, with women in CEE countries exhibiting greater educational disparities in DFLE than their male counterparts. Meanwhile, the findings for Western European countries were more mixed. Drawing upon prior research, we know that men are more prone to experiencing life-threatening conditions, whereas women are more susceptible to impairing but rarely fatal conditions (Crimmins et al., 2019). Moreover, it has been acknowledged that the relationship between education and health is influenced by the institutional framework in place (Beckfield et al., 2015; Ross & Mirowsky, 2010). Therefore, the combination of biological and social differences between men and women, along with the institutional and epidemiological context of countries, is likely to contribute to the gendered patterns of educational inequalities observed. Nonetheless, while there is a clear educational gradient, the educational returns to DFLE is heterogeneous across countries. For instance, compared to the low- and medium-educated groups, DFLE was much higher for the high educated in Czechia (men only) and Estonia. Conversely, in the case of Czech women DFLE was substantially lower for the low-educated group compared to those who were medium- and high educated. While it is difficult to find a precise explanation for the fact the lowest educated in CEE have a lower DFLE than in Western Europe, the cumulative effects of exposures tied to social disadvantages are likely to play a role, as mentioned earlier in the discussion.

Notable heterogeneity was found across other findings. Slovenia was shown to have exceptionally high DFLE among high-educated men and women, even compared to most Western European countries. However, due to the small sample size for this country, the confidence intervals are wide, meaning that these estimates should be interpreted with caution. Austria was also found to have wide educational disparities. Especially the low- and medium-educated underperformed in DFLE compared to other Western European countries. In line with our findings, recent reports have also indicated a high prevalence of limitations in daily activities and chronic conditions (European Commission, 2022; OECD, 2019), as well as a large proportion of life spent in an unhealthy state in Austria (Welsh et al., 2021). Denmark was also shown to have low DFLE values, particularly among women, which could be attributed to an exceptionally high prevalence of smoking among the interwar cohort in that country (Lindahl-Jacobsen et al., 2016).

5.1. Methodological strengths and limitations

Our study proposed combining two estimation approaches: discrete-time modeling and simulation for estimating multivariate life tables, and the Sullivan method for estimating health expectancies using SHARE data. Using SHARE as a sole data source in estimating educational gradient in DFLE is not only an original alternative to the existing methods, it can also offer insights into DFLE trends for countries that currently cannot provide differential mortality statistics, which is a crucial component of DFLE estimation. We also estimated health and life expectancies for countries that are otherwise under-represented in the literature, which is especially relevant in light of the persisting European East-West health divide.

Furthermore, unlike vital statistics, the SHARE survey provides comprehensive information on the health and demographic and socio-economic characteristics of individuals. Thus, SHARE data can be used to investigate dimensions other than education, which can help to advance our understanding of health disparities across and between CEE and Western European countries. At the same time, we acknowledge several data-related limitations. Klotz et al. (2017) have extensively discussed the validity concerns associated with survey data, such as the under-representation of institutionalized populations and the possible misrepresentation of differential mortality between subgroups. Indeed,

household surveys like SHARE usually do not cover institutionalized populations, which might result in a selectively healthier sample. The proportion of institutionalized individuals varies across countries, adding complexity to cross-country comparisons (Permanyer & Spijker, 2020). Additionally, research indicates that respondents in SHARE panel are positively selected in regards to education and physical health, factors that significantly impact sample attrition and mortality (Schulz & Doblhammer, 2011). However, Muszynska-Spielauer and Spielauer (2022) analyzing a larger number (and more recent) waves found only negligible differences in the effects of panel attrition on the education-specific prevalence of disability and DFLE estimates. Nevertheless, future research, especially focusing on cross-sectional estimates of health expectancies would benefit from including weights that account for variations in educational composition between the survey sample and the general population (Spitzer, 2020), as well as across different countries (Sauerberg, 2021).

The use of the Global Activity Limitation Indicator (GALI) as a measure of health has limitations due to potential differences across cultures in self-reported disability (Permanyer & Spijker, 2020). Additionally, in 2008, the translations of the question into different languages was revised. To address this issue, sensitivity analyses were conducted using only waves 4–8 and an alternative health indicator (limitations in at least one basic activity of daily living) (see supplementary material Fig. A1-A3). Finally, it is important to note that the selection of survey may significantly impact the prevalence of GALI, educational disparities in GALI (Rubio-Valverde et al., 2019).

Albeit with some exceptions, the SHARE data produced population-level life expectancy estimates that were similar to the Human Mortality Database estimates. The biannual nature of the panel and the exclusion of institutionalized populations should be considered when evaluating the quality of these estimates. Furthermore, our findings on the association between education and health were consistent with the results of previous research showing that longevity and health tend to be positively associated with education (Montez & Friedman, 2014). Our findings of relative differences in disability-free life expectancy by gender and education, as well as generally lower DFLE and relatively larger educational inequalities within CEE countries as compared to other European countries fall in line with the previous literature. For instance, Permanyer and Bramajo (2022) showed that CEE countries have been underperforming in health-adjusted life expectancy for decades, while Solé-Auró and Gumà (2022) provided evidence of more pronounced educational inequalities in transitioning to poor health in CEE. Although the study period and the age range for reporting remaining years of disability-free life vary significantly (Majer et al., 2011; Mäki et al., 2013), s (Majer et al., 2011; Mäki et al., 2013), the relative differences in disability-free life expectancy by gender and education are larger in countries considered belonging to the CEE region.

6. Conclusions

The study found substantial health disparities by education in both Central and Eastern European and Western European countries, with the largest cross-country differences being observed among the low-educated. Although the differences in disability-free life expectancy between the high and the low educational groups were not necessarily larger in CEE countries, the relative disparities by educational attainment were more significant due to the generally lower level of life expectancy in these countries. In addition, the educational inequalities in CEE countries were found to be more pronounced for women than for men, whereas the results for Western European countries were inconclusive. These findings underline the need for further research on how individual factors impact health that also considers the contextual environment in which people live their lives. The results of this study may advance our understanding of the factors that drive educational health inequalities, and the effectiveness of national policies designed to address them.

Authors' contributions

All authors contributed to the study conception and design. Data management: DS; Methodology: DS, AL; Formal analysis and investigation: DS; Visualization: DS; Writing - original draft preparation: DS; Writing - review and editing: DS, AL, JS; all authors read and approved the final manuscript.

Funding

Angelo Lorenti was supported by grants to the Max Planck – University of Helsinki Center from the Max Planck Society, Jane and Aatos Erkkö Foundation, Faculty of Social Sciences at the University of Helsinki, and Cities of Helsinki, Vantaa and Espoo.

Jeroen J.A. Spijker was supported by the research project “The health of people in advanced ages: the analysis of comorbidity, multiple causes of death and gender and socioeconomic inequalities in health (COMORHEALTHSES)” (IP: JS), funded by the Spanish Ministry of Science and Innovation (PID2020-113934RB-I00), the European Research Council (ERC-2019-COGAgreement No 864616, HEALIN) and the Catalan Government under the CERCA programme.

Ethical statement regarding the use of secondary data

This study was conducted with harmonized data from the Survey of Health and Retirement in Europe. No ethical approval was needed for conducting the research.

After filling in a user statement, every person with scientific affiliation can download the data for free as long as the data are used for purely scientific purposes.

The SHARE study is subject to continuous ethics review. During Waves 1 to 4, SHARE was reviewed and approved by the Ethics Committee of the University of Mannheim. Wave 4 and the continuation of the project were reviewed and approved by the Ethics Council of the Max Planck Society. In addition, the country implementations of SHARE were reviewed and approved by the respective ethics committees or institutional review boards whenever this was required. The numerous reviews covered all aspects of the SHARE study, including sub-projects and confirmed the project to be compliant with the relevant legal norms and that the project and its procedures agree with international ethical standards.

Declaration of competing interest

None declared.

Data availability

All study code is fully replicable and available in the following GitHub repository: https://github.com/DonataStonkute/dfle_50.

This paper uses data from SHARE Waves 1, 2, 4, 5, 6, and 8 (DOIs: 10.6103/SHARE.w1.800, 10.6103/SHARE.w2.800, 10.6103/SHARE.w4.800, 10.6103/SHARE.w5.800, 10.6103/SHARE.w6.800, 10.6103/SHARE.w8.800) see Börsch-Supan et al. (2013) for methodological details.(1) The SHARE data collection has been funded by the European Commission, DG RTD through FP5 (QLK6-CT-2001-00360), FP6 (SHARE-I3: RII-CT-2006-062193, COMPARE: CIT5-CT-2005-028857, SHARELIFE: CIT4-CT-2006-028812), FP7 (SHARE-PREP: GA N°211909, SHARE-LEAP: GA N°227822, SHARE M4: GA N°261982, DASISH: GA N°283646) and Horizon 2020 (SHARE-DEV3: GA N°676536, SHARE-COHESION: GA N°870628, SERISS: GA N°654221, SSHOC: GA N°823782, SHARE-COVID19: GA N°101015924) and by DG Employment, Social Affairs & Inclusion through VS 2015/0195, VS 2016/0135, VS 2018/0285, VS 2019/0332, and VS 2020/0313. Additional funding from the German Ministry of Education and Research, the Max Planck Society for the Advancement of Science, the U.S. National Institute on

Aging (U01_AG09740-13S2, P01_AG005842, P01_AG08291, P30_AG12815, R21_AG025169, Y1-AG-4553-01, IAG_BSR06-11, OGH4_04-064, HHSN271201300071C, RAG052527A) and from various national funding sources is gratefully acknowledged (see www.share-project.org).

This analysis uses data or information from the Harmonized Survey of Health, Ageing and Retirement in Europe (SHARE) dataset and Codebook, Version F as of February 2023 developed by the Gateway to Global Aging Data. The development of the Harmonized SHARE data was funded by the National Institute on Aging (R01 AG030153, RC2 AG036619, 1R03AG043052). For more information, please refer to www.g2aging.org.

Acknowledgements

DS gratefully acknowledges the resources made available by the International Max Planck Research School for Population, Health and Data Science (IMPRS-PHDS).

The authors acknowledge Lara Bister and Sneha Sarah Mani (International Max Planck Research School for Population, Health and Data Science) for their helpful comments on early versions of this manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ssmph.2023.101470>.

References

- Avendano, M., Glymour, M. M., Banks, J., & Mackenbach, J. P. (2009). Health disadvantage in US adults aged 50 to 74 years: A comparison of the health of rich and poor Americans with that of Europeans. *American Journal of Public Health*, *99*(3), 540–548.
- Beckfield, J., Bamba, C., Eikemo, T. A., Huijts, T., McNamara, C., & Wendt, C. (2015). An institutional theory of welfare state effects on the distribution of population health. *Social Theory & Health*, *13*(3), 227–244.
- Bergmann, M., Kneip, T., De Luca, G., & Scherpenzeel, A. (2019). *Survey participation in the Survey of Health, Ageing and Retirement in Europe (SHARE), Wave 1-7. Based on Release 7.0.0. SHARE Working Paper Series 41-2019*. Munich: MEA, Max Planck Institute for Social Law and Social Policy.
- Börsch-Supan, A. M., Brandt, M., Hunkler, C., Kneip, T., Korbmacher, J., Malter, F., Schaas, B., Stuck, S., & Zuber, S. (2013). Data Resource Profile: The Survey of Health, Ageing and Retirement in Europe (SHARE). *International Journal of Epidemiology*. <https://doi.org/10.1093/ije/dyt088>
- Börsch-Supan, A. (2022a). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 1; Release Version: 8.0. 0; SHARE-ERIC. Dataset*. <https://doi.org/10.6103/SHARE.w1.800>
- Börsch-Supan, A. (2022b). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 2; Release Version: 8.0. 0; SHARE-ERIC. Dataset*. <https://doi.org/10.6103/SHARE.w2.800>
- Börsch-Supan, A. (2022c). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 4; Release Version: 8.0. 0; SHARE-ERIC. Dataset*. <https://doi.org/10.6103/SHARE.w4.800>
- Börsch-Supan, A. (2022d). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 5; Release Version: 8.0. 0; SHARE-ERIC. Dataset*. <https://doi.org/10.6103/SHARE.w5.800>
- Börsch-Supan, A. (2022e). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 6; Release Version: 8.0. 0; SHARE-ERIC. Dataset*. <https://doi.org/10.6103/SHARE.w6.800>
- Börsch-Supan, A. (2022f). *Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 8; Release Version: 8.0. 0; SHARE-ERIC. Dataset*. <https://doi.org/10.6103/SHARE.w8.800>
- Brown, D. C., Hayward, M. D., Montez, J. K., Hummer, R. A., Chiu, C.-T., & Hidajat, M. M. (2012). The significance of education for mortality compression in the United States. *Demography*, *49*(3), 819–840.
- Cambois, E., Solé-Auró, A., Brønnum-Hansen, H., Egidi, V., Jagger, C., Jeune, B., ... Robine, J.-M. (2016). Educational differentials in disability vary across and within welfare regimes: A comparison of 26 European countries in 2009. *Journal of Epidemiology & Community Health*, *70*(4), 331–338.
- Cameron, A. C., & Trivedi, P. K. (2005). *Microeconometrics: Methods and applications*. Cambridge university press.
- Carlson, P. (1998). Self-perceived health in East and west Europe: Another European health divide. *Social Science & Medicine*, *46*(10), 1355–1366.
- Carlson, P. (2004). The European health divide: A matter of financial or social capital? *Social Science & Medicine*, *59*(9), 1985–1992.
- Christensen, K., Doblhammer, G., Rau, R., & Vaupel, J. W. (2009). Ageing populations: The challenges ahead. *Lancet*, *374*(9696), 1196–1208.

- Crimmins, E. M., Kim, J. K., & Solé-Auró, A. (2011). Gender differences in health: Results from SHARE, ELSA and HRS. *The European Journal of Public Health*, 21(1), 81–91.
- Crimmins, E. M., Shim, H., Zhang, Y. S., & Kim, J. K. (2019). Differences between men and women in mortality and the health dimensions of the morbidity process. *Clinical Chemistry*, 65(1), 135–145.
- Cutler, D. M., & Lleras-Muney, A. (2010). Understanding differences in health behaviors by education. *Journal of Health Economics*, 29(1), 1–28.
- Eikemo, T. A., & Bambra, C. (2008). The welfare state: A glossary for public health. *Journal of Epidemiology & Community Health*, 62(1), 3–6.
- Eikemo, T. A., Huisman, M., Bambra, C., & Kunst, A. E. (2008). Health inequalities according to educational level in different welfare regimes: A comparison of 23 European countries. *Sociology of Health & Illness*, 30(4), 565–582.
- European Commission. (2022). *State of health in the EU: Companion report 2022*. Publications Office of the European Union Luxembourg.
- Eurostat. (2022). *Life expectancy by age and sex [DEMO_MLEXPEEC]*.
- Geyer, S., Hemström, Ö., Peter, R., & Vågerö, D. (2006). Education, income, and occupational class cannot be used interchangeably in social epidemiology. Empirical evidence against a common practice. *Journal of Epidemiology & Community Health*, 60(9), 804–810.
- Haggard, S., & Kaufman, R. R. (2009). *The Eastern European welfare state in comparative perspective*. Springer.
- HMD. Human Mortality Database Max Planck Institute for Demographic Research (Germany), University of California, Berkeley (USA), and French Institute for Demographic Studies (France). Available at: www.mortality.org.
- Jagger, C., Crimmins, E. M., Saito, Y., Yokota, R. T. D. C., Van Oyen, H., & Robine, J.-M. (2020). *International handbook of health expectancies*. Springer.
- Jagger, C., Gillies, C., Moscone, F., Cambois, E., Van Oyen, H., Nusselder, W., & Robine, J.-M. (2008). Inequalities in healthy life years in the 25 countries of the European union in 2005: A cross-national meta-regression analysis. *Lancet*, 372(9656), 2124–2131.
- Jagger, C., Weston, C., Cambois, E., Van Oyen, H., Nusselder, W., Doblhammer, G., ... Robine, J.-M. (2011). Inequalities in health expectancies at older ages in the European union: Findings from the survey of health and retirement in Europe (SHARE). *Journal of Epidemiology & Community Health*, 65(11), 1030–1035.
- Kiely, K. M., Brady, B., & Byles, J. (2019). Gender, mental health and ageing. *Maturitas*, 129, 76–84.
- Klotz, J., Göllner, T., & Austria, S. (2017). *Estimating differential mortality from EU-SILC longitudinal data: A feasibility study* (Vol. 4). FACTAGE project report D.
- Lindahl-Jacobsen, R., Rau, R., Jeune, B., Canudas-Romo, V., Lenart, A., Christensen, K., & Vaupel, J. W. (2016). Rise, stagnation, and rise of Danish women's life expectancy. *Proceedings of the National Academy of Sciences*, 113(15), 4015–4020.
- Luy, M., & Minagawa, Y. (2014). Gender gaps-Life expectancy and proportion of life in poor health. *Health Reports*, 25(12), 12.
- Mackenbach, J. P. (2017). Nordic paradox, southern miracle, eastern disaster: Persistence of inequalities in mortality in Europe. *The European Journal of Public Health*, 27(suppl_4), 14–17.
- Mackenbach, J. P., Kulháňová, I., Bopp, M., Deboosere, P., Eikemo, T. A., Hoffmann, R., ... Menvielle, G. (2015). Variations in the relation between education and cause-specific mortality in 19 European populations: A test of the “fundamental causes” theory of social inequalities in health. *Social Science & Medicine*, 127, 51–62.
- Mackenbach, J. P., Valverde, J. R., Artink, B., Bopp, M., Brønnum-Hansen, H., Deboosere, P., ... Martikainen, P. (2018). Trends in health inequalities in 27 European countries. *Proceedings of the National Academy of Sciences*, 115(25), 6440–6445.
- Majer, I. M., Nusselder, W., Mackenbach, J., & Kunst, A. (2011). Socioeconomic inequalities in life and health expectancies around official retirement age in 10 Western-European countries. *Journal of Epidemiology & Community Health*, 65(11), 972–979.
- Mäki, N., Martikainen, P., Eikemo, T., Menvielle, G., Lundberg, O., Östergren, O., ... Mackenbach, J. P. (2013). Educational differences in disability-free life expectancy: A comparative study of long-standing activity limitation in eight European countries. *Social Science & Medicine*, 94, 1–8.
- Minagawa, Y. (2013). Inequalities in healthy life expectancy in Eastern Europe. *Population and Development Review*, 39(4), 649–671.
- Montez, J. K., & Friedman, E. M. (2014). Educational attainment and adult health: Under what conditions is the association causal? *Social Science & Medicine*, 127, 1–7, 1982.
- Montez, J. K., Zajacova, A., Hayward, M. D., Woolf, S. H., Chapman, D., & Beckfield, J. (2019). Educational disparities in adult mortality across US states: How do they differ, and have they changed since the mid-1980s? *Demography*, 56(2), 621–644.
- Montez, J. K., Zhang, W., Zajacova, A., & Hamilton, T. G. (2018). Does college major matter for women's and men's health in midlife? Examining the horizontal dimensions of educational attainment. *Social Science & Medicine*, 198, 130–138.
- Muszynska-Spielauer, M., & Spielauer, M. (2022). *The effect of sample attrition in the EU statistics on income and living conditions on the estimates of eurostat's healthy life years*.
- Nusselder, W., Looman, C., Van Oyen, H., Robine, J.-M., & Jagger, C. (2010). Gender differences in health of EU10 and EU15 populations: The double burden of EU10 men. *European Journal of Ageing*, 7, 219–227.
- OECD. (2019). *Health at a glance*. <https://doi.org/10.1787/4dd50c09-en>, 2019.
- Oksuzyan, A., Juel, K., Vaupel, J. W., & Christensen, K. (2008). Men: Good health and high mortality. Sex differences in health and aging. *Ageing-Clinical & Experimental Research*, 20, 91–102.
- Permanyer, I., & Bramajo, O. (2022). Increased longevity in Europe: Adding years to life or life to years? *Perspectives Demographiques*, 28, 1–4.
- Permanyer, I., & Spijker, J. (2020). *Socioeconomics and the macro-and micro-level determinants of global health inequality*. *Handbook of Global Health* (pp. 1–36).
- Pinho-Gomes, A.-C., Vassallo, A., Carcel, C., Peters, S., & Woodward, M. (2022). Gender equality and the gender gap in life expectancy in the European Union. *BMJ Global Health*, 7(2), Article e008278.
- Preston, S. H., Heuveline, P., & Guillot, M. (2001). *Demography. Measuring and modeling population processes* (Vol. 27). Blackwell Publishers.
- Robine, J.-M., & Jagger, C. (2003). Creating a coherent set of indicators to monitor health across Europe: The euro-REVES 2 project. Euro-REVES Group *The European Journal of Public Health*, 13(suppl_3), 6–14.
- Robine, J.-M., Romieu, I., & Cambois, E. (1999). Health expectancy indicators. *Bulletin of the World Health Organization*, 77(2), 181.
- Romero-Ortuno, R., Fouweather, T., & Jagger, C. (2014). Cross-national disparities in sex differences in life expectancy with and without frailty. *Age and Ageing*, 43(2), 222–228.
- Ross, C. E., Masters, R. K., & Hummer, R. A. (2012). Education and the gender gaps in health and mortality. *Demography*, 49(4), 1157–1183.
- Ross, C. E., & Mirowsky, J. (2010). Why education is the key to socioeconomic differentials in health. *Handbook of medical sociology*, 6, 33–51.
- Rubio-Valverde, J. R., Nusselder, W. J., & Mackenbach, J. P. (2019). Educational inequalities in global activity limitation indicator disability in 28 European countries: Does the choice of survey matter? *International Journal of Public Health*, 64, 461–474.
- Sauerberg, M. (2021). The impact of population's educational composition on Healthy Life Years: An empirical illustration of 16 European countries. *SSM-Population Health*, 15, Article 100857.
- Schulz, A., & Doblhammer, G. (2011). *Longitudinal research with the second wave of share: Representativeness of the longitudinal sample and the mortality follow-up*.
- Shkolnikov, V. M., Andreev, E. M., Jasilionis, D., Leinsalu, M., Antonova, O. I., & McKee, M. (2006). The changing relation between education and life expectancy in central and eastern Europe in the 1990s. *Journal of Epidemiology & Community Health*, 60(10), 875–881.
- Siegrist, J., & Marmot, M. (2004). Health inequalities and the psychosocial environment—two scientific challenges. *Social Science & Medicine*, 58(8), 1463–1473.
- Solé-Auró, A., Beltrán-Sánchez, H., & Crimmins, E. M. (2015). Are differences in disability-free life expectancy by gender, race, and education widening at older ages? *Population Research and Policy Review*, 34, 1–18.
- Solé-Auró, A., & Gumà, J. (2022). Healthy aging patterns in Europe: A multistate health transition approach. *Journal of Population Ageing*, 1–23.
- Spitzer, S. (2020). Biases in health expectancies due to educational differences in survey participation of older Europeans: It's worth weighting for. *The European Journal of Health Economics*, 21(4), 573–605.
- Statistics, U. I. S. (2012). *International standard classification of education: ISCED 2011*. UNESCO Institute for Statistics Montreal.
- Sullivan, D. F. (1971). A single index of mortality and morbidity. *HSMHA Health Reports*, 86(4), 347.
- Tawiah, R., Jagger, C., Anstey, K. J., & Kiely, K. M. (2021). Trends in disability-free life expectancy at age 50 years in Australia between 2001 and 2011 by social disadvantage. *Journal of Epidemiology & Community Health*, 75(11), 1056–1062.
- Teachman, J. D., & Hayward, M. D. (1993). Interpreting hazard rate models. *Sociological Methods & Research*, 21(3), 340–371.
- Valverde, J. R. R., Mackenbach, J. P., & Nusselder, W. J. (2021). Trends in inequalities in disability in Europe between 2002 and 2017. *Journal of Epidemiology & Community Health*, 75(8), 712–720.
- Van Oyen, H., Cox, B., Jagger, C., Cambois, E., Nusselder, W., Gilles, C., & Robine, J.-M. (2010). Gender gaps in life expectancy and expected years with activity limitations at age 50 in the European union: Associations with macro-level structural indicators. *European Journal of Ageing*, 7, 229–237.
- Van Oyen, H., Nusselder, W., Jagger, C., Kolip, P., Cambois, E., & Robine, J.-M. (2013). Gender differences in healthy life years within the EU: An exploration of the “health-survival” paradox. *International Journal of Public Health*, 58(1), 143–155.
- Van Oyen, H., Van der Heyden, J., Perenboom, R., & Jagger, C. (2006). Monitoring population disability: Evaluation of a new global activity limitation indicator (GALI). *Sozial-und Präventivmedizin*, 51(3), 153–161.
- Welsh, C. E., Matthews, F. E., & Jagger, C. (2021). Trends in life expectancy and healthy life years at birth and age 65 in the UK, 2008–2016, and other countries of the EU28: An observational cross-sectional study. *The Lancet Regional Health-Europe*, 2, Article 100023.