


Original Russian text [www.bionet.nsc.ru/vogis/](http://www.bionet.nsc.ru/vogis/)

## Starch metabolism in potato *Solanum tuberosum* L.

E.M. Sergeeva , K.T. Larichev, E.A. Salina, A.V. Kochetov


Institute of Cytology and Genetics of the Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia  
 [sergeeva@bionet.nsc.ru](mailto:sergeeva@bionet.nsc.ru)

**Abstract.** Starch is a major storage carbohydrate in plants. It is an important source of calories in the human and animal diet. Also, it is widely used in various industries. Native starch consists of water-insoluble semicrystalline granules formed by natural glucose polymers amylose and amylopectin. The physicochemical properties of starch are determined by the amylose:amylopectin ratio in the granule and degrees of their polymerization and phosphorylation. Potato *Solanum tuberosum* L. is one of the main starch-producing crops. Growing industrial needs necessitate the breeding of plant varieties with increased starch content and specified starch properties. This task demands detailed information on starch metabolism in the producing plant. It is a complex process, requiring the orchestrated work of many enzymes, transporter and targeting proteins, transcription factors, and other regulators. Two types of starch are recognized with regard to their biological functions. Transitory starch is synthesized in chloroplasts of photosynthetic organs and degraded in the absence of light, providing carbohydrates for cell needs. Storage starch is synthesized and stored in amyloplasts of storage organs: grains and tubers. The main enzymatic reactions of starch biosynthesis and degradation, as well as carbohydrate transport and metabolism, are well known in the case of transitory starch of the model plant *Arabidopsis thaliana*. Less is known about features of starch metabolism in storage organs, in particular, potato tubers. Several issues remain obscure: the roles of enzyme isoforms and different regulatory factors in tissues at various plant developmental stages and under different environmental conditions; alternative enzymatic processes; targeting and transport proteins. In this review, the key enzymatic reactions of plant carbohydrate metabolism, transitory and storage starch biosynthesis, and starch degradation are discussed, and features specific for potato are outlined. Attention is also paid to the known regulatory factors affecting starch metabolism.

Key words: potato; *Solanum tuberosum*; starch; amylose; amylopectin; synthesis; degradation.

**For citation:** Sergeeva E.M., Larichev K.T., Salina E.A., Kochetov A.V. Starch metabolism in potato *Solanum tuberosum* L. *Vavilovskii Zhurnal Genetiki i Seleksii* = *Vavilov Journal of Genetics and Breeding*. 2022;26(3):250-263. DOI 10.18699/VJGB-22-32

## Метаболизм крахмала у картофеля *Solanum tuberosum* L.

E.M. Сергеева , К.Т. Ларичев, Е.А. Салина, А.В. Кочетов

Федеральный исследовательский центр Институт цитологии и генетики Сибирского отделения Российской академии наук, Новосибирск, Россия  
 [sergeeva@bionet.nsc.ru](mailto:sergeeva@bionet.nsc.ru)

**Аннотация.** Крахмал – основной запасной углевод растений и важный источник калорий в рационе питания человека и животных, широко применяется также для ряда промышленных нужд. Нативный крахмал представляет собой не растворимые в воде полукристаллические гранулы, сформированные молекулами амилозы и амилопектина, которые являются природными полимерами глюкозы. Физико-химические свойства крахмала определяются соотношением амилозы и амилопектина в грануле, степенью их полимеризации и фосфорилирования. Одно из основных растений-производителей крахмала – картофель *Solanum tuberosum* L. Растущие потребности рынка диктуют необходимость получения сортов с повышенным содержанием и заданными свойствами крахмала, необходимым условием чего является получение детальной информации о процессе его метаболизма в организме растения. Процесс метаболизма крахмала сложен и представляет собой согласованную работу множества ферментов, транспортных и направляющих белков, транскрипционных и других регуляторных факторов. По принципу выполняемой биологической функции крахмал делится на два типа: транзитный, который синтезируется в хлоропластах фотосинтезирующих органов и распадается при отсутствии освещения, обеспечивая клетку углеводами; и запасной, который синтезируется и хранится в амилопластах запасующих органов (зерен, клубней). Основные ферментативные реакции биосинтеза и деградации крахмала, а также транспорта и метаболизма углеводов хорошо изучены на транзитном крахмале модельного объекта *Arabidopsis thaliana*; об особенностях метаболизма крахмала в запасующих органах, в частности клубнях картофеля, известно несколько меньше. При этом ряд вопросов остается открытым: недостаточно изучены участие различных изоформ ферментов и влияние на них

регуляторных факторов в зависимости от ткани и стадии развития растения, а также условий внешней среды; реализация альтернативных путей ферментативных превращений; участие направляющих и транспортных белков. В данном обзоре рассмотрены ключевые ферментативные реакции углеводного обмена, биосинтеза и деградации транзиторного и запасного крахмала, общие для растений, и отмечены особенности, характерные для картофеля; уделено внимание известным регуляторным факторам, влияющим на метаболизм крахмала.

Ключевые слова: картофель; *Solanum tuberosum*; крахмал; амилоза; амилопектин; синтез; деградация.

## Introduction

Starch is the main storage carbohydrate in plants. It constitutes up to 85 % of the dry matter of their edible parts: cereal grains (maize *Zea mays* L., rice *Oryza sativa* L., wheat *Triticum* spp., barley *Hordeum vulgare* L., etc.), potato tubers *Solanum tuberosum* L., edible roots (cassava *Manihot esculenta* Crantz, sweet potato *Ipomoea batatas* (L.) Lam., and yam *Dioscorea alata* L.), sago palm stems *Metroxylon sagu* Rottb., plantain fruit *Musa* spp. (Zeeman et al., 2010; Santana, Meireles, 2014). Starch provides a great portion of calories for human and animal nutrition. In addition, it is a natural reproducible and biodegradable material used in non-food industry, e. g., in the production of fabric, paper, drugs, and plastics.

Chemically, starch is a mixture of amylose and amylopectin. These natural glucose polymers form water-insoluble semicrystalline granules. Amylopectin consists of highly branched glucan molecules, the linear regions of which are formed by  $\alpha$ -1,4-glycosidic bonds, whereas the branching points are formed by  $\alpha$ -1,6-bonds. Amylose is a practically linear polymer with few branches. Amylopectin molecules constitute about 75–80 % of starch weight. They form the structural framework of the granule, consisting of repetitive amorphous and semicrystalline lamellae. Amylose molecules are dispersed in the semicrystalline amylopectin matrix (Zeeman et al., 2010; Tetlow, Bertoft, 2020).

The amylose:amylopectin ratio determines the dietetic, physicochemical, and functional starch properties essential for particular industries. The starch present in food is classified into glycemic and resistant. Being readily digestible in the small intestine, amylopectin increases the glycemic potential of starch. In contrast, higher amylose contents make starch more resistant (Li et al., 2008). Resistant starch is less degradable by amylases in the small intestine. It serves as substrate for microbes in the large intestine to produce short-chain fatty acids, which exert local antiinflammatory and antitumor effects (Birt et al., 2013). Also, physicochemical (gelatinization and retrogradation) and functional (swelling and viscosity) properties are taken into consideration in certain applications. These properties are determined

genetically: by size and morphology of starch granules, amylose:amylopectin ratio, glucan branching, and glucan phosphorylation (Visser et al., 1991; Schwall et al., 2000; Hofvander et al., 2004; Khlestkin et al., 2017).

Potato (*Solanum tuberosum* L.) ranks fourth among starch-producing crops in the world, next to maize, cassava, and wheat. Potato starch differs from cereal starches in a variety of important features. Potato amylose and amylopectin have higher degrees of polymerization and phosphorylation; therefore, potato starch is more suitable for bioplastic production (Hofvander et al., 2004; Reyniers et al., 2020). In response to the increasing commercial demand, the global production of potato starch steadily increases: 3.7 million tons in 2018 and 3.9 million tons in 2020 (<https://www.researchandmarkets.com/reports/5330932/potato-starch-market-global-industry-trends>). To obtain native starches with specified properties and to increase the overall amount of starch per plant are topical tasks in potato breeding.

The key enzymes in starch biosynthesis (see the Table) and their genes have been studied in detail in model plants (*Arabidopsis thaliana* L.) and in crops, including potato (Streb, Zeeman, 2012; Van Harsselaar et al., 2017; Slugina, Kochieva, 2018).

Starch is produced by the polymerization of ADP-glucose, catalyzed by granule-bound (GBSS) and soluble (SS) starch synthases. Other enzymes involved are the starch branching enzyme (SBE) and the debranching enzyme (DBE).

Starch metabolism is a component of general carbohydrate metabolism. It is essential for plant functions: growth, development, and stress response. The enzymes and genes associated with the metabolism of starch and other plant carbohydrates have been extensively studied for three decades. A considerable body of data on the location of these genes in plant genomes and their expression has been accumulated. Seventy-five *S. tuberosum* genes have been mapped on the reference genome, and the expression patterns of 64 genes in leaves and tubers have been studied (Van Harsselaar et al., 2017; Slugina, Kochieva, 2018). It has been shown that starch metabolism genes experienced numerous duplications and produced paralogs by sub- and neofunctionaliza-

## Enzymes involved in starch metabolism

Enzyme	EC no.	Izozyme	Function
<b>Carbohydrate metabolism</b>			
Sucrose synthase	EC:2.4.1.13	SuSy1, SuSy2, SuSy3, SuSy4, SuSy5, SuSy6, SuSy7	Reversible cleavage of sucrose to fructose and UDP-glucose (Stein, Granot, 2019))
UDP-glucose pyrophosphorylase	EC:2.7.7.9	UGPase1, UGPase2	Reversible conversion of UDP-glucose and pyrophosphate to glucose-1-phosphate and UDP (Kleczkowski et al., 2010)
Phosphoglucoisomerase	EC:5.3.1.9	PGI, PGI-like1, PGI-like2	Reversible conversion of fructose-6-phosphate to glucose-6-phosphate (Yu et al., 2000)
Phosphoglucomutase	EC:5.4.2.2	PGM1, PGM2.1, PGM2.2	Reversible conversion of glucose-6-phosphate to glucose-1-phosphate (Yu et al., 2000)
ADP-glucose pyrophosphorylase: large and small subunits	EC:2.7.7.27	AGPL1, AGPL2, AGPL3	Reversible synthesis of ADP-glucose and pyrophosphate from glucose-1-phosphate and ATP (Geigenberger et al., 1999)
	EC:2.7.7.27	AGPS1.1, AGPS1.2, AGPS2	
Inorganic pyrophosphatase	EC:3.6.1.1	PPase, PPase-like	Pyrophosphate cleavage to orthophosphate (George et al., 2010)
<b>Starch granule synthesis</b>			
Granule-bound starch synthase	EC:2.4.1.242	GBSS1	Amylose synthesis by forming $\alpha$ -1,4-glycosidic bonds (Pfister, Zeeman, 2016)
Soluble starch synthases	EC:2.4.1.21	SS1, SS2, SS3, SS4, SS5, SS6	SS1, SS2, SS3: amylopectin synthesis by forming $\alpha$ -1,4-glycosidic bonds (Pfister, Zeeman, 2016). SS4, SS5: granule initiation (Helle et al., 2018; Tetlow, Bertoft, 2020). SS6: associated with starch granules in potato tubers, function unknown (Helle et al., 2018)
Starch-branching enzymes	EC:2.4.1.18	SBE1.1, SBE1.2, SBE2, SBE3	SBE2, SBE3: branching of amylose and amylopectin molecules by forming $\alpha$ -1,6-glycosidic bonds (Tetlow, Bertoft, 2020). The SBE1 function is unknown (Van Harselaar et al., 2017)
Debranching enzymes $\alpha$ - (1 $\rightarrow$ 6) glucan hydrolase starch-debranching enzyme (DBE). Isoamylase	EC:3.2.1.68	ISA1.1, ISA1.2, ISA2, ISA3	Hydrolyse $\alpha$ -1,6-glycosidic bonds of amylopectin. ISA1 and ISA2 are involved in amylopectin synthesis and the formation of its semicrystalline structure (Hennen-Bierwagen et al., 2012). The ISA1/ISA2 complex participates in the regulation of starch granule formation (Bustos et al., 2004). ISA3 participates in starch degradation (Streb, Zeeman, 2012; Pfister, Zeeman, 2016)
<b>Starch degradation</b>			
Glucan water dikinase	EC:2.7.9.4	GWD	Glucan phosphorylation at position C6 (Ritte et al., 2006)
Phosphoglucan water dikinase	EC:2.7.9.5	PWD	Glucan phosphorylation at position C3 (Ritte et al., 2006)

## End of the Table

Enzyme	EC no.	Izozyme	Function
Phosphoglucan phosphatases Like SEX Four; Starch Excess 4	EC:3.1.3.48	LSF1, LSF2 SEX4, SEX4-like	Glucan dephosphorylation. SEX4 dephosphorylates positions C3 and C6 of glycoside residues. LSF2 dephosphorylates position C3 (Hejazi et al., 2010; Santelia et al., 2011)
$\alpha$ -Amylase	EC:3.2.1.1	AMY1.1, AMY1.2, AMY2 (AMY23), AMY3, AMY3-like	AMY3: hydrolysis of $\alpha$ -1,4-glycosidic bonds of amylose and amylopectin (Yu et al., 2005). AMY2 (AMY23): may be involved in the cold-induced sweetening of starch (Hou et al., 2017)
$\beta$ -Amylase	EC:3.2.1.2	BAM1, BAM2, BAM3.1, BAM3.2, BAM4, BAM6.1, BAM6.2, BAM6.3, BAM7, BAM9	BAM1 and BAM3: cleavage of $\alpha$ -1,4-glycosidic bonds in amylose and amylopectin; maltose formation (Fulton et al., 2008). BAM1, BAM9 may be involved in the cold-induced sweetening of starch (Hou et al., 2017)
$\alpha$ -Glucan phosphorylase	EC:2.4.1.1	PHO1a, PHO1b, PHO2a, PHO2b	Glucose-1-phosphate production by maltooligosaccharide degradation; participation in starch granule growth (Pfister, Zeeman, 2016)
4- $\alpha$ -Glucanotransferase, disproportionating enzyme	EC:2.4.1.25	DPE1, DPE2	Maltooligosaccharide degradation to glucose (Critchley et al., 2001). Involved in granule initiation (Tetlow, Bertoft, 2020)

tion during evolution. The paralogous genes, which encode different enzyme isoforms, show tissue- and/or stage-specific expression patterns (Van Harsselaar et al., 2017; Qu et al., 2018; López-González et al., 2019). Various factors affect the expression of starch metabolism genes: circadian rhythms, photoperiod, levels of plant hormones and sugars, and stressing factors (drought and cold) (López-González et al., 2019). Although the key enzymatic reactions in starch biosynthesis and degradation are well known, there are many unclear points concerning alternative enzymatic processes and their localization inside the cell, roles of particular isozymes in starch metabolism in different organs at different developmental stages, and the influence of regulatory factors.

By now, functions of many proteins involved in starch metabolism in potato have been identified (see the Table). Most enzymes have isoforms with partially overlapping functions (Van Harsselaar et al., 2017). Translocator proteins essential for transporting metabolites through plastid membranes are important for starch metabolism as well. They also have multiple forms: adenylate translocators NTT1 and NTT2 (Tjaden et al., 2001), glucose transporter pGlcT1 (Cho et al., 2011), glucose 6-phosphate translocator (isoforms GPT1.1, GPT1.2, GPT2.1, and GPT2.2) (Kammerer et al., 1998),

maltose transporter MEX1 (Cho et al., 2011), and triose phosphate translocator (TPT, TPT-like) (Flügge et al., 1989).

The amylose:amylopectin ratio in plants can be modified by raising lines carrying certain alleles of genes involved in starch synthesis. Such accessions were obtained in cereals; for instance, in maize *Z. mays*. The *amylose extender* (*ae*<sup>-</sup>) mutation is associated with the loss of the activity of the starch-branching enzyme SBEIIb. Starch in plants with the *ae*<sup>-</sup> phenotype is enriched with amylose, and its amylopectin chains are longer (Stinard et al., 1993). The maize phenotype whose starch has practically no amylose is named *waxy*, and it is determined by a mutation in the gene for granule-bound starch synthase *GBSSI*. Its endosperm is gluey (Hossain et al., 2019).

However, the breeding of potato varieties with specified starch properties is complicated by its autotetraploid genome. The only amylose-free potato cultivar Eliane was obtained by mutation-assisted breeding (Muth et al., 2008). Genetic engineering and genome editing aimed at the modification of key starch biosynthesis genes produced plants with expected phenotypes: amylose-free starch (knockout and knockdown of the *GBSSI* gene), amylose-enriched starch (knockdown of



both *SBE1* and *SBE2*), and starch with modified amylopectin properties (editing of *SBE1* and/or *SBE2*) (Visser et al., 1991; Schwall et al., 2000; Hofvander et al., 2004; Andersson et al., 2006, 2017; Tuncel et al., 2019).

In some cases, plants with a desired phenotype acquired additional traits. For example, the increase in amylose content obtained by antisense suppression of the genes for starch-branching enzymes *SBE1* and *SBE2* was accompanied by a decrease in starch content, formation of smaller granules, and larger tubers (Hofvander et al., 2004). Apparently, changes in certain starch metabolism steps may affect the overall carbohydrate metabolism in the plant. An association study revealed genetic loci associated with starch content and productivity (tuber weight), whereas the functions of some detected genes were unknown at all, and some other genes were involved in signaling and regulation: transcriptional and posttranscriptional (Schönhals et al., 2017). Part of the detected SNPs exerted antagonistic effects on potato productivity and starch content (Schönhals et al., 2017). Thus, the investigation of regulation pathways of starch metabolism genes is important for improving potato quality and productivity.

### Carbohydrate metabolism in potato plants

Two starch forms are recognized with regard to biologic function: transitory and storage. Transitory starch is synthesized and accumulated in chloroplasts of photosynthetic organs (leaves) in the daytime and degraded in darkness to provide nutrients for the cell. Storage starch is synthesized in amyloplasts (nonphotosynthetic plastids) of storage organs (e. g., potato tubers) and stored there for a long time to be utilized in the preparation for sprouting (Zeeman et al., 2010; Streb, Zeeman, 2012).


The main difference in carbohydrate metabolism between cells of leaves and storage organs is in the sources of carbohydrates and ATP required for starch synthesis and enzymatic reactions. In leaves, they can form in the same cells that produce transitory starch, and in storage organs, they are imported from photosynthetic ones.

During photosynthesis, chloroplasts produce ATP and fix atmospheric carbon dioxide by the Calvin–Benson cycle, in which triose phosphate is produced as an intermediate (Streb, Zeeman, 2012). Part of triose phosphate molecules remain in the chloroplast stroma to serve as carbohydrate material for transitory starch synthesis. The rest is transported to cytosol by triose phosphate translocator TPT (Flügge et al., 1989). In chloroplasts, a series of enzymatic reactions converts triose phosphate to glucose-1-phosphate (G1P). Then ADP-glucose pyrophosphorylase (AGPase) converts G1P to ADP-glucose, the main substrate for starch synthesis (see the Figure).

As triose phosphate molecules are exported to cytosol, they are converted to sucrose, which is then delivered to storage organs through phloem and apoplast to be a carbohydrate material for storage starch synthesis. Sucrose is transported into cells of storage organs either by sucrose transporter proteins or, after being hydrolyzed by invertase to glucose and fructose, by hexose transporters (Ruan, 2014). There are two pathways to cleave sucrose in cell cytoplasm: saccharolytic, catalyzed by sucrose synthase SuSy, or hydrolytic, catalyzed by invertase Inv. Invertase irreversibly cleaves sucrose to glucose and fructose, and SuSy catalyzes the reversible cleavage to fructose and UDP-glucose (Stein, Granot, 2019). The predominant pathway depends on the tuber development stage. At the beginning of growth, at a high cell division rate, the hydrolytic pathway prevails, and the saccharolytic pathway steps forward at the starch accumulation stage (Appeldoorn et al., 1997). The SuSy-catalyzed pathway is important for the rate of starch accumulation in potato. It has been shown that a decrease in SuSy activity reduces starch content in mature tubers (Zrenner et al., 1995; Baroja-Fernández et al., 2009). Seven *SuSy* isoforms have been predicted in potato, and the *SuSy4* gene had tissue-specific expression in growing tubers (Van Harsselaar et al., 2017). It is likely that some sucrose synthase isoforms are involved in sucrose transport through phloem, as observed in *A. thaliana* (Yao et al., 2020).

The subsequent steps, catalyzed by fructokinase FK, hexokinase HK, and UDP-glucose pyrophosphorylase, produce a pool of phosphorylated hexoses in cytosol. They can be reversibly interconverted by cytosolic isoforms of phosphoglucoisomerase PGI2 and phosphoglucomutase PGM2 (Yu et al., 2000; Kleczkowski et al., 2010). One of the phosphorylated hexoses, glucose-6-phosphate, is transported to amyloplasts by membrane glucose-6-phosphate/phosphate-translocator GPT (Kammerer et al., 1998). In amyloplasts, G6P is utilized for ADP-glucose production. The rate of starch biosynthesis in potato tubers directly depends on G6P transport (Tauberger et al., 2000; Fernie et al., 2002).

ATP is also imported to storage tissues from photosynthetic ones. It is delivered inside amyloplasts by plastid adenylate translocator NTT. It is known that even a slight decrease in NTT activity reduces the overall starch content in potato tubers (Tjaden et al., 2001), whereas the combination of NTT and GPT overexpression increases it (Zhang L. et al., 2008). There may be an alternative route of hexose transport to amyloplasts: direct import of glucose-1-phosphate (G1P) and its utilization in ADP-glucose synthesis. There is evidence that this route acts in potato, although the corresponding transporter



Starch metabolism in potato photosynthesizing (leaves) and storage (tubers) organs.

Blue arrows indicate the starch biosynthesis and sugar metabolism pathways; black arrows, the starch degradation pathway. Starch degradation in leaves and starch biosynthesis in tubers are shown in more detail. The red arrow indicates the alternative pathway of ADP-glucose synthesis in cytosol and subsequent transport to amyloplasts performed by BT1 (Brittle1-like transporter). Carbohydrates are shown in black: Fru, fructose; F6P, fructose-6-phosphate; Glc, glucose; G1P, glucose-1-phosphate; G6P, glucose-6-phosphate; S6P, sucrose-6-phosphate; TrioseP, triose phosphate; ADP-Glc, ADP-glucose; UDP-Glc, UDP-glucose. Enzymes and invertase inhibitor (in blue): AcInv, vacuolar acid invertase; AGPase, ADP-glucose pyrophosphorylase; ALD, aldolase; AMY,  $\alpha$ -amylase; BAM,  $\beta$ -amylase; FK, fructokinase; DPE, 4- $\alpha$ -glucanotransferase; FBP, fructose-1,6-bisphosphatase; GBSS, granule-bound starch synthase; GWD,  $\alpha$ -glucan water dikinase; HK, hexokinase; ISA, isoamylase; InvInh, invertase inhibitor; Inv, invertase; PGI, phosphoglucomutase; PGM, phosphoglucomutase; PHO,  $\alpha$ -glucan phosphorylase; PPase, inorganic pyrophosphatase; PWD, phosphoglucan water dikinase; SBE, starch-branching enzyme; SEX, LSF, phosphoglucan phosphatases; SPP, sucrose phosphate phosphatase; SPS, sucrose phosphate synthase; SS, starch synthase; SuSy, sucrose synthase; UGPase, UDP-glucose pyrophosphorylase. Transporter proteins (in green): GPT, glucose-6-phosphate translocator; NTT, ATP-ADP antiporter; MEX, maltose transporter; TPT, triose phosphate/phosphate translocator; GlcT, glucose transporter; BT1L, Brittle1-like transporter.

protein is not known yet (Fettke et al., 2010). Two candidate G1P transporters through plasma membrane were recently found in *A. thaliana* (Malinova et al., 2020).

ADP-glucose acts as the substrate for starch biosynthesis. It is produced in a reversible reaction catalyzed by ADP-glucose pyrophosphorylase (AGPase) in the stroma of chloroplasts and amyloplasts. AGPase syn-

thesizes ADP-glucose and pyrophosphate (PPi) from G1P and ATP. AGPase is a heterotetramer consisting of two large and two small subunits, AGPL and AGPS. Its activity is essential for starch synthesis in potato tubers (Geigenberger et al., 1999). Inorganic pyrophosphatase (PPase) degrades pyrophosphate to orthophosphate (George et al., 2010). The plastid PPase isoform con-

tributes much to starch accumulation in potato tubers. Lines knocked down for the *StpsPPase* gene had lower contents of starch, in particular, amylose, and smaller granules. That study also recorded elevated amounts of starch biosynthesis intermediates: pyrophosphate, glucose, fructose, hexose phosphates, and, unexpectedly, ADP-glucose. The increase in ADP-glucose content indicates that pyrophosphate does not affect the direction of the AGPase-catalyzed reaction in potato. Thus, the mechanism by which PPase participates in starch synthesis in potato tubers is still to be understood (Andersson et al., 2018).

An alternative cytosolic pathway of ADP-glucose synthesis, catalyzed by SuSy and UGPase, acts in cereals (monocots). It is essential for grain growth. ADP-glucose is transported to amyloplast by the Brittle1-like transporter protein (BT1) (Bowsher et al., 2007). The homolog protein of BT1 (StBT1) has been found in *S. tuberosum* plant. However, there is no evidence for ADP-glucose transport through the amyloplast membrane. The StBT1 protein performs unidirectional transport of AMP, ADP, and ATP (Leroch et al., 2005).

### Starch granule synthesis

Transitory starch synthesis in leaf chloroplasts and storage starch synthesis in tuber amyloplasts follow basically the same route. Amylose and amylopectin synthesis is performed by 16 key enzymes belonging to the following groups: starch synthases, starch-branching enzymes, and starch-debranching enzymes (see the Table). Most enzymes exist as isoforms, the functions of which may partly overlap (Pfister, Zeeman, 2016; Van Harsseelaar et al., 2017).

Starch synthases catalyze the formation of glycosidic bonds by transferring the glucose residue of ADP-glucose to the nonreducing end of the glucose polymer. They are subdivided into granule-bound (GBSS) and soluble (SS) starch synthases. The former synthesize long chains, mainly in amylose, and long chain fragments in amylopectin. The latter include a series of isoforms: SS1, SS2, SS3, SS4, SS5, and SS6. Of them, SS1, SS2, and SS3 synthesize chains of various lengths in amylopectin (Pfister, Zeeman, 2016). The SS4 isoform performs a special function among starch synthases, as it initiates starch granule formation (Tetlow, Bertoft, 2020). One or two large starch granules instead of five to seven wild-type small ones were found in *A. thaliana* plants with the knocked out *ss4* gene (Roldán et al., 2007). The functions of SS5 and SS6 are still vague. The C end of the SS5 protein lacks the conservative fragment characteristic of starch synthases, which has catalytic domain GT1, although the protein has the conservative glucan-binding site. Probably, SS5 is involved in starch granule

initiation, as it has been shown that the loss of SS5 from *A. thaliana* reduces the granule number in leaves (Abt et al., 2020). The SS6 isoform and its gene were found in potato in recent years (Van Harsseelaar et al., 2017), and the role of this enzyme is unknown. It may participate in granule growth, as it is directly bound to it; in addition, it bears conservative motifs XXGGL and KXGGL, characteristic of glycosyl transferase domains of starch synthases GT1 and GT5, respectively (Helle et al., 2018).

Starch granule initiation was an obscure issue for a long time. The studies reported by now concern transitory starch initiation in the model plant *A. thaliana*, but it seems that our notion of some key steps in granule formation may be extended to other plant species (Mérida, Fettke, 2021). As mentioned above, SS4 is the main granule-initiating enzyme, and SS5 and SS6 also take part in the process. Maltooligosaccharides, probably forming in the degradation of starch polyglucans by amylases, are the substrate (Mérida, Fettke, 2021). The steric interaction of starch synthases, substrate molecules, and the growing granule is driven by the PTST2 and PTST1 proteins, targeting to starch. They are associated with starch synthases SS4 and GBSS1, respectively. Also, they contain a carbohydrate-binding domain (Seung et al., 2015, 2017). Note that PTST2 is not found in potato tubers, and this fact indicates that the starch granule initiation processes in *A. thaliana* and potato differ (Helle et al., 2018). A heteromultimeric complex of isoamylases ISA1 and ISA2 has been shown to influence starch granule initiation in potato tubers. By all appearances, isoamylases suppress the formation of new starch granules by disrupting the formation of soluble glucan molecules in chloroplast stroma (Bustos et al., 2004).

As the chains of amylose and amylopectin molecules are elongated by starch synthases SS1, SS2, and SS3, starch-branching enzymes SBE attach side branches to them (Pfister, Zeeman, 2016). Starch-branching enzymes cleave  $\alpha$ -1,4-glycosidic bonds of polyglucans, synthesized by starch synthases, and attach short chains to the so-called acceptor chain by forming  $\alpha$ -1,6-glycosidic bonds. The starch-branching enzymes of potato have three isoforms: SBE1.1, SBE1.2, SBE2, and SBE3 (formerly designated as SBE1) (see the Table) (Van Harsseelaar et al., 2017). Thus, forms referred to in other papers as SBE1 and SBE2 are designated as SBE3 and SBE2 according to the notation of Van Harsseelaar et al. SBE3 produces mainly long side chains, and SBE2 produces short amylopectin chains (Tetlow, Bertoft, 2020). The roles of SBE1.1 and SBE1.2 in starch production are unknown, but studies of *A. thaliana* demonstrate a pleiotropic effect of SBE1 on plant growth and development. Transformants overexpressing SBE1 were



white-colored and low. They had a longer life cycle and produced fewer seeds than control plants (Wang X. et al., 2010). The joint action of different isoforms affects amylopectin structure. Experiments with potato plants with the knocked out genes *sbe3* and/or *sbe2* (designated by the experimenters as *sbe1* and *sbe2*) demonstrate that *sbe3* inactivity results in the formation of starch with longer amylopectin chains and lower branching level. The knockout of *sbe2* with active *sbe3* did not affect the amylopectin structure much, but the number of starch granules in potato tubers increased and size decreased (Tuncel et al., 2019).

Debranching enzymes (DBE) are another group of enzymes involved in the formation of the amylopectin structure (see the Table). They reconstruct branched glucans into easier crystallizable forms, which is essential for granule formation (Pfister, Zeeman, 2016). Debranching enzymes include isoamylases (ISA), which catalyze the hydrolysis of  $\alpha$ -1,6-glycosidic amylopectin bonds and remove excessive branching. Potato isoamylases include isoforms ISA1, ISA2, and ISA3. The ISA1 and ISA2 proteins can form heteromultimers, capable of more efficient removal of long outer chains of amylopectin (Hussain et al., 2003). The ISA3 isoenzyme is important for starch degradation, as it cleaves short outer chains of glucans (Streb et al., 2008). Transgenic potato plants with lower expression of the *isa1*, *isa2*, and *isa3* genes had significantly less starch in developing tubers, whereas the starch contents in leaves did not change. The plants also had fewer and larger granules and higher sucrose contents, probably resulting from the increase in the overall granule surface and easier access for degrading enzymes (Ferreira et al., 2017).

In addition to starch synthases, branching and debranching enzymes, the synthesis of starch granules involves  $\alpha$ -glucan phosphorylases. Their plastid (PHO1) and cytoplasmic (PHO2) isoforms catalyze the reversible transfer of the glycosyl group of glucose-1-phosphate to the nonreducing end of the chain of an  $\alpha$ -1,4-bound glucan (Pfister, Zeeman, 2016). The PHO2 enzyme is involved in carbohydrate metabolism in cytoplasm, and PHO1 contributes to starch synthesis and degradation in plastids. It has been shown that at lower temperatures starch synthesis in potato tubers can also follow the phosphorylase pathway with G1P as the substrate (Fettke et al., 2012).

### Starch granule degradation

Degradation is an intrinsic part of the metabolism of starch and carbohydrates in general, although it has been studied much poorer than starch biosynthesis. The degradation pathways of transitory starch have been investigated in most detail in leaves of the model

plant *A. thaliana*. The knowledge of starch degradation in potato tubers is limited to cold-induced sweetening and sprouting. The main steps of starch degradation are the release of soluble glucan from starch granules, glucan conversion to linear forms (maltooligosaccharides), maltooligosaccharide hydrolysis to maltose, and subsequent maltose metabolism in the cell. Starch degradation is performed by a broad range of enzymes:  $\alpha$ - and  $\beta$ -amylases, isoamylase,  $\alpha$ -glucan water dikinase (GWD), phosphoglucan water dikinase (PWD),  $\alpha$ -glucan phosphorylase, phosphoglucan phosphatase, and 4- $\alpha$ -glucanotransferase (see the Table).

Starch granule degradation is initiated by GWD and PWD. They phosphorylate glucans at positions C6 and C3 of glucose residues, making them more hydrophilic and allow  $\alpha$ -,  $\beta$ -, and isoamylases access to them (see the Table) (Ritte et al., 2006; Streb, Zeeman, 2012). The phosphorylation by GWD seems to play the key role in starch degradation in potato tubers and leaves (Claassen et al., 1993; Orzechowski et al., 2021). Tubers of transgenic potato plants with lower expression of the *StGWD* gene were less prone to starch degradation at low temperatures (Lorberth et al., 1998).

The next step of starch granule degradation is glucan hydrolysis by amylases. Potato  $\alpha$ - and  $\beta$ -amylases include many isoforms, and functions of some of them are not known in detail (see the Table) (Van Harsselaar et al., 2017). By extrapolating data on *A. thaliana*, we suppose that  $\beta$ -amylases BAM1 and BAM3 hydrolyze linear fragments of amylose and amylopectin, and the degradation of branched fragments demands the debranching enzyme DBE (ISA3 in potato) (Hussain et al., 2003; Fulton et al., 2008; Pfister, Zeeman, 2016). With knocked down *StBAM3*, starch content in potato leaves was higher than in the wild genotype (Scheidig et al., 2002). Cold-induced sweetening in potato tubers is also affected by some amylase species:  $\alpha$ -amylase AMY2 (AMY23) and  $\beta$ -amylases BAM1 and BAM9. The supposed function of BAM1 and BAM9 is starch degradation in plastids, and AMY2 is likely to degrade phytoglycogen in cytosol (Hou et al., 2017). An alternative pathway of starch degradation is observed in *A. thaliana*. It is initiated by  $\alpha$ -amylase AMY3, which releases linear and branched glucans from starch granules, and these glucans are then hydrolyzed by  $\beta$ - and isoamylases (see the Figure) (Streb et al., 2008).

Alongside starch glucan hydrolysis by amylases, the glucans are dephosphorylated by phosphoglucan phosphatases SEX4 (Starch Excess) and LSF2 (LIKE SEX FOUR2), first described in *A. thaliana*. These processes are interrelated: phosphorylation by dikinases increases granule solubility and makes them accessible for amylases, whereas phosphate moieties may hamper



hydrolysis (Hejazi et al., 2010; Santelia et al., 2011). Reduction of *SEX4* or *LSF2* activities in potato inhibited starch degradation in leaves. Starch content in tubers remained unchanged, and granules were smaller and less phosphorylated (Samodien et al., 2018).

The cooperation of dikinases, amylases, and phosphatases produces a pool of soluble maltooligosaccharides (linear glucans). Maltooligosaccharides are degraded by two pathways: hydrolytic, by  $\beta$ -amylases, or phosphorytic, by  $\alpha$ -glucan phosphorylase *PHO1* (Weise et al., 2006; Fulton et al., 2008). The end product of the phosphorytic pathway is G1P, which can be utilized in metabolism inside the plastid. Also, glucose can be produced by 4- $\alpha$ -glucanotransferase *DPE1* (DisProportionating Enzyme) and exported to cytosol by glucose transporter *pGlcT1* (Critchley et al., 2001; Cho et al., 2011). Knockdown of the chloroplast enzyme *DPE* slows down starch degradation in potato leaves in the cold and induces maltooligosaccharide accumulation, although these effects are not observed in tubers (Lloyd et al., 2004). Cold-induced sweetening in potato tubers is accompanied by increasing  $\beta$ -amylase activity and higher maltose content (Nielsen et al., 1997).

Maltose, which is the predominant product of hydrolytic starch degradation, is exported to cytosol by the transmembrane transporter *MEX1* (Cho et al., 2011). In cytosol, maltose is processed by 4- $\alpha$ -glucanotransferase *DPE2* or phosphorylase *PHO2* to glucose or G1P, which are then converted to sucrose by the joint action of *PGI2*, *PGM2*, *HK*, *UGPase*, *SPS* (sucrose phosphate synthase), and *SPP* (sucrose phosphate phosphatase) (see the Figure) (López-González et al., 2019).

Sucrose is exported from leaf cells to storage organs; also, it is used in cell metabolism. In potato tubers, sucrose is used as a source of nutrients in sprouting, and its level controls dormancy release (Sonnewald S., Sonnewald U., 2014).

To delay sprouting, potato tubers are stored at low temperatures, 2–5 °C, and these conditions initiate cold-induced sweetening. This process involves sucrose hydrolysis by vacuolar acid invertase *AcInv*, encoded by the *Pain-1* gene, and the accumulation of reducing sugars (glucose and fructose) in tubers (see the Figure) (Sowokinos et al., 2018). Knockout of *Pain-1* resulted in lower contents of reducing sugars (Clasen et al., 2016). One of the key regulators of cold-induced sweetening is invertase inhibitor *SbAI*, which inhibits *AcInv* (McKenzie et al., 2013). It has been shown that *SbAI* can also inhibit  $\alpha$ - and  $\beta$ -amylases (*StAmy23*, *StBAM1*, and *StBAM9*), in potato tubers, thereby influencing the rate of starch degradation in cold-induced sweetening (Zhang H. et al., 2014).

## Mechanisms controlling starch metabolism

Starch metabolism requires orchestrated work of many enzymes, transporters, and targeting proteins, which implies many regulation levels: gene expression, post-transcriptional regulation, and the posttranslational regulation of enzymatic activity. The expression patterns of genes for key enzymes involved in starch metabolism are well known in various plant species, but much less is known about expression-regulating factors (López-González et al., 2019). The difficulty is that the starch-metabolizing enzymes exist as numerous isoforms, which are encoded by the corresponding number of paralogous genes. The expression patterns of these genes depend on tissue (leaves, developing seeds, or growing tubers) and developmental stage, as shown in *A. thaliana* and maize (Tsai et al., 2009; Chen et al., 2014). In potato, the tissue-specific mode of expression has been shown for *SuSy4*, *SS5*, *SBE3*, *APL3*, *PHO1a*, *PHO1b*, *GPT1.1*, *GPT2.1*, *SEX4*, and *NTT2* in tubers and for *AMY1.1*, *APL1*, and *BAM3.1* in leaves (Van Harselaar et al., 2017). A number of external and internal factors affect the expression of starch biosynthesis genes: circadian rhythms, photoperiod, and sugar content (Tiessen et al., 2002; Kötting et al., 2010). It is known that the expression rates of *GBSSI*, *LSF1*, *LSF2*, *SEX4*, and *BAM3* in *A. thaliana* leaves are governed by transcription factors depending on circadian rhythms and photoperiod, so that the demand for energy is rapidly met in response to ambient changes (Tenorio et al., 2003; Flis et al., 2016). The expression rates of the genes *GBSS*, *SuSy*, and *AGPase* respond to photoperiod in growing potato tubers, being highest in the end of the light time and lowest in the beginning. This variation is determined by the influx of photoassimilates from leaves (Geigenberger, Stitt, 2000; Ferreira et al., 2010).

The formation of the storage organ, tuber, from the stolon is an important step in potato plant development. It includes intense starch production, the formation of starch granules, and increase in metabolite flux. Tuber formation is a complex process, influenced by environmental factors (photoperiod) and a variety of signals: biochemical, hormonal, and molecular, mediated by microRNAs and transcription factors (Hannapel et al., 2017; Kondhare et al., 2021). The investigation of tuber formation contributed much to the understanding of mechanisms that regulate starch metabolism in potato tubers.

Plant hormones are an important factor influencing the expression of genes involved in starch metabolism, and their effect on tuber formation has been studied in sufficient detail. The level of abscisic acid correlates with starch accumulation in potato tubers (Borzenkova, Borovkova, 2003). Treatment of stolons with indole ace-

tic acid increased starch content in growing tubers, but a twofold increase in concentration caused the opposite effect (Wang D. et al., 2018). A correlation between the transcription rates of the genes for auxin, on the one hand, and starch biosynthesis (*PGM*, *AGPase*, *GBSS*, *SS*, and *BE*), on the other hand, was observed in the formation of cassava storage roots (Rüscher et al., 2021).

Sugars (hexoses, sucrose, and trehalose) are another group of signaling molecules influencing the expression of starch metabolism genes. Sucrose increases the expression of the *SuSy* and *AGPase* genes in potato (Salanoubat, Belliard, 1989; Müller-Röber et al., 1990). The rates of *SuSy* and *AGPase* expression are high in growing tubers, but they decrease rapidly after the separation of the tuber from the plant and, correspondingly, cease of sucrose import from photosynthesizing organs (Ferreira et al., 2010).

The differential expression of starch biosynthesis genes was detected at various tuber development stages (Ferreira et al., 2010; Van Harsselaar et al., 2017). The expression rate of the *SS4* gene was elevated at the stolon stage, and it lowered with tuber growth, confirming the role of this starch synthase in granule initiation (Ferreira et al., 2010). Also, tuber growth was accompanied by an increase in the expression rate of sucrose synthase *SuSy4* and decrease in the expression of cell wall invertase *cw-Inv*. These changes point to transition to the sucrose synthase-mediated pathway of sucrose degradation. Genes for glucose-6-phosphate translocator *GPT*, adenylate translocator *NTT*, ADP-glucose pyrophosphorylase (*AGPase*), starch synthases, and starch-branching enzymes increased their expression with tuber growth. Of this group, the isogenes *SuSy4*, *SBE3*, and *GPT2.1* demonstrated just tuber-specific expression (Ferreira et al., 2010; Van Harsselaar et al., 2017). Coexpression analysis was employed to investigate the mechanisms of molecular regulation of gene activity, and transcription factors *LOB*, *TIFY5a*, and *WRKY4* were found to be associated with the expression of the *SuSy4* and *GPT2.1* genes (Van Harsselaar et al., 2017). Analysis of coexpression networks for starch biosynthesis genes of seven plant species (*Arabidopsis thaliana*, cassava *Manihot esculenta*, millet *Panicum virgatum*, maize *Zea mays*, rice *Oryza sativa*, barley *Hordeum vulgare*, and sweet potato *Ipomoea batatas*) revealed the involvement of 24 transcription factors (López-González et al., 2019).

Little is known about mechanisms regulating starch metabolism genes at the posttranscriptional step. Post-transcriptional regulation involves a variety of factors, including RNA-binding proteins (RBPs), microRNAs, and alternative splicing, so that plants can rapidly reprogram their transcriptomes in response to external and internal factors. Photoperiod significantly influ-

ences microRNA expression patterns in the growth and development of potato tubers. It has been shown that differentially expressing microRNAs are targeted to genes coding for transcription factors and RNA-binding regulatory proteins *StGRAS*, *StTCP2/4*, and *StPTB6* (Kondhare et al., 2018).

Posttranslational regulation is the next step of protein activity control. It is mediated by allosteric regulation, in which an effector molecule is bound to a noncatalytic site of the enzyme, altering its conformation, catalytic properties, and, thereby, its specificity and interaction with other proteins (Zeeman et al., 2010). Allosteric regulation involves protein phosphorylation and the formation of multimeric complexes and disulfide bridges (Kötting et al., 2010; Zeeman et al., 2010). Many starch-metabolizing enzymes assume the phosphorylated state: *PGI*, *PGM1*, *AGPase*, *SS3*, *GWD1*, *GWD2*, *DPE2*, *AMY3*, *BAM1*, *BAM3*, *LDA*, *pGlcT*, and *MEX1* (Kötting et al., 2010). ADP-glucose pyrophosphorylase (*AGPase*) is a clear example of allosterically regulated potato enzyme. It is activated by 3-phosphoglyceric acid and inhibited by inorganic phosphate (Sowokinos, Preiss, 1982). Depending on the redox state in the cell, *AGPase* can be reversibly inactivated by the formation of disulfide bridges between small subunits of the heterotetramer (Ballicora et al., 2000).

Enzymes can aggregate into complexes known as metabolons (Sweetlove, Fernie, 2013). Complexes formed by starch biosynthesis enzymes were found in the endosperm of growing cereal seeds; in particular, *SSIII*, *SSIIa*, *SBEIIa*, and *SBEIIb* form a protein complex (Tetlow et al., 2008). Protein complexes of *PTST2* and *SS4* form in the initiation of starch granules in *A. thaliana* leaves (Seung et al., 2015, 2017). Potato isoamylases *ISA1* and *ISA2* form a heterotetrameric complex, which controls starch granule formation (Bustos et al., 2004).

## Conclusions

The investigation of starch metabolism in potato plants, particularly, starch biosynthesis and degradation in tubers, is topical in connection with the growing demand for potato starch in industry. A large body of information on key enzymes for starch and carbohydrate metabolism in various crops and the model species *A. thaliana* has been accumulated in the past three decades. Although the starch biosynthesis scheme is basically the same in different species, there are significant variations associated with different sets of isozymes, features of their functions, metabolite transport pathways (e. g., ADP-glucose transport through plastid membranes in cereals), and the existence of intricate and multileveled regulation, governed by external (photoperiod and temperature) and internal (plant hormones, metabolites,

microRNA, and regulatory proteins) factors. Isogenes encoding six starch synthase isoforms, seven sucrose synthases, nine  $\beta$ -amylases, and three to five for each of the starch-branching and other enzymes were identified in the potato genome (Van Harsselaar et al., 2017).

The functions of many isoforms, including the majority of  $\alpha$ - and  $\beta$ -amylases, are still unknown. Some isogenes (*SuSy4*, *SS5*, *SBE3*, *APL3*, *PHO1a*, *PHO1b*, *GPT1.1*, *GPT2.1*, *SEX4*, and *NTT2*) demonstrate tuber-specific expression and activity variation at various stages of tuber formation (Ferreira et al., 2010; Van Harsselaar et al., 2017). Isoenzymes AMY23, BAM1, BAM9 are specifically involved in starch degradation and carbohydrate metabolism in cold-induced sweetening (Hou et al., 2017). Also, the action of various factors on starch accumulation during tuber development has been shown: transcription factors *LOB*, *TIFY5a*, and *WRKY4*; plant hormones (auxin and abscisic acid); sugars; and microRNAs, the contents of which may mediate the effect of photoperiod. However, the functions of many isoenzymes and proteins involved in the regulatory and directing functions in starch metabolism in potato plants are poorly explored. To resolve this issue, modern methods are proposed: combined analysis of the metabolome and transcriptome inside a single cell or tissue (López-González et al., 2019). Bottom-up proteomics also seems promising in search for new components (Helle et al., 2018). For example, the analysis of 36 proteins associated with potato starch granules revealed, in addition to already known starch metabolism enzymes, targeting and regulatory proteins described in *A. thaliana*: *PTST1* (Protein Targeting to Starch), *ESV1* (Early StarVation1), and *LESV* (Like *ESV*). Also, Kunitz-type proteinase inhibitor and enzymes involved in redox regulation (thioredoxin *TRX* and glutathione peroxidase *GPX*) were found (Helle et al., 2018). Detailed information on all components involved in starch metabolism and on their interactions, including their behavior under varying ambient conditions, is essential for raising potato varieties with high performance and specified starch properties.

## References

- Abt M., Pfister B., Sharma M., Eicke S., Bürgy L., Neale I., Seung D., Zeeman S.C. STARH SYNTHASE5, a noncanonical starch synthase-like protein, promotes starch granule initiation in *Arabidopsis*. *Plant Cell*. 2020;32(8):2543-2565. DOI 10.1105/tpc.19.00946.
- Andersson M., Melander M., Pojmark P., Larsson H., Bülow L., Hofvander P. Targeted gene suppression by RNA interference: an efficient method for production of high-amylose potato lines. *J. Biotechnol.* 2006;123:137-148. DOI 10.1016/j.jbiotec.2005.11.001.
- Andersson M., Turesson H., Arrivault S., Zhang Y., Fält A.-S., Fernie A.R., Hofvander P. Inhibition of plastid PPase and NTT leads to major changes in starch and tuber formation in potato. *J. Exp. Bot.* 2018;69(8):1913-1924. DOI 10.1093/jxb/ery051.
- Andersson M., Turesson H., Nocolia A., Fält A.S., Samuelsson M., Hofvander P. Efficient targeted multiallelic mutagenesis in tetraploid potato (*Solanum tuberosum*) by transient CRISPR-Cas9 expression in protoplasts. *Plant Cell Rep.* 2017;36:117-128. DOI 10.1007/s00299-016-2062-3.
- Appeldoorn N.J.G., de Bruijn S.M., Koot-Gronsveld E.A.M., Visser R.G.F., Vreugdenhil D., van der Plas L.H.W. Developmental changes of enzymes involved in conversion of sucrose to hexose-phosphate during early tuberisation of potato. *Planta*. 1997;202:220-226. DOI 10.1007/s004250050122.
- Ballicora M.A., Frueauf J.B., Fu Y., Schurmann P., Preiss J. Activation of the potato tuber ADP-glucose pyrophosphorylase by thioredoxin. *J. Biol. Chem.* 2000;275(2):1315-1320. DOI 10.1074/jbc.275.2.1315.
- Baroja-Fernández E., Muñoz F.J., Montero M., Etxeberria E., Sesma M.T., Ovecka M., Bahaji A., Ezquer I., Li J., Prat S., Pozueta-Romero J. Enhancing sucrose synthase activity in transgenic potato (*Solanum tuberosum* L.) tubers results in increased levels of starch, ADPglucose and UDPglucose and total yield. *Plant Cell Physiol.* 2009;50(9):1651-1662. DOI 10.1093/pcp/pcp108.
- Birt D.F., Boylston T., Hendrich S., Jane J.L., Hollis J., Li L., McClelland J., Moore S., Phillips G.J., Rowling M., Schalinske K., Scott M.P., Whitley E.M. Resistant starch: promise for improving human health. *Adv. Nutr.* 2013;4(6):587-601. DOI 10.3945/an.113.004325.
- Borzenkova R.A., Borovkova M.P. Developmental patterns of phytohormone content in the cortex and pith of potato tubers as related to their growth and starch content. *Russ. J. Plant Physiol.* 2003;50:119-124. DOI 10.1023/A:1021957022595.
- Bowsher C.G., Scrase-Field E.F.A.L., Esposito S., Emes M.J., Tetlow I.J. Characterization of ADP-glucose transport across the cereal endosperm amyloplast envelope. *J. Exp. Bot.* 2007;58(6):1321-1332. DOI 10.1093/jxb/erl297.
- Bustos R., Fahy B., Hylton C.M., Seale R., Nebane N.M., Edwards A., Martin C., Smith A.M. Starch granule initiation is controlled by a heteromultimeric isoamylase in potato tubers. *Proc. Natl. Acad. Sci. USA*. 2004;101(7):2215-2220. DOI 10.1073/pnas.0305920101.
- Chen J., Zeng B., Zhang M., Xie S., Wang G., Hauck A., Lai J. Dynamic transcriptome landscape of maize embryo and endosperm development. *Plant Physiol.* 2014;166(1):252-264. DOI 10.1104/pp.114.240689.
- Cho M.H., Lim H., Shin D.H., Jeon J.S., Bhoo S.H., Park Y.I., Hahn T.R. Role of the plastidic glucose translocator in the export of starch degradation products from the chloroplasts in *Arabidopsis thaliana*. *New Phytologist*. 2011;190(1):101-112. DOI 10.1111/j.1469-8137.2010.03580.x.
- Claassen P.A.M., Budde M.A.W., Van Calker M.H. Increase in phosphorylase activity during cold-induced sugar accumulation in potato tubers. *Potato Res.* 1993;36:205-217. DOI 10.1007/BF02360529.
- Clasen B.M., Stoddard T.J., Luo S., Demorest Z.L., Li J., Cedrone F., Tibebe R., Davison S., Ray E.E., Daulhac A., Coffman A., Yabandith A., Retterath A., Haun W., Baltes N.J., Mathis L., Voytas D.F., Zhang F. Improving cold storage and processing traits in potato through targeted gene knockout. *Plant Biotechnol. J.* 2016;14(1):169-176. DOI 10.1111/pbi.12370.
- Critchley J.H., Zeeman S.C., Takaha T., Smith A.M., Smith S.M. A critical role for disproportionating enzyme in starch breakdown is revealed by a knock-out mutation in *Arabidopsis*. *Plant J.* 2001;26(1):89-100. DOI 10.1046/j.1365-313x.2001.01012.x.
- Fernie A.R., Swiedrych A., Tauberger E., Lytovchenko A., Trethewey R.N., Willmitzer L. Potato plants exhibiting combined antisense repression of cytosolic and plastidial isoforms of phosphoglucomutase surprisingly approximate wild type with respect to the rate of starch synthesis. *Plant Physiol. Biochem.* 2002;40(11):921-927. DOI 10.1016/S0981-9428(02)01457-2.
- Ferreira S.J., Senning M., Fischer-Stettler M., Streb S., Ast M., Neuhaus H.E., Zeeman S.C., Sonnewald S., Sonnewald U. Simultaneous silencing of isoamylases ISA1, ISA2 and ISA3 by multi-target



- RNAi in potato tubers leads to decreased starch content and an early sprouting phenotype. *PLoS One*. 2017;12(7):e0181444. DOI 10.1371/journal.pone.0181444.
- Ferreira S.J., Senning M., Sonnewald S., Kessling P.M., Goldstein R., Sonnewald U. Comparative transcriptome analysis coupled to X-ray CT reveals sucrose supply and growth velocity as major determinants of potato tuber starch biosynthesis. *BMC Genomics*. 2010;11:93. DOI 10.1186/1471-2164-11-93.
- Fettke J., Albrecht T., Hejazi M., Mahlow S., Nakamura Y., Steup M. Glucose 1-phosphate is efficiently taken up by potato (*Solanum tuberosum*) tuber parenchyma cells and converted to reserve starch granules. *New Phytol*. 2010;185(3):663-675. DOI 10.1111/j.1469-8137.2009.03126.x.
- Fettke J., Leifels L., Brust H., Herbst K., Steup M. Two carbon fluxes to reserve starch in potato (*Solanum tuberosum* L.) tuber cells are closely interconnected but differently modulated by temperature. *J. Exp. Bot*. 2012;63(8):3011-3029. DOI 10.1093/jxb/ers014.
- Flis A., Sulpice R., Seaton D.D., Ivakov A.A., Liput M., Abel C., Millar A.J., Stitt M. Photoperiod-dependent changes in the phase of core clock transcripts and global transcriptional outputs at dawn and dusk in *Arabidopsis*. *Plant Cell Environ*. 2016;39(9):1955-1981. DOI 10.1111/pce.12754.
- Flügge U.I., Fischer K., Gross A., Sebald W., Lottspeich F., Ecker-skorn C. The triose phosphate-3-phosphoglycerate-phosphate translocator from spinach chloroplasts: nucleotide sequence of a full-length cDNA clone and import of the *in vitro* synthesized precursor protein into chloroplasts. *EMBO J*. 1989;8(1):39-46.
- Fulton D.C., Stettler M., Mettler T., Vaughan C.K., Li J., Francisco P., Gil M., Reinhold H., Eicke S., Messerli G., Dorken G., Halliday K., Smith A.M., Smith S.M., Zeeman S.C.  $\beta$ -AMYLASE4, a non-catalytic protein required for starch breakdown, acts upstream of three active  $\beta$ -amylases in *Arabidopsis* chloroplasts. *Plant Cell*. 2008;20(4):1040-1058. DOI 10.1105/tpc.107.056507.
- Geigenberger P., Muller-Rober B., Stitt M. Contribution of adenosine 5'-diphosphoglucose pyrophosphorylase to the control of starch synthesis is decreased by water stress in growing potato tubers. *Planta*. 1999;209(3):338-345. DOI 10.1007/s004250050641.
- Geigenberger P., Stitt M. Diurnal changes in sucrose, nucleotides, starch synthesis and AGPS transcript in growing potato tubers that are suppressed by decreased expression of sucrose phosphate synthase. *Plant J*. 2000;23:795-806. DOI 10.1046/j.1365-313x.2000.00848.x.
- George G.M., van der Merwe M.J., Nunes-Nesi A., Bauer R., Fernie A.R., Kossmann J., Lloyd J.R. Virus-induced gene silencing of plastidial soluble inorganic pyrophosphatase impairs essential leaf anabolic pathways and reduces drought stress tolerance in *Nicotiana benthamiana*. *Plant Physiol*. 2010;154(1):55-66. DOI 10.1104/pp.110.157776.
- Hannapel D.J., Sharma P., Lin T., Banerjee A.K. The multiple signals that control tuber formation. *Plant Physiol*. 2017;174(2):845-856. DOI 10.1104/pp.17.00272.
- Hejazi M., Fettke J., Kötting O., Zeeman S.C., Steup M. The Laforin-like dual-specificity phosphatase SEX4 from *Arabidopsis* hydrolyzes both C6- and C3-phosphate esters introduced by starch-related dikinases and thereby affects phase transition of  $\alpha$ -glucans. *Plant Physiol*. 2010;152(2):711-722. DOI 10.1104/pp.109.149914.
- Helle S., Bray F., Verbeke J., Devassine S., Courseaux A., Facon M., Tokarski C., Rolando C., Szydłowski N. Proteome analysis of potato starch reveals the presence of new starch metabolic proteins as well as multiple protease inhibitors. *Front. Plant Sci*. 2018;9:746. DOI 10.3389/fpls.2018.00746.
- Hennen-Bierwagen T.A., James M.G., Myers A.M. Involvement of debranching enzymes in starch biosynthesis. In: Tetlow I.J. (Ed.) *Essential Reviews in Experimental Biology*. Vol. 5. The Synthesis and Breakdown of Starch. Society for Experimental Biology, London, 2012;179-215.
- Hofvander P., Andersson M., Larsson C.T., Larsson H. Field performance and starch characteristics of high-amylose potatoes obtained by antisense gene targeting of two branching enzymes. *Plant Biotechnol. J*. 2004;2:311-320. DOI 10.1111/j.1467-7652.2004.00073.x.
- Hossain F., Chhabra R., Devi E.L., Zunjare R.U., Jaiswal S.K., Muthusamy V. Molecular analysis of mutant *granule-bound starch synthase-1 (waxy1)* gene in diverse waxy maize inbreds. *3 Biotech*. 2019;9(1):3. DOI 10.1007/s13205-018-1530-6.
- Hou J., Zhang H., Liu J., Reid S., Liu T., Xu S., Tian Z., Sonnewald U., Song B., Xie C. Amylases StAmy23, StBAM1 and StBAM9 regulate cold-induced sweetening of potato tubers in distinct ways. *J. Exp. Bot*. 2017;68(9):2317-2331. DOI 10.1093/jxb/erx076.
- Hussain H., Mant A., Seale R., Zeeman S., Hinchliffe E., Edwards A., Hylton C., Bornemann S., Smith A.M., Martin C., Bustos R. Three isoforms of isoamylase contribute different catalytic properties for the debranching of potato glucans. *Plant Cell*. 2003;15(1):133-149. DOI 10.1105/tpc.006635.
- Kammerer B., Fischer K., Hilpert B., Schubert S., Gutensohn M., Weber A., Flügge U.I. Molecular characterization of a carbon transporter in plastids from heterotrophic tissues: the glucose 6-phosphate/phosphate antiporter. *Plant Cell*. 1998;10(1):105-117. DOI 10.1105/tpc.10.1.105.
- Khlestkin V.K., Peltek S.E., Kolchanov N.A. Target genes for development of potato (*Solanum tuberosum* L.) cultivars with desired starch properties. *Selskokhozyaystvennaya Biologiya = Agricultural Biology*. 2017;52(1):25-36. DOI 10.15389/agrobiology.2017.1.25eng.
- Kleczkowski L.A., Kunz S., Wilczynska M. Mechanisms of UDP-glucose synthesis in plants. *Crit. Rev. Plant Sci*. 2010;29(4):191-203. DOI 10.1080/07352689.2010.483578.
- Kondhare K.R., Malankar N.N., Devani R.S., Banerjee A.K. Genome-wide transcriptome analysis reveals small RNA profiles involved in early stages of stolon-to-tuber transitions in potato under photoperiodic conditions. *BMC Plant Biol*. 2018;18:284. DOI 10.1186/s12870-018-1501-4.
- Kondhare K.R., Patil A.B., Giri A.P. Auxin: an emerging regulator of tuber and storage root development. *Plant Sci*. 2021;306:110854. DOI 10.1016/j.plantsci.2021.110854.
- Kötting O., Kossmann J., Zeeman S.C., Lloyd J.R. Regulation of starch metabolism: the age of enlightenment? *Curr. Opin. Plant Biol*. 2010;13(3):321-329. DOI 10.1016/j.pbi.2010.01.003.
- Leroch M., Kirchnerberger S., Haferkamp I., Wahl M., Neuhaus H.E., Tjaden J. Identification and characterization of a novel plastidic adenine nucleotide uniporter from *Solanum tuberosum*. *J. Biol. Chem*. 2005;280(18):17992-8000. DOI 10.1074/jbc.M412462200.
- Li L., Jiang H.X., Campbell M., Blanco M., Jane J.L. Characterization of maize amylose-extender (*ae*) mutant starches. Part I: Relationship between resistant starch contents and molecular structures. *Carbohydr. Polym*. 2008;74(3):396-404. DOI 10.1016/j.carbpol.2008.03.012.
- Lloyd J.R., Blennow A., Burhenne K., Kossmann J. Repression of a novel isoform of disproportionating enzyme (stDPE2) in potato leads to inhibition of starch degradation in leaves but not tubers stored at low temperature. *Plant Physiol*. 2004;134(4):1347-1354. DOI 10.1104/pp.103.038026.
- López-González C., Juárez-Colunga S., Morales-Eliás N.C., Tiessen A. Exploring regulatory networks in plants: transcription factors of starch metabolism. *Peer. J*. 2019;7:e6841. DOI 10.7717/peerj.6841.
- Lorberth R., Ritte G., Willmitzer L., Kossmann J. Inhibition of a starch-granule-bound protein leads to modified starch and repression of cold sweetening. *Nat. Biotechnol*. 1998;16(5):473-477. DOI 10.1038/nbt0598-473.
- Malinova I., Kössler S., Orawetz T., Matthes U., Orzechowski S., Koch A., Fettke J. Identification of two *Arabidopsis thaliana* plasma membrane transporters able to transport glucose 1-phosphate. *Plant Cell Physiol*. 2020;61(2):381-392. DOI 10.1093/pcp/pcz206.
- McKenzie M.J., Chen R.K., Harris J.C., Ashworth M.J., Brummell D.A. Post-translational regulation of acid invertase activity by vacuolar invertase inhibitor affects resistance to cold-induced sweetening of potato tubers. *Plant Cell Environ*. 2013;36(1):176-185. DOI 10.1111/j.1365-3040.2012.02565.x.
- Mérida A., Fettke J. Starch granule initiation in *Arabidopsis thaliana* chloroplasts. *Plant J*. 2021;107(3):688-697. DOI 10.1111/tbj.15359.



- Müller-Röber B.T., Kossmann J., Hannah L.C., Willmitzer L., Sonnewald U. One of two different ADP-glucose pyrophosphorylase genes from potato respond strongly to elevated levels of sucrose. *Mol. Gen. Genet.* 1990;224:136-146. DOI 10.1007/BF00259460.
- Muth J., Hartje S., Twyman R.M., Hofferbert H.R., Tacke E., Prüfer D. Precision breeding for novel starch variants in potato. *Plant Biotechnol. J.* 2008;6(6):576-584. DOI 10.1111/j.1467-7652.2008.00340.x.
- Nielsen T.H., Deiting U., Stitt M.  $\beta$ -amylase in potato tubers is induced by storage at low temperature. *Plant Physiol.* 1997;113(2):503-510. DOI 10.1104/pp.113.2.503.
- Orzechowski S., Sitnicka D., Grabowska A., Compart J., Fettke J., Zdunek-Zastocka E. Effect of short-term cold treatment on carbohydrate metabolism in potato leaves. *Int. J. Mol. Sci.* 2021;22(13):7203. DOI 10.3390/ijms22137203.
- Pfister B., Zeeman S.C. Formation of starch in plant cells. *Cell Mol. Life Sci.* 2016;73(14):2781-2807. DOI 10.1007/s00018-016-2250-x.
- Qu J., Xu S., Zhang Z., Chen G., Zhong Y., Liu L., Zhang R., Xue J., Guo D. Evolutionary, structural and expression analysis of core genes involved in starch synthesis. *Sci. Rep.* 2018;8:12736. DOI 10.1038/s41598-018-30411-y.
- Reyniers S., Ooms N., Gomand S.V., Delcour J.A. What makes starch from potato (*Solanum tuberosum* L.) tubers unique: a review. *Compr. Rev. Food Sci. Food Saf.* 2020;19(5):2588-2612. DOI 10.1111/1541-4337.12596.
- Ritte G., Heydenreich M., Mahlow S., Haebel S., Kötting O., Stepup M. Phosphorylation of C6- and C3-positions of glucosyl residues in starch is catalysed by distinct dikinases. *FEBS Lett.* 2006;580(20):4872-4876. DOI 10.1016/j.febslet.2006.07.085.
- Roldán I., Watebled F., Lucas M.M., Delvallé D., Planchot V., Jiménez S., Pérez R., Ball S., D'Hulst C., Mérida A. The phenotype of soluble starch synthase IV defective mutants of *Arabidopsis thaliana* suggests a novel function of elongation enzymes in the control of starch granule formation. *Plant J.* 2007;49:492-504. DOI 10.1111/j.1365-313X.2006.02968.x.
- Ruan Y.L. Sucrose metabolism: gateway to diverse carbon use and sugar signaling. *Annu. Rev. Plant Biol.* 2014;65:33-67. DOI 10.1146/annurev-arplant-050213-040251.
- Rüscher D., Corral J.M., Carluccio A.V., Klemens P.A.W., Gisel A., Stavolone L., Neuhaus H.E., Ludewig F., Sonnewald U., Zierer W. Auxin signaling and vascular cambium formation enable storage metabolism in cassava tuberous roots. *J. Exp. Bot.* 2021;72(10):3688-3703. DOI 10.1093/jxb/erab106.
- Salanoubat M., Belliard G. The steady-state level of potato sucrose synthase mRNA is dependent on wounding, anaerobiosis and sucrose concentration. *Gene.* 1989;84:181-185. DOI 10.1016/0378-1119(89)90153-4.
- Samodien E., Jewell J.F., Loedolf B., Oberlander K., George G.M., Zeeman S.C., Damberger F.F., van der Vyver C., Kossmann J., Lloyd J.R. Repression of *Sex4* and *Like Sex Four2* orthologs in potato increases tuber starch bound phosphate with concomitant alterations in starch physical properties. *Front. Plant Sci.* 2018;9:1044. DOI 10.3389/fpls.2018.01044.
- Santana A.L., Meireles M.A.A. New starches are the trend for industry applications: a review. *Food Public Health.* 2014;4(5):229-241. DOI 10.5923/j.fph.20140405.04.
- Santelia D., Kötting O., Seung D., Schubert M., Thalmann M., Bischof S., Meekins D.A., Lutz A., Patron N., Gentry M.S., Alain F.H., Zeeman S.C. The phosphoglucan phosphatase Like Sex Four2 dephosphorylates starch at the C3-position in *Arabidopsis*. *Plant Cell.* 2011;23(11):4096-4111. DOI 10.1105/tpc.111.092155.
- Scheidig A., Fröhlich A., Schulze S., Lloyd J.R., Kossmann J. Downregulation of a chloroplast-targeted  $\beta$ -amylase leads to a starch-excess phenotype in leaves. *Plant J.* 2002;30(5):581-591. DOI 10.1046/j.1365-313x.2002.01317.x.
- Schönhals E.M., Ding J., Ritter E., Paulo J.M., Cara N., Tacke E., Hofferbert H.R., Lübeck J., Strahwald J., Gebhardt C. Physical mapping of QTL for tuber yield, starch content and starch yield in tetraploid potato (*Solanum tuberosum* L.) by means of genome wide genotyping by sequencing and the 8.3 K SolCAP SNP array. *BMC Genomics.* 2017;18:642. DOI 10.1186/s12864-017-3979-9.
- Schwall G.P., Safford R., Westcott R.J., Jeffcoat R., Tayal A., Shi Y.C., Gidley M.J., Jobling S.A. Production of very-high-amylose potato starch by inhibition of SBE A and B. *Nat. Biotechnol.* 2000;18(5):551-554. DOI 10.1038/75427.
- Seung D., Boudet J., Monroe J., Schreier T.B., David L.C., Abt M., Lu K.J., Zanella M., Zeeman S.C. Homologs of PROTEIN TARGETING TO STARCH control starch granule initiation in Arabidopsis leaves. *Plant Cell.* 2017;29(7):1657-1677. DOI 10.1105/tpc.17.00222.
- Seung D., Soyk S., Coiro M., Maier B.A., Eicke S., Zeeman S.C. PROTEIN TARGETING TO STARCH is required for localising GRANULE-BOUND STARCH SYNTHASE to starch granules and for normal amylose synthesis in Arabidopsis. *PLoS Biol.* 2015;13(2):e1002080. DOI 10.1371/journal.pbio.1002080.
- Slugina M.A., Kochieva E.Z. The use of carbohydrate metabolism genes for potato (*Solanum tuberosum* L.). *Selskokhozyaystvennaya Biologiya = Agricultural Biology.* 2018;53(3):450-463. DOI 10.15389/agrobiology.2018.3.450eng.
- Sonnewald S., Sonnewald U. Regulation of potato tuber sprouting. *Planta.* 2014;239(1):27-38. DOI 10.1007/s00425-013-1968-z.
- Sowokinos J.R., Hayes R.J., Thill C.A. Coordinated regulation of cold induced sweetening in tetraploid potato families by isozymes of UDP-glucose pyrophosphorylase and vacuolar acid invertase. *Am. J. Potato Res.* 2018;95:487-494. DOI 10.1007/s12230-018-9653-1.
- Sowokinos J.R., Preiss J. Pyrophosphorylases in *Solanum tuberosum*: III. Purification, physical, and catalytic properties of ADPglucose pyrophosphorylase in potatoes. *Plant Physiol.* 1982;69(6):1459-1466. DOI 10.1104/pp.69.6.1459.
- Stein O., Granot D. An overview of sucrose synthases in plants. *Front. Plant Sci.* 2019;10:95. DOI 10.3389/fpls.2019.00095.
- Stinard P.S., Robertson D.S., Schnable P.S. Genetic isolation, cloning, and analysis of a mutator-induced, dominant antimorph of the maize *amylose extender1* locus. *Plant Cell.* 1993;5(11):1555-1566. DOI 10.1105/tpc.5.11.1555.
- Streb S., Delatte T., Umhang M., Eicke S., Schorderet M., Reinhardt D., Zeeman S.C. Starch granule biosynthesis in *Arabidopsis* is abolished by removal of all debranching enzymes but restored by the subsequent removal of an endoamylase. *Plant Cell.* 2008;20(12):3448-3466. DOI 10.1105/tpc.108.063487.
- Streb S., Zeeman S.C. Starch metabolism in Arabidopsis. *Arabidopsis Book.* 2012;10:e0160. DOI 10.1199/tab.0160.
- Sweetlove L.J., Fernie A.R. The spatial organization of metabolism within the plant cell. *Annu. Rev. Plant Biol.* 2013;64:723-746. DOI 10.1146/annurev-arplant-050312-120233.
- Tauberger E., Fernie A.R., Emmermann M., Renz A., Kossmann J., Willmitzer L., Trethewey R.N. Antisense inhibition of plastidial phosphoglucomutase provides compelling evidence that potato tuber amyloplasts import carbon from the cytosol in the form of glucose-6-phosphate. *Plant J.* 2000;23(1):43-53. DOI 10.1046/j.1365-313x.2000.00783.x.
- Tenorio G., Orea A., Romero J.M., Mérida A. Oscillation of mRNA level and activity of granule-bound starch synthase I in *Arabidopsis* leaves during the day/night cycle. *Plant Mol. Biol.* 2003;51(6):949-958. DOI 10.1023/a:1023053420632.
- Tetlow I.J., Beisel K.G., Cameron S., Makhmoudova A., Liu F., Bresolin N.S., Wait R., Morell M.K., Emes M.J. Analysis of protein complexes in wheat amyloplasts reveals functional interactions among starch biosynthetic enzymes. *Plant Physiol.* 2008;146(4):1878-1891. DOI 10.1104/pp.108.116244.
- Tetlow I.J., Bertoft E. A review of starch biosynthesis in relation to the building block-backbone model. *Int. J. Mol. Sci.* 2020;21:7011. DOI 10.3390/ijms21197011.
- Tiessen A., Hendriks J.H.M., Stitt M., Branscheid A., Gibon Y., Faré E.M., Geigenberger P. Starch synthesis in potato tubers is regulated by post-translational redox modification of ADP-glucose pyrophosphorylase: a novel regulatory mechanism linking starch

- synthesis to the sucrose supply. *Plant Cell*. 2002;14:2191-2213. DOI 10.1105/tpc.003640.
- Tjaden J., Möhlmann T., Kampfenkel K., Neuhaus E. Altered plastidic ATP/ADP-transporter activity influences potato (*Solanum tuberosum* L.) tuber morphology, yield and composition of tuber starch. *Plant J*. 2001;16:531-540. DOI 10.1046/j.1365-313x.1998.00317.x.
- Tsai H.-L., Lue W.-L., Lu K.-J., Hsieh M.-H., Wang S.-M., Chen J. Starch synthesis in Arabidopsis is achieved by spatial cotranscription of core starch metabolism genes. *Plant Physiol*. 2009;151(3):1582-1595. DOI 10.1104/pp.109.144196.
- Tuncel A., Corbin K.R., Ahn-Jarvis J., Harris S., Hawkins E., Smedley M.A., Harwood W., Warren F.J., Patron N.J., Smith A.M. Cas9-mediated mutagenesis of potato starch-branching enzymes generates a range of tuber starch phenotypes. *Plant Biotechnol. J*. 2019;17(12):2259-2271. DOI 10.1111/pbi.13137.
- Van Harselaar J.K., Lorenz J., Senning M., Sonnewald U., Sonnewald S. Genome-wide analysis of starch metabolism genes in potato (*Solanum tuberosum* L.). *BMC Genomics*. 2017;18(1):37. DOI 10.1186/s12864-016-3381-z.
- Visser R.G., Somhorst I., Kuipers G.J., Ruys N.J., Feenstra W.J., Jacobsen E. Inhibition of the expression of the gene for granule-bound starch synthase in potato by antisense constructs. *Mol. Gen. Genet*. 1991;225(2):289-296. DOI 10.1007/BF00269861.
- Wang D., Cheng L., Wang Y., Zhang F. Comparative proteomic analysis of potato (*Solanum tuberosum* L.) tuberization *in vitro* regulated by IAA. *Am. J. Potato Res.* 2018;95(4):395-412. DOI 10.1007/s12230-018-9640-6.
- Wang X., Xue L., Sun J., Zuo J. The *Arabidopsis BE1* gene, encoding a putative glycoside hydrolase localized in plastids, plays crucial roles during embryogenesis and carbohydrate metabolism. *J. Integr. Plant Biol.* 2010;52:273-288. DOI 10.1111/j.1744-7909.2010.00930.x.
- Weise S.E., Schrader S.M., Kleinbeck K.R., Sharkey T.D. Carbon balance and circadian regulation of hydrolytic and phosphorolytic breakdown of transitory starch. *Plant Physiol*. 2006;141(3):879-886. DOI 10.1104/pp.106.081174.
- Yao D., Gonzales-Vigil E., Mansfield S.D. Arabidopsis sucrose synthase localization indicates a primary role in sucrose translocation in phloem. *J. Exp. Bot.* 2020;71(6):1858-1869. DOI 10.1093/jxb/erz539.
- Yu T.S., Lue W.L., Wang S.M., Chen J. Mutation of *Arabidopsis* plastid phosphoglucose isomerase affects leaf starch synthesis and floral initiation. *Plant Physiol*. 2000;123(1):319-326. DOI 10.1104/pp.123.1.319.
- Yu T.S., Zeeman S.C., Thorneycroft D., Fulton D.C., Dunstan H., Lue W.L., Hegemann B., Tung S.Y., Umemoto T., Chapple A., Tsai D.L., Wang S.M., Smith A.M., Chen J., Smith S.M.  $\alpha$ -Amylase is not required for breakdown of transitory starch in *Arabidopsis* leaves. *J. Biol. Chem.* 2005;280(11):9773-9779. DOI 10.1074/jbc.M413638200.
- Zeeman S.C., Kossmann J., Smith A.M. Starch: its metabolism, evolution, and biotechnological modification in plants. *Annu. Rev. Plant Biol.* 2010;61:209-234. DOI 10.1146/annurev-arplant-042809-112301.
- Zhang H., Liu J., Hou J., Yao Y., Lin Y., Ou Y., Song B., Xie C. The potato amylase inhibitor gene *SbAI* regulates cold-induced sweetening in potato tubers by modulating amylase activity. *Plant Biotechnol. J*. 2014;12(7):984-993. DOI 10.1111/pbi.12221.
- Zhang L., Häusler R.E., Greiten C., Hajirezaei M.R., Haferkamp I., Neuhaus H.E., Flügge U.I., Ludewig F. Overriding the co-limiting import of carbon and energy into tuber amyloplasts increases the starch content and yield of transgenic potato plants. *Plant Biotechnol. J*. 2008;6(5):453-464. DOI 10.1111/j.1467-7652.2008.00332.x.
- Zrenner R., Salanoubat M., Willmitzer L., Sonnewald U. Evidence of the crucial role of sucrose synthase for sink strength using transgenic potato plants (*Solanum tuberosum* L.). *Plant J*. 1995;7(1):97-107. DOI 10.1046/j.1365-313x.1995.07010097.x.

#### ORCID ID

E.M. Sergeeva orcid.org/0000-0001-8234-3063  
E.A. Salina orcid.org/0000-0001-8590-847X  
A.V. Kochetov orcid.org/0000-0003-3151-5181

**Acknowledgements.** This work was supported by State Budgeted Project 0259-2021-0012.

**Conflict of interest.** The authors declare no conflict of interest.

Received October 13, 2021. Revised December 20, 2021. Accepted December 20, 2021.