

Supplemental Material

Applied Microbiology and Biotechnology

Distinct alterations of gut microbiota between viral and non-viral-related hepatocellular carcinoma

Thananya Jinato^{1,2}, Songtham Anuntakarun¹, Nantawat Sathawiwat¹, Natthaya Chuaypen^{1*},
Pisit Tangkijvanich^{1*}

¹Center of Excellence in Hepatitis and Liver Cancer, Department of Biochemistry, Faculty of Medicine, Chulalongkorn University, Bangkok, 10330, Thailand

²Doctor of Philosophy Program in Medical Sciences, Graduate Affairs, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

*Correspondence:

Pisit Tangkijvanich, M.D., Center of Excellence in Hepatitis and Liver Cancer, Faculty of medicine, Chulalongkorn University, Bangkok, 10330, Thailand.

pisittkvn@yahoo.com; Tel: +66 2 256 4482

*Co-correspondence:

Natthaya Chuaypen, PhD. Center of Excellence in Hepatitis and Liver Cancer, Faculty of medicine, Chulalongkorn University, Bangkok, 10330, Thailand.

natthaya.c@chula.ac.th; natthaya.ch56@gmail.com; Tel: + 66 2 256 4482

Supplemental Material




Fig. S1 Alpha diversity indices based on (a) Chao1, (b) Simpson, and (c) Shannon compared between HBV-HCC, HCV-HCC, and NBNC-HCC




Fig. S2 Top 50 relative abundance at the genus level in healthy, Viral-HCC and NBNC-HCC