

5-Fluorouracil-induced apoptosis in colorectal cancer cells is caspase-9-dependent and mediated by activation of protein kinase C- δ

NIZAR M. MHAIDAT¹, MOHAMMED BOUKLIHACENE¹ and RICK F. THORNE²

¹Department of Clinical Pharmacy, Jordan University of Science and Technology, Irbid 22110, Jordan;

²Department of Cancer Research Unit, Newcastle University, Newcastle, NSW 2300, Australia

Received October 21, 2013; Accepted May 13, 2014

DOI: 10.3892/ol.2014.2211

Abstract. Elucidation of the molecular mechanisms by which 5-fluorouracil (5-FU) induces apoptosis is required in order to understand the resistance of colorectal cancer (CRC) cells to 5-FU. In the current study, 5-FU-induced apoptosis was assessed using the propidium iodide method. Involvement of protein kinase C (PKC) was assessed by evaluating the extent of their activation in CRC, following treatment with 5-FU, using biochemical inhibitors and western blot analysis. The results revealed that 5-FU induces varying degrees of apoptosis in CRC cells; HCT116 cells were identified to be the most sensitive cells and SW480 were the least sensitive. In addition, 5-FU-induced apoptosis was caspase-dependent as it appeared to be initiated by caspase-9. Furthermore, PKC ϵ was marginally expressed in CRC cells and no changes were observed in the levels of cleavage or phosphorylation following treatment with 5-FU. The treatment of HCT116 cells with 5-FU increased the expression, phosphorylation and cleavage of PKC δ . The inhibition of PKC δ was found to significantly inhibit 5-FU-induced apoptosis. These results indicated that 5-FU induces apoptosis in CRC by the activation of PKC δ and caspase-9. In addition, the levels of PKC δ activation may determine the sensitivity of CRC to 5-FU.

Introduction

Colorectal cancer (CRC) is the third most common malignancy in the populations of developed western countries and represents the third leading cause of cancer-associated mortality in the USA (1). Although the incidence of CRC varies by ~20-fold worldwide, in Jordan, CRC is the most common type

of cancer in males and the second, following breast cancer, in females (2). Once the disease spreads to distant sites, it is usually incurable using the current systemic treatment options, including chemotherapy. This is largely due to the resistance of cancer cells to apoptosis. Understanding and overcoming the resistance of cancer cells to apoptosis may simplify the identification of novel therapeutic targets and the development of novel treatment options.

The chemotherapeutic agent, 5-fluorouracil (5-FU), induces cytotoxic effects via targeting the metabolism of RNA bases (3). It is predominantly administered in the treatment of various types of cancer, including breast, aerodigestive tract and head cancer. In CRC, 5-FU monopolizes a place of choice even in individuals with advanced metastatic disease (4,5). The response rate associated with 5-FU monotherapy is only 10-15%, however, in a small number of cases the response rate was >50% when associated with other types of medication (6-8).

The protein kinase C (PKC) family includes 11 distinct members, which share a similar serine (Ser)/threonine (Thr) structure (9,10). These polypeptide isoenzymes are classically classified into three distinct groups according to the second messengers to which they respond. Conventional PKCs, PKC α , β 1, β 2 and γ , may be activated by diacylglycerol (DAG) and calcium (Ca²⁺). The activation of novel PKCs, PKC δ , θ , η and ϵ , is DAG-dependent and Ca²⁺-independent. The activation of atypical PKCs, PKC ι/λ and ζ , is independent of these second messengers (11). PKC ϵ is an antiapoptotic enzyme, which promotes cell proliferation and resistance to chemotherapeutic agents (12), whereas PKC δ is classified among the proapoptotic PKCs as its cleavage and activation promotes apoptotic cell death (13).

Materials and methods

Cell lines. The human CRC cell lines, SW480, SW620, HCT116 and HT29, were obtained from Dr Rick F. Thorne (University of Newcastle, Newcastle, NSW, Australia) and were cultured in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum.

Antibodies and other reagents. 5-FU was obtained from Ebewe Pharma (Unterach, Austria), stored at room

Correspondence to: Dr Nizar M. Mhaidat, Department of Clinical Pharmacy, Faculty of Pharmacy, Jordan University of Science and Technology, 19 Ramtha Street, Irbid 22110, Jordan
E-mail: nizarm@just.edu.jo

Key words: protein kinase C, colorectal cancer, 5-fluorouracil, apoptosis

temperature, dissolved in dimethyl sulfoxide and prepared as a stock solution of 200 μ M immediately prior to use. The pan-caspase inhibitor, Z-VAD-fmk; the caspase-2-specific inhibitor, z-VDVAD-fmk; the caspase-3-specific inhibitor, z-DEVAD-fmk; the caspase-8-specific inhibitor, z-IETD-fmk; and the caspase-9-specific inhibitor, z-LEHD-fmk, were purchased from R&D Systems (Minneapolis, MN, USA). The general PKC inhibitor, bisindolylmaleimide I (GF109203X), and the specific PKC δ inhibitor, rottlerin, were purchased from Calbiochem (La Jolla, CA, USA), while the cell-permeable-specific PKC ϵ inhibitor, epsilon-V1-2 inhibitor Cys-conjugated, was obtained from AnaSpec Inc. (Fremont, CA, USA). The mouse anti-human monoclonal antibodies directed against caspase-2, -3, -8 and -9, propidium iodide (PI) and 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) were obtained from Sigma-Aldrich (St. Louis, MO, USA). The rabbit polyclonal antibodies, anti-PKC ϵ , p-PKC, PKC δ and p-PKC δ Ser 645, and the mouse monoclonal antibody directed against p-PKC δ , Thr 505, were obtained from Santa Cruz Biotechnology Inc. (Santa Cruz, CA, USA).

Cell viability analysis. The MTT assay was performed as previously described to assess the viability of the CRC cells following treatment with 5-FU (14).

Apoptosis. The quantification of the apoptotic cells was obtained by measuring the sub-G1 population of apoptotic cells using flow cytometry following staining with PI as previously described (15).

Western blot analysis. Changes in protein expression induced by 5-FU were assessed by western blot analysis as previously described (15).

Statistical analysis. Data are presented as the mean \pm standard error. The statistical significance of intergroup differences in normally distributed continuous variables was determined using Student's t-test. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

5-FU induces the caspase-dependent apoptosis of human CRC cells. To examine the antitumor potential of 5-FU, cell viability analysis was performed using the MTT assay on a panel of CRC cell lines. The cells were incubated with a wide range of 5-FU concentrations (0, 1, 10, 25, 50, 100 and 200 μ M) for 72 h. As shown in Fig. 1, 5-FU induced growth inhibition in a dose-dependent manner in all CRC cells, to different degrees. The optimal concentration was 50 μ M. HCT116 cells were the most sensitive cells followed by HT29 and SW620, and the least sensitive cell line was SW480. In addition, kinetic analysis revealed that 5-FU (50 μ M) induced a significant growth inhibition following 72 h of incubation (Fig. 2).

To investigate whether the 5-FU-induced reduction in cell viability is due to the induction of apoptotic cell death, the cellular and molecular apoptotic events were evaluated following treatment with 5-FU (50 μ M) for 72 h. The results showed different degrees of CRC sensitivity to 5-FU-induced




Figure 1. 5-FU inhibits CRC cell growth. HCT116, SW480, HT29 and SW620 human CRC cell lines were treated with a wide range of 5-FU prior to assess cell viability using the 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide assay. Data are presented as the mean \pm standard error of three individual experiments. 5-FU, 5-fluorouracil; CRC, colorectal cancer.




Figure 2. 5-Fluorouracil (FU) induces time-dependent cell growth inhibition in colorectal cancer cells. HCT116 and SW480 cells were treated with 5-FU (50 μ M) for different time periods. The cell viability was determined using the 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide assay. Data are representative of three individual experiments.




Figure 3. 5-FU-induced apoptosis in a panel of human CRC cell lines. CRC cells were treated with 5-FU (50 μ M) for 72 h. The apoptotic cell fractions were determined using the propidium iodide method. Data are representative of three independent experiments. 5-FU, 5-fluorouracil; CRC, colorectal cancer.

apoptosis. HCT116 represented the most sensitive cell line, HT29 and SW620 were intermediately sensitive, and SW480 cells were the least sensitive (Fig. 3).

To assess the involvement of caspases in 5-FU-induced apoptosis, HCT116 and SW480 cells were pretreated with the




Figure 4. 5-Fluorouracil (FU) induces caspase-dependent apoptosis. HCT116 and SW480 cells were pretreated with the pan-caspase inhibitor, Z-VAD-fmk (20 μ M), for 2 h prior to the addition of 5-FU for an additional 72 h. Apoptosis was measured using the propidium iodide method using flow cytometry. Data are presented as the mean \pm standard error of three individual experiments. *P<0.05 vs. 5-FU treatment group.




Figure 5. Effect of specific caspase inhibitors on 5-FU-induced apoptosis. HCT116 cells were pretreated with the pan-caspase inhibitor, z-VAD-fmk (20 μ M); the caspase-2 inhibitor, z-VDVAD-fmk (30 μ M); the caspase-3 inhibitor, z-DEVAD-fmk (30 μ M); the caspase-8 inhibitor, z-IETD-fmk (30 μ M); and the caspase-9 inhibitor, z-LEHD-fmk (30 μ M) 2 h prior to the addition of 5-FU for an additional 72 h. Apoptosis was measured using the propidium iodide method using flow cytometry. Data are presented as the mean \pm standard error of three individual experiments. *P<0.05 vs. 5-FU treatment group. 5-FU, 5-fluorouracil.

pan-caspase inhibitor, Z-VAD-fmk (20 μ M), and the specific inhibitors of caspase-2, z-VDVAD-fmk (30 μ M); caspase-3, z-DEVAD-fmk (30 μ M); caspase-8, z-IETD-fmk (30 μ M); and caspase-9, Z-LEHD-fmk (30 μ M). The inhibitors were added 2 h prior to the addition of 5-FU (50 μ M) for an additional 72 h.

As shown in Fig. 4, the pretreatment with the pan-caspase inhibitor, Z-VAD-fmk, was found to significantly inhibit the 5-FU-induced apoptosis in HCT116 cells, which indicates that 5-FU induces a caspase-dependent apoptosis. In addition, these results indicate that the 5-FU-induced apoptosis in HCT116 cells was also significantly inhibited by the specific caspase-9 inhibitor, Z-LEHD-fmk. The inhibition of caspase-2, -3 and -8 exhibited a minimal effect on 5-FU-induced apoptosis.

The kinetics of caspase activation by 5-FU was investigated by western blot analysis (Fig. 6). The cleavage of caspase-9




Figure 6. 5-FU induces caspase-9-dependent apoptosis. HCT116 and SW480 cells were treated with 5-FU (50 μ M) for the indicated time points (0, 16, 24 and 36 h). Whole cell lysates were subjected to western blot analysis and western blot analysis of the GAPDH levels was included to demonstrate that equivalent amounts of protein were loaded in each lane. Data are representative of two individual experiments. 5-FU, 5-fluorouracil; GAPDH, glyceraldehyde-3-phosphate dehydrogenase.




Figure 7. Effect of PKC inhibition on 5-FU-induced apoptosis. HCT116 and SW480 cells treated with 5-FU were pretreated with or without PKC I, bisindolylmaleimide I (5 μ M), 2 h prior to the addition of 5-FU (50 μ M) for an additional 72 h. Apoptosis was measured using the propidium iodide method using flow cytometry. Data are presented as the mean \pm standard error of three individual experiments. 5-FU, 5-fluorouracil; PKC, protein kinase C; I, inhibitor.

was evident at 16 h in the HCT116 cells, while the cleavage of the other caspases was not detected in the HCT116 or the SW480 cell lines. These findings demonstrated that caspase-9 is the initial caspase in 5-FU-induced apoptosis.

5-FU-induced apoptosis is mediated by PKC δ . To assess the involvement of PKC activation in 5-FU-induced apoptosis, the HCT116 and SW480 cell lines were pretreated with the pan-PKC inhibitor, bisindolylmaleimide I (2.5 or 5 μ M; data not shown for 2.5 μ M), the specific PKC ϵ inhibitor, epsilon-VI-2 inhibitor cys-conjugated (10 μ M), and the specific PKC δ inhibitor, rottlerin (5 μ M). The inhibitors were added 2 h prior to the addition of 5-FU for an additional 72 h.

As shown in Fig. 7, pretreatment with the pan-PKC inhibitor reduced the 5-FU-induced apoptosis in HCT116 and




Figure 8. Effect of PKC δ and PKC ϵ inhibition on 5-FU-induced apoptosis. HCT116 and SW480 cells were pretreated with PKC ϵ I, epsilon-VI-2 inhibitor cys-conjugated (10 μ M), or PKC δ I, rottlerin (5 μ M), 2 h prior to the addition of 5-FU for an additional 72 h. The apoptotic percentage was measured using the propidium iodide method. Data are presented as the mean \pm standard error of three individual experiments. * P <0.05 vs. 5-FU treatment group. 5-FU, 5-fluorouracil; PKC, protein kinase C; I, inhibitor.




Figure 9. 5-FU induces PKC δ activation in the HCT116 cell line. HCT116 and SW480 cells were treated with 5-FU (50 μ M) for the indicated time points (0, 16, 24 and 36 h). Whole cell lysates were subjected to western blot analysis and western blot analysis of the GAPDH levels was included to demonstrate that equivalent amounts of protein were loaded in each lane. Data are representative of two individual experiments. 5-FU, 5-fluorouracil; PKC, protein kinase C; GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

did not affect the response in SW480. This may indicate that the PKC pathway is activated by 5-FU in CRC cells, however, does not provide information on the implicated isoform(s). The pretreatment of the two cell lines with the specific PKC ϵ inhibitor did not sensitize the cells to the 5-FU-induced apoptosis. By contrast, the pretreatment of HCT116 with rottlerin significantly inhibited 5-FU-induced apoptosis in HCT116 cells (Fig. 8). In addition, the kinetics of PKC ϵ and PKC δ activation in HCT116 and SW480 cells prior to and following exposure to 5-FU were investigated. The results shown in Fig. 9 demonstrate that the full-length PKC ϵ was particularly small in the two cell lines prior to and following treatment, and did

not undergo cleavage or phosphorylation following treatment with 5-FU. Additionally, it appeared that the full-length PKC δ was gradually upregulated and concomitantly cleaved to its active form following exposure to 5-FU in HCT116, although not in SW480. In the sensitive cells, the phosphorylation of PKC δ at Thr 505 preceded that which occurred at Ser 645 and was downregulated. Taken together, these results may indicate that PKC ϵ is not involved in the resistance of CRC cells to 5-FU and that 5-FU induces PKC δ activation in CRC cells.

Discussion

In the present study, molecular and biochemical experiments were conducted to evaluate the *in vitro* cytotoxic effect of 5-FU against a panel of CRC cells. The cytotoxicity of 5-FU appeared to be due to the induction of apoptosis, which was identified by the PI assay. The results revealed that HCT116 was the most sensitive cell line followed by HT29 and SW620, while SW480 cells were the least sensitive.

The caspases, a family of cysteine proteases, are major mediators of the execution phase of apoptosis; possibly by direct activation of the death receptor or following mitochondrial changes (16,17). Caspase-9 and -8 are generally considered to be the initiator caspases in chemotherapy-induced apoptosis. Caspase-2 is unique in the family of caspases as it is the only caspase that may be involved in the initiation and execution of apoptosis (18-20). Thus, various studies have indicated that it is caspase-2, and not caspase-9, that initiates the DNA damage-induced apoptosis (19,21,22).

In the present study, kinetic analysis of caspase cascade activation using western blot analysis with specific caspase inhibitors revealed that the activation of caspase-9 is the initiating event, which establishes caspase-9 as the apical caspase in the 5-FU-induced apoptosis. This was consistent with previous reports indicating that caspase-9 may act as an initiator caspase in cisplatin-induced apoptosis (23,24).

The activation of the PKC ϵ isoform has been reported to be antiapoptotic in various cellular systems, including lung and prostate cancer cells (25,26). Whereas, its overexpression has been found to inhibit the apoptosis of melanoma (27) and glioma (11) cells. In the current study, the full-length PKC ϵ was found to be particularly small in the two cell lines prior to and following treatment with 5-FU, and was not found to undergo cleavage or phosphorylation following treatment with 5-FU, which indicated that PKC ϵ is not involved in the resistance of CRC to 5-FU-induced apoptosis.

The cleavage and activation of PKC δ has been reported as a response to various apoptotic stimuli, including radiation, oxidative stress and chemotherapeutic agents (13,28). These stimuli may activate the enzyme by phosphorylation and cleavage (26). In the present study, 5-FU was found to induce the activation of PKC δ in HCT116 cells, however, not in the SW480 cell line.

The kinetics of PKC δ activation following treatment with 5-FU in CRC cells was studied by western blot analysis using specific antibodies against PKC δ , as well as p-PKC δ (F-7) that targets the phosphorylated Thr 505 of PKC δ (the activation loop motif), and p-PKC (Ser 645), which targets the phosphorylated Ser 645 of PKC δ (the turn motif). The phosphorylation of PKC δ in the activation loop preceded that which occurred at

the turn motif and was downregulated. Similarly, inactivation of PKC δ implied the initial dephosphorylation at Thr 505 prior to that at Ser 645. In the present study, it is hypothesized that PKC δ activation following treatment with 5-FU may occur in a stepwise manner, initially requiring phosphorylation of the activation loop, followed by phosphorylation of the turn motif.

Although the current study demonstrates that the specific phosphorylations of PKC δ at the sites of the activation loop and the turn motif (critical steps prior to the effective activation of the protein), previous studies have indicated that these phosphorylations are not a prerequisite for the enzymatic activity of PKC δ (29), and that the only structural modification enabling PKC δ to exert its proapoptotic activity is its cleavage to the fully active 40-kDa catalytic fragment (CF) (30). In HCT116 cells, the current study showed that 5-FU increased the expression of full-length PKC δ and induced the cleavage of the enzyme to its CF. These findings may indicate that 5-FU induces PKC δ activation in CRC cells.

PKC δ promotes apoptosis by acting on different signaling pathways, including the mitogen-activated protein kinase (MAPK) signaling pathway and its members; p38 (31), extracellular signal-regulated kinase (32) and c-Jun N-terminal kinase (13). Furthermore, previous studies have shown that the c-Abl-PKC δ -p38 MAPK signaling pathway may trigger the mitochondrial apoptotic signaling pathway by activating the proapoptotic Bcl-2 proteins, Bax and Bak (33,34). Furthermore, caspase-9 is activated during the early stages of the intrinsic apoptotic signaling pathway, immediately following the formation of the apoptosome, although predominantly prior to all other caspases (35). Therefore, in the present study it was hypothesized that when there is a cross-link between the 5-FU-induced PKC δ activation and the caspase-dependent apoptosis that is induced by 5-FU in CRC cells, it may be via the c-Abl-PKC δ -p38 MAPK signaling pathway.

In conclusion, the results of the present study indicate that PKC ϵ is not involved in the resistance of CRC cells to 5-FU chemotherapy, and that 5-FU may induce apoptosis by activating PKC δ and caspase-9. Our ongoing studies will be directed towards elucidating the exact sequence of events, which are triggered following the activation of PKC δ by 5-FU in CRC and to evaluate the signaling of caspase-9 activation.

Acknowledgements

The authors would like to acknowledge the Jordan University of Science & Technology (Irbid, Jordan) for providing financial support (grant no. 60-2012).

References

- Jemal A, Siegel R, Ward E, *et al*: Cancer statistics, 2008. *CA Cancer J Clin* 58: 71-96, 2008.
- Al-Tarawneh M, Khatib S and Arqub K: Cancer incidence in Jordan, 1996-2005. *East Mediterr Health J* 16: 837-845, 2010.
- Longley DB, Harkin DP and Johnston PG: 5-fluorouracil: mechanisms of action and clinical strategies. *Nat Rev Cancer* 3: 330-338, 2003.
- Willett CG, Czito BG and Bendell JC: Radiation therapy in stage II and III rectal cancer. *Clin Cancer Res* 13: 6903-6908, 2007.
- Goldberg RM, Rothenberg ML, Van Cutsem E, *et al*: The continuum of care: a paradigm for the management of metastatic colorectal cancer. *Oncologist* 12: 38-50, 2007.
- Loehrer PJ Sr, Turner S, Kubilis P, *et al*: A prospective randomized trial of fluorouracil versus fluorouracil plus cisplatin in the treatment of metastatic colorectal cancer: a Hoosier Oncology Group trial. *J Clin Oncol* 6: 642-648, 1988.
- Douillard JY, Cunningham D, Roth AD, *et al*: Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *Lancet* 355: 1041-1047, 2000.
- Giacchetti S, Perpoint B, Zidani R, *et al*: Phase III multicenter randomized trial of oxaliplatin added to chronomodulated fluorouracil-leucovorin as first-line treatment of metastatic colorectal cancer. *J Clin Oncol* 18: 136-147, 2000.
- Newton AC: Protein kinase C: structural and spatial regulation by phosphorylation, cofactors, and macromolecular interactions. *Chem Rev* 101: 2353-2364, 2001.
- Nishizuka Y: Protein kinase C and lipid signaling for sustained cellular responses. *FASEB J* 9: 484-496, 1995.
- Basu A, Lu D, Sun B, *et al*: Proteolytic activation of protein kinase C-epsilon by caspase-mediated processing and transduction of antiapoptotic signals. *J Biol Chem* 277: 41850-41856, 2002.
- Akita Y: Protein kinase C-epsilon (PKC-epsilon): its unique structure and function. *J Biochem* 132: 847-52, 2002.
- Reyland ME, Anderson SM, Matassa AA, Barzen KA and Quissell DO: Protein kinase C delta is essential for etoposide-induced apoptosis in salivary gland acinar cells. *J Biol Chem* 274: 19115-19123, 1999.
- Mhaidat NM, Zhang XD, Allen J, *et al*: Temozolomide induces senescence but not apoptosis in human melanoma cells. *Br J Cancer* 97: 1225-1233, 2007.
- Mhaidat NM, Wang Y, Kiejda KA, Zhang XD and Hersey P: Docetaxel-induced apoptosis in melanoma cells is dependent on activation of caspase-2. *Mol Cancer Ther* 6: 752-761, 2007.
- Ashkenazi A and Dixit VM: Death receptors: signaling and modulation. *Science* 281: 1305-1308, 1998.
- Jaattela M: Escaping cell death: survival proteins in cancer. *Exp Cell Res* 248: 30-43, 1999.
- Tinel A and Tschopp J: The PIDDosome, a protein complex implicated in activation of caspase-2 in response to genotoxic stress. *Science* 304: 843-846, 2004.
- Lin CF, Chen CL, Chang WT, *et al*: Sequential caspase-2 and caspase-8 activation upstream of mitochondria during ceramideand etoposide-induced apoptosis. *J Biol Chem* 279: 40755-40761, 2004.
- Baliga BC, Read SH and Kumar S: The biochemical mechanism of caspase-2 activation. *Cell Death Differ* 11: 1234-1241, 2004.
- Vakifahmetoglu H, Olsson M, Orrenius S and Zhivotovsky B: Functional connection between p53 and caspase-2 is essential for apoptosis induced by DNA damage. *Oncogene* 25: 5683-5692, 2006.
- Guo Y, Srinivasula SM, Druilhe A, Fernandes-Alnemri T and Alnemri ES: Caspase-2 induces apoptosis by releasing proapoptotic proteins from mitochondria. *J Biol Chem* 277: 13430-13437, 2002.
- Basu A, Woollard MD and Johnson CL: Involvement of protein kinase C-delta in DNA damage-induced apoptosis. *Cell Death Differ* 8: 899-908, 2001.
- Slee EA, Harte MT, Kluck RM, *et al*: Ordering the cytochrome c-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. *J Cell Biol* 144: 281-292, 1999.
- Ding L, Wang H, Lang W and Xiao L: Protein kinase C-epsilon promotes survival of lung cancer cells by suppressing apoptosis through dysregulation of the mitochondrial caspase pathway. *J Biol Chem* 277: 35305-35313, 2002.
- McJilton MA, Van Sikes C, Wescott GG, *et al*: Protein kinase Cepsilon interacts with Bax and promotes survival of human prostate cancer cells. *Oncogene* 22: 7958-7968, 2003.
- Gillespie S, Zhang XD and Hersey P: Variable expression of protein kinase C epsilon in human melanoma cells regulates sensitivity to TRAIL-induced apoptosis. *Mol Cancer Ther* 4: 668-676, 2005.
- Matassa AA, Carpenter L, Biden TJ, Humphries MJ and Reyland ME: PKCdelta is required for mitochondrial-dependent apoptosis in salivary epithelial cells. *J Biol Chem* 276: 29719-29728, 2001.
- Stempka L, Girod A, Müller HJ, *et al*: Phosphorylation of protein kinase Cdelta (PKCdelta) at threonine 505 is not a prerequisite for enzymatic activity. Expression of rat PKCdelta and an alanine 505 mutant in bacteria in a functional form. *J Biol Chem* 272: 6805-6811, 1997.

30. Emoto Y, Kasaki H, Manome Y, Kharbanda S and Kufe D: Activation of protein kinase C δ in human myeloid leukemia cells treated with 1-beta-D-arabinofuranosylecytosine. *Blood* 87: 1990-1996, 1996.
31. Gomel R, Xiang C, Finniss S, *et al*: The localization of protein kinase C δ in different subcellular sites affects its proapoptotic and antiapoptotic functions and the activation of distinct downstream signaling pathways. *Mol Cancer Res* 5: 627-639, 2007.
32. Basu A and Tu H: Activation of ERK during DNA damage-induced apoptosis involves protein kinase C δ . *Biochem Biophys Res Commun* 334: 1068-1073, 2005.
33. Owens TW, Valentijn AJ, Upton JP, *et al*: Apoptosis commitment and activation of mitochondrial Bax during anoikis is regulated by p38MAPK. *Cell Death Differ* 16: 1551-1562, 2009.
34. Choi SY, Kim MJ, Kang CM, *et al*: Activation of Bak and Bax through c-abl-protein kinase C δ -p38 MAPK signaling in response to ionizing radiation in human non-small cell lung cancer cells. *J Biol Chem* 281: 7049-7059, 2006.
35. Vucic D, Dixit VM and Wertz IE: Ubiquitylation in apoptosis: a post-translational modification at the edge of life and death. *Nat Rev Mol Cell Biol* 12: 439-452, 2011.