



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Management of mediastinal parathyroid adenoma via minimally invasive thoracoscopic surgery: Case report



Saulat Hasnain Fatimi^a, Hina Inam^a, Farida Karim Chagan^b, Usama Khalid Choudry^{c,*}

^a Department of Cardiothoracic Surgery, Aga Khan University Hospital, Pakistan

^b Department of Ophthalmology, Aga Khan University Hospital, Pakistan

^c Department of Post Graduate Medical Education, Aga Khan University Hospital, Karachi, Pakistan

ARTICLE INFO

Article history:

Received 22 May 2017

Received in revised form 27 August 2017

Accepted 28 August 2017

Available online 8 September 2017

Keywords:

Case report

VATS

Ectopic parathyroid adenoma

Idiopathic hypercalcemia

ABSTRACT

INTRODUCTION: The most common cause of chronic hypercalcemia is primary hyperparathyroidism (PHPT). However, owing to the diverse presentation of hypercalcemia, the diagnosis often goes unnoticed culminating as a continuum of recurrence of symptoms. Nephrolithiasis, decreased bone mineral density and peptic ulcer disease are the main clinical sequelae. Among the causes of PHPT 80% are caused by parathyroid adenomas (PA). However, only rarely, these adenomas are found ectopically.

PRESENTATION OF CASE: We present the case of a 66-year-old female with a history of recurrent renal stones and peptic ulcer disease. She was found to have elevated serum calcium and PTH levels. However, subsequent high resolution CT scan of chest and neck failed to demonstrate any abnormality. Therefore, an anterior planar Technetium-99m-sestamibi (MIBI) scintigraphy scan using a single-tracer was done and it identified ectopic anterior mediastinal parathyroid adenoma. The patient was successfully managed with video-assisted thoracoscopic surgery and excision of the mass with follow up calcium level monitoring.

DISCUSSION: An elevated calcium level should prompt a thorough workup, as sometimes it's the only clue to the unrelated and diversified systemic manifestations of hypercalcemia. Hyperparathyroidism due to ectopic adenoma is quite rare and possess a diagnostic and management challenge.

CONCLUSION: Symptomatic hypercalcemia and high level of PTH without local PA should alert physicians to search for ectopic locations through imaging. VATS is a safe and effective minimally invasive procedure for the resection of ectopic mediastinal PA and it should be considered as the first line approach for resection of these ectopic tumors.

© 2017 Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The following case report has been reported from Aga Khan University Hospital which is an internationally recognized teaching hospital and a tertiary care centre based in Pakistan, in accordance with the SCARE guidelines for case reports [1]. Parathyroid adenomas are the most common cause of PTHP [2]. However, the prevalence of ectopic PA is about 2%–43% in anatomical series and reaches upto 16% and 14% in primary and secondary hyperparathyroidism, respectively [3]. Among the various ectopic locations, mediastinal ectopic PAs are extremely rare tumors, constituting 1–2% of all PAs [4]. Due to the rare possibility of ectopic PA, there is a diagnostic and management improbability, which may delay the diagnosis and subsequent surgery due to difficulties in locating the ectopic PAs. Therefore, it is essential for physicians to keep a high index of suspicion while coming across symptomatic malignant

hypercalcemia with raised PTH levels. An accurate pre-operative localization is the key to successful surgery for an ectopic PA. The two most commonly employed imaging modalities to locate PAs include ultrasonography (US) and 99mTc-MIBI scintigraphy scan having a combined sensitivity of 88%–95% [3]. CT and MRI are alternative options with sensitivities of 65% and 78% respectively. The resection of mediastinal ectopic PAs has been traditionally done via median sternotomy and thoracotomy. However, Video assisted thoracoscopic surgery (VATS) has proved to be a safe and successful approach for management of mediastinal ectopic PA in our experience.

2. Case report

A 66-year old female housewife (BMI: 24.1 kg/m²), from a remote city of Pakistan, presented to the surgical OPD with the presenting complaint of intermittent flank pain for the past two years with a history of recurrent renal stones. She had undergone a percutaneous nephrolithotomy (PCNL) 14 years back for nephrolithiasis secondary to calcium oxalate stones. She later had a right extra-

* Corresponding author.

E-mail addresses: uk.choudry@hotmail.com, harry.potter811@hotmail.com (U.K. Choudry).




Fig. 1. a. 99mTc-MIBI scintigraphy scan demonstrating PTH adenomas. b. 99mTc-MIBI scintigraphy scan demonstrating ectopic PTH adenomas in anterior mediastinum.

corporeal shock wave lithotripsy (ESWL) for a residual stones 2 years ago. CT scan of the kidneys showed non-obstructing bilateral renal calculi along with cholelithiasis in the gall bladder. Patient also had multiple gastroenterology clinic visits with the complaints of postprandial bloating and recurrent oral ulcers. Bone densitometry showed marked osteopenia. Her serum calcium level was found elevated at 12.6 mg/dl. The constellation of her symptoms and the elevated calcium levels warranted further workup. Serum PTH was 403 pg/ml (16–87) and serum phosphorus was 2.2 mg/dl (2.5–4.5) respectively; she was diagnosed with primary hyperparathyroidism, aka, PTH dependent hyperparathyroidism causing malignant hypercalcemia. In order, to rule out the possibility of parathyroid adenoma, a high resolution CT scan of chest and neck where carried out, but, there was no abnormal enlargement of the parathyroid gland present, however, 99mTc-MIBI scintigraphy scan was done that showed a well defined area of abnormal tracer deposition in anterior mediastinum just below the manubriosternal junction confirming the diagnosis of mediastinal ectopic PA, furthermore, parathyroid adenomas were outlined over lower pole of right thyroid in anterior mediastinum as well. (Fig. 1a and b).

Pre-operatively, injection pamidronate was administered three week prior to surgery. Repeat chemistry showed a serum calcium level of 10.2 mg/dl. She, subsequently, underwent VATS where right lower parathyroid tissue was identified and excised. The histopathology report established parathyroid parenchyma composed of chief and oxyphil cells with intervening fibroadipose tissue. There was no evidence of malignancy. The ectopic PA was

excised as a whole from anterior mediastinum just behind the sternum and right inferior parathyroidectomy was also carried out. Postoperatively, serum calcium level was repeated every 12 hourly and calcium with vitamin D supplements were administered. Follow up chemistry panel showed a serum calcium in decreasing trend from 9.2–8.2 mg/dl. Her PTH levels also remarkably dropped to 23 pg/dl. She had a complete recovery. Upon return to clinic her serum calcium level and PTH level were found within the normal ranges; 8.9 mg/dl and 86.90 pg/dl respectively. Post operative MR imaging also showed no residual tumor. Patient remained asymptomatic at 6 month follow up as well.

3. Discussion

Symptomatic hypercalcemia can present with several different clinical manifestations and possess a difficulty in diagnosis. Elevated calcium levels should always alert the physician and it should be followed by a thorough endocrinology workup, especially PTH levels. Our patient had complaints involving different systems including GI, renal and musculoskeletal. However, the only clue to the endocrine issue was the elevated serum calcium level, which unfortunately remained unnoticed for a long time, leading to the development of symptoms involving multiple systems. The link between malignant hypercalcemia and PHPT is now completely established as per the recent literature available [22].

The incidence of PHPT is approximately 25 per 100,000 in the general population that increases with age [5]. PHPT is classified as being caused by a single adenoma (75%–85%), hyperplasia (10%–20%), multiple adenomas (4%–5%) or carcinoma (1%) giving the histopathological characteristics and the number of involved glands [5]. The superior and inferior parathyroid glands are derivatives of the third and fourth pharyngeal pouches, respectively. During embryogenesis, these glands descend along with the thyroid gland, sharing the same route, while detaching from the pharyngeal apparatus. However, if they fail to descend, they are found high in the neck under the mandible as undescended parathyroids. On the other hand, if they descend too far, they are found within the mediastinum as ectopic parathyroids [4]. The base of the tongue upto the mediastinum can be regarded as potential positions for ectopic PAs [6] Ectopic parathyroid gland adenoma is a rare tumor, particularly causing PHPT. The clinical and laboratory features of patients with primary hyperparathyroidism due to ectopic adenomas have been reported to be more severe than those with localized parathyroid adenomas, since the former remained undetected and may manifest with higher calcium levels and more frequent primary hyperparathyroidism related bone disease more frequently [7].

The localization and subsequent treatment of an ectopic PA can usually be challenging. There are various imaging modalities available for the diagnosis include ultrasonography, CT scan, magnetic resonance imaging (MRI), positron emission tomography (PET), single photon emission tomography (SPECT) and Technetium (Tc99m-Sestamibi) scintigraphy scan. Among these, U/S is the most widely utilized modality due to its low cost and easy availability. High-resolution ultrasound can accurately localize adenomas relative to the thyroid gland. However, its ability to detect abnormalities depends on the experience and skill of the operator, and therefore, its sensitivity in localization of enlarged parathyroid glands varies greatly (44%–87%) [3]. Recently, scintigraphy scan is increasingly being utilized for the localization of ectopic PA with reported sensitivity of up to 90% [8]. High-resolution ultrasound and parathyroid scintigraphy both have their own advantages. The former is good at localizing adenomas relative to the thyroid gland, whereas the latter is usually reserved for detecting adenomas in

multi-nodular thyroid disease and at ectopic sites, whose incidence is up to 20% among all patients with suspicion of PHTP [9,17].

SPECT is combined with sestamibi scintigraphy to provide three-dimensional imaging. These hybrid imaging techniques with SPECT/CT combining scintigraphic datasets (SPECT) and anatomical (CT) has become increasingly valuable over the last few years [3]. Akram et al. in their case series found a study of 16 patients, in which SPECT/CT identified 39% more lesions compared with SPECT imaging alone. SPECT/CT permitted higher reader confidence in localization, especially for mediastinal adenomas [18].

The combination of MIBI with US raises the sensitivity to 78%–96% [19] and with CT or MRI may raise the sensitivity and specificity to 100% [20]. These additional combinations are moderately cost effective and are indicated as a means for routine preoperative localization of ectopic parathyroid adenomas, especially in cases with faint 99mTc-MIBI scans [21].

The conventional approach for the management of PHPT due to PA is neck exploration followed by excision of the tumor via median sternotomy or thoracotomy. However, ectopic PAs have been renowned as the most frequent reason for a failed cervical exploration [10]. Consequently, precise preoperative localization is the key to successful surgical removal. The choice of the surgical technique greatly relies on the anatomical site of the ectopic PAs. Before the introduction of VATS to thoracic surgery, mediastinal ectopic PAs were resected by a lateral thoracotomy or a median sternotomy [11]. These surgical interventions are associated with a risk of injury to phrenic and recurrent laryngeal nerve, laceration of brachiocephalic vein, delayed wound healing, wound infection, mediastinitis and even death. The size of surgical incisions is often large and aesthetically unsound for the patient [12].

VATS offers significant advantages over both procedures and is being more frequently employed now [13]. It allows better visualization of the tumor, a shorter operative time, shorter tube dwell time and shorter hospital stay with the superior cosmetic result compared to sternotomy and thoracotomy scars as concluded by Amer et al. The study reported seven cases of VATS resection of mediastinal PA from Southampton General Hospital, UK between 2004–2009. The median hospital stay was two days and six out of seven patients had normalized serum calcium levels. There was minimal post surgical morbidity, however VATS was converted to open thoracotomy in one patient on account of bleeding from the azygos vein due to excessive traction [4]. Furthermore, several case series have reported VATS to be a feasible and safe approach for Mediastinal PA with an overall cure rate of 98–100% [14]. Tcherveniakov et al., reported the successful removal of parathyroid adenoma in four cases of ectopic MPA using VATS without any complications; all four cases had previous unsuccessful explorations through the cervical route [15].

In accordance with the apparent benefits of VATS compared to the conventional procedures, it should be considered as the procedure of choice for ectopic mediastinal PAs, wherever the surgical expertise are available. The unfrequent side effects of VATS reported include hemorrhage, subcutaneous emphysema, recurrent pneumothorax and pulmonary edema [16].

4. Conclusion

Symptomatic hypercalcemia and high level of PTH without PA in the neck should alert physicians to search for ectopic locations through combination of imaging techniques. The mediastinum should be cautiously investigated since it is the foremost location for ectopic PA after the neck. VATS is a safe and effective minimally invasive procedure for the resection of ectopic mediastinal PA. It should be considered as the first line approach for resection of these ectopic tumors.

Conflicts of interest

None.

Funding

None.

Ethical approval

Not Required.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent will be made available for review by the Editor-in-Chief of this journal on request.

Authors contribution

1. Saulat Hasnain Fatmi – Data Analysis, Interpretation, Manuscript Drafting.
2. Hina Inam – Study Concept, Study Design, Data Collection, Manuscript Writing.
3. Farida Karim – Study Concept, Data Analysis, Proof Reading.
4. Usama Khalid Choudry – Critical Review of Literature, Data Interpretation, Proof reading.

Guarantor

Dr. Saulat Hasnain Fatmi Associate professor, Department of Cardiothoracic surgery. Aga Khan University Hospital.

References

- [1] R.A. Agha, A.J. Fowler, A. Saetta, I. Barai, S. Rajmohan, D.P. Orgill, for the SCARE Group, The SCARE statement: consensus-based surgical case report guidelines, *Int. J. Surg.* 34 (2016) 180–186.
- [2] D. Naik, K.F. Jebasingh, G.B. Ramprasath, M.J. Paul, Video assisted thoroscopic surgery (VATS) for excision of an ectopic anterior mediastinal intra-thymic parathyroid adenoma, *J. Clin. Diagn. Res.* 10 (June (6)) (2016) PD22.
- [3] W. Zhou, M. Chen, A case report of mediastinal ectopic parathyroid adenoma presented as parathyroid crisis localized by SPECT/CT, *Medicine* 95 (October (41)) (2016).
- [4] K. Amer, A.Z. Khan, D. Rew, N. Lagattolla, N. Singh, Video assisted thoroscopic excision of mediastinal ectopic parathyroid adenomas: a UK regional experience, *Ann. Cardiothorac. Surg.* 4 (October (4)) (2015) 527–534.
- [5] H.T. Kojidi, N. Vagharimehr, S. Mohseni, M. Pajouhi, M.R. Mohajeri-Tehrani, Unusual ectopic parathyroid adenoma: a case report, *Acta Med. Iran.* 54 (September (8)) (2016) 547–550.
- [6] G. Nousseios, P. Anagnostis, K. Natsis, Ectopic parathyroid glands and their anatomical, clinical and surgical implications, *Exp. Clin. Endocrinol. Diabetes* 120 (2012) 604–610.
- [7] V. Mendoza, C. Ramirez, A.E. Espinoza, et al., Characteristics of ectopic parathyroid glands in 145 cases of primary hyperparathyroidism, *Endocr. Pract.* 16 (2010) 977–981.
- [8] Y. Krausz, L. Bettman, L. Guralnik, et al., Technetium-99m-MIBI SPECT/CT in primary hyperparathyroidism, *World J. Surg.* 30 (2006) 76–83.
- [9] K.K. Wong, L.M. Fig, M.D. Gross, B.A. Dwamena, Parathyroid adenoma localization with 99mTc-sestamibi SPECT/CT: a meta-analysis, *Nucl. Med. Commun.* 36 (April (4)) (2015) 363–375.
- [10] C.F. Russell, A.J. Edis, D.A. Scholz, et al., Mediastinal parathyroid tumors: experience with 38 tumors requiring mediastinotomy for removal, *Ann. Surg.* 193 (1981) 805–809.
- [11] N.J. Downey, J.A. McGuigan, S.J. Dolan, et al., Median sternotomy for parathyroid adenoma, *Ir. J. Med. Sci.* 168 (1999) 13–16.
- [12] J. Hu, K.Y. Ngiam, R. Parameswaran, Mediastinal parathyroid adenomas and their surgical implications, *Ann. R. Coll. Surg. Engl.* 97 (4) (2015) 259–261.
- [13] S.M. Said, S.D. Cassivi, M.S. Allen, et al., Minimally invasive resection for mediastinal ectopic parathyroid glands, *Ann. Thorac. Surg.* 96 (2013) 1229–1233.

- [14] P.F. Alesina, D. Moka, J. Mahlstedt, M.K. Walz, Thoracoscopic removal of mediastinal hyperfunctioning parathyroid glands: personal experience and review of the literature, *World J. Surg.* 32 (2) (2008) 224–231.
- [15] P. Tcherveniakov, A. Menon, R. Milton, K. Papagiannopoulos, M. Lansdown, J.A. Thorpe, Video-assisted mediastinoscopy (VAM) for surgical resection of ectopic parathyroid adenoma, *J. Cardiothorac. Surg.* 2 (2007) 41.
- [16] A. Imperatori, N. Rotolo, M. Gatti, E. Nardecchia, L. De Monte, V. Conti, et al., Peri-operative complications of video-assisted thoracoscopic surgery (VATS), *Int. J. Surg.* 6 (Suppl. 1) (2008) S78–S81.
- [17] G. Akerstrom, J. Malmaeus, R. Bergstrom, Surgical anatomy of human parathyroid glands, *Surgery* 95 (1984) 14–21.
- [18] K. Akram, J.A. Parker, K. Donohoe, G. Kolodny, Role of single photon emission computed tomography/computed tomography in localization of ectopic parathyroid adenoma: a pictorial case series and review of the current literature, *Clin. Nucl. Med.* 34 (August (8)) (2009) 500–502.
- [19] C.R. McHenry, What's new in general surgery: endocrine surgery, *J. Am. Coll. Surg.* 195 (2002) 364–371.
- [20] I. Zerizer, A. Parsai, Z. Win, et al., Anatomical and functional localization of ectopic parathyroid adenomas: 6-year institutional experience, *Nucl. Med. Commun.* 32 (2011) 496–502.
- [21] D.M. Elaraj, R.S. Sippel, S. Lindsay, et al., Are additional localization studies and referral indicated for patients with primary hyperparathyroidism who have negative sestamibi scan results? *Arch. Surg.* 145 (2010) 578–581.
- [22] A. Saif, Primary hyperparathyroidism presenting with acute pancreatitis and asymptomatic bone involvement, *Clin. Cases Min. Bone Metab.* 12 (May (2)) (2015) 199.

Open Access

This article is published Open Access at [sciencedirect.com](https://www.sciencedirect.com). It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.