

Differential patterns of ophiostomatoid fungal communities associated with three sympatric *Tomicus* species infesting pines in south-western China, with a description of four new species

Hui Min Wang¹, Zheng Wang¹, Fu Liu¹, Cheng Xu Wu¹, Su Fang Zhang¹,
Xiang Bo Kong¹, Cony Decock², Quan Lu¹, Zhen Zhang¹

1 Key Laboratory of Forest Protection, National Forestry and Grassland Administration; Research Institute of Forest Ecology, Environment and Protection, Chinese Academy of Forestry, Beijing 100091, China

2 Mycothèque de l'Université Catholique de Louvain (BCCM/MUCL), Earth and Life Institute, Microbiology, B-1348 Louvain-la-Neuve, Belgium

Corresponding author: Zhen Zhang (zhangzhen@caf.ac.cn); Quan Lu (luquan@caf.ac.cn)

Academic editor: Kevin Hyde | Received 27 December 2018 | Accepted 9 March 2019 | Published 9 April 2019

Citation: Wang HM, Wang Z, Liu F, Wu CX, Zhang SF, Kong XB, Decock C, Lu Q, Zhang Z (2019) Differential patterns of ophiostomatoid fungal communities associated with three sympatric *Tomicus* species infesting pines in south-western China, with a description of four new species MycoKeys 50: 93–133. <https://doi.org/10.3897/mycokeys.50.32653>

Abstract

Bark beetles and their associated fungi, which cause forest decline and sometimes high mortality in large areas around the world, are of increasing concern in terms of forest health. Three *Tomicus* spp. (*T. brevipilosus*, *T. minor* and *T. yunnanensis*) infect branches and trunks of *Pinus yunnanensis* and *P. kesiya* in Yunnan Province, in south-western China. *Tomicus* spp. are well known as vectors of ophiostomatoid fungi and their co-occurrence could result in serious ecological and economic impact on local forest ecosystems. Nonetheless, knowledge about their diversity, ecology, including pathogenicity and potential economic importance is still quite rudimentary. Therefore, an extensive survey of ophiostomatoid fungi associated with these *Tomicus* species infesting *P. yunnanensis* and *P. kesiya* was carried out in Yunnan. Seven hundred and seventy-two strains of ophiostomatoid fungi were isolated from the adult beetles and their galleries. The strains were identified based on comparisons of multiple DNA sequences, including the nuclear ribosomal large subunit (LSU) region, the internal transcribed spacer regions 1 and 2, together with the intervening 5.8S gene (ITS) and the partial genes of β -tubulin (*TUB2*), elongation

factor 1 α (*TEF1-a*) and calmodulin (*CAL*). Phylogenetic analyses were performed using maximum parsimony (MP) as well as maximum likelihood (ML). Combinations of culture features, morphological characters and temperature-dependent growth rates were also employed for species identification. Eleven species belonging to five genera were identified. These included six known species, *Esteya vermicola*, *Leptographium yunnanense*, *Ophiostoma brevipilosi*, *O. canum*, *O. minus* and *O. tingens* and four novel taxa, described as *Graphilbum anningense*, *O. aggregatum*, *Sporothrix pseudoabietina* and *S. macroconidia*. A residual strain was left unidentified as *Ophiostoma* sp. 1. The overall ophiostomatoid community was by far dominated by three species, representing 87.3% of the total isolates; in decreasing order, these were *O. canum*, *O. brevipilosi* and *O. minus*. Furthermore, the ophiostomatoid community of each beetle, although harbouring a diversity of ophiostomatoid species, was differentially dominated by a single fungal species; *Ophiostoma canum* was preferentially associated with and dominated the ophiostomatoid community of *T. minor*, whereas *O. brevipilosi* and *O. minus* were exclusively associated with and dominated the ophiostomatoid communities of *T. brevipilosus* and *T. yunnanensis*, respectively. Eight additional species, representing the remaining 12.7% of the total isolates, were marginal or sporadic. These results suggested that sympatric *Tomicus* populations are dominated by distinct species showing some level of specificity or even exclusivity.

Keywords

Esteya vermicola, *Graphilbum*, *Leptographium*, *Ophiostoma*, species-specific association, *Sporothrix*, taxonomy

Introduction

Associations between insects and microorganisms are increasingly recognised as one of the major issues in forest ecology and forest health around the world (Wingfield et al. 2016). Many bark beetles are well known as tree pests causing various levels of tree mortality and forest decline in large areas of the world, mostly in temperate areas (Jankowiak 2006, Wingfield et al. 2017). These bark beetles are well known vectors of variably pathogenic fungi, forming symbiosis-like relationships (Six 2003, Lu et al. 2009).

The pine shoot beetles, *Tomicus* Latreille (syn. *Blastophagus* Eichhoff, *Myelophilus* Eichhoff, Scolytidae, Coleoptera), are destructive insects with a range spanning the Eurasian pine forests, seriously affecting tree growth and causing a great threat to the forest ecosystems (Kirkendall et al. 2008, Lieutier et al. 2015). Currently, eight species are recorded worldwide, i.e. *T. armandii* Li and Zhang (Li et al. 2010), *T. brevipilosus* Eggers, *T. destruens* Wollaston, *T. minor* Hartig, *T. pilifer* Spessivtsev, *T. piniperda* L., *T. puellus* Reitter, and *T. yunnanensis* Kirkendall and Faccoli (Kirkendall et al. 2008). They all occur in China except *T. destruens* and five of them, viz. *T. armandii*, *T. brevipilosus*, *T. minor*, *T. pilifer* and *T. yunnanensis*, are sympatric in forests of the Yunnan Province (Li et al. 1997, 2010, Kirkendall et al. 2008; Ye 2011). *Tomicus brevipilosus*, *T. minor* and *T. yunnanensis* have overlapping geographical distribution, host range and infection periods. They aggregately infect branches and trunks of two indigenous pines, *Pinus yunnanensis* and *P. kesiya* (Li et al. 1997, 2006, Chen et al. 2009, 2010, Lu et al. 2012, 2014), causing locally extensive tree decline or mortality (Ye and Dang 1986, Ye 1991, 2011). Since the 1980s, damage caused by these bark beetles has resulted in losses of more than 93,000 m³ of pinewood (Ji et al. 2007).

Generally, two or three pine shoot beetles co-occur underneath the bark or in shoots of a single host tree, either simultaneously but with spatially isolated galleries or successively, during differential infesting peaks. Spatial and chorological differentiation would reduce competition between beetles, but their co-occurrence also could enhance cooperation (Lu et al. 2012, Chen et al. 2015). *Tomicus yunnanensis* is considered to be the most aggressive species in Yunnan, causing primary infestations of healthy *P. yunnanensis* trees and eventually tree death (Ye and Lieutier 1997, Kirkendall et al. 2008, Chen et al. 2010, 2015, Lu et al. 2014). Although *T. brevipilosus* is able to infect healthy trees, it preferably colonises trunks already infested by *T. yunnanensis* or both *T. yunnanensis* and *T. minor* (Chen et al. 2010, 2015). *Tomicus minor* is often regarded as a secondary, opportunist species infesting trees already weakened by *T. yunnanensis* or/and *T. brevipilosus* (Ye and Ding 1999, Lieutier et al. 2003, Chen et al. 2009).

Pine shoot beetles such as *T. piniperda*, *T. minor* and *T. destruens* are commonly associated with ophiostomatoid fungi (Masuya et al. 1999, Kim et al. 2005, Jankowiak 2006, 2008). Fifteen ophiostomatoid fungi were reported associated with *T. piniperda* in Europe (Mathiesen 1950, Lieutier et al. 1989, Gibbs and Inman 1991, Solheim and Långström 1991, Jankowiak 2006, Jankowiak and Bilański 2007) and 11 were documented in eastern Asia (Japan and Korea) (Masuya et al. 1999, Kim et al. 2005). *Ophiostoma minus* was shown to be the dominant species associated with *T. piniperda* in Europe and Japan (Mathiesen 1950, Lieutier et al. 1989, Gibbs and Inman 1991, Masuya et al. 1999, Jankowiak 2006). *Leptographium wingfieldii* was shown to be the strongest pathogenic one (Gibbs and Inman 1991) in Europe. *Tomicus minor* also infests various pines in Europe and Asia. Fifteen (Mathiesen-Käärik 1953, Masuya et al. 1999, Jankowiak 2008) and 11 (Masuya et al. 1999) ophiostomatoid species have been reported to be associated with this beetle species in Europe and Japan, respectively. *Ophiostoma canum* was recorded as a frequent/dominant species in association with *T. minor*, both in Europe and Japan (Mathiesen 1950, 1951, Rennerfelt 1950, Francke-Grosmann 1952, Masuya et al. 1999) but seems to represent a weak pathogen to *P. sylvestris* (Solheim et al. 2001). Additionally, six ophiostomatoid fungi were documented associated with *T. destruens* in Europe (Lieutier 2002, Sabbatini Peverieri et al. 2006, Ben Jamaa et al. 2007).

Despite the fact that *Tomicus* spp. have caused serious losses to forest ecosystems in south-western China, there are no systematic studies of their ophiostomatoid associates but only a few sporadic reports. So far, nine ophiostomatoid species have been reported as being associated with *Tomicus* spp. in Yunnan. Six species (*Leptographium yunnanense*, *Ophiostoma ips*, *O. minus*, *O. quercus*, *S. abietina* and *S. nebularis*) were recorded to be associated with *T. yunnanensis* (Ye et al. 2000, Zhou et al. 2000, 2013, Chang et al. 2017). Two species (*Graphilbum fragrans* and *O. tingens*) were recorded as being associated with *T. minor* (Zhou et al. 2013, Pan et al. 2017), whereas only a single species (*O. brevipilosus*) was recorded as being associated with *T. brevipilosus* (Chang et al. 2017). Amongst them, *L. yunnanense* was the first species newly described from the area (Zhou et al. 2000) and is likely the most virulent one (Liao and Ye 2004, Gao et al. 2017). Until now, the relative abundance with which these fungi occur, their host (pine and beetle) relationships, and their pathogenicity remain unknown.

The symbiosis between bark beetles and ophiostomatoid fungi enhances their pathogenicity. The fitness of bark beetle populations may depend in part on the degree of the fungal partners' pathogenicity and the resulting weakening of the tree (Christiansen et al. 1987, Kirisits 2004, Linnakoski et al. 2012), although this has been questioned by some (Six and Wingfield 2011). Therefore, the question remains whether there is any link between the differential aggression of the pine shoot beetles and the differential virulence of their fungal associates, especially in circumstances where various beetle species co-exist.

The aim of this study was to describe the diversity of ophiostomatoid fungal communities associated with three pine shoot beetles and their galleries infesting *P. yunnanensis* and *P. kesiya* in forest ecosystems of Yunnan Province. We also analysed the degree of beetle/ophiostomatoid fungi specificity. Such studies will enable us to understand the aggressive nature of the beetles and the pathogenicity of the associated fungi and the interactions, ultimately helping to address the current situation of ceaseless outbreaks and rapid expansion of the pests.

Materials and methods

Sample collection and fungus isolation

Samples of galleries in bark and shoots and adults of *Tomicus* spp. were collected from *P. yunnanensis* and *P. kesiya* at five sites in Yunnan Province (Fig. 1, Table 1) from December 2016 to March 2017. Beetles were placed individually in sterilised Eppendorf tubes and their galleries were placed in sterile envelopes and stored at 4°C until processed within one week.

Isolations from beetles and their galleries were carried out on 2% malt extract agar (MEA: 20 g Biolab malt extract, 20 g Biolab agar and 1 000 ml deionised water) with 0.05% NaClO added, in 9-cm Petri dishes as described by Seifert et al. (1993). Hyphal tips of emerging colonies were cut and transferred to MEA plates in order to obtain pure strains. The strains were grown routinely on 2% MEA at 25 °C. Representative

Table 1. Basic information on the sample collection plots in China.

Location	Host	Insect vector	longitude/latitude	altitude(m)	No. of examined samples
Xiangyun, Yunnan	<i>Pinus yunnanensis</i>	<i>Tomicus yunnanensis</i> , <i>T. minor</i>	25°21'25.8"N, 100°51'49"E	2255.4	447
Puer, Yunnan	<i>P. kesiya</i>	<i>T. brevipilosus</i> , <i>T. minor</i>	22°56'36.1"N, 101°14'36.7"E	1400.7	346
Qujing, Yunnan	<i>P. yunnanensis</i>	<i>T. yunnanensis</i> , <i>T. minor</i> , <i>T. brevipilosus</i>	25°28'51"N, 103°46'32"E	2068.2	102
Anning, Yunnan	<i>P. yunnanensis</i>	<i>T. yunnanensis</i> , <i>T. minor</i> , <i>T. brevipilosus</i>	24°53'32"N, 102°24'23"E	1939.9	138
Yuxi, Yunan	<i>P. yunnanensis</i>	<i>T. yunnanensis</i> , <i>T. minor</i>	24°18'23"N, 102°34'37"E	1908.1	85




Figure 1. **A** Map showing the 11 species of ophiostomatoid fungi detected from Yunnan Province, China **B, D** disease symptoms on *Pinus yunnanensis* and *P. kesiya* trees infested by *Tomicus* spp. (*T. yunnanensis*, *T. minor* and *T. brevipilosus*) and ophiostomatoid fungi **C, G, H** exposed branches of *Tomicus* spp. on *P. yunnanensis* and *P. kesiya* **E, F, I–K** galleries of *Tomicus* spp. on *P. yunnanensis* and *P. kesiya*.

cultures of each morphotype were deposited in the China Forestry Culture Collection Center (CFCC, part of the National Infrastructure of Microbial Resources) and the culture collection of the Chinese Academy of Forestry (CXY) (Table 2).

Morphology and growth studies

Morphological characterisation of both the sexual and asexual reproduction forms was performed on 2% MEA media incubated 3–6 weeks at 25 °C in the dark. Slide cultures were made to observe all microscopic characters (sexual/asexual structures) using a BX51 OLYMPUS microscope with differential interference contrast. Fifty measurements were made of each relevant structure and the ranges were calculated. Standard

Table 2. Representative strains of the ophiostomatoid fungi associated with three *Tomicus* spp. in Yunnan Province, China, and three *E. vermicola* strains used in this study.

Group	Taxon	Strain no.	Host	Location	Beetle	LSU	ITS/ITS2-LSU5	GenBank no.		
								BT	EF	CAL
A	<i>Esteya vermicola</i>	CFCC52625 (CXY1893)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325143	–	MH697597	MH605999	–
		ATCC74485	Japanese black pine	Taiwan, China	<i>Bursaphelenchus xylophilus</i>	–	–	–	GQ995674	–
		CNU120806	soil	Korea	saprophytic nematodes	EU627684	–	FJ490553	GQ995671	–
B	<i>Graphilbum anningense</i>	CBS 115803	oak	Czech Republic	<i>Scolytus intricatus</i>	–	–	FJ490552	GQ995672	–
		CFCC52631 (CXY1939)	<i>P. yunnanensis</i>	Anning	<i>T. yunnanensis</i>	MH325162	MH555903	MH683595	–	–
		CFCC52632 (CXY1940)	<i>P. yunnanensis</i>	Anning	<i>T. yunnanensis</i>	MH325164	MH555901	MH683596	–	–
		CFCC52633 (CXY1944)	<i>P. yunnanensis</i>	Anning	<i>T. minor</i>	MH325163	MH555902	MH683597	–	–
		CFCC52619 (CXY1897)	<i>P. keisiya</i>	Ninger	<i>T. brevipilosus</i>	MH325138	MH487721	MH603933	MH606000	–
C	<i>Leptographium yunnanense</i>	CFCC52620 (CXY1900)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325139	MH487724	MH603934	MH606001	–
		CFCC52621 (CXY1904)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325140	MH487726	MH603935	MH606003	–
		CFCC52622 (CXY1908)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325142	MH487725	MH603938	MH606002	–
		CFCC52623 (CXY1917)	<i>P. keisiya</i>	Puer	<i>T. brevipilosus</i>	MH325137	MH487723	MH603936	MH606004	–
		CFCC52624 (CXY1925)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325141	MH487722	MH603937	MH606005	–
D	<i>Ophiostoma brevipilosi</i>	CFCC52596 (CXY1828)	<i>Pinus keisiya</i>	Puer	<i>T. brevipilosus</i>	MH325134	MH555904	MH619527	–	–
		(CXY1806) CFCC52597	<i>P. keisiya</i>	Puer	<i>T. brevipilosus</i>	MH325135	MH555905	MH619528	–	–
		CFCC52598 (CXY1808)	<i>P. keisiya</i>	Puer	<i>T. brevipilosus</i>	MH325136	MH555906	MH619529	–	–

E	<i>O. canum</i>	CFCC52601 (CXY1858)	<i>P. yunnanensis</i>	Xiangyun	<i>T. minor</i>	MH325151	MH555889	MH619521	-
		CFCC52602 (CXY1848)	<i>P. yunnanensis</i>	Xiangyun	<i>T. minor</i>	MH325152	MH555890	MH619522	-
		CFCC52603 (CXY1857)	<i>P. yunnanensis</i>	Xiangyun	<i>T. minor</i>	MH325153	MH555891	MH619523	-
F	<i>O. aggregatum</i>	CFCC52615 (CXY1876)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325146	MH555894	MH603927	-
		CFCC52616 (CXY1875)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325145	MH555893	MH603929	-
		CFCC52617 (CXY1874)	<i>P. keisya</i>	Puer	<i>T. minor</i>	MH325147	MH555895	MH603928	-
G	<i>O. minus</i>	CFCC52606 (CXY1885)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325154	MH578163	MH619524	-
		CFCC52607 (CXY1877)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325155	MH578164	MH619525	-
		CFCC52608 (CXY1881)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325156	MH578165	MH619526	-
H	<i>O. tingens</i>	CFCC52611 (CXY1866)	<i>P. yunnanensis</i>	Xiangyun	<i>T. minor</i>	MH325148	MH578166	MH603931	-
		CFCC52612 (CXY1865)	<i>P. yunnanensis</i>	Xiangyun	<i>T. minor</i>	MH325149	MH578167	MH603932	-
		CFCC52613 (CXY1868)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325150	MH578168	MH603930	-
I	<i>Ophiostoma</i> sp. 1	CFCC52618 (CXY1936)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325144	MH555892	MH683600	-
J	<i>Sporothrix macroconidia</i>	CFCC52628 (CXY1894)	<i>P. yunnanensis</i>	Xiangyun	<i>T. yunnanensis</i>	MH325157	MH555898	MH697594	MH592598
		CFCC52629 (CXY1895)	<i>P. keisya</i>	Ninger	<i>T. brevipilosus</i>	MH325158	MH555899	MH697595	MH592599
		CFCC52630 (CXY1896)	<i>P. keisya</i>	Ninger	<i>T. brevipilosus</i>	MH325159	MH555900	MH697596	MH592600
K	<i>S. pseudobietina</i>	CFCC52626 (CXY1937)	<i>P. yunnanensis</i>	Qujing	<i>T. minor</i>	MH325160	MH555896	MH683598	MH592601
		CFCC52627 (CXY1938)	<i>P. yunnanensis</i>	Qujing	<i>T. minor</i>	MH325161	MH555897	MH683599	MH592602

Species names in bold are species newly described in this study.

CFCC: China Forestry Culture Collection Center, Beijing, China;

CXY (Culture Xingyao): Culture collection of the Research Institute of Forest Ecology, Environment and Protection, Chinese Academy of Forestry.

Sequences missing data are indicated by [-]

deviation (*SD*), minimum (min) and maximum (max) measurements are presented as (min–) (mean–*SD*) – (mean+*SD*) (–max).

The optimal growth temperature of the various strains was determined by placing a 5-mm (diam.) plug from an actively growing fungal colony upside down at the centre of an MEA plate. For each strain, three replicates were incubated at temperatures ranging from 5 to 35 °C at five-degree intervals, for 8d. The diameter of each colony was measured daily. Culture characters were recorded on MEA incubated at 25 °C for 8 d and 20 d. Colour descriptions were made by reference to Rayner (1970).

DNA extraction and sequencing

DNA was extracted from actively growing mycelium scraped from seven-day-old cultures using sterile scalpels and transferred to 2 ml Eppendorf tubes. DNA extraction and purification were performed using the Invisorb Spin Plant Mini Kit (Invitex, Berlin, Germany), following the manufacturer's protocols.

DNA sequences were determined for six gene regions: the nuclear ribosomal large subunit region (LSU), the internal transcribed spacer regions 1 and 2, including the intervening 5.8S gene (ITS), as well as segments of the β -tubulin (*TUB2*), elongation factor 1 α (*TEF1-a*) and calmodulin (*CAL*) genes. DNA fragments were amplified using the primer pairs LROR/LR5 (Vilgalys and Hester 1990), ITS1/ITS4 (White et al. 1990), ITS3/LR3 (Vilgalys and Hester 1990, White et al. 1990), Bt2a/Bt2b (Glass and Donaldson 1995), EF1/EF2 (Jacobs et al. 2004) and CL1/CL2a (Zhang et al. 2015), respectively. PCR reactions were conducted in 25 μ l volumes (2.5 mM MgCl₂, 1 \times PCR buffer, 0.2 mM dNTP, 0.2 mM of each primer and 2.5 U Taq-polymerase enzyme). PCR amplifications were carried out in a thermocycler (Applied Biosystems, Foster City, California, USA). The reaction conditions for these six gene regions were similar to those described in the references used for primer design. PCR products were cleaned with an MSB Spin PCR apace Kit (250), following the manufacturer's instructions.

Phylogenetic analyses

BLAST searches for the obtained sequences were performed in NCBI GenBank and published sequences of closely related species were downloaded. Alignments of the genes were made using MAFFT 7.0 (Kato and Standley 2013) and the E-INS-i strategy and edited manually in MEGA 5.2 (Tamura et al. 2011). Phylogenetic analyses were performed using maximum parsimony (MP) as well as maximum likelihood (ML).

ML analyses were implemented using RAxML v. 7.0.3 (Stamatakis 2006), under the GTR-GAMMA model. Support for the nodes was estimated from 1 000 bootstrap replicates. The results were subsequently exported to Figtree v.1.4.2 to visualise the trees.

MP analyses were implemented in PAUP* 4.0b10 (Swofford 2003). The most parsimonious trees were identified by a heuristic search of 1 000 random addition sequence replicates, using the tree-bisection-recognition (TBR) algorithm for branch

swapping. Branch support was assessed by 1 000 bootstrap replicates. Tree length (TL), consistency index (CI), retention index (RI), rescaled consistency index (RC) and homoplasy index (HI) were used to evaluate the trees.

Results

Fungal isolation and sequence comparisons

Three *Tomicus* species occurred on *P. yunnanensis* and *P. kesiya* in the areas studied, either independently or concomitantly in individuals of the host trees (Fig. 1). In total, 772 strains of ophiostomatoid fungi (*Hyalorhinochlaediella*-like, *Ophiostoma*, *Pesotum*-like, *Leptographium*-like and *Sporothrix*-like) were isolated from 223 adult beetles (20% of the strains) and 890 galleries (80% of the strains). Galleries or adults of *T. yunnanensis* yielded 297 strains whereas 247 strains were retrieved from galleries or adults of *T. minor* and 228 strains from galleries or adults of *T. brevipilosus* (Table 3).


The LSU sequence was used to search for preliminary affinities using the BLASTn search option in GenBank. As a result, these strains were found to be distributed over 5 genera and 11 tentative species/groups (A–K) (Table 2).

Phylogenetic analyses

The degrees of polymorphism of LSU, ITS, *TUB2*, *TEF1-a* and *CAL* make them variably suitable for genus or species discrimination amongst ophiostomatoid fungi. The LSU sequence is a suitable marker to infer the generic affinities (de Beer and Wingfield 2013,

Table 3. Strain numbers of various ophiostomatoid fungi obtained from three *Tomicus* spp. and their galleries collected in Yunnan Province.

Group	Fungi species	<i>Tomicus yunnanensis</i>	<i>T. minor</i>	<i>T. brevipilosus</i>	Total no. strains/samples
A	<i>Ophiostoma brevipilosi</i>	0	0	224	224
B	<i>O. canum</i>	52	201	0	253
C	<i>O. minus</i>	197	0	0	197
D	<i>O. tingens</i>	4	26	0	30
E	<i>O. aggregatum</i>	3	2	0	5
F	<i>Ophiostoma</i> sp. 1	1	0	0	1
G	<i>Leptographium yunnanense</i>	30	0	2	32
H	<i>Esteya vermicola</i>	1	0	0	1
I	<i>Sporothrix pseudoabietina</i>	4	15	0	19
J	<i>S. macroconidia</i>	1	0	2	3
K	<i>Graphilbum anningense</i>	4	3	0	7
	Total no. strains	297	247	228	772
	Total no. samples	455	324	339	1118



de Beer et al. 2013a, 2016); it allowed confirming the preliminary placement of our strains based on morphological characters (Fig. 2). The ITS region would be useful to place strains within the *Ophiostoma s. l.* complex, but the degree of polymorphism does not allow distinguishing species. Usually, *TUB2*, *TEF1-a* and *CAL* regions are better markers to identify and, where appropriate, to show the genetic diversity within ophiostomatoid fungi (Zipfel et al. 2006, de Beer and Wingfield 2013, de Beer et al. 2016).

On the basis of the LSU blast searches, one to six strains of each tentative species (A–K) were selected for sequencing of five additional DNA markers (ITS, ITS2-LSU, *TUB2*, *TEF1-a* and *CAL*) to infer more accurate identification and phylogenetic affinities. Six sequence datasets (LSU, ITS, ITS2-LSU, *TUB2*, *TEF1-a* and *CAL*) were generated for a total of 31 representative strains (Table 2) and the sequences were deposited in GenBank. Resulting alignments were deposited in TreeBASE (submission no: 24032). The topologies generated by the ML and MP analyses were highly concordant and the ML phylograms are presented for all the individual genes, incorporating nodal supports of both the ML and MP analyses.

The LSU dataset consisted of 109 sequences, 11 sequences obtained in this study and 98 downloaded from GenBank. The phylogenetic analyses confirmed the morphology-based placement of our strains into *Esteya*, *Graphilbum*, *Leptographium*, *Ophiostoma* and *Sporothrix* (Fig. 2).

Group A consisted of a single strain. LSU-based phylogenetic analysis showed this strain to be close to *E. vermicola* (Fig. 2). *TUB2* and *TEF1-a* data analysis confirmed the strain's close affinities to *E. vermicola* (Fig. 3a, b), that could justify conspecificity.

Group B strains nested within the *Graphilbum* lineage in the LSU-based phylogenetic analysis (Fig. 2). Phylogenetic analysis based on LSU, ITS and *TUB2* concordantly showed that the group B strains formed a single, well-supported clade related to but distinct from *Gra. rectangulosporium* and *Gra. microcarpum* (Fig. 4a, b); this would warrant its recognition as a distinct, undescribed species.

Group C strains were shown to belong to the *Leptographium* lineage in the LSU-based phylogenetic analysis (Fig. 2). The ITS2-LSU dataset consisted of six of our own sequences and 49 reference sequences downloaded from GenBank. Within the *Leptographium* lineage, group C strains nested in the *L. lundbergii*-complex; they were related to *L. yunnanense*, *L. lundbergii* and *L. conjunctum* (Fig. 5a). *TUB2*- and *TEF1-a* based analysis confirmed their close affinities with *L. yunnanense*, although forming a slightly divergent clade (Fig. 5b, c). *TUB2* and *TEF1-a* sequences of group C strains showed some polymorphisms, which could be considered as falling within the natural diversity of *L. yunnanense*.

Figure 2. Phylograms obtained from ML analysis of LSU sequences, showing fungal associates with pines infected by *Tomicus yunnanensis*, *T. minor* and *T. brevipilosus* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with 'T'; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 743 positions, including gaps.




Figure 3. Phylograms obtained from ML analysis of β -tubulin **A** and elongation factor **B** sequences of *Esteya*, showing fungal associates with pines infected by *Tomicus yunnanensis* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with ‘T’; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 320 (**A**), 856 (**B**) positions, including gaps.




Figure 4. Phylograms obtained from ML analysis of ITS sequences **A** and β -tubulin sequences **B** of *Graphilbum* showing fungal associates with pines infected by *Tomicus yunnanensis* and *T. minor* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with ‘T’; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 515 (**A**), 481 (**B**) positions, including gaps.

The six strains from groups D to I nested within the *Ophiostoma* lineage based on the LSU phylogenetic tree (Fig. 2). The ITS dataset comprised species from all lineages discovered in this study. Analysis of this dataset yielded the phylograms shown in Fig. 6. Sixteen ITS sequences generated in this study were compared with 61 sequences




Figure 5. Phylograms obtained from ML analysis of ITS2-28S **A** β -tubulin **B** and elongation factor **C** sequences of *Leptographium*, showing fungal associates with pines infected by *Tomicus yunnanensis* and *T. brevipilosus* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with ‘T’; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 641 (**A**), 358 (**B**), 639 (**C**) positions, including gaps.

retrieved from GenBank, representing the major groups of *Ophiostoma* (de Beer and Wingfield 2013, Linnakoski et al. 2016).

The ITS- and *TUB2*-based phylogenetic inferences (Figs 6, 7a, b) showed that the strains of groups D and E nested within the *O. clavatum*- and *O. piceae*-complex (de Beer and Wingfield 2013, Yin et al. 2016, Linnakoski et al. 2016), in which they were positioned in the near vicinity of the *O. brevipilosus* and *O. canum* clades, respectively. From these results, and considering their morphological features, we concluded that the strains of groups D and E are conspecific with *O. brevipilosus* and *O. canum*, respectively.




Figure 6. Phylograms obtained from ML analysis of ITS sequences of *Ophiostoma*, showing fungal associates with pines infected by *Tomicus yunnanensis*, *T. minor* and *T. brevispilosus* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with 'T'; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 633 positions, including gaps.




Figure 7. Phylograms obtained from ML analysis of β -tubulin sequences of *Ophiostoma* **A, B, D, E** and ITS sequences of *O. minus*-complex **C** showing fungal associates with pines infected by *Tomicus yunnanensis*, *T. minor* and *T. brevopilosus* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with 'T'; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 455 (**A**), 430 (**B**), 541 (**C**), 378 (**D**), 423 (**E**) positions, including gaps.




Figure 8. Phylograms obtained from ML analysis of ITS sequences of *Sporothrix*, showing fungal associates with pines infected by *Tomicus yunnanensis*, *T. minor* and *T. brevipilosus* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with ‘T’; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 546 positions, including gaps.




Figure 9. Phylograms obtained from ML analysis of β -tubulin **A, C** and calmodulin **B, D** sequences of *Sporothrix*, showing fungal associates with pines infected by *Tomicus yunnanensis*, *T. minor* and *T. brevipilosus* in Yunnan Province, China. Novel sequences obtained in this study are printed in bold type. Bootstrap values $\geq 70\%$ for ML and MP are indicated above branches. Bootstrap values $< 70\%$ are indicated by the symbol *. Strains representing ex-type sequences are marked with ‘T’; ML, maximum likelihood; MP, maximum parsimony and the final alignment of 284(**A**), 622(**B**), 260(**C**), 675(**D**) positions, including gaps.

In the ITS-based phylogenetic analysis, strains of groups G and I were grouped with the *O. minus* complex (Fig. 6). ITS- and *TUB2*-based phylogenetic analyses consistently showed that group G strains formed a well-supported subclade between the North American and European subclades within the *O. minus* lineage (Fig. 7c, d). The strains of group G are therefore identified as *O. minus*. The ITS- and *TUB2*-based phylogenetic analyses consistently showed that the single strain of group I formed a branch that is related to, but distinct from the *O. minus*, *O. kryptum* and *O. olgensis* clades (Figs 6, 7d). Hence, this strain is interpreted as belonging to a distinct, undescribed *Ophiostoma*.

The remaining two groups (F and H) were not placed in any defined complex. Phylogenetic analyses, based on ITS and *TUB2* sequences, consistently showed that the group H strains clustered in the near vicinity of the *O. tingens* clade whereas group F strains formed a clade related to, but distinct from the *O. macrosporum* and *O. tingens* clades (Figs 6, 7e). Thus, the strains in group H should be identified as *O. tingens* whereas the strains of group F represent an undescribed *Ophiostoma*.

Strains of groups J and K nested within the *Sporothrix* lineage in LSU-based phylogenetic analysis (Fig. 2). The phylograms resulting from the analyses of individuals are shown in Fig. 8 (ITS), Fig. 9a, c (*TUB2*) and Fig. 9b, d (*CAL*).

The ITS-based analyses showed that group K strains belonged to the *S. gossypina*-complex whereas the group J strains were not placed in any species complex as defined by de Beer et al. (2016) (Fig. 8). Both groups formed independent, well-supported clades in ITS-, *TUB2*- and *CAL*-based phylogenetic analyses (Figs 8, 9). It could be deduced from results of multiple phylogenies that both groups represent novel species.

Morphology and taxonomy

From a morphological perspective, strains of groups D, E and G appeared, overall, concordant with the descriptions or our own observations of reference strains, namely of *O. brevipilosi*, *O. canum* and *O. minus*, respectively. However, although strains of groups A, C, and H are phylogenetically close to *E. vermicola*, *L. yunnanense* and *O. tingens*, respectively, justifying, for the time being, conspecificity, their phenotype deviated slightly from published descriptions and/or our own observation of type material. The description of these species is extended. Strains of groups B, F, J and K revealed unique combinations of phenotypes, allowing morphological distinction from their closest phylogenetic relatives; consequently, they are described below as new species. The strain of the stand-alone group I also may represent an undescribed species; however, we refrain from describing it for the time being, waiting for more material to become available.

Taxonomy

***Esteya vermicola* J.Y. Liou, J.Y. Shih & Tzean, Mycol. Res. 103(2): 243. 1999.**

MycoBank MB450702

Fig. 10

Description. Sexual form: unknown.

Asexual form: *Hyalorhinocladiella*-like. *Conidiophores* mononematous, micronematous; *conidiophorous cells* solitary, integrated, flask-shaped, with an inflated base (3.6–) 4.6–6.1 (–7.1) μm in diam., the fertile hyphoid part (9.1–) 12.2–19.0 (–22.5) \times (1.4–) 1.9–3.1 (–4.7) μm , often crooked due to successive conidial development; *conidia*




Figure 10. **A–H** Morphological characters of *Esteya vermicola* **A, B** upper and reverse of cultures on 2% MEA 20 d after inoculation **C–H** conidiogenous cells with lunate conidia **I–M** the cuticle of a nematode attached by many lunate conidia. Scale bars: 20 μm (**I, K, L**); 10 μm (**C–H, J, M**).

1-celled, asymmetrically ellipsoidal in face view, concave, lunate in side view, with a layer of adhesive mucus on the concave surface, ending slightly apiculate, hyaline, smooth, $(8.0\text{--}10\text{--}12\text{--}13.1) \times (3.3\text{--}3.4\text{--}4.5\text{--}5.1) \mu\text{m}$, containing an ovoid endospore-like structure.

Culture characteristics. Colonies on 2% MEA in the dark reaching 31 mm in diam. in 8 days at 25 °C, growth rate up to 5 mm/day at the fastest; colony margin smooth. Mycelium compact, somewhat floccose in the margin, white at first, gradually discolouring to greyish-green, eventually dark green. Optimal growth temperature 25 °C, growth at 5 °C and 35 °C.

Known substrate and host. Galleries of *Tomicus yunnanensis* in *Pinus yunnanensis*.

Known insect vector. *Tomicus yunnanensis*.

Known distribution. Yunnan Province, China.

Specimen examined. CHINA, Yunnan, *Tomicus yunnanensis* galleries in *Pinus yunnanensis*, Dec. 2016, HM Wang, CFCC 52625 = CXY 1893.

Note. *Esteya vermicola* is known only from an asexual, *Hyalorhinocladiella*-like state producing lunate and bacilliform conidia (Liou et al. 1999, Kubátová et al. 2000, Wang et al. 2009, 2014) that we also observed in various strains of *E. vermicola* with a different origin (Taiwan, Korea, Czech Republic). Our strain was identified as *E. vermicola* based on phylogenetic inferences and morphological characters. However, our strain differed from previous descriptions (Liou et al. 1999) in having only lunate conidia *in vitro*. The size of the lunate conidia of our strains (mostly 10 - 12 × 3.4 - 4.5 µm) was similar to that reported for *E. vermicola*, *viz.* 9.9–11.9 × 3.4–4.5 µm vs 8.2–11.1 × 3.5–3.7 µm (Taiwan, Liou et al. 1999), 9.3–12.4 × 3.0–3.2 µm (Czech Republic, Kubátová et al. 2000), 7.7–12.1 × 3.0–3.8 µm (Korea, Wang et al. 2009) or 8.7–11.9 × 3.0–3.6 µm (Brazil, Wang et al. 2014).

This is the first report of *E. vermicola* from continental China. The species was originally isolated from Japanese black pine infected by the pinewood nematode *Bursaphelenchus xylophilus*, in Taiwan (Liou et al. 1999). Since then, its distribution range has been extended to Japan and Korea, Europe (Czech Republic, Italy) and both North (USA) and South America (Brazil) (Liou et al. 1999, Kubátová et al. 2000, Wang et al. 2009, 2014, Li et al. 2018). This species is associated with various vectors, including the pinewood nematode, *Oxoplatypus quadridentatus* and the bark beetle *Scolytus intricatus*. It was isolated also from wooden packaging material infested by *Bursaphelenchus rainulfi*.

***Graphilbum anningense* H. Wang, Q. Lu & Z. Zhang, sp. n.**

Mycobank MB828884

Fig. 11

Etymology. ‘*anningense*’ (Latin), referring to the type locality.

Type. CHINA, Yunnan, *Tomicus yunnanensis* galleries in *Pinus yunnanensis*, Apr. 2017, HM Wang, holotype CXY 1939, culture ex-holotype CFCC 52631 = CXY 1939.

Description. Sexual form: unknown.

Asexual forms: *Pesotum*-like and *Hyalorhinocladiella*-like. *Pesotum*-like conidiophores abundant on 2% MEA, macronematous, synnematous, (150–) 210–293 (–336) µm long including conidiogenous apparatus, the base dark brown, slightly widened, (6.7–)




Figure 11. Morphological characters of *Graphilbum anningense* sp. n. **A, B** Upper and reverse of cultures on 2% MEA 8 d after inoculation **C, D, G** conidiogenous cells of Pesotum-like macronematal asexual state and conidia **E, F, H** conidiogenous cells of *Hyalorhinocladiella*-like asexual state and conidia. Scale bars: 50 μm (**C**); 20 μm (**D**); 10 μm (**E–H**).

7.9–18.8 (–29.0) μm wide anchored in the media by brown rhizoid-like hyphae, the apex slightly enlarging, fan-shaped; *conidiogenous cells* hyaline, thin-walled, aseptate, (15.3–) 21.0–35.5 (–42) \times (0.7–) 1.1–1.9 (–2.3) μm ; *conidia* 1-celled, clavate, ellipsoid to ovoid with truncate base and rounded apex, hyaline, smooth, (3.1–) 3.6–6.3 (–9.7) \times (1.4–) 1.6–2.2 (–2.5) μm .

Hyalorhinocladiella-like: *conidiogenous cells* macronematous or semi-macronematous, mononematous, hyaline, simple or loosely branched, thin-walled, aseptate, (4.5–) 10.8–29.0 (–47) \times (1.5–) 1.7–2.3 (–2.6) μm ; *conidia* hyaline, clavate to ellipsoid, with obtuse ends, 1-celled, aseptate, smooth, (3.7–) 4.5–6.4 (–9.0) \times (1.4–) 1.7–2.3 (–2.9) μm .

Culture characteristics. Colonies on 2% MEA in the dark reaching 90 mm in diam. in 6 days at 25 $^{\circ}\text{C}$, growth rate up to 19.5 mm/day at the fastest; colony margin smooth. Mycelium superficial to flocculose or floccose, hyaline; reverse hyaline to pale yellowish. Optimal growth temperature 30 $^{\circ}\text{C}$, slow growth at 40 $^{\circ}\text{C}$, no growth at 5 $^{\circ}\text{C}$.

Known substrate and hosts. Galleries of *Tomicus yunnanensis* and *T. minor* in *Pinus yunnanensis*.

Known insect vectors. *Tomicus yunnanensis*, *T. minor*.

Known distribution. Yunnan Province, China.

Additional specimens examined. CHINA, Yunnan, *Tomicus yunnanensis*, *T. minor* galleries in *Pinus yunnanensis*, Apr. 2017, HM Wang, CFCC 52632 = CXY 1940, CFCC 52633 = CXY 1944.

Note. *Graphilbum anningense* is characterised by a *Pesotum*-like and a *Hyalorhinocladiella*-like asexual state. It is phylogenetically closely related to *Gra. rectangulosporium*. However, *Gra. rectangulosporium* produced a sexual state *in vitro* (Ohtaka et al. 2006) which has not been observed in *Gra. anningense*. Other morphologically similar species include *Gra. fragrans*, *Gra. crescericum*, *Gra. kesiyae* and *Gra. puerense*. *Graphilbum kesiyae* and *Gra. puerense* also produce a *Pesotum*-like and a *Hyalorhinocladiella*-like asexual state. *Graphilbum anningense* and *Gra. kesiyae* differ by the size of their synnemata, whose length ranges do not overlap, *viz.* 210–293 µm and 112.5–173 µm long (Harrington et al. 2001), respectively. They also differ by their optimal growth temperature, respectively 30°C and 25°C. The synnemata of *Gra. puerense*, 206–357 µm long (Chang et al. 2017), are marginally longer than those of *Gra. anningense*. *Graphilbum fragrans* and *Gra. crescericum* produce only a *Leptographium*-like and/or a *Hyalorhinocladiella*-like asexual state *in vitro* (Harrington et al. 2001, Chang et al. 2017).

Graphilbum anningense was isolated from galleries of *T. yunnanensis* and *T. minor* infesting *P. yunnanensis*. Previously, *Gra. fragrans* had been reported from *T. yunnanensis* infesting *P. yunnanensis* and from *Pissodes* spp. infesting *Tsuga dumosa* and *P. armandii* in China (Paciura et al. 2010, Zhou et al. 2013). *Graphilbum kesiyae* and *Gra. puerense* were isolated from galleries of *Polygraphus aterrimus*, *Po. szemaoensis* and *Ips acuminatus* infesting *P. kesiya* (Chang et al. 2017). Although the geographic distribution of these four *Graphilbum* species overlaps, their hosts and vectors are nevertheless, as far as it is known, different (Chang et al. 2017).

***Leptographium yunnanense* X.D. Zhou, K. Jacobs, M.J. Wingf. & M. Morelet, Mycoscience 41(6): 576. 2000.**

MycoBank MB 466542

Fig. 12

Description. Sexual form: unknown.

Asexual form: *Leptographium*-like. *Conidiophores* occurring singly or in groups of up to three, arising from the superficial mycelium, erect, macronematous, mononematous, (93.5–) 159–412 (–544) µm long, without rhizoid-like structures; *stipes* simple, cylindrical, not constricted at septa, 1–6-septate, pale olivaceous at the base, (12–) 19.0–128 (–245) × (3.3–) 4.1–6.1 (–7.3) µm; *conidiogenous apparatus* (33.0–) 65.5–119.5 (–168.0) µm long (high), with 2 to 3 series of cylindrical branches; *primary branches* hyaline to pale olivaceous, smooth, cylindrical, 2–3 septate, (11.5–) 18.2–37.7 (–56.0) µm long and (3.0–) 3.7–5.9 (–7.7) µm wide; *secondary branches* hyaline, 0–2 septate, (10.3–) 14.5–30.0 (–50.1) µm long, (2.8–) 3.4–5.5 (–7.3) µm wide; *conidiogenous cells* discrete, 2–3 per branch, cylindrical, (10.2–) 13.2–29.6 (–57.4) × (2.2–) 2.9–3.9 (–4.4) µm; *conidia* 1-celled, oblong to obovoid with truncate bases, hyaline, (5.8–) 7.0–10.4 (–13.0) × (2.9–) 3.6–5.3 (–6.4) µm.




Figure 12. Morphological characters of *Leptographium yunnanense* **A, B** upper and reverse of cultures on 2% MEA 8 d after inoculation **D, I** conidiophore on 2% MEA **C, E–H** conidiogenous cells of *Leptographium*-like asexual state and conidia. Scale bars: 50 μm (**D, I**); 10 μm (**C, E–H**).

Culture characteristics. Colonies on 2% MEA medium fast growing in the dark, reaching 76 mm in diam. in 8 days at 25 °C, growth rate up to 20 mm/day at the fastest; colony margin smooth. Hyphae submerged in agar with aerial mycelium, greenish-olivaceous to olivaceous, smooth, straight; reverse hyphae umber-brown to dark olivaceous. Optimal growth temperature 25 °C, slow growth at 5 °C and 30 °C.

Known substrate and hosts. *Tomicus yunnanensis* and its galleries in *Pinus yunnanensis*, galleries of *T. brevipilosus* in *P. kesiya*.

Known insect vectors. *Tomicus brevipilosus*, *T. yunnanensis*.

Known distribution. Yunnan Province, China.

Specimens examined. CHINA, Yunnan, adults of *Tomicus yunnanensis* and their galleries in *Pinus yunnanensis*, *Tomicus brevipilosus* galleries in *P. kesiya*. Apr. 2017, HM Wang, CFCC 52619 = CXY 1897, CFCC 52620 = CXY 1900, CFCC 52621 = CXY 1904, CFCC 52622 = CXY 1908, CFCC 52623 = CXY 1917, CFCC 52624 = CXY 1925.

Note. The sole reproductive structure formed on MEA in *L. yunnanense* is a *Leptographium*-like state. Our strains were identified as *L. yunnanense*, based on phylogenetic evidence and secondarily, on morphological features. However, our strains slightly deviated from *L. yunnanense* in having longer conidiophores, mainly 159–412 μm vs mostly 74–227 (–233) μm (Zhou et al. 2000) or 80–240 μm (Yamaoka et al. 2008). Furthermore, our strains grew faster than reported for the species, 76 mm vs 44 mm in 8 days at 25 °C (Zhou et al. 2000).

Although our strains were slightly genetically and morphologically divergent, we are of the opinion that they enter into the current *L. yunnanense* species concept (e.g. *sensu* Zhou et al. 2000). Yamaoka et al. (2008) showed the genetic diversity of *L. yunnanense* in Yunnan to be higher than in other places, that which is confirmed by the present study.

Leptographium yunnanense was originally described from Yunnan Province with only an asexual state (Zhou et al. 2000). Subsequently, mating of strains from different origins (Thailand, China and Japan) yielded the sexual state, which is formed by neckless ascocarps and cucullate ascospores (Yamaoka et al. 2008).

Leptographium yunnanense was the third most abundant species associated with *T. yunnanensis* in our study. A few strains also were isolated from *T. brevipilosus* infesting *P. kesiya* and none from *T. minor*.

***Ophiostoma aggregatum* H. Wang, Q. Lu & Z. Zhang, sp. n.**

MycoBank MB828885

Fig. 13

Etymology. ‘*aggregatum*’ (Latin), reflects to the conidiophores aggregated in clusters.

Type. CHINA, Yunnan, from *Tomicus minor* galleries in *Pinus yunnanensis*, Dec. 2016, HM Wang, holotype CXY 1876, culture ex-holotype CFCC 52615 = CXY 1876.

Description. Sexual form: unknown.

Asexual form: *Leptographium*-like. *Conidiophores* macronematous, mononematous, gathered in groups up to 5, (28.5–) 34–45.5 (–52) μm long; *stipes* cylindrical, 1–2 septate, not constricted at septa, umber-brown to dark olivaceous, (6.3–) 7.3–14.5 (–18) μm long \times (2.2–) 3.1–4.6 (–5.8) μm wide. *Conidiogenous apparatus* (22–) 26.5–31 (–34) μm long, with 2–3 series of cylindrical branches; primary branches olivaceous, smooth, cylindrical all over, (5.9–) 7.2–13.5 (–20.5) \times (3–) 3.3–4.2 (–4.6) μm ; *conidiogenous cells* discrete, 2–3 per branch, aseptate, cylindrical, hyaline to pale umber, (5.8) 7.2–12.1 (–18.5) \times (2.1–) 2.8–4.0 (–4.7) μm ; *conidia* 1-celled, globose, elliptical with truncate bases, hyaline to pale umber, (4.0–) 4.8–5.9 (–6.3) \times (3.1–) 4.0–5.0 (–5.6) μm .

Culture characteristics. Colonies on 2% MEA fast growing in the dark, reaching 90 mm in diam. in 8 days at 25 °C, growth rate up to 13 mm/day at the fastest; colony margin smooth. Hyphae submerged and aerial, umber-brown to dark olivaceous, flocculose or floccose; reverse hyphae umber-brown to dark olivaceous. Optimal growth temperature 25 °C, able to grow at 5 °C and 30 °C. No growth at 35 °C.

Known substrate and hosts. Galleries of *Tomicus yunnanensis* and *T. minor* in *Pinus yunnanensis*.




Figure 13. Morphological characters of *Ophiostoma aggregatum* sp. n. **A, B** upper and reverse of cultures on 2% MEA 8 d after inoculation **C** conidiomata on 2% MEA (bar = 50 µm) **D–H** conidiogenous cells of *Leptographium*-like asexual state and conidia. Scale bars: 20 µm (**C**); 10 µm (**D–H**).

Known insect vectors. *Tomicus minor*, *T. yunnanensis*.

Known distribution. Yunnan Province, China.

Additional specimens examined. CHINA, Yunnan, from *Tomicus yunnanensis* and *T. minor* galleries in *Pinus yunnanensis*, Dec. 2016, Apr. 2017, HM Wang, CFCC 52616 = CXY 1875, CFCC 52617 = CXY 1874.

Note. *Ophiostoma aggregatum* produced a single asexual, *Leptographium*-like state *in vitro*. This species is phylogenetically closely related to *O. macrosorum*, *O. tingens*,

O. floccosum, *O. tapionis* and *O. piliferum* in LSU-, ITS- and *TUB2*-based phylogenetic inferences. *Ophiostoma aggregatum* and *O. tingens* are shown to be sympatric in Yunnan pine forest; both taxa were isolated from galleries and adults of *T. minor* and *T. yunnanensis* infesting *P. yunnanensis* (Table 2). *Ophiostoma tingens* was also reported from *T. minor* infesting *P. yunnanensis* in Yunnan (Pan et al. 2017).

Ophiostoma aggregatum and *O. tingens* differ in their asexual state. *Ophiostoma aggregatum* only produces a *Leptographium*-like state. Inversely, the asexual states of *O. tingens* are variable. Our strains produced a *Pesotum*-like and a *Sporothrix*-like state whereas previously, Francke-Grosmann (1952) and de Beer et al. (2013b) reported a *Hyalorhinocladiella*- to *Raffaelea*-like state in European strains. The origin of this variability and its importance for taxonomy is uncertain.

Ophiostoma macrosporum, *O. floccosum*, *O. tapionis* and *O. piliferum* also differ from *O. aggregatum* by their asexual state. *Ophiostoma macrosporum* and *O. floccosum* produce a *Pesotum*-like asexual state, *O. tapionis* a *Hyalorhinocladiella*-like state and *O. piliferum* produces a *Sporothrix*-like state (Francke-Grosmann 1952, Upadhyay 1981, Yamaoka et al. 2004, Linnakoski et al. 2008).

Ophiostoma macrosporum and *O. tingens* were both originally described in *Trichosporium* as *T. tingens* var. *macrosporum* and *T. tingens* (Lagerberg 1927, Francke-Grosmann 1952). Batra (1967) transferred these two species into *Ambrosiella*. It is only recently that the morphological characteristics were found to agree with those of *Ophiostoma* (de Beer et al. 2013b). *Ophiostoma macrosporum* has been reported from various *Pinus* spp. (including *P. sylvestris*) infected by *Ips acuminatus* in Europe (Francke-Grosmann 1952, Batra 1967).

***Ophiostoma tingens* (Lagerb. & Melin) Z.W. de Beer & M.J. Wingf., Svensk Skogs-vårdsförening Tidskr. 25:233. 1927.**

MycoBank: MB801091

Fig. 14

Description. Sexual form: unknown.

Asexual forms: *Pesotum*-like and *Sporothrix*-like. *Pesotum*-like: *conidiophores* macronematous, synnematosus; synnemata simple, anchored into the substrate by brown rhizoid-like hyphae, (333–) 344–584 (–684) μm long including *conidiogenous apparatus*, the base dark brown, slightly widened, (16.7–) 17–50.5 (–65.5) μm wide, the apex cream-coloured or pale brown, slightly widening; *conidia* hyaline, globose to elliptical, 1-celled, smooth, (2.7–) 3.6–7.2 (–8.0) \times (2.8–) 4.3–6.1 (–7.0) μm .

Sporothrix-like: *conidiophores* semi-macronematous, mononematous, hyaline, simple or loosely branched, smooth, bearing terminal denticulate *conidiogenous cells* (8.3–) 15.6–30.0 (–42.5) \times (1.1–) 1.7–3.1 (–4.7) μm ; *conidia* hyaline, globose to elliptical, obovoid with pointed bases and rounded apices, 1-celled, smooth, (2.6–) 4.0–6.8 (–8.7) \times (2.2–) 3.5–5.5 (–7.4) μm .




Figure 14. Morphological characters of *Ophiostoma tingens* **A,B** upper and reverse of cultures on 2% MEA 20 d after inoculation **C–G** conidiogenous cells of *Sporothrix*-like asexual state and conidia **H–J** conidiogenous cells of *Pesotum*-like macronematal asexual state and conidia. Scale bars: 10 µm (**C–H**); 50 µm (**I, J**).

Culture characteristics. Colonies on 2% MEA medium slow growing in the dark, reaching 39 mm in diam. in 8 days at 25 °C, growth rate up to 5 mm/day at the fastest; colony margin anomalous. Hyphae appressed to flocculose, black; reverse hyphae also black. Optimal growth temperature 25 °C, no growth at 5 °C and 30 °C.

Known substrate and hosts. Galleries of *Tomicus yunnanensis* and *T. minor* in *Pinus yunnanensis*.

Known insect vectors. *Tomicus yunnanensis*, *T. minor*.

Known distribution. Yunnan Province, China; Europe.

Specimens examined. CHINA, Yunnan, from *Tomicus minor* and *T. yunnanensis* galleries in *Pinus yunnanensis*, Feb. 2017, Nov. 2016, HM Wang, CFCC 52611 = CXY 1866, CFCC 52612 = CXY 1865, CFCC 52613 = CXY 1868.

Note. Our strains of *O. tingens* were identified based on phylogenetic affinities and morphological features. (cf. above under note for *O. aggregatum*.)

Ophiostoma tingens has been reported from sapwood of various *Pinus* spp. (including *P. sylvestris*) infested by *T. minor*, *T. piniperda* and *Ips sexdentatus* in Europe (Francke-Grosmann 1952, Batra 1967, Jankowiak 2008). The species was recorded in Yunnan Province in China in 2017, associated with *T. minor* infesting *P. yunnanensis* (Pan et al. 2017).

***Sporothrix macroconidia* H. Wang, Q. Lu & Z. Zhang, sp. n.**

MycoBank: MB828886

Fig. 15

Etymology. ‘*macroconidia*’ (Latin), referring to the large conidia of this fungus.

Type. CHINA, Yunnan, from *Tomicus yunnanensis* galleries in *Pinus yunnanensis*, Dec. 2016, collected by HM Wang, holotype CXY 1894, culture ex-holotype CFCC 52628 = CXY 1894.

Description. Sexual form: unknown.

Asexual form: *Sporothrix*-like. *Conidiophores* semi-macronematous, mononematous; *conidiogenous cells* hyaline, simple or loosely branched, thin-walled, aseptate, bearing denticles forming a rachis (4.1–) 11.0–24.5 (–36.5) × (1.4–) 2.1–3.4 (–4.9) μm; *conidia* hyaline, cylindrical, ellipsoid to ovoid, 1-celled, smooth, (3.6–) 4.8–7.4 (–9.9) × (2.5–) 3.2–4.9 (–9.9) μm, solitarily or aggregating in slimy masses.

Culture characteristics. Colonies on 2% MEA medium slow growing in the dark, reaching 34 mm in diam. in 8 days at 25 °C, growth rate up to 5 mm/day at the fastest; colony margin smooth. Hyphae appressed to flocculose, white; reverse hyaline to pale yellowish. Optimal growth temperature 25 °C, little growth at 5 °C and 35 °C.

Known substrates and hosts. Galleries of *Tomicus yunnanensis* and *T. brevipilosus* in *Pinus yunnanensis* and *P. kesiya*.

Known insect vectors. *Tomicus yunnanensis*, *T. brevipilosus*.

Known distribution. Yunnan Province, China.

Additional specimens examined. CHINA, Yunnan, from *Tomicus brevipilosus* galleries in *Pinus kesiya*, Dec. 2016, Jan. 2017, HM Wang, CFCC 52629 = CXY 1895, CFCC 52630 = CXY 1896.

Note. *Sporothrix macroconidia* is closely related to *O. valdivianum*, *S. bragantina*, *S. brunneoviolacea* and *S. fumea* in phylogenetic analyses inferred from LSU, ITS, *TUB2* and *CAL* DNA sequence data. It differs from these species by its conidia, which are larger than those of the other four species, mostly 4.8–7.4 × 3.2–4.9 μm and 4–6 × 2 μm in *O. valdivianum* (Butin and Aquilar 1984), 4–6 × 2–2.5 μm in *S. bragantina* (Pfenning and Oberwinkler 1993), 3–7 × 1.5–3 μm in *S. brunneoviolacea* (Madrid et al. 2010) and 1.5–2.0 × 0.5–1.0 μm in *S. fumea* (Nkuekam et al. 2012). In addition, a sexual state was observed *in vitro* for *O. valdivianum*, *S. bragantina* and *S. fumea*, which was not observed in *S. macroconidia* and *S. brunneoviolacea*.

Sporothrix macroconidia was found associated with *T. yunnanensis* infesting *P. yunnanensis* and with *T. brevipilosus* infesting *P. kesiya*. The other four similar species have




Figure 15. Morphological characters of *Sporothrix macroconidia* sp. n. **A, B** Upper and reverse of cultures on 2% MEA 20 d after inoculation **C–H** conidiogenous cells of *Sporothrix*-like asexual state and conidia. Scale bars: 10 μm (**C–H**).

very different ecology and known geographic distributions. *Sporothrix fumea* was isolated from *Eucalyptus cloeziana* infested by *Phoracantha* beetles in South Africa (Nkuekam et al. 2012), whereas *O. valdivianum*, *S. bragantina* and *S. brunneoviolacea* were obtained from soil or *Nothofagus* in Europe and South America (Butin and Aquilar 1984, Pfenning 1993, Madrid et al. 2010).

***Sporothrix pseudoabietina* H. Wang, Q. Lu & Z. Zhang, sp. n.**

Mycobank: MB828887

Fig. 16

Etymology. ‘*pseudoabietina*’ (Latin), referring to the phylogenetic affinities to *S. abietina*.

Type. CHINA, Yunnan, from *T. minor* galleries in *P. yunnanensis*, Apr. 2017, HM Wang, holotype CXY 1937, culture ex-holotype CFCC 52626 = CXY 1937.

Description. Sexual form perithecial: on 2% MEA, *perithecia* superficial or partially immersed, with a globose base extending into a cylindrical neck, often terminated by ostiolar hyphae; bases (85–) 110–152 (–168) μm diam., black, the outer layer with dark brown hyphal ornamentation; apical neck mild to dark brown at the




Figure 16. Morphological characters of *Sporothrix pseudoabietina* sp. n. **A, B** upper and reverse of cultures on 2% MEA 20 d after inoculation **C, D** ostiolar hyphae present **E, F** perithecium **G** ascospores of sexual state **H–I** conidiogenous cells of *Sporothrix*-like asexual state and conidia. Scale bars: 20 µm (**C, D**); 50 µm (**E, F**); 10 µm (**G–I**).

base, pale brown to pale yellow or hyaline toward the apex, straight or slightly curved, (172–) 560–985 (–1039) µm long, (37–) 41–62 (–78) µm wide at the base, (9.3–) 12.5–17.5 (–20) µm wide at the apex; *ostiolar* hyphae numerous, hyaline, divergent, (19.5–) 21.5–38.0 (–43) µm long; *asci* not seen; *ascospores* hyaline, 1-celled, orange-shaped in lateral view, ellipsoid in face view, circular in polar view, (2.9 –) 3.4–4.4 (–5.3) × (0.8–) 1.0–1.5 (–1.9) µm, without mucilaginous sheath.

Asexual form: *Sporothrix*-like. *Conidiophores* semi-macronematous to mononematous; *conidiogenous cells* hyaline, simple or loosely branched, smooth, bearing denticles disposed in a dense rachis (16.0–) 20.5–30.5 (–34.5) × (1.2–) 1.6–2.0 (–2.3) µm; *conidia* 1-celled, clavate, ellipsoid to ovoid, hyaline, (3.0–) 4.0–7.0 (–9.0) × (1.0–) 1.1–3.1 (–4.8) µm.

Culture characteristics. Colonies on 2% MEA slow growing in the dark, reaching 23 mm in diam. in 8 days at 25 °C, growth rate up to 2.5 mm/day at the fastest; colony margin smooth. Hyphae appressed to flocculose or floccose, white; reverse hyaline to pale yellowish. Optimal growth temperature 25 °C; very slow growth at 35 °C; no growth at 5 °C.

Known substrate and hosts. Galleries of *Tomicus yunnanensis* and *T. minor* in *Pinus yunnanensis*.

Known insect vectors. *Tomicus yunnanensis*, *T. minor*.

Known distribution. Yunnan Province, China.

Additional specimen examined. CHINA, Yunnan, *Tomiscus minor* galleries in *Pinus yunnanensis*, Apr. 2017, HM Wang, CFCC 52627 = CXY 1938.

Note. *Sporothrix pseudoabietina* is characterised by a perithecial sexual form and a *Sporothrix*-like asexual state. Multiple phylogenetic inferences (LSU, ITS, *TUB2* and *CAL*) showed that *S. pseudoabietina* belonged to the *S. gossypina* complex, in which it is closely related to *S. abietina*. However, it can be distinguished from this species, based on both morphological and physiological features. The conidia of *S. pseudoabietina* ($4.0\text{--}7.0 \times 1.1\text{--}3.1 \mu\text{m}$) are wider than those of *S. abietina* ($4\text{--}7.5 \times 1\text{--}2 \mu\text{m}$) (Marmolejo and Butin 1990). Perithecia are known from *S. abietina* but only on natural substrates and not *in vitro* on artificial media, contrary to those from *S. pseudoabietina*. The perithecial neck in *S. pseudoabietina* is much longer than that of *S. abietina*, viz. mostly 560–985 μm and 450–650 μm , respectively. Ostiolar hyphae of *S. abietina* and *S. pseudoabietina* also differ in number, numerous vs 7–10 and size, mostly 13–19 μm and in *S. pseudoabietina* 21.5–38.0 μm (Fig. 11c, d). In addition, no growth of *S. abietina* was observed at 35 °C, but *S. pseudoabietina* can grow at 35 °C.

The hosts and geographic distributions of *S. pseudoabietina* and *S. abietina* are also very different. *Sporothrix pseudoabietina* was found associated with *T. minor* and *T. yunnanensis* infecting *P. yunnanensis*, whereas *S. abietina* was reported from *Abies vejari* attacked by *Pseudohylesinus* sp. in Mexico (Marmolejo and Butin 1990).

Discussion

In this study, 772 strains of ophiostomatoid fungi were isolated from galleries and adults of three pine shoot beetles, *T. brevipilosus*, *T. minor* and *T. yunnanensis*, inhabiting *P. yunnanensis* and *P. kesiya* in forests in Yunnan Province, south-western China. Multiple phylogenetic analyses and morphological features allowed the identification of 11 species from 5 genera. Six species corresponded to known taxa (*E. vermicola*, *L. yunnanense*, *O. brevipilosi*, *O. canum*, *O. minus* and *O. tingens*), whereas four species are proposed as new, *Gra. anningense*, *O. aggregatum*, *S. pseudoabietina* and *S. macroconidia*. A single strain remained unnamed.

The global ophiostomatoid fungal communities, associated with these three *Tomiscus* species in pine forest, were dominated by far by three species, which are, in decreasing order of isolates, *O. canum*, *O. brevipilosi* and *O. minus*. Furthermore, these three ophiostomatoid species are not equally associated with the three *Tomiscus* species but show variable degrees of preference or specificity.

Overall, *O. canum* was the most frequently isolated species in our study (253 out of the 772 strains). It was preferably (79.4% of the *O. canum* strains) isolated from galleries and adults of *T. minor*, infesting both *P. yunnanensis* and *P. kesiya* (Table 3) and dominated the ophiostomatoid community associated with this beetle (81.4%, 201 strains of *O. canum* out of 247 strains in the community, Table 3).

This is the first report of this species in China. It was previously reported in eastern Asia but only in Japan (Masuya et al. 1999). *Ophiostoma canum* was also shown to be

the dominant species associated with *T. minor*, both in Europe and Japan (Masuya et al. 1999, Jankowiak 2008). In addition, this species was found in association with other bark beetles in Finland and Russia, e.g. *Hylastes brunneus*, *Hylurgops palliatus*, *Ips typographus*, *Pityogenes chalcographus* and *Trypodendron lineatum* (Linnakoski et al. 2010). The close association between *O. canum* and *T. minor* appears stable over an extensive geographical distribution and tree host range, indicating likely intimate relationships.

Ophiostoma brevipilosi represented the second most frequently isolated species in our survey (224 out of 772 strains), occurring exclusively in galleries and adults of *T. brevipilosus*, dominating this beetle's ophiostomatoid community (98.2%, 224 strains of *O. brevipilosi* out of 228 strains in the community, Table 3). The occurrence or fitness of *O. brevipilosi* is therefore strongly linked to the presence of *T. brevipilosus*.

Ophiostoma brevipilosi was described originally from Yunnan, based on six strains, all isolated from *T. brevipilosus* (Chang et al. 2017). It belongs to the recently defined *O. clavatum* complex (Linnakoski et al. 2016). It is only known from this area of south-western China.

Ophiostoma minus was the third most frequently isolated species overall (197 strains out of 772), occurring exclusively in galleries and adults of *T. yunnanensis* infesting *P. yunnanensis*, dominating this beetle ophiostomatoid community (66.3%, 197 strains of *O. minus* out of 297 strains in the community, Table 3).

Ophiostoma minus, first reported as a blue-stain agent in Europe (Munch 1907), is a widely distributed species, also recorded from North America and East Asia (Japan and China) (Hedgcock 1906, Gorton and Webber 2000, Gorton et al. 2004, Lu et al. 2009, Linnakoski et al. 2010). It infests various pines and is transported by various bark beetles. This species was predominantly associated with *T. piniperda* in Europe (Jankowiak 2006) and Japan (Masuya et al. 1999) and with the southern pine beetle, *Dendroctonus frontalis*, in the southern states of the USA (Klepzig 1998, Gorton and Webber 2000, Gorton et al. 2004).

Ophiostoma minus was deemed to have two allopatric populations, viz. a North American and a Eurasian population (Gorton et al. 2004). In ITS/*TUB2* phylogenetic inferences, the North American and Eurasian populations of *O. minus* were resolved as two closely related clades (Gorton et al. 2004, Lu et al. 2009). ITS and *TUB2*-based phylogenetic inferences (Fig. 7c, d) also resolved our strains as a third distinct clade, which could thus be interpreted as a third allopatric population. The question of translating these populations into a Linnaean taxonomic rank, however, remains open.

Tomicus yunnanensis galleries and adult beetles harboured the highest diversity of ophiostomatoid fungi; ten of the 11 species identified were isolated from galleries and adults of this beetle. Three species were exclusively found with this beetle (*O. minus*, *E. vermicola*, *Ophiostoma* sp. 1). By comparison, galleries and adults of *T. minor* and of *T. brevipilosus* yielded less species; five species were isolated from *T. minor*, none of which was associated exclusively with this beetle and three species from *T. brevipilosus*, of which one was exclusive, *O. brevipilosi*. Five species are shared by both *T. yunnanensis* and *T. minor* and two species by both *T. yunnanensis* and *T. brevipilosus*, but none by *T. minor* and *T. brevipilosus* and also none by all three pine shoot beetles (Table 3, Fig. 17).




Figure 17. Venn diagram showing overlaps of the ophiostomatoid fungal communities associated with three pine shoot beetles.

The ectosymbiosis between bark beetles and fungi is widespread and diverse. Some fungi are highly specific and associated with a single beetle species, forming a ‘species-specific association’ (Six and Paine 1999, Six 2012), while others can be associated with many vectors (Kostovcik et al. 2014). The species-specific associations include, for instance, *Ips typographus* and *Endoconidiophora polonica*, *I. cembrae* and *End. laricicola* (Harrington et al. 2002) or *I. subelongatus* and *End. fujiensis* (Marin et al. 2005, Meng et al. 2015). The present study showed that species-specific associations might occur with various sympatric beetles that share the same niche. The association of *T. brevipilosus* and *O. brevipilosus* seems to be species-specific in the pine forest of Yunnan, where both taxa are, so far, endemic. In the pine forest of Yunnan, the Chinese ‘population’ of *O. minus* is also specifically associated with *T. yunnanensis*, whereas the two other *O. minus* ‘populations’ are associated, at least preferably, with *Dendroctonus frontalis* and *T. piniperda* (Gorton et al. 2004, Jankowiak 2006). The genetically distinct ‘populations’ might originate from both

the allopatric distribution and vector specificity and both factors could support recognition of three distinct taxa. In the pine forest of Yunnan, the association of *O. canum* with *T. minor* is preferential but not exclusive.

Up to now, no data have been provided proving the pathogenicity of these ophiostomatoid species to both indigenous pines, except for *L. yunnanense* (Liao and Ye 2004, Gao et al. 2017). Pathogenicity tests have been done by artificial inoculation of the dominant species into seedlings of the two pines. The results preliminarily showed that the virulence of *O. minus* and *O. brevopilosi* was significantly stronger than that of *O. canum*. This is similar to the relative aggressive nature of the three *Tomicus* species. Thus, we suspect there might be some link between beetle aggression and fungus virulence (Christiansen et al. 1987, Kirisits 2004).

Conclusions

This study provides evidence for the diversity of ophiostomatoid fungi associated with *T. yunnanensis*, *T. minor* and *T. brevopilosus* in Yunnan pine forest in south-western China. Eleven species were identified, of which four were new to science. The diversity is the highest in the galleries and adults of *T. yunnanensis* and the poorest in the galleries and adults of *T. brevopilosus*.

Three species, namely *O. brevopilosi*, *O. canum* and *O. minus*, dominate the ophiostomatoid communities; each is associated predominantly with one species of *Tomicus*, namely *T. brevopilosus*, *T. minor* and *T. yunnanensis*, respectively. In this regard, this study has revealed differential associations between beetles living sympatrically, concomitantly or sequentially, in the same ecological niche, which indicates a certain level of specificity of the relationships between the fungi and the beetles. However, the parameters behind these (partial) species-specific relationships remain unknown.

Increased study of the biodiversity, biogeography and ecology of ophiostomatoid fungi in China, in particular of those associated with *Tomicus* spp., would facilitate comparison with well-known species associated with other *Tomicus* spp. in other neighbouring or distant geographical areas, e.g. in European countries, Japan and Korea and allow a better understanding of the occurrence and mechanisms behind the outbreak of infections, enabling the development of effective management methods to alleviate the subsequent plant losses.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (Project No.: 31770693 and 31770682). We are very grateful to Shuangcheng Li and Hongxun Wang for their help in field survey and collection. Cony Decock gratefully acknowledges the financial support received from the Belgian State–Belgian Federal Science Policy through the BCCM programme.

References

- Batra LR (1967) Ambrosia fungi: a taxonomic revision, and nutritional studies of some species. *Mycologia* 59: 976–1017. <https://doi.org/10.2307/3757271>
- Ben Jamaa ML, Lieutier F, Yart A, et al. (2007) The virulence of phytopathogenic fungi associated with the bark beetles *Tomicus piniperda* and *Orthotomicus erosus* in Tunisia. *For. Pathol.* 37: 51–63. <https://doi.org/10.1111/j.1439-0329.2007.00478.x>
- Butin H, Aquilar M (1984) Blue-Stain Fungi on Nothofagus from Chile-Including Two New Species of *Ceratocystis* Ellis and Halst. *Journal of Phytopathology* 109(1): 80–89. <https://doi.org/10.1111/j.1439-0434.1984.tb04233.x>
- Chang R, Duong TA, Taerum SJ, et al. (2017) Ophiostomatoid fungi associated with conifer-infesting beetles and their phoretic mites in Yunnan, China. *MycKeys* 28: 19. <https://doi.org/10.3897/mycokeys.28.21758>
- Chen P, Li L, Liu H (2009) Interspecific competition between *Tomicus yunnanensis* and *T. minor* (Col., Scolytidae) during shoot-feeding period in Yunnan of China. *Journal of West China Forestry Science* 38(3): 52–60.
- Chen P, Li L, Liu H (2010) Host preference and competition in bark beetles, *Tomicus yunnanensis* and *T. minor* in breeding period. *Journal of West China Forestry Science* 39(1) 15–20.
- Chen P, Lu J, Haack RA, Ye H (2015) Attack pattern and reproductive ecology of *Tomicus brevipilosus* (Coleoptera: Curculionidae) on *Pinus yunnanensis* in south-western china. *Journal of Insect Science* 15(1). <https://doi.org/10.1093/jisesa/iev014>
- Christiansen E, Waring RH, Berryman AA (1987) Resistance of conifers to bark beetle attack: searching for general relationships. *Forest Ecology and Management* 22: 89–106. [https://doi.org/10.1016/0378-1127\(87\)90098-3](https://doi.org/10.1016/0378-1127(87)90098-3)
- de Beer ZW, Duong TA, Wingfield MJ (2016) The divorce of *Sporothrix* and *Ophiostoma*: solution to a problematic relationship. *Studies in mycology* 83: 165–191. <https://doi.org/10.1016/j.simyco.2016.07.001>
- de Beer ZW, Wingfield MJ (2013) Emerging lineages in the Ophiostomatales. In: KA Seifert, ZW de Beer, MJ Wingfield. *The Ophiostomatoid Fungi: Expanding Frontiers*. Utrecht, The Netherlands CBS, 21–46.
- de Beer ZW, Seifert KA, Wingfield MJ (2013a) The ophiostomatoid fungi: Their dual position in the sordariomycetes. In: Seifert KA, de Beer ZW, Wingfield MJ (Eds) *The ophiostomatoid fungi: Expanding frontiers*. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands, 1–19.
- de Beer ZW, Seifert KA, Wingfield MJ (2013b) A nomenclator for ophiostomatoid genera and species in the Ophiostomatales and Microascales. In: Seifert KA, de Beer ZW, Wingfield MJ (Eds) *The ophiostomatoid fungi: Expanding frontiers*. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands, 245–322.
- Francke-Grosmann H (1952) Über die Ambrosiazucht der beiden Kiefernborckenkäfer *Myelophilus minor* Htg. und *Ips acuminatus* Gyll. *Medd. Statens Skogforskningsinst* 41: 1–52
- Gao X, Zhou X, Wang H, Kong, et al. (2017) Influence of severe drought on the resistance of *Pinus yunnanensis*, to a bark beetle-associated fungus. *Forest Pathology* 47(4): e12345. <https://doi.org/10.1111/efp.12345>

- Gibbs JN, Inman A (1991) The pine shoot beetle *Tomicus piniperda* as a vector of blue stain fungi to windblown pine. *Forestry* 64: 239–249. <https://doi.org/10.1093/forestry/64.3.239>
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied Environmental Microbiology* 61: 1323–1330. DOI: 10.0000/PMID7747954
- Gorton C, Webber JF (2000) Reevaluation of the status of the blue-stain fungus and bark beetle associate *Ophiostoma minus*. *Mycologia* 92: 1071–1079. <https://doi.org/10.2307/3761474>
- Gorton C, Kim SH, Henricot B, et al. (2004) Phylogenetic analysis of the blue-stain fungus *Ophiostoma minus* based on partial ITS rDNA and β -tubulin gene sequences. *Mycological Research* 108: 759–765. <https://doi.org/10.1017/S0953756204000012>
- Harrington TC, McNew D, Steimel J, et al. (2001) Phylogeny and taxonomy of the *Ophiostoma piceae* complex and the Dutch elm disease fungi. *Mycologia* 111–136. <https://doi.org/10.2307/3761610>
- Harrington TC, Pashenova NV, McNew DL, et al. (2002) Species delimitation and host specialization of *Ceratocystis laricicola* and *C. polonica* to Larch and Spruce. *Plant Disease* 86: 418–422. <https://doi.org/10.1094/PDIS.2002.86.4.418>
- Hedgcock GG (1906) Studies upon some chromogenic fungi which discolor wood. *Missouri Botanical Garden Annual Report* 17: 59–114. <https://doi.org/10.2307/2400089>
- Jacobs K, Bergdahl DR, Wingfield MJ, et al. (2004) *Leptographium wingfieldii* introduced into North America and found associated with exotic *Tomicus piniperda* and native bark beetles. *Mycol Res* 108(4): 411–418. <https://doi.org/10.1017/S0953756204009748>
- Jankowiak R (2006) Fungi associated with *Tomicus piniperda* in Poland and assessment of their virulence using Scots pine seedlings. *Annals of forest science*, 63(7): 801–808. <https://doi.org/10.1051/forest:2006063>
- Jankowiak R (2008) Fungi associated with *Tomicus minor* on *Pinus sylvestris* in Poland and their succession into the sapwood of beetle-infested windblown trees. *Canadian journal of forest research* 38(10): 2579–2588. <https://doi.org/10.1139/X08-101>
- Jankowiak R, Bilański P (2007) Fungal flora associated with *Tomicus piniperda* L. in an area close to a timber yard in southern Poland. *Journal of applied entomology* 131(8): 579–584. <https://doi.org/10.1111/j.1439-0418.2007.01194.x>
- Ji M, Dong XQ, Liu HP, et al. (2007) Preliminary Study on Remote Sensing Detection of Yunnan Pine Forest Damaged by *Tomicus piniperda*. *Journal of West China Forestry Science* 36(1): 87–90.
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular biology and evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kim JJ, Lim YW, Breuil C, et al. (2005) A new *Leptographium* species associated with *Tomicus piniperda* infesting pine logs in Korea. *Mycological research* 109(3): 275–284. <https://doi.org/10.1017/S0953756204002060>
- Kirkendall LR, Faccoli M, Ye H (2008) Description of the Yunnan shoot borer, *Tomicus yunnanensis* Kirkendall and Faccoli sp. n.(Curculionidae, Scolytinae), an unusually aggressive pine shoot beetle from southern China, with a key to the species of *Tomicus*. *Zootaxa* 1819: 25–39.

- Kirisits T (2004) Fungal associates of European bark beetles with special emphasis on the ophiostomatoid fungi. In: Lieutier F, Day KR, Battisti A, Grégoire JC, Evans HF (Eds) *Bark and Wood Boring Insects in Living Trees in Europe, A Synthesis*. Kluwer Academic Publishers, The Netherlands, 181–235. https://doi.org/10.1007/978-1-4020-2241-8_10
- Klepzig KD (1998) Competition between a biological control fungus, *Ophiostoma piliferum*, and symbionts of the southern pine beetle. *Mycologia* 90: 69–75. <https://doi.org/10.1080/00275514.1998.12026880>
- Kubátová A, Novotný D, Prášil K, et al. (2000) The nematophagous hyphomycete *Esteya vermicola* found in the Czech Republic. *Czech Mycology* 52(3): 227–235.
- Kostovcik M, Bateman CC, Kolarik M, et al. (2014) The ambrosia symbiosis is specific in some species and promiscuous in others: evidence from community pyrosequencing. *Isme Journal* 9(1): 126. <https://doi.org/10.1038/ismej.2014.115>
- Lagerberg, T (1927) Biological and practical researches into blueing in pine and spruce. *Sven Skogsvardsforen Tidskr* 25, 145–272.
- Li LS, Wang HL, Chai X, Shu N (1997) Study on the Biological Characteristics of *Tomicus piniperda* and Its Damage. *Yunnan Forestry Science and Technology* 2: 1–7.
- Li LS, Liu HP, Chen P, et al. (2006) Comparison study on ecological characteristics of *Tomicus piniperda* and *T. minor*. *Journal of West China Forestry Science* 35(1): 1–5.
- Li X, Zhang Z, Wang HB, et al. (2010) *Tomicus armandii* Li & Zhang (Curculionidae, Scolytinae), a new species from China. *Zootaxa* 2572:57–64.
- Li Y, Huang YT, Kasson MT, et al. (2018) Specific and promiscuous ophiostomatalean fungi associated with Platypodinae ambrosia beetles in the southeastern United States. *Fungal Ecology* 35: 42–50. <https://doi.org/10.1016/j.funeco.2018.06.006>
- Liao ZY, Ye H (2004) Action mechanisms of phytotoxin produced by *Leptographium yunnanense* associated with *Tomicus*. *Forest Pest and Disease* 23(1): 20–22.
- Lieutier F (2002) Mechanisms of resistance in conifers and bark beetle attack strategies. In: Wagner MR, Clancy KM, Lieutier F, Paine TD (Eds) *Mechanisms and Deployment of Resistance in Trees to Insects*. Kluwer, Dordrecht, 31–75. https://doi.org/10.1007/0-306-47596-0_2
- Lieutier F, Ye H, Yart A (2003) Shoot damage by *Tomicus* sp. (Coleoptera: Scolytidae) and effect on *Pinus yunnanensis* resistance to subsequent reproductive attacks in the stem. *Agric. Forest Entomol* 5: 227–233. <https://doi.org/10.1046/j.1461-9563.2003.00182.x>
- Lieutier F, Yart A, Garcia J, et al. (1989) Champignons phytopathogènes associés à deux coléoptères scolytidae du pin sylvestre (*Pinus sylvestris* L.) et étude préliminaire de leur agressement envers l'hôte, *Ann. Sci. For.* 46: 201–216 <https://doi.org/10.1051/forest:19890301> [in French with English summary].
- Lieutier F, Langstrom B, Faccoli M (2015) The genus *Tomicus*. In F. E. Vega and R. W. Hofstetter (eds.), *Bark beetles: biology and ecology of native and invasive species*. Elsevier, Amsterdam, The Netherlands, 371–426. <https://doi.org/10.1016/B978-0-12-417156-5.00010-1>
- Linnakoski R, Jankowiak R, Villari C, Kirisits T, Solheim H, et al. (2016). The *Ophiostoma clavatum* species complex: a newly defined group in the Ophiostomatales including three novel taxa. *Antonie van Leeuwenhoek* 109: 987–1018. <https://doi.org/10.1007/s10482-016-0700-y>

- Linnakoski R, Beer ZWD, Niemelä P, et al. (2012) Associations of conifer-infesting bark beetles and fungi in Fennoscandia. *Insects* 3(1): 200–227. <https://doi.org/10.3390/insects3010200>
- Linnakoski R, de Beer ZW, Ahtiainen J, et al. (2010) *Ophiostoma* spp. associated with pine-and spruce-infesting bark beetles in Finland and Russia. *Persoonia: Molecular Phylogeny and Evolution of Fungi* 25: 72. <https://doi.org/10.3390/insects3010200>
- Linnakoski R, de Beer ZW, Rousi M, et al. (2008) Fungi, including *Ophiostoma karelicum* sp. n. associated with *scolytus ratzeburgi* infesting birch in finland and russia. *Mycological Research* 112(12): 1475–1488. <https://doi.org/10.1016/j.mycres.2008.06.007>
- Liou JY, Shih JY, Tzean SS (1999) *Esteya*, a new nematophagous genus from Taiwan, attacking the pinewood nematode (*Bursaphelenchus xylophilus*). *Mycological Research* 103(2): 242–248. <https://doi.org/10.1017/S0953756298006984>
- Lu M, Zhou XD, De Beer ZW, et al. (2009) *Ophiostomatoid* fungi associated with the invasive pine-infesting bark beetle, *Dendroctonus valens*, in China. *Fungal Diversity* 38: 133–145.
- Lu J, Zhao T, Ye H, et al. (2014) The shoot-feeding ecology of three *Tomicus* species in Yunnan Province, southwestern China. *Journal of Insect Science* 14: 37. <https://doi.org/10.1093/jis/14.1.37>
- Lu RC, Wang HB, Zhang Z, et al. (2012) Coexistence and competition between *Tomicus yunnanensis* and *T. minor* (coleoptera: scolytinae) in yunnan pine. *Psyche A Journal of Entomology* 2012(3): S17. <https://doi.org/10.1155/2012/185312>
- Madrid H, Gené J, Cano J, et al. (2010) *Sporothrix brunneoviolacea* and *Sporothrix dimorphospora*, two new members of the *Ophiostoma stenoceras-Sporothrix schenckii* complex. *Mycologia* 102(5): 1193–1203. <https://doi.org/10.3852/09-320>
- Marmolejo JG, Butin H (1990) New conifer-inhabiting species of *Ophiostoma* and *Ceratocystiopsis* (Ascomycetes, Microascales) from Mexico. *Sydowia* 42: 193–199.
- Marin M, Preisig O, Wingfield BD, et al. (2005) Phenotypic and DNA sequence data comparisons reveal three discrete species in the *Ceratocystis polonica* species complex. *Mycological Research* 109(10): 1137–1148. <https://doi.org/10.1017/S095375620500362X>
- Masuya H, Kaneko S, Yamaoka Y, et al. (1999) Comparisons of ophiostomatoid fungi associated with *Tomicus piniperda* and *T. minor* in Japanese red pine. *Journal of Forest Research* 4(2): 131–135. <https://doi.org/10.1007/BF02762237>
- Mathiesen A (1950) Über einige mit Borkenkäfern assoziierten Bläuepilze in Schweden. *Oikos* 2: 275–308. <https://doi.org/10.2307/3564798>
- Mathiesen A (1951) Einige neue Ophiostoma-Arten in Schweden. *Sven. Bot. Tidskr* 45: 203–232.
- Mathiesen-Käärrik A (1953) Eine Übersicht über die gewöhnlichsten mit Borkenkäfern assoziierten Bläuepilze in Schweden und einige für Schweden neue Bläuepilze, Meddn. St. Skogforsk Inst, 431–74.
- Meng XJ, Lu Q, Liu XW, et al. (2015) The species specific associations between *Ips subelongatus* and ophiostomatoid fungi. *Acta Ecologica Sinica* 35(2): 313–323.
- Munch E (1907) Die Blaufaule des Nadelholzes. *Naturwiss. Ztschr. F. Forst .U.Landw* 5: 531–573.

- Nkuekam G K, de Beer Z W, Wingfield M J, et al. (2012) A diverse assemblage of *Ophiostoma* species, including two new taxa on eucalypt trees in South Africa. *Mycological progress* 11(2): 515–533. <https://doi.org/10.1007/s11557-011-0767-9>
- Ohtaka N, Masuya H, Yamaoka Y, et al. (2006) Two new *Ophiostoma* species lacking conidial states isolated from bark beetles and bark beetle-infested *Abies* species in Japan. *Botany* 84(2): 282–293. <https://doi.org/10.1139/b05-164>
- Paciura D, Zhou XD, de Beer ZW, et al. (2010) Characterisation of synnematosus bark beetle-associated fungi from China, including *Graphium carbonarium* sp. n. *Fungal Diversity* 40(1): 75–88. <https://doi.org/10.1007/s13225-009-0004-x>
- Pan Y, Chen P, Lu J, et al. (2017) First report of blue-stain in *Pinus yunnanensis* caused by *Ophiostoma tingers* associated with *Tomicus minor* in china. *Journal of Plant Pathology* 99(3): 805.
- Pfenning L, Oberwinkler F (1993) *Ophiostoma bragantinum* n. sp., a possible teleomorph of *Sporothrix inflata*, found in Brazil. *Mycotaxon* 46: 381–385.
- Rayner RW (1970) A mycological colour chart. CMI and British Mycological Society, Kew, Surrey, UK.
- Rennerfelt E (1950) Über den Zusammenhang zwischen dem Verblauendes Holzes und den Insekten. *Oikos* 2: 120–137. <https://doi.org/10.2307/3564666>
- Seifert KA, Webber JF, Wingfield MJ (1993) Methods for studying species of *Ophiostoma* and *Ceratocystis*. *Ceratocystis* and *Ophiostoma*: taxonomy, ecology and pathogenicity. Minnesota, U.S.A., The American Phytopathological Society Press, 255–259.
- Sabbatini Peverieri G, Capretti P, Tiberi R (2006) Associations between *Tomicus destruens* and *Leptographium* spp. in *Pinus pinea* and *P. pinaster* stands in Tuscany, central Italy. *For. Pathol* 36, 14–20. <https://doi.org/10.1111/j.1439-0329.2006.00427.x>
- Six DL (2003) Bark beetle–fungus symbiosis. In: Bourtzis K, Miller TA (Eds) *Insect Symbiosis*. CRC Press, Washington D.C., 97–114. <https://doi.org/10.1201/9780203009918.ch7>
- Six DL (2012) Ecological and Evolutionary Determinants of Bark Beetle–Fungus Symbioses. *Insects* 3: 339–366. <https://doi.org/10.3390/insects3010339>
- Six DL, Paine TDA (1999) Phylogenetic comparison of ascomycete mycangial fungi and *Dendroctonus* bark beetles (Coleoptera: Scolytidae). *Ann Entomol Soc Am* 92: 159–166. <https://doi.org/10.1093/aesa/92.2.159>
- Six DL, Wingfield MJ (2011) The role of phytopathogenicity in bark beetle–fungus symbioses: a challenge to the classic paradigm. *Annu Rev Entomol* 56: 255–272. <https://doi.org/10.1146/annurev-ento-120709-144839>
- Solheim H, Långström B (1991) Blue-stain fungi associated with *Tomicus piniperda* in Sweden and preliminary observations on their pathogenicity, *Ann. Sci. For.* 48: 149–156. <https://doi.org/10.1051/forest:19910203>
- Solheim H, Krokene P, Långström B (2001) Effects of growth and virulence of associated blue-stain fungi on host colonization behavior of the pine shoot beetles *Tomicus minor* and *T. piniperda*. *Plant Pathol* 50: 111–116. <https://doi.org/10.1046/j.1365-3059.2001.00541.x>

- Stamatakis A (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690. <https://doi.org/10.1093/bioinformatics/btl446>
- Swofford DL (2003) PAUP*: phylogenetic analysis using parsimony, Version 4. Sinauer Associates, Sunderland, MA. <https://doi.org/10.1007/BF02198856>
- Tamura K, Peterson D, Peterson N, et al. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28: 2731–2739. <https://doi.org/10.1093/molbev/msr121>
- Upadhyay HP (1981) A monograph of *Ceratocystis* and *Ceratocystiopsis*. Univ. Georgia Press. <https://doi.org/10.1007/BF02858713>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Wang CY, Fang ZM, Wang Z, et al. (2009) High infection activities of two *Esteya vermicola* isolates against pinewood nematode. *African Journal of Microbiology Research* 3(10): 581–584. <https://doi.org/10.1007/s12275-007-0122-7>
- Wang X, Wang T, Wang J, et al. (2014) Morphological, molecular and biological characterization of *Esteya vermicola*, a nematophagous fungus isolated from intercepted wood packing materials exported from Brazil. *mycoscience* 55(5): 367–377. <https://doi.org/10.1016/j.myc.2014.01.002>
- Wingfield M J, Garnas J R, Hajek A, et al. (2016) Novel and co-evolved associations between insects and microorganisms as drivers of forest pestilence. *Biological Invasions* 18(4): 1045–1056. <https://doi.org/10.1007/s10530-016-1084-7>
- Wingfield MJ, Barnes I, De Beer ZW, et al. (2017) Novel associations between ophiostomatoid fungi, insects and tree hosts: current status–future prospects. *Biological Invasions*: 1–14. <https://doi.org/10.1007/s10530-017-1468-3>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols: a guide to methods and application. Academic Press, San Diego, USA, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Yamaoka Y, Masuya H, Ohtaka N, et al. (2004) Three new *Ophiostoma* species with *Pesotum* anamorphs associated with bark beetles infesting *Abies* species in Nikko, Japan. *Mycoscience* 45(4): 277–286. <https://doi.org/10.1007/S10267-004-0179-7>
- Yamaoka Y, Chung WH, Masuya H, et al. (2008) The teleomorph of *Leptographium yunnanense*, discovered in crosses among isolates from Thailand, China, and Japan. *Mycoscience* 49(4): 233–240. <https://doi.org/10.1007/S10267-008-0412-X>
- Yin M, Wingfield MJ, Zhou X, et al. (2016) Multigene phylogenies and morphological characterization of five new *Ophiostoma* spp. associated with spruce-infesting bark beetles in China. *Fungal biology* 120(4): 454–470. <https://doi.org/10.1016/j.funbio.2015.12.004>
- Ye H (1991) On the bionomy of *Tomicus piniperda* (L.) (Coleoptera: Scolytidae) in the Kunming region of China. *Journal of Applied Entomology* 112: 366–369. <https://doi.org/10.1111/j.1439-0418.1991.tb01069.x>

- Ye H, Zhou X, Lu J, Yang L, Ding H (2000) A preliminary study on the occurrence and pathogenicity of the fungi associated with *Tomicus*. Forest Research, Beijing 13: 451–454.
- Ye H (2011) *Tomicus yunnanensis*. Kunming: Yunnan Science and Technology Publishing Press.
- Ye H, Dang CL (1986) The study on the feature of the bark beetle (*Blastophagus piniperda* L.) injuring the pine tree (*Pinus yunnanensis*). Journal of Yunnan University 8(2): 218–221.
- Ye H, F Lieutier (1997) Shoot aggregation by *Tomicus piniperda* L (Col: Scolytidae) in Yunnan, southwestern China. Ann. Forest Sci 54: 635–641. <https://doi.org/10.1051/forest:19970704>
- Ye H, Ding XS (1999) Impacts of *Tomicus minor* on distribution and reproduction of *Tomicus piniperda* (Col., Scolytidae) on the trunk of the living *Pinus yunnanensis* trees. J. Appl. Entomol 123: 329–333. <https://doi.org/10.1046/j.1439-0418.1999.00353.x>
- Zhang Y, Hagen F, Stielow B, et al. (2015) Phylogeography and evolutionary patterns in *Sporothrix* spanning more than 14 000 human and animal case reports. Persoonia 35: 1–20. <https://doi.org/10.3767/003158515X687416>
- Zhou XD, de Beer ZW, Wingfield MJ (2013) Ophiostomatoid fungi associated with conifer infecting bark beetles in China. In: Seifert KA, de Beer ZW, Wingfield MJ (Eds) Ophiostomatoid fungi: Expanding frontiers. CBS, Utrecht, The Netherlands, 91–98.
- Zhou XD, Jacobs K, Wingfield MJ, et al. (2000) A new *Leptographium* species associated with *Tomicus piniperda* in South Western China. Mycoscience 41: 573–578. <https://doi.org/10.1007/BF02460923>
- Zipfel RD, Beer ZW de, Jacobs K, et al. (2006) Multigene phylogenies define *Ceratocystiopsis* and *Grosmannia* distinct from *Ophiostoma*. Studies in Mycology 55: 75–97. <https://doi.org/10.3114/sim.55.1.75>