

# Aberrant internal carotid artery presenting as a retrotymppanic vascular mass

Simon Nicolay<sup>1,2</sup>, Bert De Foer<sup>1</sup>, Anja Bernaerts<sup>1</sup>, Joost Van Dinther<sup>3</sup> and Paul M Parizel<sup>2</sup>

Acta Radiologica Short Reports  
3(10) 1–3  
© The Foundation Acta Radiologica  
2014  
Reprints and permissions:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/2047981614553695  
arr.sagepub.com



## Abstract

We report a case of a young woman with an aberrant right internal carotid artery (ICA) presenting as a retrotymppanic reddish mass. This variant of the ICA represents the collateral pathway that is formed as a result of an embryological agenesis of the cervical segment of the ICA. The embryonic inferior tympanic artery is recruited to bypass the absent carotid segment. This hypertrophied vessel may be seen otoscopically and wrongfully considered to be a vascular middle ear tumor. Informing the otorhinolaryngologist of this important vascular variant not only obviates biopsy but also helps in careful preoperative planning of eventual middle ear procedures.

## Keywords

Head/Neck, vascular, computed tomography (CT), magnetic resonance angiography (MRA) arteries, ear

Date received: 2 September 2009; accepted: 9 September 2014

## Introduction

Underdevelopment of the internal carotid artery (ICA) can occur in different degrees. In order to maintain sufficient blood flow to the brain the body will recruit collateral arteries. The collateral route may vary depending on the extent and the location of the agenesis.

In case of partial absence of the cervical segment (C1) of the ICA a bypass via the embryonic arteries that run through the temporal bone will develop, causing a pseudo-tumoral vascular mass lesion in the middle ear. Computed tomography (CT) or cone beam CT (CBCT) can easily demonstrate the vascular anomaly and provide the otolaryngologist this valuable information. Magnetic resonance angiography (MRA) can be used to confirm the diagnosis of this vascular variant.

## Case report

A 21-year-old woman, with no relevant medical history, consulted the otorhinolaryngologist with complaints of right-sided hearing loss. Audiometry showed slight conductive hearing loss and neurological

examination was normal. Otoscopy of the right ear revealed a reddish structure behind the tympanic membrane. A tentative diagnosis of a vascular middle ear tumor was made. Computed tomography (CT) scan of the temporal bone was subsequently performed, showing a soft tissue lesion against the promontory. There was, however, also a remarkable difference in size between the vertical segment of the vertical petrous part of both internal carotid arteries (ICA), the right being much smaller (Fig. 1a and b). The latter was also found to run more laterally, bulging into the tympanic cavity – causing the soft tissue mass against the promontory – before joining the horizontal petrous segment

<sup>1</sup>Department of Radiology, GZA Hospital Sint-Augustinus, Wilrijk, Belgium


<sup>2</sup>Department of Radiology, Antwerp University Hospital (UZA) and University of Antwerp (UA), Belgium

<sup>3</sup>European Institute for ORL, GZA Hospital Sint-Augustinus, Wilrijk, Belgium

### Corresponding author:

Simon Nicolay, Department of Radiology, Antwerp University Hospital (UZA), Wilrijkstraat 10, 2650 Edegem, Belgium.  
Email: simon.nicolay@gmail.com





**Fig. 1.** Axial non-enhanced CT images at the level of the condylar process of the mandible (a, b) and at the level of the hypotympanon (c, d). A reduced-caliber vertical segment of the petrous portion of the right ICA is noted (a) in comparison to the normal sized left ICA (b). (c) The right ICA also has a more lateral course, bulging out in the hypotympanon giving rise to a soft tissue mass lesion against the promontory (arrow) before reaching the horizontal segment of the petrous ICA (asterisk). (d) No such artery is found connecting to the left horizontal segment of the petrous ICA (asterisk).

of the ICA (Fig. 1c and d). A right-sided partial agenesis of the ICA with collateral flow through the embryonic inferior tympanic artery was suspected. An MR imaging (MRI) examination with MRA sequences was performed to confirm this diagnosis. On unenhanced three-dimensional (3D) time of flight (TOF) MRA the difference between the normal-sized left ICA and the thin, more laterally running right ICA can be nicely depicted (Fig. 2).

Based on the imaging findings diagnosis of a partial agenesis of the right ICA with collateral circulation through the inferior tympanic artery, known as aberrant ICA, was made.

## Discussion

Aberrant ICA is a rare congenital condition, with a debated pathogenesis. In the embryological stadium the third of the so-called aortic arches forms the proximal cervical portion of the ICA, while the dorsal aorta

constitutes the distal portion. In aberrant ICA, also known as partial agenesis of the ICA, the presumed pathogenesis is regression or underdevelopment of the cervical part of the ICA at the skull base (1). In order to maintain adequate blood flow to the brain, an attempt to bypass this underdeveloped segment by using a collateral route is done. In most cases of underdevelopment of the ICA the circle of Willis is used to accommodate collateral flow. However, flow can also be provided by persistent embryonic vessels or branches originating from the external carotid artery (ECA). When there is agenesis of the cervical part of the ICA (C1), as in this case, the small arteries of the middle ear will be recruited to reach the horizontal petrous ICA segment. These vessels are the inferior tympanic artery, an embryonic branch of the ECA, and the caroticotympanic artery, a branch of the horizontal petrous part of the ICA. The enlarged inferior tympanic artery enters the tympanic cavity through the inferior tympanic canalculus, passes lateral to the cochlear promontory



**Fig. 2.** Unenhanced 3D TOF MRA of the intra-cranial arteries nicely shows the smaller CI segment of the right ICA running more laterally in comparison to the normal-sized left ICA, which follows a more medial course. Note also the coincident finding of an absence of the AI segment of the right anterior cerebral artery.

bulging into the middle ear cavity, eventually anastomosing with the caroticotympanic artery to reach the horizontal petrous ICA segment (2,3). The intratympanic course of the aberrant ICA may suggest a vascular mass on otoscopic examination. The glomus tympanicum is the most frequent hypervascular middle ear tumor with a typical location against the promontory (4). However, in case of a glomus tympanicum, the size and aspect of the vertical and horizontal intrapetrous segment of the carotid artery is completely normal. There is also no communication between the mass lesion and the horizontal intrapetrous carotis segment in case of a glomus tumor.

Though in most cases asymptomatic, symptoms such as pulsatile tinnitus or conductive hearing loss may sometimes be seen in aberrant ICA but these are very non-specific. Aberrant ICA can therefore not reliably be discerned from vascular tumors or other vascular malformations like an aneurysm or a hemangioma (5), based on clinical findings alone (3). A biopsy or surgical exploration of a presumed hypervascular mass can be complicated by life-threatening hemorrhage or neurological deficit. Evaluation of a possible vascular mass in the middle ear should therefore be done by imaging. Even more, to avoid surgical injury, CT scan of the temporal bone should ideally be performed before any middle ear surgery. Several features suggestive of an aberrant ICA can be seen on CT or CBCT. These include – as demonstrated – the absence of the vertical

segment of the ICA, distention of the inferior tympanic canaliculus, an aberrant ICA passing laterally through the tympanic space and dehiscence of the bone layer covering the tympanic segment of the ICA (6). On angiography a lateral and superior location of the carotid genu in the middle ear on the antero-posterior projection is typical (7). The ICA also runs laterally to the so-called vestibular line, a line drawn vertical from the lateral border of the vestibule (8). Since MRA is non-invasive and provides excellent visualization of the intracranial and extracranial circulation, this technique is to be preferred over conventional angiography (9).

In conclusion, an aberrant ICA is a rare vascular abnormality that may mimic a hypervascular middle ear mass lesion. Biopsy or surgical exploration of this pseudolesion can have devastating consequences. Therefore evaluation of a possible vascular middle ear mass should always be done by temporal bone CT. MRA can be used to confirm the diagnosis.

### Conflict of interest

None declared.

### References

1. Roll JD, Urban MA, Larson TC, et al. Bilateral aberrant internal carotid arteries with bilateral persistent stapediaal arteries and bilateral duplicated internal carotid arteries. *Am J Neuroradiol* 2003;24:762–765.
2. Endo K, Maruyama Y, Tsukatani T, et al. Aberrant internal carotid artery as a cause of objective pulsatile tinnitus. *Auris Nasus Larynx* 2006;33:447–450.
3. Verbist B, Hermans R, Devlies F, et al. Quiz case of the month. Partial agenesis of the internal carotid artery with collateral circulation through an enlarged inferior tympanic artery. *Eur Radiol* 1996;6:939–940.
4. Botma M, Kell RA, Bhattacharya J, et al. Aberrant internal carotid artery in the middle-ear space. *J Laryngol Otol* 2000;114:784–787.
5. Anderson JM, Stevens JC, Sundt TM, et al. Ectopic internal carotid artery seen initially as middle ear tumor. *JAMA* 1983;249:2228–2230.
6. Lo WW, Solti-Bohman LG, McElveen JT Jr. Aberrant carotid artery. Radiologic diagnosis with emphasis on high-resolution computed tomography. *Radiographics* 1985;5:985–993.
7. Davis WL, Harnsberger HR. MR Angiography of an aberrant internal carotid artery. *Am J Neuroradiol* 1991; 12:1225.
8. Lapayowker MS, Liebman EP, Ronis ML, et al. Presentation of the internal carotid artery as a tumor of the middle ear. *Radiology* 1971;98:293–297.
9. Caldas J, Iffenecker C, Attal P, et al. Anomalous vessel in the middle ear: the role of CT and MR angiography. *Neuroradiology* 1998;407:48–51.