

Research Article

Electrophilic PPAR γ Ligands Attenuate IL-1 β and Silica-Induced Inflammatory Mediator Production in Human Lung Fibroblasts via a PPAR γ -Independent Mechanism

Christopher M. Hogan,¹ Thomas H. Thatcher,¹ Ramil E. Sapinoro,^{2,3}
Michael N. Gurell,¹ Heather E. Ferguson,^{2,3} Stephen J. Pollock,^{2,3} Carolyn Jones,⁴
Richard P. Phipps,^{2,3} and Patricia J. Sime^{1,2,3}

¹ Division of Pulmonary and Critical Care Medicine, University of Rochester, 601 Elmwood Avenue, P.O. Box 692, Rochester, NY 14642, USA

² Department of Environmental Medicine, University of Rochester, Rochester, NY 14642, USA

³ Lung Biology and Disease Program, University of Rochester, Rochester, NY 14642, USA

⁴ Department of Surgery, University of Rochester, Rochester, NY 14642, USA

Correspondence should be addressed to Patricia J. Sime, patricia.sime@urmc.rochester.edu

Received 23 November 2010; Revised 4 March 2011; Accepted 25 March 2011

Academic Editor: Paul Drew

Copyright © 2011 Christopher M. Hogan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Acute and chronic lung inflammation is associated with numerous important disease pathologies including asthma, chronic obstructive pulmonary disease and silicosis. Lung fibroblasts are a novel and important target of anti-inflammatory therapy, as they orchestrate, respond to, and amplify inflammatory cascades and are the key cell in the pathogenesis of lung fibrosis. Peroxisome proliferator-activated receptor gamma (PPAR γ) ligands are small molecules that induce anti-inflammatory responses in a variety of tissues. Here, we report for the first time that PPAR γ ligands have potent anti-inflammatory effects on human lung fibroblasts. 2-cyano-3, 12-dioxoolean-1, 9-dien-28-oic acid (CDDO) and 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15d-PGJ₂) inhibit production of the inflammatory mediators interleukin-6 (IL-6), monocyte chemoattractant protein-1 (MCP-1), COX-2, and prostaglandin (PG)E₂ in primary human lung fibroblasts stimulated with either IL-1 β or silica. The anti-inflammatory properties of these molecules are not blocked by the PPAR γ antagonist GW9662 and thus are largely PPAR γ independent. However, they are dependent on the presence of an electrophilic carbon. CDDO and 15d-PGJ₂, but not rosiglitazone, inhibited NF- κ B activity. These results demonstrate that CDDO and 15d-PGJ₂ are potent attenuators of proinflammatory responses in lung fibroblasts and suggest that these molecules should be explored as the basis for novel, targeted anti-inflammatory therapies in the lung and other organs.

1. Introduction

Inflammation is associated with many diseases of the lung and can result from immunologic injury, infection, and inhalation of particulate matter. Diseases strongly associated with pulmonary inflammation include asthma, chronic obstructive pulmonary disease (COPD), and silicosis. Inflammation is also associated with an increased susceptibility to developing lung cancers and other malignancies [1–4]. Aside from glucocorticoids, few effective anti-inflammatory agents exist. In this regard, it is important to investigate new anti-inflammatory targets.

Peroxisome proliferator-activated receptor gamma (PPAR γ) has emerged as an important potential anti-inflammatory target. PPAR γ is a ligand-activated nuclear receptor that binds a variety of endogenous lipids and lipid-derived compounds. Upon ligand binding, PPAR γ heterodimerizes with the retinoid X receptor (RXR), translocates to the nucleus, and regulates expression of genes containing PPAR γ response elements (PPREs) [5]. PPAR γ ligands regulate key cellular processes including differentiation, proliferation, adipogenesis, and insulin sensitization [6, 7]. PPAR γ ligands have also been shown to attenuate inflammation in many tissues including skin, liver, kidney, and lung [8–11]. Several

types of natural PPAR γ ligands exist, including prostaglandins such as 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15d-PGJ₂), and fatty acids such as lysophosphatidic acid and 15S-hydroxyicosatetraenoic acid [12–15]. The thiazolidinedione (TZD) class of drugs, including rosiglitazone and pioglitazone, are synthetic PPAR γ agonists that are used as insulin sensitizers for type II diabetes [16, 17]. 2-cyano-3, 12-dioxoolean-1, 9-dien-28-oic acid (CDDO) is a novel synthetic triterpenoid PPAR γ ligand currently in clinical trials for treatment of several different forms of cancer [18, 19].

Rosiglitazone, CDDO and 15d-PGJ₂ all tightly bind to PPAR γ [12, 20, 21], activate PPAR γ -dependent transcription [22, 23], and regulate adipogenesis via a PPAR γ -dependent mechanism [12, 24, 25]. However, CDDO and 15d-PGJ₂ also have effects which can be mediated through PPAR γ -independent pathways [26–28]. 15d-PGJ₂ and CDDO contain α/β -unsaturated ketone rings with electrophilic carbons susceptible to forming covalent bonds with cellular proteins through Michaelis addition reactions [18, 29, 30]. Rosiglitazone and other synthetic TZDs lack these electrophilic centers. We have recently demonstrated the importance of these electrophilic carbons in preventing TGF- β -induced myofibroblast differentiation [26]. Understanding the molecular pathways modulated by PPAR γ ligands will shed light on their potential therapeutic contribution(s) in the control of pulmonary inflammation.

In addition to their structural role, fibroblasts in the lung act as sentinel cells with significant effector roles in orchestrating and amplifying inflammatory cascades. They become activated when exposed to inflammatory stimuli and produce inflammatory mediators such as IL-6, monocyte chemoattractant protein-1 (MCP-1), cyclooxygenase-2 (COX-2), and PGE₂ [31–34]. We hypothesized that PPAR γ ligands would exhibit anti-inflammatory effects in human lung fibroblasts, and tested this hypothesis using IL-1 β , a potent proinflammatory cytokine, and silica, an inhaled particulate with strong pro-inflammatory effects on lung fibroblasts [32, 33]. Here, we report for the first time that PPAR γ ligands inhibit the inflammatory response of human lung fibroblasts, and do so via a largely PPAR γ -independent pathway dependent on a strong electrophilic center.

2. Material and Methods

2.1. Cells and Cell Culture. Primary human lung fibroblasts were derived from tissue explants obtained from patients undergoing surgical resection for benign hamartoma. This is an abnormal but noncancerous growth within the lung, it is not an inflammatory or fibrotic disease. The tissue pieces used to obtain the fibroblasts were taken from a region of the resected tissue that was most distal to the hamartoma that was anatomically and histologically normal [35]. Patient samples were obtained with approval of the Institutional Review Board of the University of Rochester. These cells are morphologically consistent with fibroblasts [36]. They express collagen and vimentin, and they do not express CD45, factor VIII, or cytokeratin. Fibroblasts were cultured


in minimum essential media (MEM, Life Technologies, Gaithersburg, Md, USA) supplemented with 10% fetal bovine serum (FBS, Sigma Aldrich, St. Louis, Mo, USA), 2 mM L-glutamine, penicillin (100 units/mL), streptomycin (100 μ g/mL), and amphotericin (0.25 μ g/mL) (Life Technologies) at 37°C in 7% CO₂, as previously described [26]. Cells were used at passages 6–12.

2.2. Reagents. PPAR γ agonists and related compounds rosiglitazone, 9,10-dihydro-15-deoxy- $\Delta^{12,14}$ -PGJ₂ (CAY10410), GW9662, and prostaglandin-A₁ (PGA₁) were from Cayman Chemical (Ann Arbor, MI). 2-cyano-3, 12-dioxoolean-1, 9-dien-28-oic acid (CDDO) was obtained from the NIH-RAID Program and Reata Pharmaceuticals (Dallas, Tex, USA) and 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15d-PGJ₂) was from Biomol (Plymouth Meeting, PA). These compounds were prepared as 10 mM stocks in DMSO and added to cell cultures to the final concentrations indicated. Control wells (media only) were supplemented with DMSO to the same final concentration (0.2%) as treated wells. One hour after pretreatment with PPAR γ ligands, inflammatory stimulants interleukin-1 β (IL-1 β , R&D Systems, Minneapolis, Minn, USA) and amorphous silica (a generous gift of Dr. David R. Hemenway, University of Vermont) were added to the cell cultures at the final concentrations indicated for 24 hours. Amorphous silica was prepared by baking for 2 hours at 180°C prior to addition to MEM.

2.3. Cytotoxicity Assays. Cell viability was assessed by 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) assay [37]. Fibroblasts were plated in triplicate at a density of 5,000 cells per well in 96 well plates and treated with TGF- β and PPAR γ agonists for 24 h at the indicated concentrations. MTT was added for the final 4 hours. Production of the colored reaction product was measured at 560 nm, and the results were normalized to the negative control wells. LDH activity in culture medium was measured by a commercial assay (Sigma).

2.4. Prostaglandin and Cytokine Assays. Primary human lung fibroblasts (100,000 cells/well) were plated in six-well plates (Falcon/Becton Dickson, Franklin Lakes, Nj, USA), serum starved for 48 hours, and treated with IL-1 β or silica and/or PPAR γ agonists as described. PGE₂ was measured in harvested supernatants using a commercially available competitive enzyme immunoassay (EIA) (Cayman Chemical) [38]. IL-6 and CCL2/MCP-1 in harvested supernatants were determined by ELISA according to the manufacturer's instructions (R&D Systems, Minneapolis, Minn, USA).

2.5. Western Blots for Cyclooxygenase-2 (COX-2). Total cellular protein extracts were prepared from lung fibroblast cultures with 10% Nonidet P-40 (NP-40) lysis buffer supplemented with a protease inhibitor cocktail (Sigma). Lysates were clarified by centrifugation and proteins were quantitated by the bicinchoninic acid (Pierce, Rockford, IL). Typically, 5 μ g of total solubilized cellular protein was separated by 10% SDS-PAGE under reducing conditions



EC₅₀ (μM)

| | IL-6 | MCP-1 |
|----------------------|------|-------|
| 15d-PGJ ₂ | 3.5 | 1 |
| CDDO | 0.6 | 0.1 |
| Rosi | >20 | 5 |

(e)

FIGURE 1: PPAR γ ligands attenuate IL-6 and MCP-1 production by human lung fibroblasts induced by IL-1 β or crystalline silica. Primary human lung fibroblasts were pretreated with the indicated concentrations of rosiglitazone (Rosi), CDDO or 15d-PGJ₂ (PGJ₂) for 1 hour and then cotreated with 1 ng/mL IL-1 β (a, b) or 10 μ g/mL crystalline silica (c, d) for 24 hours. Supernatants were harvested and cytokine concentrations were determined by ELISA. Results are mean \pm standard error for quadruplicate wells and are representative of 2 independent experiments that yielded similar results. (e) The EC₅₀ was determined from the given data and averaged for each cytokine. Data were analyzed by one-way ANOVA. * = $P < .05$ compared to stimulus alone (MEM).

and transferred to nitrocellulose membranes. Blots were probed with a COX-2 specific monoclonal antibody (Cayman Chemical) using GAPDH (Abcam, Cambridge, Mass, USA) as a loading control. Proteins were visualized with

enhanced chemiluminescence (Western Lightning, Perkin-Elmer, Wellesley, MA) [38] and densitometry of the resulting bands was performed using Kodak Molecular Imaging Software (Rochester, NY) normalized to the loading control.




FIGURE 2: The PPAR γ ligands are not overtly cytotoxic at the doses used. (a) Primary human lung fibroblasts were treated for 24 hours with the indicated concentrations of CDDO, or 15d-PGJ₂ (PGJ₂), and LDH released into the media was measured by commercial LDH activity assay. There were no significant differences between any of the treatment groups compared to MEM control. Results are mean \pm standard deviation for triplicate wells and are representative of 2 independent experiments that yielded similar results. (b) Primary human lung fibroblasts were plated in a 96-well plate and treated with the indicated concentrations of CDDO or 15d-PGJ₂ (PGJ₂). Cell viability was determined after 24 hours by MTT assay. The results shown are the mean \pm standard deviation of quadruplicate wells and are normalized to untreated control wells.

2.6. COX-2 Immunofluorescence. Fibroblasts were cultured in eight-well chamber slides (5×10^4 cells/well) in serum-free MEM for 48 hours before treatment. Some cells were pretreated with PPAR γ agonists for one hour prior to IL-1 β (1 ng/mL) treatment for the indicated duration. Cells maintained in serum-free MEM were used as negative controls. Cells were washed in PBS and fixed and permeabilized for 15 minutes at room temperature in BD Cytotfix/Cytoperm solution (BD Biosciences, San Diego, Calif, USA). The cells were then washed with BD Perm/Wash buffer (BD Biosciences) in this step and all future washes to maintain




FIGURE 3: CDDO and 15d-PGJ₂ attenuate IL-1 β -induced COX-2 production in human lung fibroblasts. Primary human lung fibroblasts were pretreated with 20 μ M rosiglitazone (Rosi), 1 μ M CDDO, or 5 μ M 15d-PGJ₂ (PGJ₂) for 1 hour and then cotreated with 1 ng/mL of IL-1 β for 24 hours. Protein lysates were harvested and Western blot analysis was performed by probing for protein expression of COX-2 and GAPDH (for normalization). (a) Representative samples are shown. (b) Quadruplicate samples were analyzed by densitometry and normalized to GAPDH. COX-2 expression was significantly reduced in CDDO and 15d-PGJ₂ treated cells compared to IL-1 β alone (* $P < .001$). CDDO and 15d-PGJ₂ were significantly more potent than rosiglitazone ($\dagger P < .001$). Results are mean \pm standard deviation for quadruplicate wells and are representative of 3 independent experiments that yielded similar results. Data were analyzed by ANOVA.

permeabilization. Nonspecific sites were blocked with 5% normal goat serum (Life Technologies) in BD Perm/Wash buffer for 15 minutes at room temperature or 4 $^{\circ}$ C overnight. Monoclonal mouse anti-human COX-2 antibody was diluted in 1% normal goat serum in BD Perm/Wash buffer and incubated with the cells overnight at 4 $^{\circ}$ C. Cells were washed and Alexa Fluor 488 (green-) tagged goat anti-mouse IgG (Invitrogen) diluted in 1% normal goat serum in BD Perm/Wash buffer was added to the cells for one hour at room temperature. Cells were washed a coverslipped with ProLong Antifade with DAPI (Invitrogen).

2.7. NF- κ B Luciferase Assay. We generated human lung fibroblast strains that stably expressed a mammalian codon-optimized firefly luciferase reporter gene under the transcriptional control of NF- κ B response elements using NF- κ B Cignal Lenti Reporter lentiviral particles (SABiosciences Corporation, Frederick, Md, USA) at an MOI = 25. Two days post-infection, growth medium was removed and replaced with growth medium containing 1 μ g/mL puromycin.




FIGURE 4: Immunofluorescence of human lung fibroblasts demonstrates CDDO and 15d-PGJ₂ attenuate IL-1 β -induced COX-2 expression. Fibroblasts were pretreated with PPAR γ agonists for 1 hour prior to IL-1 β (1 ng/mL) treatment for 24 hours. Cells were fixed and permeabilized and probed for COX-2 protein. COX-2 was visualized with Alexa Fluor 488 (green) and cell nuclei with DAPI (blue). (a) MEM control; (b) IL-1 β ; (c) IL-1 β + 20 μ M rosiglitazone; (d) IL-1 β + 1 μ M CDDO; (e) IL-1 β + 5 μ M 15d-PGJ₂; (f) IL-1 β + 5 μ M CAY10410. IL-1 β -induced COX-2 was inhibited by CDDO (d) and 15d-PGJ₂ (e).

Puromycin-resistant clones were identified and expanded for propagation. To investigate NF- κ B activity, primary human lung fibroblasts expressing NF- κ B-Luc (3,000 cells/well) were plated in 96-well plates, serum starved for 48 hours and treated with IL-1 β and/or PPAR γ agonists as described.

Cells were lysed in 1x Passive Lysis Buffer and mixed with Luciferase Assay Reagent II as instructed by the manufacturer (Promega, Madison, Wis, USA). Equal protein quantities were used in luciferase assays; results were reported in relative light units.




FIGURE 5: PPAR γ ligands attenuate IL-1 β -induced PGE₂ production in human lung fibroblasts. Primary human lung fibroblasts were pretreated with 20 μ M rosiglitazone (Rosi), 1 μ M CDDO, or 5 μ M 15d-PGJ₂ (PGJ₂) and then cotreated with IL-1 β for 24 hours as previously described. Supernatants were harvested, and PGE₂ was determined by EIA. PGE₂ production is significantly reduced in PPAR γ ligand-treated fibroblasts compared to IL-1 β alone (* P < .001). CDDO and 15d-PGJ₂ were significantly more potent than rosiglitazone ($\dagger P$ < .05). Results are mean \pm standard deviation for quadruplicate wells and are representative of 3 independent experiments that yielded similar results. Data were analyzed by ANOVA.

2.8. *Statistics.* All data are expressed as mean \pm SD. A one-way analysis of variance (ANOVA) with Tukey post-test were used to establish statistical significance. Results were considered significant if P < .05.

3. Results

3.1. *PPAR γ Ligands Inhibit IL-1 β -Induced Inflammatory Cytokine Production in Human Lung Fibroblasts.* To determine the efficacy of selected PPAR γ ligands in inhibiting production of inflammatory mediators in lung fibroblasts, primary human lung fibroblasts were pretreated with rosiglitazone, CDDO, or 15d-PGJ₂ for 1 hour and then cotreated with a powerful pro-inflammatory stimulus, IL-1 β (1 ng/mL), for 24 hours. IL-1 β strongly induced the pro-inflammatory mediators IL-6 and MCP-1 (Figure 1). Rosiglitazone, CDDO, and 15d-PGJ₂ all show dose-dependent inhibition of cytokine production and significantly inhibited the release of these mediators. We also investigated an alternative inflammatory stimulus, crystalline silica, which we have previously reported is a potent pro-inflammatory stimulus in human lung fibroblasts [32]. Silica also strongly induced the production of IL-6 and MCP-1, which was inhibited by the PPAR γ ligands in a dose-dependent manner (Figures 1(c) and 1(d)). Interestingly, all 3 ligands were 4-5-fold more effective at blocking MCP-1 than IL-6 (Figure 1(e)).

It should be noted that the maximum dose used for each compound is the highest dose that does not cause overt cytotoxicity (Figure 2 and data not shown). Rosiglitazone is

at least 10-fold less effective than CDDO or 15d-PGJ₂ and is a very poor inhibitor of IL-6 release even at the maximum tolerable dose of 20 μ M.

3.2. *PPAR γ Ligands Inhibit IL-1 β -Induced Upregulation of COX-2 and PGE₂.* To further evaluate the potential anti-inflammatory actions of PPAR γ ligands, lung fibroblasts were treated with IL-1 β and PPAR γ ligands, and expression of COX-2 was determined by Western blot. As expected, IL-1 β strongly induces COX-2 (Figure 3). CDDO and 15d-PGJ₂ significantly inhibited IL-1 β -induced COX-2 production. However, rosiglitazone failed to suppress COX-2 induction in human lung fibroblasts.

We also performed immunofluorescence to confirm the induction of COX-2 protein and to localize the expression of COX-2 in IL-1 β -treated lung fibroblasts. COX-2-specific staining demonstrated that untreated fibroblasts expressed minimal COX-2 at baseline (Figure 4(a)). Upon treatment with IL-1 β , cytoplasmic COX-2 production was markedly upregulated (Figure 4(b)). Rosiglitazone failed to suppress IL-1 β -induced COX-2 upregulation (Figure 4(c)). In contrast, however, both CDDO and 15d-PGJ₂ suppressed cytoplasmic expression of COX-2 in IL-1 β -treated fibroblasts (Figures 4(d) and 4(e)). CAY10410, a structural analogue of 15d-PGJ₂, did not suppress IL-1 β -induced COX-2 upregulation (Figure 4(f)), suggesting structural differences between the ligands account for their anti-inflammatory actions.

COX-2 mediates the first step in the conversion of arachidonic acid to prostaglandins. The immunomodulatory prostaglandin PGE₂, a product of this reaction, was measured in lung fibroblast culture supernatants following treatment with PPAR γ ligands and IL-1 β . Consistent with the COX-2 results, CDDO and 15d-PGJ₂ inhibited IL-1 β -induced production of PGE₂ by greater than 90% compared to controls treated with IL-1 β alone (Figure 5). Rosiglitazone also inhibited IL-1 β -induced production of PGE₂, but was less effective than CDDO and 15d-PGJ₂.

3.3. *Suppression of Inflammatory Mediators by PPAR γ Ligands in Human Lung Fibroblasts Occurs via a PPAR γ -Independent Mechanism.* We used a pharmacological approach to determine whether the anti-inflammatory actions of PPAR γ ligands are dependent on or independent of PPAR γ . GW9662 is an irreversible PPAR γ antagonist that covalently binds to a cysteine residue in the ligand binding site of PPAR γ [39]. GW9662 inhibits PPAR γ agonist-driven adipogenesis, which is a completely PPAR γ -dependent process [24]. Primary human lung fibroblasts were pretreated for 4 hours with GW9662 and one hour with CDDO or 15d-PGJ₂, followed by IL-1 β . IL-1 β strongly induced IL-6, MCP-1 and PGE₂ compared to MEM control (Figures 6(a)–6(c)). As previously shown, CDDO and 15d-PGJ₂ significantly inhibited the IL-1 β -induced production of these inflammatory mediators. GW9662 did not reverse the suppressive effects of CDDO and 15d-PGJ₂ ligands on cytokine and PGE₂ production (Figures 6(a)–6(c)). This indicates that PPAR γ is not essential for the anti-inflammatory effects of these ligands, and that PPAR γ independent pathways are therefore likely important.




FIGURE 6: The PPAR γ antagonist GW9662 does not inhibit the anti-inflammatory effects of CDDO and 15d-PGJ₂. Primary human lung fibroblasts were pretreated with 1 μ M GW9662 for 3 hours, then PPAR γ ligands were added for 1 hour, then 1 ng/mL IL-1 β was added for 24 hours. Supernatants were harvested and proinflammatory mediators were measured by ELISA or EIA. (a) IL-6, (b) MCP-1, (c) PGE₂. Pretreatment with GW9662 did not significantly alter the attenuation of pro-inflammatory mediator production by CDDO or 15d-PGJ₂ alone. Results are mean \pm standard deviation for quadruplicate wells and are representative of 2 independent experiments that yielded similar results.

3.4. A Strong Electrophilic Center Is Important for PPAR γ Ligand-Mediated Suppression of Inflammation in Human Lung Fibroblasts. CDDO and 15d-PGJ₂ contain α/β -unsaturated ketone rings with electrophilic carbons that can form covalent bonds with free sulfhydryls in cellular proteins [40, 41]. CAY10410 (9,10-dihydro-15-deoxy- $\Delta^{12,14}$ -PGJ₂) is a structural analog of 15d-PGJ₂ that lacks the unsaturated ketone containing the electrophilic carbon. To investigate the importance of the electrophilic center in suppressing inflammatory endpoints, we compared the ability of 15d-PGJ₂ and CAY10410 to inhibit the pro-inflammatory effects of IL-1 β on human lung fibroblasts. CAY10410 treatment resulted in a small reduction in IL-1 β -induced IL-6 production that was not statistically significant, and a 60% reduction in MCP-1 production compared to 98% inhibition by 15d-PGJ₂ (Figure 7).

To further investigate the importance of the electrophilic center, we tested another prostaglandin that is also a potent electrophile, PGA₁. PGA₁ was partially effective at inhibiting IL-1 β -induced production of IL-6 and completely effective in blocking MCP-1 production (Figure 7).

3.5. A Strong Electrophilic Center Is Important for PPAR γ Ligand-Mediated Suppression of NF- κ B in Human Lung Fibroblasts. To better understand the mechanism involved in PPAR γ ligand-mediated immune suppression, we investigated the effect of PPAR γ ligands on the activation of NF- κ B, a transcription factor that regulates the expression of numerous pro-inflammatory mediators. Primary human lung fibroblasts were transfected with an NF- κ B luciferase reporter construct, and treated with PPAR γ ligands and IL-1 β . CDDO, 15d-PGJ₂, and PGA₁, but not CAY10410 or rosiglitazone, significantly decreased IL-1 β -induced NF- κ B luciferase activity (Figure 8).

4. Discussion

Inflammation is associated with many diseases of the lung including asthma, chronic obstructive pulmonary disease (COPD), silicosis, and lung cancer [1–4]. Aside from glucocorticoids, there are few effective anti-inflammatory therapies; therefore, the development of novel therapies that




FIGURE 7: A strong electrophilic carbon is necessary for PPAR γ ligand-mediated attenuation of inflammation in IL-1 β -treated human lung fibroblasts. Primary human lung fibroblasts were pretreated with 1 μ M CDDO, 5 μ M 15d-PGJ₂ (PGJ₂), 5 μ M CAY10410, (CAY) or 15 μ M PGA₁ for 1 hour and then co-treated with 1 ng/mL of IL-1 β for 24 hours. Supernatants were harvested, and cytokine concentrations were measured by ELISA. (a) CDDO, 15d-PGJ₂, and PGA₁, but not CAY10410, significantly inhibited production of IL-6 ($*P < .01$). (b) CDDO, 15d-PGJ₂, PGA₁, and CAY10410 all significantly reduced MCP-1 production ($*P < .01$). CDDO, 15d-PGJ₂, and PGA₁ were significantly more potent than CAY10410 ($\dagger P < .01$). Results are mean \pm standard deviation for quadruplicate wells and are representative of 3 independent experiments.

have the potential to alleviate pulmonary diseases associated with inflammatory etiologies is a priority.

PPAR γ ligands are receiving increasing attention as potential anti-inflammatory therapeutics because of their anti-inflammatory properties in a variety of tissues in vivo and cells in vitro [42]. The anti-inflammatory effects of PPAR γ ligands have not previously been reported in human lung fibroblasts, a sentinel cell of inflammatory cascades in the lung [31, 34, 43, 44]. Here, we report that PPAR γ ligands have potent anti-inflammatory effects in human lung fibroblasts exposed to divergent inflammatory stimuli, and that the mechanism is largely PPAR γ -independent.

To induce a pro-inflammatory response in human lung fibroblasts, we used two different inflammatory stimuli. IL-1 β is an acute phase inflammatory cytokine, while silica is a particulate that has potent proinflammatory effects when inhaled and is capable of causing both acute and chronic inflammatory lung disease [32, 33]. Both IL-1 β and




FIGURE 8: CDDO and PGA₁ inhibit IL-1 β -induced NF- κ B transcriptional activity in human lung fibroblasts. Primary human lung fibroblasts were transfected with a luciferase reporter, then pretreated with 20 μ M rosiglitazone (Rosi), 1 μ M CDDO, 5 μ M 15d-PGJ₂ (PGJ₂), 5 μ M CAY10410 (CAY), or 15 μ M PGA₁ and cotreated with of IL-1 β for 24 hours as described. NF- κ B-dependent luciferase activity was measured in lysates as described, normalized to protein concentration, and expressed as relative light units (RLU). NF- κ B activation is significantly reduced in PPAR γ ligand-treated fibroblasts compared to IL-1 β alone ($*P < .01$). Results are mean \pm standard deviation for quadruplicate wells and are representative of 2 independent experiments that yielded similar results.

silica induced the inflammatory mediators IL-6 and MCP-1, which were inhibited by CDDO, rosiglitazone, and 15d-PGJ₂ (Figure 1). Interestingly, rosiglitazone was much less effective at inhibiting IL-6 and MCP-1, with an EC₅₀ 5–10-fold higher than 15d-PGJ₂ and at least 30-fold higher than CDDO. CDDO and 15d-PGJ₂, but not rosiglitazone, also blocked upregulation of COX-2 and PGE₂ (Figures 3 and 4). This is in agreement with our previous finding that rosiglitazone is less effective than CDDO or 15d-PGJ₂ at inhibiting the pro-fibrotic effects of TGF- β in lung fibroblasts [26], and suggests that there are significant differences in the mechanism of action between rosiglitazone and CDDO and 15d-PGJ₂.

Rosiglitazone, CDDO and 15d-PGJ₂ all tightly bind the PPAR γ receptor [12, 20, 21], activate PPAR γ -dependent transcription [22, 23], and promote adipogenesis via a solely PPAR γ -dependent mechanism [12, 25]. However, in addition to stimulating PPAR γ -dependent transcriptional changes, CDDO and 15d-PGJ₂ are reported to have effects that are mediated through PPAR γ -independent pathways [26, 28, 45]. To determine whether CDDO, and 15d-PGJ₂ might be acting via a PPAR γ -independent mechanism, we used a pharmacological approach to block PPAR γ . GW9662 is an irreversible competitive PPAR γ antagonist that covalently binds to a cysteine residue in the ligand binding

domain of PPAR γ [46]. GW9662 is a highly effective inhibitor of PPAR γ -dependent processes including differentiation of osteoclasts and activation of hepatic stellate cells [47, 48]. We have previously reported that rosiglitazone, CDDO and 15d-PGJ₂ drive the differentiation of fibroblasts to adipocytes. GW9662 at 1 μ M completely inhibits this effect, demonstrating that this compound is effective at blocking the PPAR γ -dependent actions of these PPAR γ ligands [24]. Here, GW9662 did not reverse the anti-inflammatory effects of CDDO and 15d-PGJ₂ (Figure 6), indicating that the anti-inflammatory effects of CDDO and 15d-PGJ₂ on human lung fibroblasts are largely independent of the PPAR γ -dependent transcriptional pathway. Rosiglitazone was such a poor inhibitor of the inflammatory effects of IL-1 β that it was not possible to show a reversal of inhibition by GW9662, which would be expected if rosiglitazone acted by a purely PPAR γ -dependent mechanism.

Comparing the chemical structures of rosiglitazone, CDDO, and 15d-PGJ₂, it is notable that CDDO and 15d-PGJ₂ have strong electrophilic carbons, whereas rosiglitazone does not. 15d-PGJ₂ has one α/β -unsaturated ketone ring with an electrophilic carbon capable of forming covalent bonds through Michael addition reactions [49], whereas CDDO has two [18, 30]. We have recently demonstrated the importance of these electrophilic carbons in preventing TGF- β -induced myofibroblast differentiation [26, 50]. We hypothesize that the electrophilic carbons of CDDO and 15d-PGJ₂ are also important for their anti-inflammatory effects. To test this hypothesis, we used CAY10410, a structural analog of 15d-PGJ₂ that lacks the α/β -unsaturated ketone, and PGA₁, another electrophilic prostaglandin. In lung fibroblasts stimulated with IL-1 β , CAY10410 did not inhibit COX-2 upregulation or IL-6 production and was half as effective as 15d-PGJ₂ at blocking MCP-1 production (Figures 4 and 7). On the other hand, PGA₁ significantly attenuated IL-6 and completely blocked production of MCP-1 (Figure 7). Because CAY10410 has an identical structure to 15d-PGJ₂ except for the electrophilic carbon, the fact that CAY10410 lacks the effects of 15d-PGJ₂ strongly suggests that the electrophilic centers present in CDDO and 15d-PGJ₂ are critical for mediating their maximal anti-inflammatory therapeutic potential. CDDO and 15d-PGJ₂, but not rosiglitazone or CAY10410, significantly inhibited IL-1 β -induced NF- κ B activity (Figure 8).

The molecular targets of CDDO and 15d-PGJ₂ in inflammation are not completely known. 15d-PGJ₂ can bind to the NF- κ B components I κ B and p65 [50]. Another candidate is the transcription factor Nrf2, which regulates anti-oxidant and anti-inflammatory pathways. CDDO and 15d-PGJ₂ activate Nrf2 in mouse cells and human cancer cells [51, 52]. However, these compounds do not activate Nrf2 in human lung fibroblasts [27, 53]. We have previously reported that CDDO activates AP-1 transcriptional activity in human lung fibroblasts [27]. However, AP-1 is a promoter, rather than an inhibitor of inflammation, and AP-1 activation leads to upregulation of IL-6 via NF- κ B [54]. We hypothesize that these electrophilic compounds suppress inflammation and activate AP-1 via different pathways, and that the anti-

inflammatory effects are stronger and override the potentially proinflammatory effects of AP-1 activation.

In addition to PPAR γ -independent effects, PPAR γ ligands have anti-inflammatory effects that are moderated via a PPAR γ -dependent mechanism. This PPAR γ -dependent mechanism can be accessed by TZDs such as rosiglitazone and pioglitazone [9, 55–57], and indeed, rosiglitazone has limited anti-inflammatory properties in this report. However, while TZDs are currently used clinically as insulin sensitizers in type 2 diabetes, they have a complex sideeffect profile including edema, weight gain, bone weakness, and potentially an increased risk of cardiovascular disease [58–60], that may limit their widespread use as anti-inflammatory therapies. Although TZDs have high binding affinity for PPAR γ they lack electrophilic centers and are thus unable to access PPAR γ -independent anti-inflammatory pathways that use this mechanism [27, 49, 61, 62]. We suggest that additional research on the PPAR γ -independent anti-inflammatory activities of CDDO and 15d-PGJ₂, including identification of additional targets beyond NF- κ B, should lead to development of novel compounds with greater specificity for the anti-inflammatory targets of PPAR γ ligands but decreased binding of PPAR γ itself, with fewer resulting side-effects. As CDDO is orally active, has a long half-life, and is currently in clinical trials as an anticancer therapy, it may be a useful platform for derivatization and further study. Further development of small compounds with strong electrophilic centers is warranted as these drugs may be effective anti-inflammatory treatments for human lung diseases.

Acknowledgments

This work was supported in part by the National Institutes of Health Clinical and Translational Sciences Award and Grants nos. HL75432-0451, HL75432-0452, DE011390, ES01247, HL095402, EY017123, EY015836, T32-HL066988, T32-HL007152, and ES07026. C. M. Hagon was supported by Award no. 5TL1RR024135 from the National Center for Research Resources. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Research Resources or National Institutes of Health.

References

- [1] P. J. Barnes, "Future advances in COPD therapy," *Respiration*, vol. 68, no. 5, pp. 441–448, 2001.
- [2] D. H. Bowden and I. Y. R. Adamson, "The role of cell injury and the continuing inflammatory response in the generation of silicotic pulmonary fibrosis," *Journal of Pathology*, vol. 144, no. 3, pp. 149–161, 1984.
- [3] J. S. Brody and A. Spira, "Chronic obstructive pulmonary disease, inflammation, and lung cancer," *Proceedings of the American Thoracic Society*, vol. 3, no. 6, pp. 535–537, 2006.
- [4] E. A. Engels, "Inflammation in the development of lung cancer: epidemiological evidence," *Expert Review of Anticancer Therapy*, vol. 8, no. 4, pp. 605–615, 2008.
- [5] I. Issemann, R. A. Prince, J. D. Tugwood, and S. Green, "The peroxisome proliferator-activated receptor:retinoid X receptor heterodimer is activated by fatty acids and fibrate

- hypolipidaemic drugs," *Journal of Molecular Endocrinology*, vol. 11, no. 1, pp. 37–47, 1993.
- [6] A. Cabrero, J. C. Laguna, and M. Vázquez, "Peroxisome proliferator-activated receptors and the control of inflammation," *Curr Drug Targets Inflamm Allergy*, vol. 1, no. 3, pp. 243–248, 2002.
 - [7] N. S. Tan, L. Michalik, B. Desvergne, and W. Wahli, "Multiple expression control mechanisms of peroxisome proliferator-activated receptors and their target genes," *Journal of Steroid Biochemistry and Molecular Biology*, vol. 93, no. 2–5, pp. 99–105, 2005.
 - [8] M. G. Belvisi, D. J. Hele, and M. A. Birrell, "Peroxisome proliferator-activated receptor gamma agonists as therapy for chronic airway inflammation," *European Journal of Pharmacology*, vol. 533, no. 1–3, pp. 101–109, 2006.
 - [9] R. Kapadia, J. H. Yi, and R. Vemuganti, "Mechanisms of anti-inflammatory and neuroprotective actions of PPAR-gamma agonists," *Frontiers in Bioscience*, vol. 13, no. 5, pp. 1813–1826, 2008.
 - [10] M. Lehrke and M. A. Lazar, "The many faces of PPAR γ ," *Cell*, vol. 123, no. 6, pp. 993–999, 2005.
 - [11] G. Rizzo and S. Fiorucci, "PPARs and other nuclear receptors in inflammation," *Current Opinion in Pharmacology*, vol. 6, no. 4, pp. 421–427, 2006.
 - [12] S. A. Kliewer, J. M. Lenhard, T. M. Willson, I. Patel, D. C. Morris, and J. M. Lehmann, "A prostaglandin J₂ metabolite binds peroxisome proliferator-activated receptor γ and promotes adipocyte differentiation," *Cell*, vol. 83, no. 5, pp. 813–819, 1995.
 - [13] T. M. McIntyre, A. V. Pontsler, A. R. Silva et al., "Identification of an intracellular receptor for lysophosphatidic acid (LPA): LPA is a transcellular PPAR γ agonist," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 1, pp. 131–136, 2003.
 - [14] F. J. Schopfer, Y. Lin, P. R. S. Baker et al., "Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor γ ligand," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 7, pp. 2340–2345, 2005.
 - [15] S. B. Shappell, R. A. Gupta, S. Manning et al., "15S-hydroxyeicosatetraenoic acid activates peroxisome proliferator-activated receptor γ and inhibits proliferation in PC3 prostate carcinoma cells," *Cancer Research*, vol. 61, no. 2, pp. 497–503, 2001.
 - [16] J. W. F. Elte and J. F. Blicklé, "Thiazolidinediones for the treatment of type 2 diabetes," *European Journal of Internal Medicine*, vol. 18, no. 1, pp. 18–25, 2007.
 - [17] A. Zieleniak, M. Wójcik, and L. A. Woźniak, "Structure and physiological functions of the human peroxisome proliferator-activated receptor γ ," *Archivum Immunologiae et Therapiae Experimentalis*, vol. 56, no. 5, pp. 331–345, 2008.
 - [18] S. Chintharlapalli, S. Papineni, M. Konopleva, M. Andreef, I. Samudio, and S. Safe, "2-Cyano-3,12-dioxoolean-1,9-dien-28-oic acid and related compounds inhibit growth of colon cancer cells through peroxisome proliferator-activated receptor γ -dependent and γ -independent pathways," *Molecular Pharmacology*, vol. 68, no. 1, pp. 119–128, 2005.
 - [19] D. M. Ray, K. M. Morse, S. P. Hilchey et al., "The novel triterpenoid 2-cyano-3,12-dioxooleana-1,9-dien-28-oic acid (CDDO) induces apoptosis of human diffuse large B-cell lymphoma cells through a peroxisome proliferator-activated receptor gamma-independent pathway," *Experimental Hematology*, vol. 34, no. 9, pp. 1202–1211, 2006.
 - [20] J. M. Lehmann, L. B. Moore, T. A. Smith-Oliver, W. O. Wilkison, T. M. Willson, and S. A. Kliewer, "An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor γ (PPAR γ)," *The Journal of Biological Chemistry*, vol. 270, no. 22, pp. 12953–12956, 1995.
 - [21] Y. Wang, W. W. Porter, N. Suh et al., "A synthetic triterpenoid, 2-cyano-3,12-dioxooleana-1,9-dien-28-oic acid (CDDO), is a ligand for the peroxisome proliferator-activated receptor γ ," *Molecular Endocrinology*, vol. 14, no. 10, pp. 1550–1556, 2000.
 - [22] H. A. Burgess, L. E. Daugherty, T. H. Thatcher et al., "PPAR γ agonists inhibit TGF- β induced pulmonary myofibroblast differentiation and collagen production: implications for therapy of lung fibrosis," *American Journal of Physiology*, vol. 288, no. 6, pp. L1146–L1153, 2005.
 - [23] A. A. Kulkarni, T. H. Thatcher, K. C. Olsen, S. B. Maggior, R. P. Phipps, and P. J. Sime, "PPAR- γ ligands repress TGF β -induced myofibroblast differentiation by targeting the PI3K/Akt pathway: implications for therapy of fibrosis," *PLoS ONE*, vol. 6, no. 1, article e15909, 2011.
 - [24] S. E. Feldon, C. W. O'Loughlin, D. M. Ray, S. Landskroner-Eiger, K. E. Seweryniak, and R. P. Phipps, "Activated human T lymphocytes express cyclooxygenase-2 and produce proadipogenic prostaglandins that drive human orbital fibroblast differentiation to adipocytes," *American Journal of Pathology*, vol. 169, no. 4, pp. 1183–1193, 2006.
 - [25] B. M. Forman, P. Tontonoz, J. Chen, R. P. Brun, B. M. Spiegelman, and R. M. Evans, "15-deoxy- Δ , -prostaglandin J₂ is a ligand for the adipocyte determination factor PPAR γ ," *Cell*, vol. 83, no. 5, pp. 803–812, 1995.
 - [26] H. E. Ferguson, A. Kulkarni, G. M. Lehmann et al., "Electrophilic peroxisome proliferator-activated receptor- γ ligands have potent antifibrotic effects in human lung fibroblasts," *American Journal of Respiratory Cell and Molecular Biology*, vol. 41, no. 6, pp. 722–730, 2009.
 - [27] H. E. Ferguson, T. H. Thatcher, K. C. Olsen et al., "Peroxisome proliferator-activated receptor- γ ligands induce heme oxygenase-1 in lung fibroblasts by a PPAR γ -independent, glutathione-dependent mechanism," *American Journal of Physiology*, vol. 297, no. 5, pp. L912–L919, 2009.
 - [28] B. Melichar, M. Konopleva, W. Hu, K. Melicharova, M. Andreeff, and R. S. Freedman, "Growth-inhibitory effect of a novel synthetic triterpenoid, 2-cyano-3,12-dioxoolean-1,9-dien-28-oic acid, on ovarian carcinoma cell lines not dependent on peroxisome proliferator-activated receptor- γ expression," *Gynecologic Oncology*, vol. 93, no. 1, pp. 149–154, 2004.
 - [29] B. Shi and M. F. Greaney, "Reversible Michael addition of thiols as a new tool for dynamic combinatorial chemistry," *Chemical Communications*, no. 7, pp. 886–888, 2005.
 - [30] N. Suh, Y. Wang, T. Honda et al., "A novel synthetic oleanane triterpenoid, 2-cyano-3,12-dioxoolean-1,9-dien-28-oic acid, with potent differentiating, antiproliferative, and anti-inflammatory activity," *Cancer Research*, vol. 59, no. 2, pp. 336–341, 1999.
 - [31] M. Jordana, B. Sarnstrand, P. J. Sime, and I. Ramis, "Immune-inflammatory functions of fibroblasts," *European Respiratory Journal*, vol. 7, no. 12, pp. 2212–2222, 1994.
 - [32] K. M. A. O'Reilly, R. P. Phipps, T. H. Thatcher, B. A. Graf, J. Van Kirk, and P. J. Sime, "Crystalline and amorphous silica differentially regulate the cyclooxygenase-prostaglandin pathway in pulmonary fibroblasts: Implications for pulmonary fibrosis," *American Journal of Physiology*, vol. 288, no. 6, pp. L1010–L1016, 2005.
 - [33] M. A. Olman, K. E. White, L. B. Ware et al., "Pulmonary edema fluid from patients with early lung injury stimulates fibroblast

- proliferation through IL-1 β -induced IL-6 expression,” *Journal of Immunology*, vol. 172, no. 4, pp. 2668–2677, 2004.
- [34] R. S. Smith, T. J. Smith, T. M. Blieden, and R. P. Phipps, “Fibroblasts as sentinel cells. Synthesis of chemokines and regulation of inflammation,” *The American Journal of Pathology*, vol. 151, no. 2, pp. 317–322, 1997.
- [35] K. M. Fries, G. D. Sempowski, A. A. Gaspari, T. Blieden, R. J. Looney, and R. P. Phipps, “CD40 expression by human fibroblasts,” *Clinical Immunology and Immunopathology*, vol. 77, no. 1, pp. 42–51, 1995.
- [36] C. J. Baglolle, S. Y. Reddy, S. J. Pollock et al., “Isolation and phenotypic characterization of lung fibroblasts,” *Methods in Molecular Medicine*, vol. 117, pp. 115–127, 2005.
- [37] T. Mosmann, “Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays,” *Journal of Immunological Methods*, vol. 65, no. 1-2, pp. 55–63, 1983.
- [38] L. Koumas, T. J. Smith, and R. P. Phipps, “Fibroblast subsets in the human orbit: thy-1 and Thy-1 subpopulations exhibit distinct phenotypes,” *European Journal of Immunology*, vol. 32, no. 2, pp. 477–485, 2002.
- [39] S. Giri, R. Rattan, A. K. Singh, and I. Singh, “The 15-deoxy- δ 12,14-prostaglandin J₂ inhibits the inflammatory response in primary rat astrocytes via down-regulating multiple steps in phosphatidylinositol 3-kinase-akt-NF- κ B-p300 pathway independent of peroxisome proliferator-activated receptor γ ,” *Journal of Immunology*, vol. 173, no. 8, pp. 5196–5208, 2004.
- [40] J. Atsmon, B. J. Sweetman, S. W. Baertschi, T. M. Harris, and L. J. Roberts, “Formation of thiol conjugates of 9-deoxy-delta 9,delta 12(E)-prostaglandin D2 and delta 12(E)-prostaglandin D2,” *Biochemistry*, vol. 29, pp. 3760–3765, 1990.
- [41] T. Shibata, T. Yamada, T. Ishii et al., “Thioredoxin as a molecular target of cyclopentenone prostaglandins,” *The Journal of Biological Chemistry*, vol. 278, no. 28, pp. 26046–26054, 2003.
- [42] K. Asada, S. Sasaki, T. Suda, K. Chida, and H. Nakamura, “Antiinflammatory roles of peroxisome proliferator-activated receptor γ in human alveolar macrophages,” *American Journal of Respiratory and Critical Care Medicine*, vol. 169, no. 2, pp. 195–200, 2004.
- [43] J. Kaufman, B. A. Graf, E. C. Leung et al., “Fibroblasts as sentinel cells: role of the CD40-CD40 ligand system in fibroblast activation and lung inflammation and fibrosis,” *Chest*, vol. 120, no. 1, supplement, pp. S53–S55, 2001.
- [44] L. Koumas, A. E. King, H. O. D. Critchley, R. W. Kelly, and R. P. Phipps, “Fibroblast heterogeneity: existence of functionally distinct Thy 1⁺ and Thy 1⁻ human female reproductive tract fibroblasts,” *American Journal of Pathology*, vol. 159, no. 3, pp. 925–935, 2001.
- [45] R. A. Derlacz, K. Hyc, M. Usarek, A. K. Jagielski, J. Drozak, and R. Jarzyna, “PPAR- γ -independent inhibitory effect of rosiglitazone on glucose synthesis in primary cultured rabbit kidney-cortex tubules,” *Biochemistry and Cell Biology*, vol. 86, no. 5, pp. 396–404, 2008.
- [46] L. M. Leesnitzer, D. J. Parks, R. K. Bledsoe et al., “Functional consequences of cysteine modification in the ligand binding sites of peroxisome proliferator activated receptors by GW9662,” *Biochemistry*, vol. 41, no. 21, pp. 6640–6650, 2002.
- [47] A. C. Bendixen, N. K. Shevde, K. M. Dienger, T. M. Willson, C. D. Funk, and J. W. Pike, “IL-4 inhibits osteoclast formation through a direct action on osteoclast precursors via peroxisome proliferator-activated receptor γ 1,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 5, pp. 2443–2448, 2001.
- [48] T. Miyahara, L. Schrum, R. Rippe et al., “Peroxisome proliferator-activated receptors and hepatic stellate cell activation,” *The Journal of Biological Chemistry*, vol. 275, no. 46, pp. 35715–35722, 2000.
- [49] J. U. Scher and M. H. Pillinger, “The anti-inflammatory effects of prostaglandins,” *Journal of Investigative Medicine*, vol. 57, no. 6, pp. 703–708, 2009.
- [50] D. S. Straus, G. Pascual, M. Li et al., “15-Deoxy- Δ -prostaglandin J₂ inhibits multiple steps in the NF- κ B signaling pathway,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 9, pp. 4844–4849, 2000.
- [51] H. R. Kim, S. Kim, E. J. Kim et al., “Suppression of Nrf2-driven heme oxygenase-1 enhances the chemosensitivity of lung cancer A549 cells toward cisplatin,” *Lung Cancer*, vol. 60, no. 1, pp. 47–56, 2008.
- [52] T. E. Sussan, T. Rangasamy, D. J. Blake et al., “Targeting Nrf2 with the triterpenoid CDDO-imidazolide attenuates cigarette smoke-induced emphysema and cardiac dysfunction in mice,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 106, no. 1, pp. 250–255, 2009.
- [53] C. J. Baglolle, P. J. Sime, and R. P. Phipps, “Cigarette smoke-induced expression of heme oxygenase-1 in human lung fibroblasts is regulated by intracellular glutathione,” *American Journal of Physiology*, vol. 295, no. 4, pp. L624–L636, 2008.
- [54] C. Viedt, J. Vogel, T. Athanasiou et al., “Monocyte chemoattractant protein-1 induces proliferation and interleukin-6 production in human smooth muscle cells by differential activation of nuclear factor- κ B and activator protein-1,” *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 22, no. 6, pp. 914–920, 2002.
- [55] A. Jacob, R. Wu, M. Zhou, and P. Wang, “Mechanism of the anti-inflammatory effect of curcumin: PPAR- γ activation,” *PPAR Research*, Article ID 89369, 2007.
- [56] A. Nakajima, K. Wada, H. Miki et al., “Endogenous PPAR γ mediates anti-inflammatory activity in murine ischemia-reperfusion injury,” *Gastroenterology*, vol. 120, no. 2, pp. 460–469, 2001.
- [57] M. D. Rollins, S. Sudarshan, M. A. Firpo et al., “Anti-inflammatory effects of PPAR- γ agonists directly correlate with PPAR- γ expression during acute pancreatitis,” *Journal of Gastrointestinal Surgery*, vol. 10, no. 8, pp. 1120–1130, 2006.
- [58] P. S. Chaggar, S. M. Shaw, and S. G. Williams, “Review article: thiazolidinediones and heart failure,” *Diabetes & Vascular Disease Research*, vol. 6, no. 3, pp. 146–152, 2009.
- [59] C. V. Rizos, M. S. Elisaf, D. P. Mikhailidis, and E. N. Liberopoulos, “How safe is the use of thiazolidinediones in clinical practice?” *Expert Opinion on Drug Safety*, vol. 8, no. 1, pp. 15–32, 2009.
- [60] P. Shah and S. Mudaliar, “Pioglitazone: side effect and safety profile,” *Expert Opinion on Drug Safety*, vol. 9, no. 2, pp. 347–354, 2010.
- [61] C. Giaginis, A. Giagini, and S. Theocharis, “Peroxisome proliferator-activated receptor- γ (PPAR- γ) ligands as potential therapeutic agents to treat arthritis,” *Pharmacological Research*, vol. 60, no. 3, pp. 160–169, 2009.
- [62] W. K. Jung, I. S. Park, S. J. Park et al., “The 15-Deoxy- Δ -prostaglandin J₂ inhibits LPS-stimulated AKT and NF- κ B activation and suppresses interleukin-6 in osteoblast-like cells MC3T3E-1,” *Life Sciences*, vol. 85, no. 1-2, pp. 46–53, 2009.