

Resting Dead Space Fraction as Related to Clinical Characteristics, Lung Function, and Gas Exchange in Male Patients with Chronic Obstructive Pulmonary Disease

This article was published in the following Dove Press journal:
International Journal of General Medicine

Ming-Lung Chuang ^{1,2}
Benjamin Yung-
Thing Hsieh ³
I-Feng Lin ⁴

¹Division of Pulmonary Medicine and Department of Internal Medicine, Chung Shan Medical University Hospital, Taichung, Taiwan, 40201, Republic of China; ²School of Medicine, Chung Shan Medical University, Taichung, Taiwan, 40201, Republic of China; ³School of Medicine, National Yang Ming University, Taipei, Taiwan, 11221, Republic of China; ⁴Institute of Public Health, National Yang Ming University, Taipei, Taiwan, 11221, Republic of China

Correspondence: Ming-Lung Chuang
Division of Pulmonary Medicine and
Department of Internal Medicine, Chung
Shan Medical University Hospital, #110,
Section 1, Chien-Kuo North Road, South
District, Taichung, 40201, Taiwan,
Republic of China
Tel +886-4-2473-9595 ext. 34718
Email yuan1007@ms36.hinet.net

I-Feng Lin
Institute of Public Health, National Yang
Ming University, 155, Sec 2, LihNong
Street, ShiPai, Taipei, Taiwan, 11221,
Republic of China
Email iflin@ym.edu.tw

Background: Measures of forced expired volume in one second % predicted (FEV₁%), residual volume to total lung capacity ratio (RV/TLC) and diffusing capacity for carbon monoxide measurements (D_LCO) are the standard lung function test for evaluating patients with chronic obstructive pulmonary disease (COPD). The dead space fraction (V_D/V_T) has been shown to be a robust marker of gas exchange abnormality. However, the use of V_D/V_T has gradually become less common. As V_D/V_T measured at rest (V_D/V_TR) has been successfully used in non-COPD conditions, it was hypothesized that in COPD the V_D/V_TR was more sensitive than the standard lung function test in correlation with clinical characteristics and gas exchange. This study aimed to test the hypothesis and to identify the variables relevant to V_D/V_TR.

Methods: A total of 46 male subjects with COPD were enrolled. Clinical characteristics included demographic data, oxygen-cost diagram (OCD), and image studies for pulmonary hypertension. The standard lung function was obtained. To calculate V_D/V_T, invasive arterial blood gas and pulmonary gas exchange (PGX) were measured. The variables relevant to V_D/V_TR were analyzed by multiple linear regression.

Results: Compared to lung function, V_D/V_TR was more frequently and significantly related to smoking, carboxyhemoglobin level, pulmonary hypertension and P_aCO₂ (all p <0.05) whereas FEV₁% was more related to lung function test, P_aO₂ and OCD score. V_D/V_TR and FEV₁% were highly related to resting gas exchange but RV/TLC and D_LCO% were not. Cigarette consumption, the equivalent for CO₂ output, arterial oxyhemoglobin saturation, and the product of tidal volume and inspiratory duty cycle were identified as the parameters relevant to V_D/V_TR with a power of 0.72.

Conclusion: Compared to lung function test, V_D/V_TR is more related to clinical characteristics and is a comprehensive marker of resting gas exchange. Further studies are warranted to provide a noninvasive measurement of V_D/V_TR.

Registration Number: MOST 106-2314-B-040-025 and CSH-2019-C-30.

Keywords: obstructive airway disease, residual volume and total lung capacity ratio, diffusing capacity of lung, dead space and tidal volume ratio, cigarette consumption, carboxyhemoglobin, pulmonary hypertension

Background

The severity of chronic obstructive pulmonary disease (COPD) is usually graded by forced expired volume in one second % predicted (FEV₁%).¹ Despite mMRC and acute exacerbation being related to the risk assessment of COPD, the power of

correlations of FEV₁% with the clinical outcomes is not robust,²⁻⁴ and lung volume such as the residual volume to total lung capacity ratio (RV/TLC) and diffusing capacity measurements (D_LCO) provide additional information.⁵⁻⁸

The dead space to tidal volume ratio at rest or at peak exercise or change during exercise ($V_D/V_{T,R}$, -P, and -C) indicating the severity of ventilation/perfusion (V/Q) mismatch⁹ can serve as a marker of physiological change in patients with COPD in a various clinical conditions.^{8,10-15} However, exercise testing is labor consuming and costly and in diseases other than COPD, the V_D/V_T has been successfully used in non-exercise conditions.¹⁶⁻²⁰

It was hypothesized that the $V_D/V_{T,R}$ was more significant than the standard lung function test in correlation with clinical characteristics and gas exchange (ie, all variables relevant to oxygen and CO₂ tensions and their differences in arterial blood and alveoli) in patients with COPD. This study aimed to test the hypothesis by correlation analysis and to identify variables of demographics, symptom scores, images for pulmonary hypertension, lung function, and gas exchange relevant to $V_D/V_{T,R}$ by multiple regression analysis.

Methods

Study Design

In this observational cross-sectional study, we measured demographic characteristics, cigarette smoking history, functional activity and pulmonary hypertension determined by image studies, and lung function in subjects with COPD at a university teaching hospital. Resting pulmonary gas exchange (PGX) and arterial blood gas (ABG) were measured simultaneously. The local Institutional Review Board of Chung Shan Medical University Hospital (CS16174) approved this study. This study was conducted in compliance with the Declaration of Helsinki.

Subjects

Subjects aged ≥ 40 years with COPD were enrolled. Anthropometric measurements, cigarette consumption, and functional activity were recorded. Male adult subjects who underwent lung function testing were enrolled if their forced expired volume in one second (FEV₁)/forced vital capacity (FVC) was < 0.7 .¹ The diagnosis of COPD was made according to the Global Initiative for Chronic Lung Disease (GOLD) criteria.¹ As few female subjects met the

criteria of COPD in Taiwan (4%),² for simplicity they were not included in this study. Subjects with a body mass index $\leq 18 \text{ kg}\cdot\text{m}^{-2}$ or $\geq 32 \text{ kg}\cdot\text{m}^{-2}$, any other chronic diseases including uncontrolled diabetes mellitus, uncontrolled hypertension, anemia (hemoglobin $< 13 \text{ g}\cdot\text{dL}^{-1}$ in males), or with laboratory findings of cardiovascular, hematological, metabolic or neuromuscular diseases or acute illnesses in the recent 1 month were not enrolled. Uncontrolled hypertension means having hypertension but no or inadequate treatment so that the blood pressure remained high (ie, $> 140/90 \text{ mm Hg}$).²¹ Signed informed consent was obtained from each participant. Some of the participants were enrolled in other studies.^{22,23}

Measurements

Demographic Data and Functional Activity

Age, height, weight, body mass index, triceps skinfold, mid-arm circumference, and cigarette consumption were recorded. The oxygen cost diagram (OCD) was used to evaluate the participants' functional activity. The OCD a 100-mm long vertical line with everyday activities listed alongside the line, above which breathlessness limited the participants. The distance from zero was measured and scored.

Image Study

Chest radiography was obtained within one month of enrolling in the study. The hila thoracic ratio (HTR), cardiac thoracic ratio, and the diameter of anterior descending pulmonary artery (ADPA) on the standing posterior-anterior chest radiograph were measured as these variables were reported to represent pulmonary hypertension.²⁴ The chest radiographs were evaluated by an experienced pulmonologist without knowing the clinical information and the average values were recorded for analysis.

Two-dimensional echocardiography was performed by an experienced cardiologist who was unaware of the clinical data and lung function. Parasternal, apical, and sub-costal studies were conducted.^{25,26}

Pulmonary Function Testing

Pre-test preparation was followed the standard guide and bronchodilators were not administered within 3 h for short-acting beta-agonists and 12 h for long-acting beta-agonists before the tests.²⁷⁻²⁹ FEV₁, FVC, TLC, RV, and D_LCO were measured using spirometry, body plethysmography, and the single-breath technique (MasterScreenTM Body; Carefusion, Wuerzburg, Germany), respectively, in accordance with the currently recommended standards by

ATS/ERS task force.^{30,31} The best of three technically satisfactory readings was used.^{30,32,33} All of the spirometry data were obtained before and after inhaling a standard dose of fenoterol HCl. Post-dose measurements were performed 15 minutes after inhalation. Static lung volume data and D_LCO data were obtained before inhaling fenoterol.

Maximum inspiratory pressure at the mouth indicating inspiratory muscle strength and maximum expiratory pressure indicating expiratory muscle strength at the mouth were measured.³⁴

Pulmonary Gas Exchange and V_D/V_T Measurement

Each subject completed PGX measured at a 2-min complete rest. VO_2 (mL/min), CO_2 output (VCO_2) (mL/min), and minute ventilation (V_E) were continuously measured and the data at the last 15 seconds were averaged and reported (MasterScreen CPX™, Carefusion, Wuerzburg, Germany). The physiological V_D/V_T was measured (V_D/V_T) using a standard formula as follows:³⁵

$$V_D/V_T = (P_aCO_2 - P_ECO_2)/P_aCO_2 - V_{Dm}/V_T \quad (1)$$

where P_aCO_2 was arterial PCO_2 and $P_ECO_2 = VCO_2/V_E \times (P_B - 47\text{mmHg})$ and P_B was barometric pressure measured daily and V_{Dm} was the dead space of mouth piece and pneumotachograph as the manufacture reported.³⁶ Artery blood samples were drawn at the brachial artery and heparinized at rest. The sample was immediately placed on ice and then analyzed for pH, PCO_2 , and PO_2 with body temperature correction (model 278, CIBA-Corning, Medfield, MA, USA).

Statistical Analysis

Data were summarized as mean \pm standard deviation. Pearson's or Spearman correlation coefficients were used when appropriate for quantifying the pair-wise relationships among the variables of interest. Multiple linear regression analysis was used to select important parameters of V_D/V_T . All possible regression algorithms were performed using the candidate variables with p values < 0.35 in univariate analysis. The Student's t -test was used for comparisons between two groups. The chi-square test or Fisher exact test was used to compare the proportions between the two groups. All statistical analyses were performed using SAS statistical software (SAS Institute Inc., Cary, NC, USA). Statistical significance was set at $p < 0.05$. Marginal significance was set at $0.05 \leq p \leq 0.1$.

Results

Fifty-seven patients were enrolled, of whom 10 were excluded because two subjects did not meet the inclusion criteria and another two met the exclusion criteria and six subjects declined to participate (Figure 1). The remaining 46 male subjects with COPD were enrolled after excluding one subject whose arterial blood gas analysis was not obtained (mean age 65.2 ± 5.8 years) (Tables 1–3). In Tables 1–3, some data were missing in one to six subjects because of technical failure or no measurements. Most of the patients had GOLD stages II and III, hyperinflation of lung and static air trapping, high V_D/V_{TR} , and impaired diffusing capacity, inspiratory muscle strength and peak flow, and hyperventilation with normoxemia and normocapnia.

Compared to the standard lung function test, V_D/V_T R was more frequently (Table 4, $p < 0.0001$) and significantly related to cigarette consumption, carboxyhemoglobin level, and pulmonary hypertension (all $p < 0.05$) and these associated factors were not related to $FEV_1\%$, RV/TLC , and $D_LCO\%$ (all $p = \text{NS}$). In contrast, $FEV_1\%$ and RV/TLC were more related to lung function as expected and $D_LCO\%$ was singly and mildly related to RV/TLC . V_D/V_T R and $FEV_1\%$ were correlated with the resting gas exchange whereas RV/TLC and $D_LCO\%$ were not.

Multiple linear regression analysis revealed that cigarette consumption and HTR were positively related to V_D/V_T R whereas the height was negatively related to V_D/V_T R (Table 5). When PGX data were added in analysis, the HTR and height were replaced by VE/VCO_2 , the product of V_T and inspiratory duty cycle (V_TIDC), and S_pO_2 where VE/VCO_2 was positively related to V_D/V_{TR} whereas V_TIDC and S_pO_2 were negatively related to V_D/V_{TR} . Cigarette consumption and the three PGX variables were highly related to V_D/V_{TR} with a power of 0.72.

Discussion

In this study, compared to $FEV_1\%$, RV/TLC , and $D_LCO\%$, V_D/V_{TR} was a unique marker for male patients with COPD (Table 4). V_D/V_{TR} was related to cigarette consumption, carboxyhemoglobin, and pulmonary hypertension, whereas the standard lung function was not. Using statistical technique for evaluation of V_D/V_{TR} , cigarette consumption, VE/VCO_2 , V_TIDC and S_pO_2 were the four contributors (Table 5, $r^2 = 0.72$).

Regarding lung function, $D_LCO\%$, KCO , and V_A/TLC have been related to V_D/V_{TR} and V_D/V_{TP} in the literature




Figure 1 Flow diagram. A total of 57 subjects with chronic obstructive pulmonary disease were screened. After excluding 10 subjects, the remaining 46 were retained for analysis after further excluding another one subject whose arterial blood gas data was not obtained.

($r = -0.20$ – -0.60 , $p < 0.05$ - <0.01) where KCO is D_LCO divided by alveolar volume (V_A).⁷ The V_A and V_A/TLC were related to lung volumes and airway obstruction, respectively (all $p < 0.05$ - <0.01),⁷ suggesting that D_LCO , KCO, and V_A are related to poor communication of airways “that would be indicative of high ventilation-perfusion areas (~alveolar dead space)”.⁷ Hence, it was expected that RV/TLC was related to V_D/V_{TR} and thus measuring V_D/V_{TR} is redundant. However, in this study, V_D/V_{TR} was not correlated with the lung function variables suggesting that V_D/V_T and lung hyperinflation, air trapping or diffusing capability might be different components in physiology.²² V_D/V_{TR} was correlated with cigarette consumption, carboxyhemoglobin, and pulmonary hypertension measured by chest radiography (Table 4). Thus, V_D/V_{TR} may offer additional information to lung function test, whereas $FEV_1\%$, RV/TLC , and $D_LCO\%$ were more related to lung function test (Table 4).

Tobacco smoking is the key preventable risk factor for the development of COPD and the pathogenesis of emphysema. The duration of cigarette consumption associated with the development of COPD is considered to be ≥ 20 pack-years.^{1,37} Emphysema with a tissue-to-airway progression pattern³⁷ was related to an increased V_D/V_T .¹⁵ Although cigarette consumption is thought to contribute to an increased V_D/V_T as found in this study, their relationship may not be necessary through the mechanism of emphysema. This was noted that RV/TLC was not related to V_D/V_{TR} in this study. Further studies are warranted to elucidate the

relationship between cigarette consumption, emphysema, and V_D/V_T . Furthermore, carboxyhemoglobin level is modestly related to cigarette smoking ($r = 0.28$, $p = 0.06$) and was mildly correlated with V_D/V_{TR} ($r = 0.35$, $p = 0.02$) in the current study. It was consistent with a previous report that carboxyhemoglobin was positively correlated with VE/VCO_2 and rapid breathing and inversely with $VO_{2max}\%$ and expandable volume excursion.³⁸ Of note, carboxyhemoglobin level and V_D/V_{TR} were not in a cause–effect relationship and that carboxyhemoglobin was not correlated with $D_LCO\%$ in the current study does not exclude that carboxyhemoglobin can affect D_LCO in normal subjects. The findings indicate that the lung pathology of COPD plays a more important role than carboxyhemoglobin in relation to D_LCO .

Regarding chest radiography, a $HTR \geq 0.36$ and diameter of the right ADPA ≥ 1.8 cm have been reported to be markers of pulmonary hypertension in COPD.²⁴ COPD is usually complicated with pulmonary hypertension in the later stages. Although there was discordance between radiographic and hemodynamic measures of pulmonary arterial pressure, HTR and diameter of the right ADPA were modestly correlated with increased V_D/V_{TR} in this study (Table 4, $r = 0.33$ and 0.34 , both $p < 0.05$).

In normal subjects, age and height have been positively related to V_D/V_{TR} whereas weight has been negatively related to V_D/V_{TR} , and V_D/V_{TR} has been reported to be smaller in females.³⁹ However, in this study, in subjects with COPD, height was negatively related to V_D/V_{TR} . The

Table 1 Demographic Data, Symptom Score, Blood Tests, Chest Radiography, and Echocardiography (n = 46)

	n	Mean	SD
Age, year	46	65.2	5.8
Height, cm	46	165.0	6.4
Weight, kg	46	60.4	11.2
Body mass index, kg/m ²	46	22.12	3.53
Smoke, py	46	42.3	19.2
OCD, cm	46	7.0	1.4
Triceps, mm	44	6.4	2.6
Mid-arm, cm	44	27.3	3.4
Borg, A.U.	46	0.05	0.19
Hemoglobin, g/dL	46	14.8	1.5
Albumin, g/dL	40	4.2	0.4
GPT, IU/L	40	21.1	16.8
Bilirubin, direct, mg/dL	42	0.25	0.29
Creatinine, mg/dL	42	1.09	0.20
Na ⁺ , mmol/L	41	140.5	2.4
K ⁺ , mmol/L	40	4.3	0.5
Sugar, mg/dL	41	98.8	14.7
Cholesterol, mg/dL	43	182.2	36.6
Hila-thoracic ratio	44	0.36	0.04
Cardiac-thoracic ratio	41	0.44	0.06
ADPA, cm	46	1.62	0.33
Apical 4 EDRV, cm ²	42	13.5	3.7
Apical 4 ESRV, cm ²	42	7.8	1.8
Subcostal 4 EDRV, cm ²	42	14.2	3.3
Subcostal 4 ESRV, cm ²	42	7.6	2.1
Right ventricle wall, mm	42	6.0	2.0

Abbreviations: A.U., absolute unit, ADPA: anterior descending pulmonary artery of the right lung, apical four chamber view, end-diastolic right ventricle area (EDRV), end-systolic right-ventricle area (ESRV); subcostal four-chamber view; long and short axes view: the right ventricle free wall thickness at an end-diastolic phase between the tricuspid annulus and the papillary muscle.

reason for this discrepancy is not clear, but it is probably due to the high correlation between height and weight in this study in an analysis a posteriori ($r = 0.51$, $p = 0.0003$). A higher weight is probably beneficial against the development of emphysema and stage 4 disease of COPD⁴⁰ and thus negatively related to V_D/V_T (Table 4, $r = -0.27$, $p = 0.08$), while sarcopenia or low lean body mass may predispose to the development of emphysema (multi-organ loss of tissue, MOLT phenotype).⁴¹ Another possible reason is that the lung pathology of COPD outweighs the contribution of height to V_D/V_T as normal subjects do.

VE/VCO_2 has been reported to be strongly related to V_D/V_T when the data involve healthy subjects and those with COPD at rest and during submaximal exercise.⁷ This is attributed to the Bohr-Enghoff equation in which VE/VCO_2 and V_D/V_T are mathematically related at a given level of P_aCO_2 . A high VE/VCO_2 and high P_{a-ET}

Table 2 Lung Function

Variables	n	Mean	SD
Total lung capacity, TLC, L	46	6.5	1.0
TLCpred, %	46	135	21
Functional residual capacity, FRC, L	46	4.8	1.0
FRCpred, %	46	162	34
Inspiratory capacity, L	46	1.7	0.5
Residual volume, RV, L	46	3.8	0.9
RV/TLC, %	46	58	9
D_LCO , mL/min/mmHg	45	15.8	5.6
D_LCO pred, %	45	69	22
Forced vital capacity, FVC, L	46	2.5	0.7
FVCpred, %	46	81	21
FEV ₁ , L	46	1.2	0.5
FEV ₁ pred, %	46	50	19
GOLD, I, II, III, IV, n (%)			
3 (6.5), 18 (39.1), 19 (41.3), 6 (13.0)			
FEV ₁ /FVC, %	46	49	13
Slow vital capacity, SVC, L	46	2.7	0.7
SVCpred, %	46	89	22
MIP, cmH ₂ O	43	68.4	18.6
MEP, cmH ₂ O	43	103.7	22.6
PEFR, L/sec	46	2.9	1.3
PEFRpred, %	46	42	20

Abbreviations: D_LCO , the diffusion capacity of the lungs for carbon monoxide; FEV₁, forced expiratory volume in one second; GOLD I-IV, global initiative for chronic obstructive lung disease stages; MIP, maximal inspiratory pressure; MEP, maximum expiratory pressure; PEFR, peak expiratory flow rate.

CO_2 explain 40–50% of high V_D/V_T in this study (Table 4, $r = 0.64$ and 0.67 , both $p < 0.0001$) compatible with that Lewis et al and Liu et al reported that V_D/V_T and $P_{a-ET}CO_2$ were highly correlated ($r = 0.76–0.88$, $p < 0.001$).^{42,43} Compared to FEV₁% in this study, V_D/V_T was more related to P_aCO_2 at rest ($r^2 = 0.24$ versus 0.10) whereas FEV₁% was more related to P_aO_2 and OCD.

Despite A-aDO₂ and S_pO₂ being marginally correlated with V_D/V_T (Table 4, $r = -0.28$ to -0.29 , both $p = 0.06$), S_pO₂ was selected in multiple regression analysis. However, in COPD even in subjects with mild severity A-aDO₂ was positively related to V_D/V_T .¹⁰ The discrepancy between these two studies in the V_D/V_T versus A-aDO₂ relationship might be due to poor correlation of V_D/V_T with P_aO_2 where A-aDO₂ was highly negatively related to P_aO_2 in this study ($r = -0.66$, $p < 0.0001$).

V_T IDC probably represents expandable tidal volume excursion in a timely manner. When both V_T and IDC are large, the mean inspiratory flow (V_T /inspiratory time) can be favorably generated by the force of the inspiratory muscles. On the other hand, when V_T is large and IDC is small, the mean inspiratory flow is extremely high and

Table 3 Cardiopulmonary Physiological Data, Arterial Blood Gases, Carboxyhemoglobin and Plasma Lactate Concentrations at Rest

Variables	n	Mean	SD
VO ₂ %predicted maximum, %	46	16	3
VO ₂ /kg, L/min/kg	46	4.1	0.8
VCO ₂ , L/min	46	0.20	0.04
Respiratory exchange ratio	46	0.83	0.05
O ₂ Pulse, mL/beat	46	3.1	0.7
VE/VCO ₂	46	53.5	10.0
VE/VO ₂	46	44.5	7.4
V _D /V _T	45	0.58	0.08
P _{a-ET} CO ₂ , mmHg	45	4.0	4.4
A-aDO ₂ , mmHg	45	18.9	1.0
S _p O ₂ , %	45	95.3	2.6
VE, L/min	46	10.7	2.1
MVV, L/min	46	36.3	16.9
VE/MVV	46	0.35	0.16
Systolic blood pressure, mmHg	46	145.0	18.6
Diastolic blood pressure, mmHg	45	79.3	9.0
Heart rate, beat/min	46	80.6	13.2
pH	45	7.4	0.03
P _a CO ₂ , mmHg	45	40.6	6.4
P _a O ₂ , mmHg	45	79.2	10.1
HCO ₃ ⁻ , mEq/L	45	24.7	1.9
Carboxyhemoglobin, %	45	1.3	0.9
Methemoglobin, %	43	0.25	0.08
Lactate, mmol/L	43	1.2	0.3

Abbreviations: VO₂, oxygen uptake; VCO₂, CO₂ output; Respiratory exchange ratio, CO₂ output and oxygen uptake ratio; O₂ Pulse, oxygen uptake divided by heart rate; V_D/V_T, dead space and tidal volume ratio; P_{a-ET}CO₂, arterial end-tidal CO₂ pressure gradient; A-aDO₂, alveolar arterial oxygen pressure gradient; VE, minute ventilation; MVV, maximal voluntary ventilation; S_pO₂, hemoglobin saturation measured by pulse oximetry.

would be not biologically plausible. V_TIDC is a variable firstly reported in the literature and its importance needs to be confirmed.

Study Limitations

The number of cases in this study was small; however, it may be inevitable for an invasive study. The small number of cases may have caused insufficient power when performing correlation coefficient analysis on V_D/V_T R. However, the sample size of 46 achieved a power of 80% to detect a difference between a correlation of 0.4 and the null (no correlation) using a two-sided test with a significance level of 0.05.²³ As the power is related to type II error, nonsignificant test results should be interpreted more conservatively. All of the participants in this study were male, so using the results applied to females should be cautious. To investigate the relationship

Table 4 Comparison of Correlation Coefficient (r) Between Resting Dead Space Fraction (V_D/V_TR) and Lung Function with Variables of Interest

	n	V _D /V _T R	FEV ₁ %	RV/ TLC	D _L CO %
		r	r	r	r
Clinical parameters					
Age	45	0.30*	0.04	0.04	0.02
Height	45	-0.40**	-0.04	-0.01	-0.17
Weight	45	-0.27 [¶]	0.16	-0.15	0.37**
Body mass index	45	-0.12	-0.16	-0.16	0.52[†]
Smoke, p-y	45	0.32*	-0.11	0.21	-0.01
OCD	45	0.08	0.34*	-0.14	0.03
Hemoglobin	45	0.29*	-0.16	-0.18	0.31*
Carboxyhemoglobin	45	0.42**	0.01	0.08	-0.16
Hila-thoracic ratio	44	0.33*	0.25 [¶]	-0.22	0.00
ADPA diameter	45	0.34*	-0.06	-0.11	0.13
Number of significant correlation, n		10 [#]	2	1	3
Lung function parameters					
FRC%	45	0.02	-0.23	0.48[†]	-0.24
RV/TLC	45	-0.04	-0.66[‡]	1	-0.31*
D _L CO%	45	0.05	0.25 [¶]	-0.31*	1
Slow vital capacity%	45	0.10	0.69[‡]	-0.72[‡]	0.12
Forced vital capacity%	45	0.02	0.75[‡]	-0.64[‡]	0.05
FEV ₁ %	45	-0.01	1	-0.66[‡]	0.25 [¶]
MIP	43	-0.19	0.13	-0.14	0.19
Post MIP	43	-0.28 [¶]	-0.07	-0.02	0.30 [¶]
Number of significant correlation, n		1 [!]	4	5	3
Gas exchange parameters					
P _{a-ET} CO ₂	45	0.67[‡]	-0.02	0.06	0.08
V _E /V _{CO} ₂	45	0.64[‡]	0.34*	-0.19	-0.05
P _a CO ₂	45	0.49[†]	-0.32*	0.10	0.18
P _a O ₂	45	-0.18	0.44**	-0.06	-0.08
AaDO ₂	45	-0.28 [¶]	-0.25 [¶]	0.02	-0.06
S _p O ₂	45	-0.29 [¶]	0.12	0.08	-0.17
Number of significant correlation, n		5	4	0	0

Notes: P_{a-ET}CO₂, V_E/V_{CO}₂, AaDO₂, S_pO₂. Fisher exact test was used for comparison of V_D/V_TR versus 3 lung function tests, # and ! indicating p = 0.0001 and 0.1, respectively; [¶]Indicating modest correlation with 0.05 < p < 0.1, whereas *, **, †, ‡Indicating significant correlation with p < 0.05, <0.01, <0.001, and <0.0001, respectively; bolded numerals highlighting a significant correlation.

Abbreviations: FEV₁, forced expired volume in one second %predicted; RV/TLC, residual volume to total lung capacity ratio; D_LCO%, diffusing capacity for carbon monoxide % predicted; p-y, packyear; OCD, oxygen cost diagram; ADPA, anterior descending pulmonary artery; FRC, functional residual capacity; SVC, slow vital capacity; MIP, maximal inspiratory capacity.

between two or more variables, it is usually to adjust known factors that influence the dependent variables. We used % predicted of lung function variables instead of using the absolute measured values because lung function variables are obviously influenced by anthropometrics. In this context, the unwanted influences by anthropometrics were avoided and highlight the important

Table 5 Risk Factors of Dead Space Fraction at Rest (V_D/V_{T-R})

Variable	Equation	r^2	p
$V_D/V_{T-R_{stat}}$	$1.046 (\pm 0.279) - 0.0042 \times \text{height} (\pm 0.0015) + 0.0014 \times \text{smoke} (\pm 0.0006) + 0.491 \times \text{HTR} (\pm 0.276)$	0.33	0.001
	$0.942 (\pm 0.251) + 0.0012 \times \text{smoke} (\pm 0.0004) + 0.0048 \times \text{VE}/\text{VCO}_2 (\pm 0.0007) - 0.472 \times \text{V}_T\text{IDC} (\pm 0.118) - 0.006 \times \text{S}_p\text{O}_2 (\pm 0.0027)$	0.72	< 0.0001

Notes: $V_D/V_{T-R_{stat}}$ is the V_D/V_{T-R} obtained using multiple linear regression, smoke refers to cigarette consumption in pack-years, HTR is the hila-thoracic ratio in decimals measured from chest radiography, with the height in centimeters, and the numbers in parenthesis are standard errors. VE/VCO_2 is the ventilatory equivalent for CO_2 output, V_TIDC is the product of tidal volume in liters, inspiratory duty cycle (IDC) in decimals, and S_pO_2 is the oxyhemoglobin saturation measured with pulse oximetry. IDC is the ratio of inspiratory time of an entire respiratory cycle.

factors. This issue has been strongly recommended in a previous report that TLC, FRC, IC, IRV, ERV, RV, FVC, SVC, and FEV1 in liters and DLCO in mL/min/mmHg were omitted from correlation analysis.²⁷ As transcutaneous PCO_2 ($P_{tc}\text{CO}_2$) is a noninvasive measurement, it warrants further study to compare the study with that using $P_{tc}\text{CO}_2$.^{7,44} Although using radioisotopes or applying the multiple inert gas elimination technique (MIGET) is the standard of measuring V/Q mismatch, these techniques are costly and the latter is not common in clinical practice whereas V_D/V_{T-R} measurement is practical and high V_D/V_{T-R} is consistent with high V/Q mismatch.⁹

Clinical Implication

Compared to lung function test, V_D/V_{T-R} was more related to cigarette consumption and pulmonary hypertension and was a comprehensive variable of resting PGX. Whether or not it can serve as a prognosticator or can be applied in smoking cessation strategy warrants further studies.

Conclusion

Compared to $\text{FEV}_1\%$, RV/TLC , and $\text{D}_L\text{CO}\%$, V_D/V_{T-R} is more related to smoking, carboxyhemoglobin, pulmonary hypertension, and P_aCO_2 whereas $\text{FEV}_1\%$ is more related to P_aO_2 and oxygen-cost diagram score and thus should be considered when evaluating patients with COPD. Further studies are warranted to provide a noninvasive measurement of V_D/V_{T-R} .

Author Contributions

MLC: initiated and designed the study, analyzed and interpreted the data, wrote and revised the manuscript, and approved the version to be published in *Inter J Gen Med*. and take responsibility and is accountable for the contents of the article. BYTH: analyzed and interpreted the data, critically reviewed the manuscript and approved the version to be published in *Inter J Gen Med*. and take

responsibility and is accountable for the contents of the article. IFL: analyzed and interpreted the data and revised and critically reviewed the manuscript and approved the version to be published in *Inter J Gen Med*. and take responsibility and is accountable for the contents of the article.

Funding

Chung Shan Medical University Hospital (CS16174).

Disclosure

The authors declare that there are no competing interests.

References

- GOLD Committees. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. disclosure forms for GOLD Committees are posted on the GOLD Website, www.goldcopd.org. 2020
- Huang TH, Hsiue TR, Lin SH, Liao XM, Su PL, Chen CZ. Comparison of different staging methods for COPD in predicting outcomes. *Eur Respir J*. 2018;51(3):pii:1700577. doi:10.1183/13993003.00577-2017
- Jones PW. Health status measurement in chronic obstructive pulmonary disease. *Thorax*. 2001;56(11):880–887. doi:10.1136/thorax.56.11.880
- Oga T, Nishimura K, Tsukino M, Sato S, Hajiro T. Analysis of the factors related to mortality in chronic obstructive pulmonary disease: role of exercise capacity and health status. *Am J Respir Crit Care Med*. 2003;167(4):544–549. doi:10.1164/rccm.200206-583OC
- Casanova C, Cote C, de Torres JP, et al. Inspiratory-to-total lung capacity ratio predicts mortality in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2005;171(6):591–597. doi:10.1164/rccm.200407-867OC
- Martinez FJ, Foster G, Curtis JL, et al. Predictors of mortality in patients with emphysema and severe airflow obstruction. *Am J Respir Crit Care Med*. 2006;173(12):1326–1334. doi:10.1164/rccm.200510-1677OC
- Mahut B, Chevalier-Bidaud B, Plantier L, et al. Diffusing capacity for carbon monoxide is linked to ventilatory demand in patients with chronic obstructive pulmonary disease. *COPD*. 2012;9(1):16–21. doi:10.3109/15412555.2011.630700
- Sue DY, Oren A, Hansen JE, Wasserman K. Diffusing capacity for carbon monoxide as a predictor of gas exchange during exercise. *N Engl J Med*. 1987;316(21):1301–1306. doi:10.1056/NEJM198705213162103
- Wasserman K, Hansen JE, Sue DY, Stringer WW, Whipp BJ. Physiology of exercise. In: Wasserman K, editor. *Principles of Exercise Testing and Interpretation*. Philadelphia: Lippincott Williams & Wilkins; 2005:10–65.

10. Elbehairy AF, Ciavaglia CE, Webb KA, et al. Pulmonary gas exchange abnormalities in mild chronic obstructive pulmonary disease. implications for dyspnea and exercise intolerance. *Am J Respir Crit Care Med*. 2015;191(12):1384–1394. doi:10.1164/rccm.201501-0157OC
11. Benditt JO, Lewis S, Wood DE, Klima L, Albert RK. Lung volume reduction surgery improves maximal O₂ consumption, maximal minute ventilation, O₂ pulse, and dead space-to-tidal volume ratio during leg cycle ergometry. *Am J Respir Crit Care Med*. 1997;156(2 Pt 1):561–566. doi:10.1164/ajrccm.156.2.9611032
12. Elbehairy AF, Webb KA, Laveneziana P, et al. Acute bronchodilator therapy does not reduce wasted ventilation during exercise in COPD. *Respir Physiol Neurobiol*. 2018;252–253:64–71. doi:10.1016/j.resp.2018.03.012
13. Rocha A, Arbex FF, Sperandio PA, et al. Excess ventilation in chronic obstructive pulmonary disease-heart failure overlap. implications for dyspnea and exercise intolerance. *Am J Respir Crit Care Med*. 2018;196(10):1264–1274. doi:10.1164/rccm.201704-0675OC
14. Wasserman K, Hansen JE, Sue DY, Casaburi R, Whipp BJ. Measurements during integrative cardiopulmonary exercise testing. In: Wasserman K, editor. *Principles of Exercise Testing and Interpretation*. Philadelphia: Lea & Febiger; 1994:53–78.
15. Yamasawa W, Tasaka S, Betsuyaku T, Yamaguchi K. Correlation of a decline in aerobic capacity with development of emphysema in patients with chronic obstructive pulmonary disease: a prospective observational study. *PLoS One*. 2015;10(4):e0125053. doi:10.1371/journal.pone.0125053
16. Beitler JR, Thompson BT, Matthay MA, et al. Estimating dead-space fraction for secondary analyses of acute respiratory distress syndrome clinical trials. *Crit Care Med*. 2015;43(5):1026–1035. doi:10.1097/CCM.0000000000000921
17. González-Castro A, Suárez-Lopez V, Gómez-Marcos V, et al. Utility of the dead space fraction (V_d/V_t) as a predictor of extubation success. *Med Intensiva*. 2011;35(9):529–538. doi:10.1016/j.medint.2011.05.016
18. Meyer K, Görndt L, Schwaibold M, et al. Predictors of response to exercise training in severe chronic congestive heart failure. *Am J Cardiol*. 1997;80(1):56–60. doi:10.1016/S0002-9149(97)00283-X
19. Rodger MA, Bredeson CN, Jones G, et al. The bedside investigation of pulmonary embolism diagnosis study: a double-blind randomized controlled trial comparing combinations of 3 bedside tests vs ventilation-perfusion scan for the initial investigation of suspected pulmonary embolism. *Arch Intern Med*. 2006;166(2):181–187. doi:10.1001/archinte.166.2.181
20. Sinha P, Calfee CS, Beitler JR, et al. Physiologic analysis and clinical performance of the ventilatory ratio in acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2019;199(3):333–341. doi:10.1164/rccm.201804-0692OC
21. Gebremichael GB, Berhe KK, Zemichael TM. Uncontrolled hypertension and associated factors among adult hypertensive patients in Ayder comprehensive specialized hospital, Tigray, Ethiopia, 2018. *BMC Cardiovasc Disord*. 2019;19(1):121. doi:10.1186/s12872-019-1091-6
22. Chuang ML. Combining dynamic hyperinflation with dead space volume during maximal exercise in patients with chronic obstructive pulmonary disease. *J Clin Med*. 2020;9(10):1127. doi:10.3390/jcm9041127
23. Chuang ML. Mechanisms affecting exercise ventilatory inefficiency-airflow obstruction relationship in male patients with chronic obstructive pulmonary disease. *Respir Res*. 2020;21(1):206. doi:10.1186/s12931-020-01463-4
24. Chetty KG, Brown SE, Light RW. Identification of pulmonary hypertension in chronic obstructive pulmonary disease from routine chest radiographs. *Am Rev Respir Dis*. 1982;126(2):338–341. doi:10.1164/arrd.1982.126.2.338
25. Bertoli L, Mantero A, Cicero SL, Alpago R, Rizzato G, Belli C. Usefulness of two-dimensional echocardiography in the assessment of right heart in chronic obstructive lung disease. *Progress Respiration Res Basel: Karger*. 1985;91–100.
26. Danchin N, Cornette A, Henriquez A, et al. Two-dimensional echocardiographic assessment of the right ventricle in patients with chronic obstructive lung disease. *Chest*. 1987;92(2):229–233. doi:10.1378/chest.92.2.229
27. Chuang ML, Lin IF. Investigating the relationships among lung function variables in chronic obstructive pulmonary disease in men. *Peer J*. 2019;7:e7829. doi:10.7717/peerj.7829
28. Chuang ML, Lin IF, Hsieh MJ. More impaired dynamic ventilatory muscle oxygenation in congestive heart failure than in chronic obstructive pulmonary disease. *J Clin Med*. 2019;8(10):E1641. doi:10.3390/jcm8101641
29. Chuang ML, Lin IF, Wasserman K. The body weight-walking distance product as related to lung function, anaerobic threshold and peak VO₂ in COPD patients. *Respir Med*. 2001;95(7):618–626. doi:10.1053/rmed.2001.1115
30. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir J*. 2005;26(2):319–338. doi:10.1183/09031936.05.00034805
31. Wanger J, Clausen JL, Coates A, et al. Standardisation of the measurement of lung volumes. *Eur Respir J*. 2005;26(3):511–522.
32. ATS/ERS S. ATS/ERS Statement on respiratory muscle testing. *Am J Respir Crit Care Med*. 2002;166(4):518–624.
33. Miller MR, Crapo R, Hankinson J, et al. General considerations for lung function testing. *Eur Respir J*. 2005;26(1):153–161. doi:10.1183/09031936.05.00034505
34. Chuang ML, Lin IF. Clinical characteristics and lung function in chronic obstructive pulmonary disease complicated with impaired peripheral oxygenation. *Intern Emerg Med*. 2014;9(6):633–640. doi:10.1007/s11739-013-0989-8
35. Wasserman K, Hansen JE, Sue DY, Stringer WW, Whipp BJ. Calculations, formulas, and examples. In: Wasserman K, editor. *Principles of Exercise Testing and Interpretation*. Philadelphia: Lippincott Williams & Wilkins; 2005:556–565.
36. Sun XG, Hansen JE, Garatachea N, Storer TW, Wasserman K. Ventilatory efficiency during exercise in healthy subjects. *Am J Respir Crit Care Med*. 2002;166(11):1443–1448. doi:10.1164/rccm.2202033
37. Young AL, Bragman FJS, Rangelov B, et al. Disease progression modelling in chronic obstructive pulmonary disease (COPD). *Am J Respir Crit Care Med*. 2020;201(3):294–302. doi:10.1164/rccm.201908-1600OC
38. Kitahara Y, Hattori N, Yokoyama A, et al. Cigarette smoking decreases dynamic inspiratory capacity during maximal exercise in patients with type 2 diabetes. *Hiroshima J Med Sci*. 2012;61(2):29–36.
39. Gläser S, Ittermann T, Koch B, et al. Influence of smoking and obesity on alveolar-arterial gas pressure differences and dead space ventilation at rest and peak exercise in healthy men and women. *Respir Med*. 2013;107(6):919–926. doi:10.1016/j.rmed.2013.02.013
40. Pinto-Plata VM, Celli-Cruz RA, Vassaux C, et al. Differences in cardiopulmonary exercise test results by american thoracic society/european respiratory society-global initiative for chronic obstructive lung disease stage categories and gender. *Chest*. 2007;132(4):1204–1211. doi:10.1378/chest.07-0593
41. Celli BR, Locantore N, Tal-Singer R, et al. Emphysema and extrapulmonary tissue loss in COPD: a multi-organ loss of tissue phenotype. *Eur Respir J*. 2018;51(2):pii: 1702146. doi:10.1183/13993003.02146-2017
42. Lewis DA, Sietsema KE, Casaburi R, Sue DY. Inaccuracy of non-invasive estimates of VD/VT in clinical exercise testing. *Chest*. 1994;106(5):1476–1480. doi:10.1378/chest.106.5.1476

43. Liu Z, Vargas F, Stansbury D, Sasse SA, Light RW. Comparison of the end-tidal arterial PCO₂ gradient during exercise in normal subjects and in patients with severe COPD. *Chest*. 1995;107(5):1218–1224. doi:10.1378/chest.107.5.1218
44. Sridhar MK, Carter R, Moran F, Banham SW. Use of a combined oxygen and carbon dioxide transcutaneous electrode in the estimation of gas exchange during exercise. *Thorax*. 1993;48(6):643–647. doi:10.1136/thx.48.6.643

International Journal of General Medicine

Dovepress

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>