

A checklist of the millipedes of Georgia, Caucasus (Diplopoda)

Mzia S. Kokhia¹, Sergei I. Golovatch²

1 Institute of Zoology, Ilia State University, K. Cholokashvili Ave., 3/5, Tbilisi 0162, Georgia **2** Institute for Problems of Ecology and Evolution, Russian Academy of Sciences, Leninsky prospekt 33, Moscow 119071, Russia

Corresponding author: Sergei I. Golovatch (sgolovatch@yandex.ru)

Academic editor: P. Stoev | Received 4 August 2017 | Accepted 9 November 2017 | Published 7 March 2018

<http://zoobank.org/7B60748C-1972-4952-A133-7F28B5B6BBD3>

Citation: Kokhia MS, Golovatch SI (2018) A checklist of the millipedes of Georgia, Caucasus (Diplopoda). In: Stoev P, Edgecombe GD (Eds) Proceedings of the 17th International Congress of Myriapodology, Krabi, Thailand. ZooKeys 741: 35–48. <https://doi.org/10.3897/zookeys.741.20042>

Abstract

The diplopod fauna of Georgia, Transcaucasia, is very rich given the country's relatively small territory, presently comprising 95 species from 42 genera, 12 families, and seven orders. Most of the Diplopoda known from Georgia are subendemics (39 species, or 38%), shared with one or more neighbouring countries, but another 33 species (33%) are strict endemics, nearly all highly localized, including 12 presumed troglobites. Several genera are likewise endemic to Georgia, including a few troglobionts. Within Georgia, the fauna of the western part (= Colchis) is particularly rich and diverse, the faunas of the central and eastern parts of the country growing increasingly depauperate inland and apparently following a rather gradual climatic aridisation gradient from west (the Black Sea coast) to east (Armenia and Azerbaijan). Much more work to include alpine and cave environments is required in order to reveal and refine the real diversity of Georgia's Diplopoda.

Keywords

Colchis, distribution, endemism, fauna, Myriapoda

Introduction

Georgia is one of the main countries in the Caucasus, lying between western Asia and Eastern Europe. It is bounded to the west by the Black Sea, to the north by Russia, to the south by Turkey, and to the southeast and east by Armenia and Azerbaijan



Figure 1. Crude geographical division of Georgia.

(Fig. 1). The area is largely montane to high montane, situated between latitudes 41° and 44°N, and longitudes 40° and 47°E. The Greater Caucasus Mountain Range, or Caucasus Major, forms the northern border of Georgia, while the southern border is bounded by the Lesser Caucasus Mountains, or Caucasus Minor. The Caucasus Major is much higher in elevation (up to more than 5,000 m a.s.l.) than the plateau-like Caucasus Minor, both being connected by the submeridional Surami (= Likhi) Mountain Range which divides Georgia into the western and central + eastern parts. Both parts are quite varied in climate and biota. Western Georgia's landscape ranges from lowland marsh-forests, swamps, and temperate rainforests within the Colchis Plain to eternal snows and glaciers, while the eastern part of the country even contains a small segment of semi-arid plains. Forests cover around 40% of Georgia's territory, while the alpine/subalpine zone accounts for approximately 10% of the land. The climate of Georgia is extremely diverse, but largely mild to warm, considering the nation's small size. There are two main climatic zones, roughly corresponding to the eastern and western parts of the country. The Greater Caucasus Mountain Range plays an important role in moderating Georgia's climate and protects the nation from the penetration of colder air masses from the north. The Lesser Caucasus Mountains partially protect the region from the influence of dry and hot air masses from the south (Bondyrev et al. 2015).

The history of diplopodological research in the Caucasus generally, and in Georgia in particular, started with the works of Victor (1839), Brandt (1840) and Karsch (1881), followed by faunistic contributions by Timotheew (1897), Attems (1898, 1899, 1901, 1903, 1907), Lignau (1903, 1907, 1911, 1915, 1924), Muralewicz (1907, 1911, 1913,

1927) and Issaev (1911). Muralewicz (1911) was the first to thoroughly review the fauna of Caucasian Myriapoda known to that date. Verhoeff (1921, 1930), Jawłowski (1929) and Lohmander (1928, 1932) had added a few more species of Caucasian millipedes before a real milestone synthesis appeared. That historical stage culminated with Lohmander's (1936) monograph which still serves, however outdated taxonomically, as one of the main sources of our knowledge of the Diplopoda of the Caucasus.

Several checklists, partly containing new faunistic records of Caucasian Diplopoda, appeared since then (Lang 1959, Kobakhidze 1964, 1965, Lokšina and Golovatch 1979, Talikadze 1984), but marked progress in the taxonomic study of millipedes in the region resumed only with contributions by Golovatch (1975, 1976a, 1976b, 1976c, 1977, 1979, 1980, 1981a, 1981b, 1984/85). Sporadic descriptions have since been upgraded to regional reviews of certain higher taxa such as genera, families and orders, with few exceptions only. These reviews mostly covered not only the Caucasus proper, but also the faunas of the adjacent parts of Turkey and Iran, e.g. the families Blaniulidae (Enghoff 1984, 1990, Golovatch and Enghoff 1990), Nemasomatidae (Enghoff 1985) and several tribes and genera of Julidae (Read 1992, Enghoff 1995, Evsyukov 2016b, Vagalinski and Lazányi 2018), all from the order Julida. The order Glomerida (Golovatch 1989a, 1989b, 1990, 1993, Golovatch and Chumachenko 2013), three orders of the subclass Colobognatha (Golovatch et al. 2015, Zuev 2017), as well as the orders Polyxenida (Short 2015, Short et al. 2018), Polydesmida (Golovatch et al. 2016, Evsyukov et al. 2016) and Chordeumatida (Antić and Makarov 2016) have also been revised in the scope of the entire Caucasian fauna, sometimes even broader. The faunas of two larger areas in Ciscaucasia have also been reviewed and updated (Evsyukov and Golovatch 2013, Evsyukov 2016a, Zuev 2014). A couple of nature reserves at the Black Sea coast of the Russian Caucasus have likewise been thoroughly surveyed for their local millipede faunas, with some data on ecology and distribution (Chumachenko 2016, Korobushkin et al. 2016).

The present paper provides an up-to-date checklist (Table 1) of the millipede fauna of Georgia, based on all available publications. It shows very considerable progress achieved since the latest lists by Kobakhidze (1964, 1965), who grossly repeated Lohmander's (1936) and added many new faunistic records, and by Talikadze (1984), who only considered the Colchidian part of the Caucasus, including the Black Sea coast area of Russia. Cave fauna has been reviewed within the entire former Soviet Union, including the Diplopoda of the Caucasus together with Georgia (Turbanov et al. 2016).

Georgia is conventionally divisible into three main parts: western, central, and eastern (Figure 1). This division is followed in the checklist below. Data are also given on the presence or absence of relevant species in the immediately neighbouring countries, including the Crimean Peninsula, as well as the distribution patterns and main literature sources. The checklist is arranged in alphabetic order per family, omitting subgeneric categories. All accepted designations are explained at the bottom of Table 1.

Results

Table 1. Checklist of the Diplopoda of Georgia, with data on species distributions, both within and beyond the country, their statuses, and the main relevant literature sources. Dp = Distribution pattern.

| Taxonomic composition | G | R | T | Ar | Az | Cr | St | Dp | Main relevant references |
|---|------|---|---|----|----|----|------|-----|---|
| Class Diplopoda | | | | | | | | | |
| Order Polyxenida | | | | | | | | | |
| Family Polyxenidae | | | | | | | | | |
| Genus <i>Polyxenus</i> Latreille, 1803 | | | | | | | | | |
| 1. <i>Polyxenus argentifer</i> Verhoeff, 1921 | G | + | | + | + | + | | AM | Short et al. 2018 |
| 2. <i>P. lagurus</i> (Linnaeus, 1758) | W, E | | | | | + | | sc | Short et al. 2018 |
| Family Lophoproctidae | | | | | | | | | |
| Genus <i>Lophoproctus</i> Pocock, 1894 | | | | | | | | | |
| 3. <i>Lophoproctus coecus</i> Pocock, 1894 | G | + | | | | + | | EM | Short 2015, Short et al. 2018 |
| Order Polyzoniida | | | | | | | | | |
| Family Hirudisomatidae | | | | | | | | | |
| Genus <i>Hirudosoma</i> Fanzago, 1881 | | | | | | | | | |
| 4. <i>Hirudosoma roseum</i> (Victor, 1839) | G | + | + | | + | | se | EM | Golovatch et al. 2015 |
| Order Siphonocryptida | | | | | | | | | |
| Family Siphonocryptidae | | | | | | | | | |
| Genus <i>Hirudicryptus</i> Enghoff & Golovatch, 1985 | | | | | | | | | |
| 5. <i>Hirudicryptus abchasicus</i> Golovatch, Esvyukov & Reip, 2015 | W | + | | | | | se | Ca | Golovatch et al. 2015, Zuev 2017 |
| Order Glomerida | | | | | | | | | |
| Family Glomeridae | | | | | | | | | |
| Genus <i>Hyleoglomeris</i> Verhoeff, 1910 | | | | | | | | | |
| 6. <i>Hyleoglomeris awchasica</i> (Brandt, 1840) | W | + | | | | | se | Ca | Golovatch 1976a, 1989b |
| 7. <i>H. specialis</i> Golovatch, 1989 | E | + | | | | | se | Ca | Golovatch 1989b |
| Genus <i>Trachysphaera</i> Heller, 1858 | | | | | | | | | |
| 8. <i>Trachysphaera costata</i> (Waga, 1857) | G | + | + | + | + | + | | EuM | Golovatch 1990, 2008 |
| 9. <i>T. fragilis</i> Golovatch, 1976 | G | | | | | | t, e | Ca | Golovatch 1990, Golovatch and Turbanov 2017 |
| 10. <i>T. minuta</i> Golovatch, 1976 | G | + | + | + | | | se | Ca | Golovatch 1990 |
| 11. <i>T. orientalis</i> Golovatch, 1976 | W | | | | | | t, e | Ca | Golovatch 1976c, 1990 |
| 12. <i>T. radiosa</i> (Lignau, 1911) | W | + | | | | | se | Ca | Golovatch 1976c, 1990 |
| 13. <i>T. solida</i> Golovatch, 1976 | W, C | | | | | | se | Ca | Golovatch 1976c, 1990, 1993 |
| Family Glomeridellidae | | | | | | | | | |
| Genus <i>Typhloglomeris</i> Verhoeff, 1898 | | | | | | | | | |
| 14. <i>Typhloglomeris lohmanderi</i> (Golovatch, 1989) | C, E | + | | + | | | se | Ca | Golovatch 1989a, 2003 |
| Order Chordeumatida | | | | | | | | | |
| Family Anthroleucosomatidae | | | | | | | | | |
| Genus <i>Acanthophorella</i> Antić & Makarov, 2016 | | | | | | | | | |
| 15. <i>Acanthophorella barjadzei</i> Antić & Makarov, 2016 | W | | | | | | t, e | Ca | Antić and Makarov 2016 |
| Genus <i>Adshardicus</i> Golovatch, 1981 | | | | | | | | | |
| 16. <i>Adshardicus strasseri</i> Golovatch, 1981 | W | | + | | | | se | Ca | Enghoff 2006, Antić and Makarov 2016 |
| Genus <i>Alpinella</i> Antić & Makarov, 2016 | | | | | | | | | |
| 17. <i>Alpinella waltheri</i> Antić & Makarov, 2016 | E | | | | | | e | Ca | Antić and Makarov 2016 |
| Genus <i>Brachychaetosoma</i> Antić & Makarov, 2016 | | | | | | | | | |
| 18. <i>Brachychaetosoma turbanovi</i> Antić & Makarov, 2016 | W | | | | | | t, e | Ca | Antić and Makarov 2016 |

| Taxonomic composition | G | R | T | Ar | Az | Cr | St | Dp | Main relevant references |
|--|------|---|---|----|----|------|-----|----|---|
| Genus <i>Caucaseuma</i> Strasser, 1970 | | | | | | | | | |
| 19. <i>Caucaseuma kelasuri</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| 20. <i>C. variabile</i> Antić & Makarov, 2016 | C | + | | | | se | Ca | | Antić and Makarov 2016 |
| Genus <i>Cryptacanthophorella</i> Antić & Makarov, 2016 | | | | | | | | | |
| 21. <i>Cryptacanthophorella manubriata</i> Antić & Makarov, 2016 | W, C | | | | | e | Ca | | Antić and Makarov 2016 |
| Genus <i>Dentatosoma</i> Antić & Makarov, 2016 | | | | | | | | | |
| 22. <i>Dentatosoma denticulatum</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| 23. <i>D. magnum</i> Antić & Makarov, 2016 | W | + | | | | se | Ca | | Antić and Makarov 2016 |
| 24. <i>D. zeraboseli</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| Genus <i>Georgiosoma</i> Antić & Makarov, 2016 | | | | | | | | | |
| 25. <i>Georgiosoma bicornutum</i> Antić & Makarov, 2016 | W | | | | | t, e | Ca | | Antić and Makarov 2016 |
| Genus <i>Herculina</i> Antić & Makarov, 2016 | | | | | | | | | |
| 26. <i>Herculina oligosagittae</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| 27. <i>H. polysagittae</i> Antić & Makarov, 2016 | C | | | | | e | Ca | | Antić and Makarov 2016 |
| Genus <i>Heterocaucaseuma</i> Antić & Makarov, 2016 | | | | | | | | | |
| 28. <i>Heterocaucaseuma longicorne</i> Antić & Makarov, 2016 | W | | | | | t, e | Ca | | Antić and Makarov 2016 |
| 29. <i>Heterocaucaseuma mauriesi</i> (Golovatch & Makarov, 2011) | W | | | | | t, e | Ca | | Golovatch and Makarov 2011, Antić and Makarov 2016 |
| Genus <i>Metamastigophorophyllum</i> Ceuca, 1976 | | | | | | | | | |
| 30. <i>Metamastigophorophyllum giljarovi</i> (Lang, 1959) | W | + | | | | se | Ca | | Antić and Makarov 2016 |
| 31. <i>M. hamatum</i> Antić & Makarov, 2016 | W | + | | | | se | Ca | | Antić and Makarov 2016 |
| 32. <i>M. lamellohirsutum</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| 33. <i>M. torsivum</i> Antić & Makarov, 2016 | G | | | | + | se | Ca | | Antić and Makarov 2016 |
| Genus <i>Paranotosoma</i> Antić & Makarov, 2016 | | | | | | | | | |
| 34. <i>Paranotosoma attemsi</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| 35. <i>P. cordatum</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| 36. <i>P. subrotundatum</i> Antić & Makarov, 2016 | W | + | | | | se | Ca | | Antić and Makarov 2016 |
| Genus <i>Pseudoflagellophorella</i> Antić & Makarov, 2016 | | | | | | | | | |
| 37. <i>Pseudoflagellophorella eskovi</i> Antić & Makarov, 2016 | C, E | | + | + | | se | Ca | | Antić and Makarov 2016 |
| 38. <i>P. mirabilis</i> Antić & Makarov, 2016 | W | | | | | e | Ca | | Antić and Makarov 2016 |
| 39. <i>P. papilioformis</i> Antić & Makarov, 2016 | E | | | | + | se | Ca | | Antić and Makarov 2016 |
| Genus <i>Ratcheuma</i> Golovatch, 1985 | | | | | | | | | |
| 40. <i>Ratcheuma excorne</i> Golovatch, 1985 | W | | | | | t, e | Ca | | Golovatch 1984/85, Antić and Makarov 2016 |
| Order Julida | | | | | | | | | |
| Family Blaniulidae | | | | | | | | | |
| Genus <i>Cibinulus</i> Verhoeff, 1927 | | | | | | | | | |
| 41. <i>Cibinulus phlepsii</i> (Verhoeff, 1897) | W | + | | | | | EuM | | Enghoff 1984, 2006 |
| Genus <i>Nopoiulus</i> Menge, 1851 | | | | | | | | | |
| 42. <i>Nopoiulus brevipilosus</i> Enghoff, 1984 | W | | | | | t, e | Ca | | Enghoff 1984 |
| 43. <i>N. densepilosus</i> Enghoff, 1984* | W | | | | + | | Ca | | Enghoff 1984, Golovatch and Enghoff 1990 |
| 44. <i>N. golovatchi</i> Enghoff, 1984 | W | + | | | | | Ca | | Enghoff 1984, 1990 |
| 45. <i>N. kochii</i> (Gervais, 1847) | G | + | + | + | + | | sc | | Enghoff 1984, Golovatch and Enghoff 1990 |

| Taxonomic composition | G | R | T | Ar | Az | Cr | St | Dp | Main relevant references |
|--|------|---|---|----|----|----|------|----|--|
| Family Nemasomatidae | | | | | | | | | |
| Genus <i>Nemasoma</i> C.L. Koch, 1847 | | | | | | | | | |
| 46. <i>Nemasoma caucasicum</i> (Lohmander, 1932) | G | + | + | + | + | | se | Ca | Enghoff 1985 |
| Family Julidae | | | | | | | | | |
| Genus <i>Amblyjulus</i> Silvestri, 1896 | | | | | | | | | |
| 47. <i>Amblyjulus adsharicus</i> Lohmander, 1936 | W | | | | | | e | Ca | Lohmander 1936 |
| 48. <i>A. georgicus</i> Lohmander, 1932 | C | | | | | | e | Ca | Lohmander 1932 |
| Genus <i>Archileucogeorgia</i> Lohmander, 1936 | | | | | | | | | |
| 49. <i>Archileucogeorgia abchasica</i> Lohmander, 1936 | W | | | | | | t, e | Ca | Lohmander 1936 |
| 50. <i>Archileucogeorgia satunini</i> Lohmander, 1936 | W | | | | | | e | Ca | Lohmander 1936 |
| Genus <i>Brachyiulus</i> Berlese, 1884 | | | | | | | | | |
| 51. <i>Brachyiulus lusitanus</i> Verhoeff, 1898* | C | | | | + | | | M | Lohmander 1936 |
| Genus <i>Catamicrophyllum</i> Verhoeff, 1900 | | | | | | | | | |
| 52. <i>Catamicrophyllum caucasicum</i> (Attems, 1901) | G | + | + | + | | | se | Ca | Lohmander 1936, Enghoff 1995 |
| Genus <i>Calyptophyllum</i> Brolemann, 1922 | | | | | | | | | |
| 53. <i>Calyptophyllum</i> sp. | W | | | | | | ? | ? | Lohmander 1936, Enghoff 1995 |
| Genus <i>Chaetoleptophyllum</i> Verhoeff, 1898 | | | | | | | | | |
| 54. <i>Chaetoleptophyllum flexum</i> Golovatch, 1979 | G | + | | | | | se | Ca | Golovatch 1979, Chumachenko 2016, Korobushkin et al. 2016 |
| Genus <i>Cylindroiulus</i> Verhoeff, 1894 | | | | | | | | | |
| 55. <i>Cylindroiulus bellus</i> (Lignau, 1903) | W? | + | | | | + | | EM | Lignau 1903, Read 1992, Chumachenko 2016 |
| 56. <i>C. crassiphyllum</i> Read, 1992 | G | | + | | | | se | Ca | Read 1992 |
| 57. <i>C. kacheticus</i> Lohmander, 1936 | E | + | | | | | se | Ca | Read 1992 |
| 58. <i>C. olgaiinna</i> Read, 1992 | W | | | | | | e | Ca | Read 1992 |
| 59. <i>C. parvus</i> Lohmander, 1928 | C, E | | | + | | | se | Ca | Read 1992 |
| 60. <i>C. placidus</i> (Lignau, 1903) | W, C | + | | | | | se | Ca | Read 1992 |
| 61. <i>C. pterophyllum</i> Read, 1992 | W, C | + | | | | | se | Ca | Read 1992, Zuev 2014 |
| 62. <i>C. quadrus</i> Read, 1992 | G | + | | | | | se | Ca | Read 1992 |
| 63. <i>C. ruber</i> (Lignau, 1903) | W | + | | | | | se | Ca | Read 1992 |
| 64. <i>C. schestoperovi</i> Lohmander, 1936 | W | + | | | | | se | Ca | Lohmander 1936, Read 1992 |
| 65. <i>C. truncorum</i> (Silvestri, 1896) | W | | | | | | | sc | Read 1992 |
| Genus <i>Grusiniulus</i> Lohmander, 1936 | | | | | | | | | |
| 66. <i>Grusiniulus redikorzevi</i> Lohmander, 1932 | C | | | | | | e | Ca | Lohmander 1936, Vagalinski and Lazányi 2018 |
| Genus <i>Julus</i> Linnaeus, 1758 | | | | | | | | | |
| 67. <i>Julus colchicus</i> Lohmander, 1936 | W | + | + | | | | se | Ca | Lohmander 1936, Enghoff 2006 |
| 68. <i>J. kubanus</i> Verhoeff, 1921 | W | + | | | | | se | Ca | Lohmander 1936, Kobakhidze 1965 |
| 69. <i>J. lindholmi</i> Lohmander, 1936 | W | | | | | | e | Ca | Lohmander 1936 |
| Genus <i>Leptoziulus</i> Verhoeff, 1894 | | | | | | | | | |
| 70. <i>Leptoziulus disparatus</i> Lohmander, 1936 | C | + | | | | | se | Ca | Lohmander 1936, Enghoff 2006 |
| 71. <i>L. tanytropis</i> (Attems, 1901) | C, E | | | + | | | | Ca | Lohmander 1936 |
| Genus <i>Leucogeorgia</i> Verhoeff, 1930 | | | | | | | | | |
| 72. <i>Leucogeorgia longipes</i> Verhoeff, 1930 | W | | | | | | t, e | | Verhoeff 1930 |
| 73. <i>L. rediviva</i> Golovatch, 1983 | W | | | | | | t, e | Ca | Golovatch 1983 |
| Genus <i>Megaphyllum</i> Verhoeff, 1894 | | | | | | | | | |
| 74. <i>Megaphyllum dioscoridis</i> (Lignau, 1915) | W | + | | | | | e | Ca | Lignau 1915, Lohmander 1936, Chumachenko 2016, Vagalinski and Lazányi 2018 |
| 75. <i>M. hercules</i> (Verhoeff, 1901) | W | + | | | | | | EM | Lazányi and Vagalinski 2013 |

| Taxonomic composition | G | R | T | Ar | Az | Cr | St | Dp | Main relevant references |
|--|------|---|---|----|----|----|----|----|---|
| 76. <i>M. spathulatum</i> (Lohmander, 1936) | W? | ? | | | | | | Ca | Lohmander 1936, Lazányi and Vagalinski 2013 |
| Genus <i>Omobrachyiulus</i> Lohmander, 1936 | | | | | | | | | |
| 77. <i>Omobrachyiulus adsharicus</i> (Lohmander, 1936) | W | | | | | | e | Ca | Lohmander 1936, Vagalinski and Lazányi 2018 |
| 78. <i>O. brachyurus</i> (Attems, 1899) | G | + | + | + | + | | | EM | Lohmander 1936, Enghoff 2006, Vagalinski and Lazányi 2018 |
| 79. <i>O. curvocaudatus</i> (Lignau, 1903) | W | + | | | | | se | Ca | Lohmander 1936, Vagalinski and Lazányi 2018 |
| 80. <i>O. divaricatus</i> (Lohmander, 1936) | G | | | + | | | se | Ca | Lohmander 1936, Vagalinski and Lazányi 2018 |
| 81. <i>O. hortensis</i> (Golovatch, 1981) | W | | | | | | e | Ca | Golovatch 1981, Vagalinski and Lazányi 2018 |
| 82. <i>O. implicitus</i> Lohmander, 1936 (= <i>O. i. ritsensis</i> (Golovatch, 1981)) | W | + | | | | | se | Ca | Lohmander 1936, Chumachenko 2016, Vagalinski and Lazányi 2018 |
| 83. <i>O. macrourus</i> (Lohmander, 1928) (= <i>O. m. abchasicus</i> (Lohmander, 1936)) | W, C | | | | | | e | Ca | Lohmander 1936, Kobakhidze 1965, Vagalinski and Lazányi 2018 |
| Genus <i>Pachyiulus</i> Berlese, 1883 | | | | | | | | | |
| 84. <i>Pachyiulus flavipes</i> (C.L. Koch, 1847) | W | | | | | + | | M | Lohmander 1936 |
| 85. <i>Pachyiulus krivolutskyi</i> Golovatch, 1977 | W | + | | | | | se | Ca | Evsukov 2016 |
| Order Polydesmida | | | | | | | | | |
| Family Paradoxosomatidae | | | | | | | | | |
| Genus <i>Oxidus</i> Cook, 1911 | | | | | | | | | |
| 86. <i>Oxidus gracilis</i> (C.L. Koch, 1847) | W | + | | | | | | sc | Lignau 1915, Lohmander 1936, Chumachenko 2016 |
| Family Polydesmidae | | | | | | | | | |
| Genus <i>Brachydesmus</i> Heller, 1858 | | | | | | | | | |
| 87. <i>Brachydesmus assimilis</i> Lohmander, 1936 | C, E | + | | | | | se | Ca | Golovatch et al. 2016 |
| 88. <i>B. furcatus</i> Lohmander, 1936 | W | + | | | | | se | Ca | Golovatch et al. 2016 |
| 89. <i>B. kalishevskyi</i> Lignau, 1915 | G | + | + | + | + | | se | Ca | Golovatch et al. 2016 |
| 90. <i>B. kvavadzei</i> Golovatch, Evsukov & Reip, 2016 | W | | | | | | e | Ca | Golovatch et al. 2016 |
| 91. <i>B. simplex</i> Golovatch, Evsukov & Reip, 2016 | W | + | | | | | se | Ca | Golovatch et al. 2016 |
| 92. <i>B. superus</i> Latzel, 1884 | W | + | | | | | | sc | Golovatch et al. 2016 |
| Genus <i>Polydesmus</i> Latreille, 1803 | | | | | | | | | |
| 93. <i>Polydesmus abchasicus</i> Attems, 1899 | W | + | | | | | se | Ca | Golovatch et al. 2016 |
| 94. <i>P. lignaui</i> Lohmander, 1936 | W | + | | | | | se | Ca | Golovatch et al. 2016 |
| 95. <i>P. mediterraneus</i> Daday, 1889 | W | | | | | + | EM | | Golovatch et al. 2016 |

Designations: G – entire Georgia; W – western Georgia; C – central Georgia; E – eastern Georgia; R – Russian Caucasus; T – Turkey; Ar – Armenia; Az – Azerbaijan; Cr – Crimean Peninsula; (+) – present; St – status; e – endemic to Georgia; se – subendemic to Georgia; t – presumed troglobiont; sc – subcosmopolitan; AM – Ancient Mediterranean; EuM – Euro-Mediterranean; M – Mediterranean; EM – eastern Mediterranean; EE – eastern European; Ca – Caucasian.

Discussion

As is evident from the above list, the millipede fauna of Georgia is, surprisingly, very diverse, especially so given the relatively small territory it covers. This is hardly surprising, because Diplopoda are largely mesophilous forest-dwellers (e.g., Kime and Golovatch 2000, Golovatch and Kime 2009). Georgia with its mostly mild climate and large

woodland areas supports the richest millipede fauna in the entire Caucasus, nearly twice as rich as neighbouring Azerbaijan (Bababekova 1996, a quite poorly compiled list) or Iran (Enghoff and Moravvej 2005), and approximately 2/3 as diverse as the fauna of the so much larger Turkey (Enghoff 2006, 135 species), for all of which rather modern country checklists are available. Although the bulk of the fauna of Georgia is represented by epigean taxa, the abundant limestone massifs, primarily those lying at the northern and northeastern peripheries of the Colchis, harbour numerous karstic caves with their own fauna. Troglobionts do account for a considerable proportion (12 species, or 14%) of Georgia's millipede species (Barjadze et al. 2015, Turbanov et al. 2016).

Western Georgia, the Colchis (Fig. 1), is especially rich in millipedes, apparently due to the moist and mild climate near the warm Black Sea, highly varied, but largely forested habitats, and abundant karst caves. The Colchidian millipede fauna is also the richest in endemics, both at the species and generic levels. The orders Chordeumatida and Julida are particularly strongly diversified in Georgia. As well the country supports also *Hirudicryptus abchasicus*, a subendemic representing one of the most relict diplopod orders, Siphonocryptida, which presently comprises only seven species in two genera and a single family (Golovatch et al. 2015, Zuev 2017). Central and eastern parts of Georgia are increasingly drier, in places even semi-arid, and the millipede fauna generally demonstrates a decline in diversity from the Black Sea coast inland, appearing to follow rather gradual climatic aridisation gradient from west to east.

Most of the Diplopoda known from Georgia are subendemics (40 species, or 42%), shared with one or more neighbouring countries, but another 33 species (34%) are strict endemics, nearly all highly localized, including 12 presumed troglobionts. Several genera are likewise endemic to Georgia, including a few troglobionts. The proportions of the remaining, more widely distributed, species are rather modest, represented by Mediterranean, Euro-Mediterranean, eastern Mediterranean, eastern European or ubiquitous elements, but even among the latter the subcosmopolitan *Nopoiulus kochii* may have originated in the Caucasus, because the remaining congeners (from all subgenera) seem to be endemic to the Caucasus region (Golovatch and Enghoff 1990).

The present checklist must be understood as temporary, far from complete, marking only the state of knowledge of diplopodological research in Georgia. Several of Lohmander's *nomina nuda* listed by Kobakhidze (1964) are thereby omitted. Much more work is required to reveal the real diversity of Georgia's Diplopoda. Discoveries and descriptions of numerous new taxa, both species and probably even genera, can still be expected in the future. Further faunistic records are necessary to refine not only the taxonomy and the above list, but also the distributions, both horizontal and vertical. Very little is known yet concerning high-montane millipedes, in particular, whether strictly alpine Caucasian/Georgian endemics exist at all, like those few recorded from the Pyrenees and Alps. Finally, cave explorations in Georgia will undoubtedly reveal many more new troglobionts, including diplopods.

Acknowledgements

Thanks are due to Boyan Vagalinski (Sofia, Bulgaria) and Megan Short (Melbourne, Australia) for kindly sharing published and unpublished information on their current research relevant to our topic, as well as acknowledgements to them and Dragan Antić (Belgrade, Serbia) and Elena Mikhajlova (Vladivostok, Russia) for their helpful reviews of the manuscript. MK is grateful for the support rendered through the Project “Georgia’s Natural and Agricultural Ecosystems: Animal Diversity, Monitoring, Biocontrol”, funded by the Ministry of Science and Education of Georgia.

References

- Antić DŽ, Makarov SE (2016) The Caucasus as a major hotspot of biodiversity: Evidence from the millipede family Anthroleucosomatidae (Diplopoda, Chordeumatida). Zootaxa 4211(1): 1–205. <https://doi.org/10.11646/zootaxa.4211.1.1>
- Attems C (1898) System der Polydesmiden. I. Theil. Denkschriften der Kaiserlichen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Classe 67: 221–482.
- Attems C (1899) Neues über paläarktische Myriopoden. Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Thiere 12: 286–336. <https://doi.org/10.5962/bhl.part.2032>
- Attems C (1901) Myriopoden. – Horvath G (Hrsg), Zoologische Ergebnisse der dritten asiatischen Forschungsreise des Grafen Eugen Zichy, Budapest 2: 275–310.
- Attems C (1903) Beiträge zur Myriopodenkunde. Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Thiere 18: 63–154.
- Attems C (1907) Myriopoden aus der Krim und dem Kaukasus von Dr. A. Stuxberg gesammelt. Arkiv för Zoologi 3(25): 1–16.
- Bababekova LA (1996) The subphylum of tracheates – Tracheata. In: Aliev SV, Kasymov EG (Eds) Animal life of Azerbaijan 2: The phylum of arthropods. Elm Publishers, Baku, 89–97. [in Russian]
- Barjadze S, Murvanidze M, Arabuli T, Mumladze L, Pkhakadze V, Djanashvili R, Salakaia M (2015) Annotated list of invertebrates of the Georgian karst caves. Georgian Academic Book, Tbilisi, 119 pp.
- Bondyrev IV, Davitashvili ZV, Singh VP (2015) Natural Resources. The Geography of Georgia, Problems and Perspectives. World Regional Geography Book Series. Springer, Cham, 227 pp. https://doi.org/10.1007/978-3-319-05413-1_11
- Brandt JF (1840) Remarques critiques sur les espèces qui composent le genre *Glomeris*, suivies de quelques observations sur leur distribution géographique. Bulletin scientifique publié par l’Académie Impériale des Sciences de Saint-Pétersbourg 7(148, 149): 37–44.
- Chumachenko YA (2016) [Millipede (Diplopoda) populations in a yew-boxtree grove in the Caucasian Nature Reserve, Russia]. Zoologicheskii Zhurnal 95(4): 406–416. <https://doi.org/10.7868/S004451341604005X> [In Russian, English summary]

- Enghoff H (1984) A revision of the Nopoziulinae, with notes on the classification of blaniulid millipedes (Diplopoda: Julida: Blaniulidae). *Senckenbergiana biologica* 64 (4–6): 393–427.
- Enghoff H (1985) The millipede family Nemasomatidae. With the description of a new genus, and a revision of *Orinisobates* (Diplopoda: Julida). *Entomologica scandinavica* 16: 27–67. <https://doi.org/10.1163/187631285X00045>
- Enghoff H (1990) A new species of *Nopoziulus* Menge 1851, subgenus *Paranopoziulus* Lohmander, 1939, from the Caucasus, with records of consubgeneric species, and the systematic position of *Thassoblaniulus* Mauriès 1985 (Diplopoda: Julida: Blaniulidae). *Senckenbergiana biologica* 70(4/6): 323–330.
- Enghoff H (1995) A revision of the Paectophyllini and Calyptophyllini: millipedes of the Middle East (Diplopoda: Julida: Julidae). *Journal of Natural History* 29: 685–786. <https://doi.org/10.1080/00222939500770241>
- Enghoff H (2006) The millipedes of Turkey (Diplopoda). *Steenstrupia* 29(2): 175–198.
- Enghoff H, Moravvej SA (2005) A review of the millipede fauna of Iran (Diplopoda). *Zoology in the Near East* 35: 61–72. <https://doi.org/10.1080/09397140.2005.10638104>
- Evsyukov AP (2016a) A new species of the millipede genus *Julus* from the Rostov-on-Don Region, southern Russia (Diplopoda: Julida: Julidae). *Arthropoda Selecta* 25(3): 241–245. http://kmkjournals.com/upload/PDF/ArthropodaSelecta/25/25_3_241_245_Evsyukov_for_Inet.pdf
- Evsyukov AP (2016b) The millipede *Pachyiulus krivolutskyi* Golovatch, 1977, the easternmost species of the eastern Mediterranean genus *Pachyiulus* Berlese, 1883, endemic to the western Caucasus (Diplopoda: Julida: Julidae). *Russian Entomological Journal* 25(3): 299–306. http://kmkjournals.com/upload/PDF/REJ/25/ent25_3_299_306_Evsyukov_for_Inet.pdf
- Evsyukov AP, Golovatch SI (2013) Millipedes (Diplopoda) from the Rostov-on-Don Region, southern Russia. *Arthropoda Selecta* 22(3): 207–215. http://kmkjournals.com/upload/PDF/ArthropodaSelecta/22/22_3%2020207_215%20Evsyukov%20Golovatch%20for%20Inet.pdf
- Evsyukov AP, Golovatch SI, Reip H (2016) The millipede genus *Strongylosoma* Brandt, 1833 in the Caucasus (Diplopoda, Polydesmida, Paradoxosomatidae). *Acta Societatis Zoologicae Bohemicae* 80: 7–16.
- Golovatch SI (1975) [Two genera of Oniscomorpha (Diplopoda) new to the USSR fauna, found in Transcaucasia, and their zoogeographical relations]. *Zoologicheskii Zhurnal* 54(10): 1566–1571. [In Russian, English summary]
- Golovatch SI (1976a) [New or little-known species of Glomeridae (Diplopoda, Oniscomorpha)]. *Zoologicheskii Zhurnal* 55(6): 931–935. [In Russian, English summary].
- Golovatch SI (1976b) [New species of polydesmidans (Polydesmida, Diplopoda) of the USSR]. *Zoologicheskii Zhurnal* 55(10): 1567–1570. [In Russian, English abstract]
- Golovatch SI (1976c) [Description of new species of *Trachysphaera* (Oniscomorpha; Diplopoda) from caves of Transcaucasia with remarks on *T. rotundata* (Lignau) 1911]. *Bulletin de la Société des Naturalistes de Moscou, Section Biologique* [Byulleten Moskovskogo obshchestva ispytateley prirody, Otdel biologicheskiy] 81(5): 30–43. [In Russian, English summary]

- Golovatch SI (1977) [New or little-known Julida (Diplopoda) of the Sataplia Nature Reserve (SSR Georgia)]. Bulletin de la Société des Naturalistes de Moscou, Section Biologique [Byulleten Moskovskogo obshchestva ispytateley prirody. Otdel biologicheskiy] 82(4): 46–51. [In Russian, English summary]
- Golovatch SI (1979) [Three genera of Diplopoda-Chilognatha new to the USSR fauna]. Zoologicheskii Zhurnal 58(3): 336–343. [In Russian, English summary]
- Golovatch SI (1980) [A new species of the millipede genus *Fioria* (Diplopoda, Platynoididae) from Transcaucasia]. Zoologicheskii Zhurnal 59(3): 458–460. [In Russian, English summary]
- Golovatch SI (1981a) A review of the cavernicolous millipede fauna (Diplopoda) of the western Caucasus. Caves of Georgia [Peshchery Gruzii], Tbilisi 9: 108–111. [In Russian]
- Golovatch SI (1981b) Some new forms of millipedes (Diplopoda) from the Caucasus. Annales Zoologici 36(5): 105–116.
- Golovatch SI (1983) Une nouvelle espèce cavernicole du genre *Leucogeorgia* de la Transcaucasie (Diplopoda, Julidae). Fragmenta Entomologica 17(1): 47–50.
- Golovatch SI (1984/85) Two new genera of cave-dwelling millipedes (Diplopoda), with remarks on the millipede fauna of West Caucasian caves. International Journal of Speleology 14(1/4): 39–50.
- Golovatch SI (1989a) Diplopoda of the Caucasus, 1. Glomeridellidae, with contributions to the fauna of Turkey. Senckenbergiana biologica 69(4/6): 405–419.
- Golovatch SI (1989b) Diplopoda of the Caucasus, 2. Glomeridae, with contributions to the fauna of Turkey. Senckenbergiana biologica 69(4/6): 421–440.
- Golovatch SI (1990) Diplopoda of the Caucasus, 3. Trachysphaeridae, with contributions to the fauna of Turkey. Senckenbergiana biologica 70(4/6): 331–358.
- Golovatch SI (1993) *Trachysphaera solida* Golovatch, 1976, a revalidated Caucasian millipede species (Diplopoda Glomerida Trachysphaeridae). Arthropoda Selecta 2(4): 19–20.
- Golovatch SI (2003) Two new species of Glomeridellidae (Diplopoda: Glomerida) from the Middle East. Arthropoda Selecta 11(4): 255–258.
- Golovatch SI (2008) On three remarkable millipedes (Diplopoda) from the Crimea, Ukraine. International Journal of Myriapodology 1: 97–110. <https://doi.org/10.1163/187525408X316767>
- Golovatch SI, Chumachenko YA (2013) The millipede *Typhloglomeris caucasica* Golovatch, 1975 found epigaeically (Diplopoda, Glomerida, Glomeridellidae). Biodiversity Data Journal 1: 1–3, e981. <https://doi.org/10.3897/BDJ.1.e981>
- Golovatch SI, Enghoff H (1990) [The millipede *Nopoiulus kochii* (Gervais, 1847) in the Caucasus (Diplopoda, Julida, Blaniulidae)]. In: Striganova BR (Ed.) Fauna of Terrestrial Invertebrates of the Caucasus. Nauka Publishers, Moscow, 114–118. [In Russian]
- Golovatch SI, Kime RD (2009) Millipede (Diplopoda) distributions: A review. Soil Organisms 81(3): 565–597.
- Golovatch SI, Makarov S (2011) Two new, easternmost species of the millipede genus *Anastigona* Silvestri, 1898 from Israel and the Caucasus (Diplopoda, Chordeumatida, Anthroleucosomatidae). Arthropoda Selecta 20(1): 11–20. http://kmkjournals.com/upload/PDF/ArthropodaSelecta/20/20_1%20011_020%20Golovatch%2C%20Makarov%20for%20Inet.pdf

- Golovatch SI, Turbanov IS (2017) The cave millipede *Trachysphaera fragilis* Golovatch, 1976, new to the fauna of Abkhazia, western Caucasus (Diplopoda: Glomerida: Glomeridae). Russian Entomological Journal 25(1): 101–102. http://kmkjournals.com/upload/PDF/REJ/26/ent26_1_101_102_Golovatch_Turbanov_for_Inet.pdf
- Golovatch SI, Evsyukov AP, Reip HS (2015) Colobognatha millipedes in the Caucasus (Diplopoda: Polyzoniida, Platydesmida, Siphonocryptida). Zootaxa 3972(2): 250–266. doi.org/10.11646/zootaxa.3972.2.6
- Golovatch SI, Evsyukov AP, Reip HS (2016) The millipede family Polydesmidae in the Caucasus (Diplopoda: Polydesmida). Zootaxa 4085(1): 1–51. <https://doi.org/10.11646/zootaxa.4085.1.1>
- Issaev VM (1911) [Observations of millipedes Diplopoda at the Black Sea coast of the Caucasus]. Travaux de la Société Impériale des Naturalistes de Saint-Pétersbourg [Trudy Imperatorskogo Sankt-Peterburgskogo obshchestva estestvoispytateley] 42(1): 292–310. [In Russian]
- Jawłowski H (1929) Über einige neue Diplopoden-Arten aus Kleinasien und Transkaukasien. Annales Musei Zoologici Polonici 8(1): 49–53.
- Karsch F (1881) Neue Juliden des Berliner Museums, als Prodromus einer Juliden-Monographie. Zeitschrift für die gesammten Naturwissenschaften 54: 1–79.
- Kime RD, Golovatch SI (2000) Trends in the ecological strategies and evolution of millipedes (Diplopoda). Biological Journal of the Linnean Society 69: 333–349.
- Kobakhidze DN (1964) [Myriopoda]. In [Animal Life of Georgia 2: Arthropoda. Institute of Zoology, SSR Georgia Academy of Sciences, Tbilisi, 186–195. [In Georgian]
- Kobakhidze DN (1965) [A list of millipedes (Diplopoda) of SSR Georgia]. Fragmenta Faunistica 11(21): 390–398. [In Russian, summaries in Polish and German]
- Korobushkin DI, Semenyuk II, Tuf IH (2016) An annotated checklist of the Chilopoda and Diplopoda (Myriapoda) of the Abrau Peninsula, northwestern Caucasus, Russia. Biodiversity Data Journal 19(4): e7308. <https://doi.org/10.3897/BDJ.4.e7308>
- Lang J (1959) [To the knowledge of millipedes (Diplopoda) of the USSR territory] // Zoologicheskii Zhurnal 38(12): 1790–1796. [In Russian, a summary in English]
- Lazányi E, Vagalinski B (2013) Redefinition of the millipede subgenus *Megaphyllum* sensu stricto Verhoeff, 1894 and neotype designation for *Megaphyllum austriacum* (Latzel, 1884) (Myriapoda: Diplopoda: Julida: Julidae). Zootaxa 3741: 55–100. <http://dx.doi.org/10.11646/zootaxa.3741.1.2>
- Lignau N (1903) [Myriapods of the Black Sea coast of the Caucasus]. Die Myriopoden am Kaukasischen Schwarzmeerofer. Mémoires de la Société Impériale des Naturalistes de la Nouvelle-Russie [Zapiski Imperatorskogo Novorossiyskogo obshchestva estestvoispytateley. Odessa] 25(1): 82–149. [In Russian and German]
- Lignau N (1907) [To the myriapod fauna of the Caucasus]. Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de Saint Pétersbourg [Ezhegodnik zoologicheskogo muzeya Imperatorskoy akademii nauk. Sankt-Peterburg] 12: 195–200.
- Lignau N (1911) Neue Beiträge zur Myriopodenfauna des Kaukasus. Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de Saint Pétersbourg [Ezhegodnik zoologicheskogo muzeya Imperatorskoy akademii nauk. Sankt-Peterburg] 16: 38–59.

- Lignau N (1915) Vielfüssler aus Abchasien. Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de Petrograd [Ezhegodnik zoologicheskogo muzeya Imperatorskoy akademii nauk. Petrograd] 19: 349–401 (for 1914).
- Lignau N (1924) Les nouveaux représentants du genre *Polyxenus*. Journal of Research Chairs in Odessa [Zhurnal nauchno-issledovatelskikh kafedr v Odesse] 1(10/11): 191–201. [In Russian and French]
- Lohmander H (1928) Neue Diplopoden aus Ukraine und dem Kaukasus. Mémoires de la Classe des Sciences Physiques et Mathématiques, Académie des Sciences de l'Ukraine [Trudy Fizichno-matematichnogo viddilu, Ukrainska akademiya nauk. Kyiv] 4(3): 529–550.
- Lohmander H (1932) Neue transkaukasische Diplopoden. 4. Aufsatz über Diplopoden aus Sowjet-Union. Zoologischer Anzeiger 98(7/8): 171–182.
- Lohmander H (1936) Über die Diplopoden des Kaukasusgebietes. Göteborgs Kungliga Vetenskaps- och Vitterhets-Samhällses Handlingar, Series 5B, 5(1): 1–196.
- Lokšina IE, Golovatch SI (1979) Diplopoda of the USSR fauna. Pedobiologia 19: 381–389.
- Muralewicz WS (1907) Zur Myriapodenfauna des Kaukasus. Zoologischer Anzeiger 31(11/12): 329–351.
- Muralewicz WS (1911) [Essay on the fauna of Myriopoda of the Caucasus]. Mitteilungen des Kaukasischen Museums [Izvestiya Kavkazskogo muzeya] 5: 1–80. [in Russian]
- Muralewicz WS (1913) [Arthropods.—Arthropoda. Chapter VII. Class of myriapods.—Myriapoda]. Proceedings of the Society for the Study of the Black Sea Coast [Trudy Obshchestva izucheniya Chernomorskogo poberezhya] 2: 215–221. [in Russian]
- Muralewicz WS (1927) [To the fauna of Myriopoda of the Caucasus]. Acta Societatis Entomologicae Stauropolitanae [Izvestiya Stavropol'skogo entomologicheskogo obshchestva] 3(1): 1–7. [in Russian]
- Read H (1992) The genus *Cylindroiulus* Verhoeff 1894 in the faunas of the Caucasus, Turkey and Iran (Myriapoda: Diplopoda: Julidae). Senckenbergiana biologica 72(4/6): 373–433.
- Short M (2015) New records of *Lophoproctus coecus* Pocock, 1894 (Diplopoda, Polyxenida, Lophoproctidae) extend the range of the genus *Lophoproctus*. ZooKeys 510: 209–222. <https://doi.org/10.3897/zookeys.510.8668>
- Short M, Vahtera V, Golovatch SI (2018) The millipede order Polyxenida in the faunas of the Crimean Peninsula and Caucasus, with notes on some other Polyxenidae. Zootaxa.
- Talikadze DA (1984) [On the millipede fauna (Diplopoda) of the Colchidan Province of the Caucasus]. Zoologicheskii Zhurnal 63(1): 124–145. [in Russian, a summary in English]
- Timotheew TE (1897) [Two new species of Diplopoda]. Bulletin de la Société des Naturalistes à l'Université Impériale de Kharkov [Trudy Obshchestva ispytateley prirody pri Imperatorskom Kharkovskom Universitete] 31: 1–19. [in Russian]
- Turbanov IS, Palatov DM, Golovatch SI (2016) The state of the art of biospeleology in Russia and other countries of the former Soviet Union: A review of cave (endogean) invertebrate fauna. 2. Arachnida – Acknowledgements. Zoologicheskii Zhurnal 95(11): 1283–1304. [in Russian; English translation: Entomological Review, 2016, 96(9): 1297–1333]. <https://doi.org/10.7868/S0044513416110064>

- Vagalinski B, Lazányi E (2018) Revision of the millipede tribe Brachyiulini Verhoeff, 1909 (Diplopoda: Julida: Julidae) with description of new taxa. Zootaxa.
- Verhoeff KW (1921) Chilognathen-Studien (91. Diplopoden-Aufsatz). Archiv für Naturgeschichte 85A: 23–80.
- Verhoeff KW (1930) Eine cavernicole Symphyognathen-Gattung aus Georgien, Transcaucasien. Mitteilungen über Höhlen- und Karstforschung 1930: 33–39.
- Victor T (1839) Insectes du Caucase et des provinces Transcaucasiennes recueillis et décrits par T. Victor. Bulletin de la Société Impériale des Naturalistes de Moscou 12: 44–67.
- Zuev RV (2014) Preliminary data on the millipedes (Diplopoda) from the Stavropol Territory, northern Caucasus, Russia. Arthropoda Selecta 23(4): 347–354. http://kmkjournals.com/upload/PDF/ArthropodaSelecta/23/23_4_347_354_Zuev_for_Inet.pdf
- Zuev RV (2017) The relict millipede *Hirudicryptus abchasicus* Golovatch, Evsyukov et Reip, 2015, new to the fauna of Russia (Diplopoda: Siphonocryptida: Siphonocryptidae). Russian Entomological Journal 26(3): 281–286.