



# HHS Public Access

Author manuscript

*Front Virol.* Author manuscript; available in PMC 2022 February 08.

Published in final edited form as:

*Front Virol.* 2021 November ; 1: . doi:10.3389/fviro.2021.753366.

## Oncogenic Viruses as Entropic Drivers of Cancer Evolution

Italo Tempera, Paul M. Lieberman\*

Program in Gene Expression and Regulation, The Wistar Institute, Philadelphia, PA, United States

### Abstract

Viral infection is an indisputable causal factor for nearly 17% of all human cancers. However, the diversity and complexity of oncogenic mechanisms raises new questions as to the mechanistic role of viruses in cancer. Classical viral oncogenes have been identified for all tumor-associated viruses. These oncogenes can have multiple oncogenic activities that may or may not be utilized in a particular tumor cell. In addition, stochastic events, like viral mutation and integration, as well as heritable host susceptibilities and immune deficiencies are also implicated in tumorigenesis. A more contemporary view of tumor biology highlights the importance of evolutionary forces that select for phenotypes better adapted to a complex and changing environment. Given the challenges of prioritizing singular mechanistic causes, it may be necessary to integrate concepts from evolutionary theory and systems biology to better understand viral cancer-driving forces. Here, we propose that viral infection provides a biological “entropy” that increases genetic variation and phenotypic plasticity, accelerating the main driving forces of cancer cell evolution. Viruses can also influence the evolutionary selection criteria by altering the tumor microenvironment and immune signaling. Utilizing concepts from cancer cell evolution, population genetics, thermodynamics, and systems biology may provide new perspectives on viral oncogenesis and identify novel therapeutic strategies for treating viruses and cancer.

### Keywords

KSHV; plasticity; hepatitis B virus; epigenetic; cancer; Merkel cancer; HPV–human papillomavirus; EBV–Epstein-Barr virus

---

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

\*Correspondence: Paul M. Lieberman [lieberman@wistar.org](mailto:lieberman@wistar.org).

#### AUTHOR CONTRIBUTIONS

PL and IT contributed to the conception, writing, figure preparation, and editing of the manuscript. Both authors contributed to the article and approved the submitted version.

**Conflict of Interest:** PL is a founder and consultant for Vironika, LLC.

The remaining author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## INTRODUCTION

Viruses have well-established causal roles in numerous human and animal cancers, collectively responsible for almost one fifth of all cancers (1, 2). Viral associated cancers are a special case of cancer biology and virology. To date, there are seven human viruses with strong epidemiological links to human cancers. These include members of the high-risk human papillomavirus (HPVs), hepatitis viruses B and C (HBV and HCV), human gammaherpesviruses (HHV4/Epstein-Barr Virus (EBV) and HHV8/Kaposi's Sarcoma-Associated Herpesvirus (KSHV), Merkel cell polyomavirus (MCPyV), and human T-cell leukemia virus I (HTLV-1). These oncoviruses represent members of vastly different families of virus, including DNA, RNA and retroviridae. Despite this species diversity, these oncoviruses are thought to share common features that enable them to drive cancer. Oncoviruses usurp key cellular pathways important for the control of cell growth and metabolism. However, many non-cancer-causing viruses perturb these pathways and have similar viral-host interactions. Consequently, it is not fully understood what features confer viruses with oncogenic potential in human populations.

Oncogenic viruses perturb numerous cellular pathways described as the hallmarks of cancers (3, 4). As expected, viral-associated cancer pathways can be readily superimposed on these cancer hallmarks (5) (Figure 1). And while the pathways of viral carcinogenesis are ultimately cellular, viruses do provide foreign genomes and gene products that create new interactions and pathways for oncogenesis. How do these viral products and viral-specific pathways work coordinately over time to overcome the many barriers to cellular carcinogenesis? What makes these seven viruses different from their non-oncogenic relatives? Here, we suggest that oncogenic viruses are unique in their ability to increase the adaptability and evolvability of infected cells, and that multiple perturbations over time enable formation of cancer cell fate choices. We suggest that a more in-depth knowledge of virus-host interactions over the time-course of cancer evolution will provide a more complete understanding of viral oncogenesis.

## A PLETHORA OF ONCOGENIC MECHANISMS

A major challenge in the field of viral oncology, and cancer biology in general, is the very large number of mechanisms and pathways that contribute to carcinogenesis. While early studies focused on one or a few viral oncogenes, we now know that viruses can promote cancers through a much greater diversity of mechanisms and pathways. Many of these mechanisms can be well-defined in the context of a particular tumor type or tissue environment. However, the diversity of mechanisms confounds the identification of single causal agent or event. Given the abundance of potential and actual mechanisms, a new challenge arises to both understand the impact of each oncogenic event and the accumulation of multiple distinct oncogenic events over time or in the context of a transient stress challenge. Here, we provide a brief overview of some classic mechanisms of viral oncogenesis, highlighting many from the human gammaherpesviruses, and argue that the multiplicity of these mechanisms makes sense only in the context of cancer cell evolution and systems biology.

## Classical Viral Oncogenes Are Not Sufficient for Cancer

Viruses have been shown to induce tumor formation through numerous and diverse mechanisms (5). From the pioneering studies of Peyton Rous in 1920s, retroviruses were found to encode oncogenes, like v-src, v-myc, v-ras, which could potentially transform normal cells to grow as tumors in animal models (6). Several of these viral oncogenes were subsequently recognized as host genes captured by viral recombination and their oncogenicity due to combinations of activating mutations and deregulated expression in infected cancer cells. DNA tumor viruses were found to encode novel oncoproteins, such as SV40 T-antigen, adenovirus E1A and E1B, and papillomavirus E6 and E7 that interact with and disable cellular tumor suppressor proteins, such as p53 and Rb (7). While these viral oncogenes may be necessary for transformation in laboratory models, they are not sufficient for tumor formation in the overwhelming majority of natural infections. In addition, some viruses, such as adenovirus, encode potent inhibitors of cellular tumor suppressors p53 and Rb, but rarely associate with human cancers (8). This is consistent with the observation that most oncogenic viruses typically cause benign infections, that only rarely lead to cancer.

## Multiple Host Targets of Viral Oncogenes

Pioneering studies identified a few cellular targets universally exploited by tumor viruses, such as p53 DNA damage surveillance and Rb cell cycle control. However, it is now known that subversion of these targets are not sufficient for viral tumorigenesis, and that additional and alternative host proteins and pathways are targeted by viral oncogenes (Figure 1). For example, the small viral oncoprotein E7 is well known to bind and degrade Rb (9), but can also interact with the Rb-associated DREAM complex (10), phosphatase PTPN1(11), histone modifying enzyme HDACs (12), stem cell promoting factors APH1B and OCT4 (13, 14), and Cullin2 to stabilize APOBEC3a (15). In addition, E7 can cooperate with another viral oncoprotein E6 to activate hTERT to overcome replicative senescence (16, 17). Viral subtypes, as well as host cell type can determine whether these different interactions are oncogenic, further demonstrating the diversity of targets for one small viral oncoprotein and its potential effects on different cancer pathways. Similarly, the MCPyV small T antigen can interact with MYCL and EP400 to alter chromatin and transcription regulatory networks implicated in cell lineage control (18), 4EBP1 affecting translational control (19), protein phosphatase 2A to affect ubiquitin ligases (20), the F-box proteins FBW7 to activate NF- $\kappa$ B signaling (21, 22), as well as binding iron-sulfur clusters (23). This promiscuous multitasking is likely to be a general feature of viral oncoproteins that target multiple cellular proteins and pathways.

Larger DNA tumor viruses, such as EBV and KSHV, encode numerous viral genes implicated in oncogenesis. Many of these oncogenes target pathways important for the tissue-specific functions of the host cell. EBV encodes two membrane oncoproteins, LMP1 and LMP2, that cooperate to immortalize primary B-lymphocytes by mimicking B-cell receptor and CD40 co-receptor (24, 25). LMP1 interacts with multiple TRAFs and TRADDs to activate NF- $\kappa$ B pathways (26), but can also interact with other proteins involved in membrane vesicle formation (27). LMP2 can interact with several different src-family kinases (28). EBV also encodes 6 nuclear antigens, EBNA-LP, 1, 2, 3a, 3b, and 3C, that are all implicated in oncogenic mechanisms (29). EBNA2 is absolutely required for

B-cell immortalization *in vitro* (30), and natural mutations in EBNA2 correlate with B-cell transformation activity (31). However, some EBV associated tumors fail to consistently express EBNA2 and LMPs. Like EBV, KSHV also encodes candidate oncogenes, including the nuclear antigen LANA that can bind host chromatin and alter p53 and Rb function, vGPCR that can induce endothelial tumors in transgenic mouse models, vCyclin that can drive cell cycle progression, vFLIP and K12 that can activate NF- $\kappa$ B and STAT3 signaling (32). In addition to these viral encoded proteins, both EBV and KSHV have numerous non-coding RNAs implicated in oncogenesis. EBV small non-coding RNA EBERs can interact with transcription factor PAX5 (33), ribosomal protein L22 (34), TLR receptors (35), and provide paracrine signals through exosome transmission (36). Numerous viral miRNAs target oncogenic pathways implicated in EBV and KSHV carcinogenesis (37, 38). EBV miRNAs are highly expressed in tumors, especially EBV-associated gastric carcinomas (EBVaGC) along with other non-coding RNAs that arise from the same genomic locations (BARTs) and have additional oncogenic potential (39, 40). Remarkably, these viral genes are expressed at variable levels and heterogeneously in most viral-associated cancers, further confounding the problem of complexity and diversity of viral oncogenic mechanisms.

### A Growing List of Viral Oncogenic Mechanisms

**Inhibition of Apoptosis**—Resistance to programmed cell death, particularly apoptosis, is among the most fundamental hallmarks of cancer and viral infection. Viruses provide numerous mechanisms to resist apoptosis (41–43). For example, EBV encodes two viral proteins, BHRF1 and BALF1, dedicated to inhibition of the Bcl2 family of pro-apoptotic factors. BHRF1 and BALF1 have some overlapping, but not completely redundant activities in the inhibition of programmed cell death mediated by Bcl2 (44, 45). In addition, EBV encodes multiple miRNAs that target the pro-apoptotic genes BIM (46) and Puma (47). KSHV also encodes numerous genes directed at the disruption of apoptosis (42), including a Bcl2 homologue ORF16 that may have mitochondrial and nuclear functions required for viral reactivation and lytic replication (48). MCPyV large T protein enhances BIRC5/survivin mRNA and protein expression to prevent caspase-mediated apoptosis (49). HBV HbX protein has been shown to have a BH3-like domain that interacts with Bcl2 and Bcl-xL to prevent apoptosis during viral replication (50). HCV non-structural protein NS5A can attenuate apoptosis by enhancing GRP78 expression and reducing ER-stress (51). HTLV-1 Tax suppresses transcription of pro-apoptotic genes Bid and Bim, while activating expression of pro-survival Bcl-2 members (52). In general, oncoviruses demonstrate numerous and diverse anti-apoptotic mechanisms, often encoding multiple, partially redundant viral genes that may be expressed heterogeneously in tumors.

**Reprogramming Host Metabolism**—Cancer cells frequently undergo a metabolic shift to aerobic glycolysis (Warburg effect) and utilize alternative metabolites, such as glutamine and serine for energy production and macromolecular biosynthesis (53). Oncogenic viruses can reprogram cellular metabolism in various ways (54). Overexpression of both HPV16 E6 and E7 promote glucose metabolism through activation of glucose transporter 1 (55). E6 was shown to stabilize HIF1A induced Warburg effect during hypoxia in keratinocytes (56, 57). E7 can bind and inhibit pyruvate kinase M2 to promote glycolysis (58). MCPyV small T has been shown to increase glucose consumption and lactate production indicative

of Warburg aerobic glycolysis (59). These changes correlate with transcriptomic changes in hypoxia, AMPK activation, and mTOR signaling. EBV infection of resting B-cells induces a hyperproliferative state that is rate limited by nucleotide metabolism (60), and EBNA2 activates a myc-dependent metabolic program to increase amino acid and nucleotide metabolism during hyperproliferation (61). EBNA2 also activates SREBP2 to promote lipid biosynthesis and fatty acid metabolism (62). KSHV miRNAs induce a metabolic shift from OXPHOS to glycolysis (63). Thus, oncogenic viruses shift cellular metabolism through multiple factors and pathways to promote infected cell fitness, similar to cancer cells.

**Modulation of the Cellular Microenvironment**—Viral-infected cells and associated tumors thrive in harsh microenvironments that reinforce viral-infected and cancer cell selection, requiring Warburg metabolism and adaptation to low oxygen and acidification (64, 65). Variations in oxygenation due to competitive crowding or vascular insufficiency can have dramatic effects on viral gene expression and cellular stress response (65). Hypoxia plays a central role in KS tumorigenesis and regulating KSHV latency (66, 67). Hypoxia inducible factors (HIF1A) can modulate KSHV oncogenes, including vIL6 and vGPCR to upregulate VEGF and angiogenesis (68, 69). Hypoxia has a strong immunosuppressive effect (70), and oncogenic viruses may induce pseudohypoxia to escape immune recognition (71). Viral induced Warburg effect can also be immunosuppressive by competing with immune cells for glucose and oxygen consumption (72). Hypoxia is known to induce EBV lytic cycle genes that have pro-survival and immune modulatory functions. EBV BRLF1 can inhibit interferon response genes IRF3 and IRF7 and interferon production (73) while BZLF1 can inhibit interferon gamma and TNF $\beta$  signaling pathways (74, 75). EBV immediate early protein BZLF1 can also induce SOCS3 to inhibit cytokine signaling (76), and the viral kinase BGLF4 can degrade TLR9 mRNA (77). Oncogenic viruses also reprogram the tumor microenvironment through production of extracellular vesicles (EV) that transmit cargo to neighboring cells (63, 78). EBV positive NPC produce EVs that transfer viral miRNAs and oncoproteins, such as LMP1, to neighboring cells (36, 79) causing a microenvironment selective for infected cell persistence (80) and immune suppression through recruitment of regulatory T-cells (81). Thus, virus infection can alter the microenvironment to promote selection of viral-infected cells that survive at the expense of uninfected and non-transformed cells.

**Attenuation of Host Immune Control**—In Darwinian terms, viruses and cancer are most limited by the predatory function of the immune system. Viral associated cancers are particularly adept at modulating immune surveillance, and are most virulent in immunosuppressed conditions, such as HIV-AIDS and solid-organ transplants. HPV and EBV associated tumors upregulate T-cell checkpoint proteins PD-L1 and PD-L2, as well as the CTLA-4 immunosuppressive pathways (82). Multiple different mechanisms act on this pathway. For EBV, EBNA2 can down regulate miR-34a to upregulation of PD-L1 in lymphoid cancers (83). LMP1 activates PD-L1 through interferon gamma pathway in NPC (84). LMP1 activates PD-L1 through NF- $\kappa$ B pathway in NKTCL (85). HPV E5 protein suppresses HLA expression and immune recognition of infected tumor cells, rendering them resistant to checkpoint immunotherapy (86). HBV sAg binds to SIGLEC-3 (CD33) on myeloid cells to induce immunosuppression (87). HCV core protein interaction with

cellular gC1qR can modulate macrophage cytokines to restrict immune targeting to HCC (88). Clearly, diverse and novel viral mechanisms function to disable the immunological barriers to cancer.

**Transcriptional Reprogramming**—Perturbations in transcription factor and gene regulatory networks are also hallmark changes in cancer. Viral immediate-early genes and oncogenes frequently target transcription factors and transcription factor networks that are fundamental to host cell differentiation and identity. E6, E7, and T-antigens are known to interact with cellular transcription factors, such as p53 and the Rb-family complexes. EBV and KSHV encode numerous other nuclear factors that alter transcription control, and these viral factors have been implicated in viral carcinogenesis. All six EBV-encoded EBNA interact with host transcriptional regulators to perturb regulatory networks in distinct and complex ways. EBNA2 can interaction with B-cell regulatory factors RBPI, EBF1, RUNX1, and PU.1 to affect cooperative DNA binding site selection (89) as well as facilitate formation of super-enhancers, such as found at the cMyc locus (90, 91). EBNA1 can activate transcription of the EBNA2 gene, while EBNA2 can auto-activate its own transcription along with that of EBNA3Cs and LMPs to change the EBV viral gene regulatory network. Similarly, these factors cooperate to regulate expression of host cell genes including the repression of tumor suppressor genes, like BIM and p16, by EBNA3C (92, 93). In contrast, KSHV encodes one major nuclear protein LANA that can affect viral and cellular transcription and chromatin structure through multiple mechanisms including direct binding to core histones H2A/H2B through its N-terminus and to GC-rich DNA through its C-terminal domain (94, 95). KSHV also encode Interferon Regulatory Factors (vIRFs) that can alter transcriptional control of cellular IRFs, but these viral factors are not typically expressed in most KSHV-associated tumor cells, unless lytic reactivation occurs (96). EBV and KSHV miRNAs and longer non-coding RNAs can also impact cellular and viral regulatory networks (97). All other oncogenic viruses have similar perturbations in host gene regulation. HBV alters the miRNA-mRNA regulatory network in HCC (98). HPV has a distinct viral gene network signature in HPV-positive head and neck squamous cell carcinoma (99). The complexity of gene regulatory networks and diversity of viral mechanisms for disrupting these networks reveals the challenges of pinpointing a single or primary causal factor.

**Epigenomic Reprogramming**—Epigenetic modifications represent an important mode of adaptive and heritable gene regulation. Persistent viral infection can impact host epigenomes in diverse ways. DNA infection in the nucleus can alter the DNA methylation patterns, frequently resulting in hypermethylation of viral and cellular genes (100). Host genome hypermethylation is detected in HPV and EBV associated carcinomas (100). EBV infection can induce host hypermethylation in a number of experimental models, including non-neoplastic gastric epithelial cells (101), telomerase immortalized keratinocytes (102), and gastric carcinoma derived AGS cells (103). Hypermethylation has been correlated with the inactivation of the dioxygenases TET1 (104) and TET2 (105) that play a role in active demethylation. For EBV, LMP1 and LMP2 can induce DNMT1 expression and subsequent methylation of cellular genes for CDH1 (E-cadherin)(106–108) and tumor suppressors p16 and p21 (109, 110). HPV is also found to alter the host epigenome,

including DNA methylation and histone modification patterning (111). Other mechanisms for host epigenetic modification in response to foreign DNA have been described, including modulation of the Sting (112) and Apobec (113) pathways.

Viruses can also induce changes in host chromosome conformation leading to the rewiring of gene regulatory circuits. EBNA-LP, EBNA2, and EBNA3C have been implicated in the reorganization of DNA regulatory loops to form super-enhancers regulating cellular oncogenes, such as *c-myc*, to drive resting B-lymphocytes into proliferating and immortalized lymphoblastoid cells (91, 114). EBNA2 has been shown to interact cooperatively with several cellular transcription factors, including EBF1 and RBPJ (89), and RUNX1 (115), and this cooperativity may explain some of the capacity to facilitate new DNA-DNA loop interactions. Episomal viruses may also influence host chromatin and histone modifications through chromosome tethering mechanism. For example, one study found EBV tethering to reinforce heterochromatic H3K9me3 silencing of neuronal genes in EBV positive BL cells (116). Another study found that EBV genome tethering caused transition of heterochromatic H3K9me3 to euchromatic H3K4me3 along with transcriptional activation of cancer-related genes in EBVaGC (117). EBV genomes were also found to transit within open chromosome territories during the switch to reactivation (118). In contrast to viral integrations, episomal tethering may be dynamic over time and provide epigenetic plasticity to both virus and host.

## COMMON THREADS: ONCOVIRAL PERSISTENCE AND PLASTICITY

### Viral Persistence at Tumor Sites

All known human tumor viruses persist, in one form or another, at the site of tumor formation. Tumor viruses can persist as chronic infections (HCV), nuclear episomes (HBV, HPV, MCPyV, EBV, KSHV) or integrated genomes in viral-associated tumor cells (HTLV-1). HCV is an unusual oncovirus in that it does not infect the cancer cell, but its long-term persistence causes inflammation conducive to cancer cell emergence. In contrast, episomal DNA tumor viruses, like EBV, KSHV and HPV have viral-specific programs dedicated to viral genome persistence in a dividing cell that can serve as a clonal outgrowth in cancer (119–121). These viruses encode proteins, such as EBNA1, LANA, and E2, dedicated to binding viral DNA and maintaining the viral genome over generations in proliferating cells. HTLV-1 persists through integration into the host genome as part of its normal life cycle (122). Integration is inherently mutagenic and there is evidence that some, albeit rare integrations are oncogenic.

Chang and Moore proposed that viral cancers arise due to aberrations in the normal productive life cycle, including genetic mutations and integrations that disrupt normal viral gene expression (123). Consistent with this, aberrant integrations are frequently observed for oncoviruses that typically persist as extrachromosomal episomes, such as HBV, HPV, and MCPyV. HBV has been found to integrate in oncogene hot-spots, such as the *TERT* (telomerase) or *KMT2B* (*MLL4*) loci (124), and HPV has been found to integrate at *ERBB2* and *PTPN13* loci (125). In each case, integration alters normal gene regulation to promote oncogenesis. Integrations can also lead to loss of viral DNA and deregulation of viral oncogenes. HPV integration with loss of viral E2 repressor protein leads to the

upregulation of viral oncogenes E6 and E7 in cervical carcinomas (126, 127). In Merkel cell carcinomas, MCPyV frequently integrates as incomplete genomes with deletions in large T and overexpression of small T (18, 128). There is some evidence that viruses can transform host cells without viral genome persistence through heritable changes in the host epigenome or regulome (129). Such “hit and run” mechanisms have been reported but are difficult to demonstrate in naturally occurring human cancers. Thus, most viral cancers are associated with long-term persistence of viral genetic material transmitted for multiple cellular generations.

### **Viral and Host Heterogeneity as Oncogenic Drivers**

Tumor cell heterogeneity is a central hallmark of cancer (130–134). Viruses can be highly heterogeneous, as well as induce a more heterogeneous phenotype in the host cells. Viral gene programs are inherently unstable with potential shifts from productive to non-productive, or latent to lytic infection cycles with variable gene expression patterns. Viral genomes may be more relaxed than host chromatin and free to move from one chromosome compartment to another. Viral genes may be activated or repressed with greater flexibility, including sporadic bursts of lytic amplification and gene expression. Oncoviruses can also increase variability in host gene expression. Viral infected tumor cells maintain a poorly differentiated state, and may toggle between the lympho-epithelial features of EBV NPC and GC, and the mesenchymal-endothelial features of KSHV infected KS spindle cells (135). This loss of fixed cell identity has been referred to as cellular plasticity. Increasing cellular plasticity provides cancer cells with the advantage of increase variation, or the capacity to adapt more rapidly to changing environmental conditions (such as hypoxia), relocate to new niches (metastasize), evade immune surveillance, and develop drug resistance (136).

Viruses can be highly heterogenous and mutate during the course of infection to increase carcinogenic risk. High and low risk subtypes of HPV may be considered a form of species heterogeneity, while integrations and deletions can be a source of mutational variation (123, 137). Tumor heterogeneity based on viral and host gene expression (including single cell RNAseq), epigenetic modifications, and immune infiltration have been observed for cancers associated with HPV (138), MCPyV (139), HBV (140), HCV (141), and HTLV-1 (142), indicating that such heterogeneity is a general rule for viral and non-viral cancers. Variations in viral gene expression and genome copy number may also account for cancer-risk. For EBV, viral latency types can contribute to genetic heterogeneity and plasticity. EBV can adopt different latency types in different host cells and tumor types. Epigenetic factors are known to regulate the different latency types, including differences in DNA methylation patterning. Among the viral genes with variable expression is the potent oncogene LMP1, involved in constitutive TNF-pathway signaling. LMP1 gene can be expressed at ranges that vary 100 fold among single cells in a population (143). Variations in LMP1 expression in NPC correlated with cellular genes linked to E2F and cellular DNA replication, as well as to changes in NF-kB and JAK/STAT signaling (144). Another variable in viral gene expression is the genome copy number (145). Viral episomes can range from 1 to several 100 copies per cell. Viral genome amplification is closely linked with over-expression of some viral genes, especially the lytic cycle genes of the gammaherpesviruses. Although a complete lytic cycle is rarely detected in viral tumors, abortive lytic cycle gene expression is likely to contribute



to viral carcinogenesis. These lytic cycle genes provide numerous potential contributions to viral carcinogenesis, including anti-apoptotic, immunomodulatory, and paracrine activities required for tumorigenesis (146).

### Temporal Heterogeneity and Cancer Cell Evolution

Temporal history of viral infection may also contribute to tumor heterogeneity (Figure 2). Viral cancers are often clonal expansions of a virally infected progenitor cell (147, 148). However, the progenitor cell is likely to have acquired somatic mutations or other aberrations enabling viral transformation. Pre-existing somatic cell mutations appear to be required for the formation of EBV-associated NPC (149). EBV infected nasopharyngeal cells undergo cell cycle arrest, unless infected cells have pre-existing mutations in cell cycle control. Gene loss from 9p21 to 3p21.3 (inactivating RASSF1A and CDKN2A1, respectively), and activation of telomerase, have been found to precede acquisition of EBV infection in the evolution of NPC (150). EBV infection further drives clonal expansion of infected cells, as has been demonstrated in NPC by examining the uniformity of the viral terminal repeat DNA (148). Subsequent genomic hypermethylation follows the infection of EBV, and then activation of NF- $\kappa$ B pathways, loss of MHC I, mutations in PI3K/MAPK, and chromatin remodeling, and subsequently TP53 and RAS, along with other mutations (151) are frequently observed in the course of NPC formation (Figure 3).

Host genetic variation, whether inherited or acquired through somatic mutation, also contribute to the risk of viral cancers. Variations in the HLA locus correlate with risk of viral cancers, suggesting that presentation of viral antigens plays a key role in immune resistance to viral-driven cancers. Genetic analysis of NPC susceptibility revealed risk loci at hTERT, CDKN2A/B, MECOM, and TNFRSF19, all of which have known roles in oncogenic pathways. Other susceptibility pathways have been linked to Notch signaling, magnesium transport (NIPAL1), EBV entry into epithelial cells (ITGB6), modulation of apoptosis (NEDD4L, BCL2L12), cAMP signaling, or DNA repair (MLH1, PRKDC) (153). Inherited mutations in magnesium channel MAGT, as found in XMEN syndrome is associated with defects in NK and T-cell control of EBV infected B-cell (154, 155), and may also contribute to risk of NPC and KSHV associated KS.

Other aberrations also contribute to variations in viral-host interaction. Rare tumors of atypical tissue types, such as EBV leiomyosarcomas, NK-T cell lymphoma, peripheral T-cell lymphoma (156) and pulmonary lympho-epithelium-like carcinoma (LELC) (157) are likely due to aberrant entry of EBV into unnatural host cells. Environmental factors, such as coinfection with HIV or malaria, can alter the immune control of viral infected cells. Thus, host cell type, immune functionality and other environmental factors can impact the course of infection and cancer progression in a temporal-dependent manner.

Stage specific effects on viral oncogenesis are observed for most oncoviruses. HPV has been shown to have different patterns of infection, integration, and gene expression at different stages of viral-associated cancers (158). Similarly, MCPyV T-antigen was found to have stage specific tumor promoting activity in a mouse model treated with defined carcinogens (159). HTLV-1 antigen expression changes in response to T-cell activity, providing evidence for viral adaptation and co-evolution with tumor cell progression (160). HBV X-protein

interaction with miRNA production impacts multiple stages of HCC through very different pathways, ranging from cell cycle control at early stages to immune suppression at later stages (161). These multifunctional viral oncoproteins can affect stage specific events in cancer cell evolution, and therefore may adapt their oncogenic activities over time.

## SYSTEMS APPROACHES TO VIRAL ONCOLOGY

### Virus-Driven Gene Regulatory Networks and Attractor States

Oncogenic viruses increase the number of possible ways a cell can propagate, survive and achieve oncogenic transformation. In terms of Darwinian evolutionary dynamics and population genetics, virus infection increases the phenotypic diversity and fitness of the population. A more heterogeneous population has a fitness advantage by adapting more rapidly to changing and stressed environments. To borrow from systems biology, a stable phenotype requires a stable gene regulatory network (GRN). GRNs are considered a thermodynamic “attractor” state (162) (Figure 4). GRNs are related to the developmental states described by Waddington (164) using an energy landscape and the canalization patterns that separate two or more distinct cell fates or GRNs. It has been proposed that cancer cells converge on a common GRN attractor state akin to the embryonic and unicellular cell states (165). Oncogenic viruses perturb major hubs in GRNs enabling greater plasticity between phenotypic states (136). Experimental validation of this concept has been provided by measuring the intrinsic plasticity of EBV positive Burkitt lymphoma cell lines (162). Viruses can also increase signal noise in a GRN [reviewed in (166)]. Viral genomes may have inherently higher “noise” than their cellular counterparts due, in part, to their relaxed epigenetic regulation, subcellular localizations, and copy number variations. Viral genomes and gene products destabilize GRNs and facilitate the transition from one attractor state to another (167). Thus, we propose that a major feature of oncogenic viruses is their ability to accelerate the rates of cancer cell evolution by increasing the genetic variability and phenotypic plasticity, and inherent cellular adaptability to changing and stressful microenvironments (168, 169).

### Viruses as “Entropic” Drivers of Cancer Evolution

In thermodynamics and statistical mechanics, terms like entropy were developed to explain the behavior of complex systems with excessively large numbers (ensembles) of microstates. In this respect, the term entropy can be used to describe the number of microstates of a complex biological system, such as the gene regulatory interactions in a viral infected tumor cell. While we can not provide a rigorous definition of biological entropy, we do suggest that viruses increase the number of possible microstates available to the host cell, and therefore may be considered a form of “genetic entropy.” In the most simplistic terms, genetic entropy may be the ability to reconfigure the genome and its programmed processes. Viruses reconfigure genomes, gene expression programs, and biochemical pathways. We further suggest that this be considered in terms of Shannon information theory, where viral genomes may be considered a source of signal noise enabling the freedom to find a lower energy state, or alternative GRN, such as the oncogenic state. Viruses can increase signal noise and alternative outputs for cellular developmental programs, and this enables transcriptional plasticity and phenotype heterogeneity associated with tumorigenesis. Efforts to quantify

cellular information and thermodynamic entropy may be useful for understanding the emergence of the cancer phenotype. In this regard, virus infection may be considered an entropic driving force for cancer (170).

### Implications for Cancer Therapy

The diversity of oncogenic mechanisms and the plasticity of cancer cells raise enormous challenges in developing precision therapies. For most cancers, early detection provides the best opportunity for effective treatment. Viral cancers have the advantage of having viral-specific targets and biomarkers. For example, early-stage NPC can be predicted from cell-free EBV DNA in plasma (171) and EBV-specific IgA (172) and can be effectively treated with radiation. However, most cancers are discovered at later stages and fail treatment or develop resistance and recurrence. This is largely attributed to tumor heterogeneity and plasticity and the emergence of resistant clonal populations. Oncogenic viruses provide a rich resource for tumor cell heterogeneity and evolutionary diversity. Elimination of persistent oncogenic viruses at early stages is ideal, but not always possible. Reduction of genetic plasticity and modulation of selection pressure may be attractive alternative approaches for treatment of viral cancers. It may also be possible to use evolutionary principles to improve dosing and timing of therapy. Additionally, it may be possible to exploit viral genetic plasticity to eliminate viral cancers. Since excessive genetic variability can be incompatible with life (173), it may be possible to amplify viral-induced plasticity through drug intervention. Treatments that increase gene regulatory noise, such as epigenetic modifiers, could provoke chaotic and lethal gene expression patterns in cancer cells that would be resisted by normal cells.

## CONCLUSIONS

Viruses are thought to be simple biological systems, yet their contributions to cancer can be fiendishly complex. If we return to the question of why some viruses cause cancers but not closely related others, we have only a partial answer. Among the common features, is that of long-term persistence of tumor virus in localized tissue compartments. Another common feature, as highlighted by Chang and Moore, is the frozen accident of the defective virus entering the wrong cell type, or cell with precancerous mutation, or acquired mutation in host or virus that blocks the natural infection and immune clearance. Host and virus genetic variations can be susceptibility factors that enable viral oncogenesis. One additional feature is the plasticity provided by chronic virus infection, and how that effects the survival options for infected tumor cells and tumor fields. Viruses co-opt and perturb numerous cellular pathways implicated in cancer. These perturbations may occur at different times (temporal heterogeneity) and in different subpopulations (spatial heterogeneity) and may be replaced by cellular oncogenic drivers at different stages of tumor evolution (interchangeability). Viral-specific cancer mechanisms may have unique features and provide new insights into cancer biology and genetic plasticity. Despite this complexity, viral cancers may be considered low hanging fruit for cancer therapeutic intervention. Prevention of virus infection and virus-specific inhibitors have been shown to diminish cancer risk, and immune targeting of viral proteins show clinical promise. Deeper understanding of the

basic mechanisms driving cancer cell evolution may also be required for more effective intervention.

Precision medicine requires discrete knowledge of the causal factors in disease. Identification of the specific drivers and treatment with selective drugs for each driver pathway is a reasonable, rational, and reductionist approach to cancer therapy. Viral cancers are likely to have different vulnerabilities than their non-viral counterparts, including mechanisms driving cellular plasticity and evolvability. Studying viral cancers may also help us to solve some of the key questions in cancer biology. What are the rate-limiting steps in cancer evolution? What are the most vulnerable nodes of a gene regulatory network for a particular type of cancer and how can we dampen genetic and environmental noise to reduce cancer cell plasticity and evolvability? Can we reverse tissue microenvironment conditions that preferentially select for cancer cell evolution? Ultimately, understanding the evolutionary and thermodynamic driving forces of virus infection and carcinogenesis will provide a more coherent conceptual framework for research and new avenues for therapy.

## FUNDING

PL received funding from NIH grants RO1 DE017336, RO1 CA117830, and RO1 CA093606. IT received funding from NIH grants R01AI130209, R01GM124449, and R21AI122035.

## GLOSSARY

<b>DREAM</b>	Dimerization partner (DP), Retinoblastoma (RB)-like, E2F and MuvB Complex
<b>EBV</b>	Epstein-Barr Virus
<b>EBVaGC</b>	Epstein-Barr Virus associated Gastric Carcinoma
<b>ER</b>	Endoplasmic reticulum
<b>EV</b>	Extracellular Vessicles
<b>GRN</b>	Gene Regulatory Network
<b>GPCR</b>	G-protein coupled receptor
<b>KSHV</b>	Kaposi's Sarcoma Associated Herpesvirus
<b>HSV</b>	Herpes Simplex Virus
<b>HTLV-1</b>	Human T-cell Leukemia Virus 1
<b>MCPyV</b>	Merkel Cell Polyomavirus
<b>NK</b>	Natural Killer
<b>HCC</b>	Hepatocellular carcinoma
<b>HDACs</b>	Histone Deacetylase
<b>HLA</b>	Human Leukocyte Antigen

<b>NPC</b>	Nasopharyngeal Carcinoma
<b>NKTCL</b>	Natural Killer-T Cell Lymphoma
<b>NS5A</b>	Non Structural Protein 5A (Influenza)
<b>OXPHOS</b>	Oxidative Phosphorylation
<b>SV40</b>	Simian Virus 40
<b>XMEN</b>	X-linked immuno-deficiency with magnesium defect, Epstein-Barr virus infection, and neoplasia

### **Viral Genes—EBV Genes:**

<b>BGLF4</b>	Kinase
<b>BALF1</b>	Anti-apoptotic
<b>BART</b>	Non-coding RNAs
<b>BHRF1</b>	Anti-apoptotic
<b>BZLF1</b>	Immediate early b-Zip protein
<b>EBNAs</b>	Epstein-Barr Nuclear Antigens (EBNA-LP, EBNA1, EBNA2, EBNA3A EBNA3B, EBNA3C)
<b>LMP1</b>	Latency Membrane Protein 1 (EBV oncogene)
<b>LMP2</b>	Latency Membrane Protein 1 (EBV oncogene)

### **KSHV Genes:**

<b>LANA</b>	Latency Associated Nuclear Antigen
<b>ORF16</b>	Open Reading Frame 16
<b>vIRFs</b>	Viral Interferon Regulatory Factors

### **Polyoma (SV40, MCPyV):**

<b>Tag</b>	Tumor Antigen
<b>LT</b>	Large T antigen
<b>ST</b>	Small T antigen

### **Adenovirus:**

<b>E1A</b>	E1B—Early 1A, 1B
------------	------------------

**HPV:**

<b>E5</b>	E6, E7—Early 5, 6, 7
<b>HBV</b>	HbX
<b>HTLV-1</b>	Tax
<b>HBZ</b>	Antisense B-zip protein

**Cellular Genes and Proteins:**

<b>APH1B</b>	Aph-1 Homolog B, Gamma-Secretase Subunit)
<b>APOBEC</b>	Apolipoprotein B mRNA Editing Catalytic protein
<b>AMPK</b>	Adenosine Monophosphate Kinase
<b>Bcl2</b>	Breakpoint cluster 2 (anti-apoptotic)
<b>Bcl-xL</b>	BCL2 like
<b>Bid</b>	BH3 Interacting Domain Death Agonist
<b>BIRC5</b>	Baculoviral IAP Repeat Containing 5/aka Survivin
<b>Bim</b>	Bcl2 like protein 11 (BCL211)
<b>CDKN2A1</b>	Cyclin-dependent kinase inhibitor 2A (aka p16)
<b>CTLA4</b>	Cytotoxic T-Lymphocyte Associated Protein 4
<b>DNMT</b>	DNA methyltransferase
<b>EP400</b>	E1A Binding Protein P400
<b>GRP</b>	Gastrin Releasing Peptide
<b>HIF1</b>	Hypoxia Inducible Factor 1
<b>IRF</b>	Interferon Regulatory Factor
<b>JAK/STAT</b>	Janus Kinase/Signal Transducer and Activator of Transcription
<b>KMT2B</b>	Lysine Methyl Transferase 2B
<b>NF-κB</b>	Nuclear Factor κB
<b>OCT4</b>	Octamer Binding Factor 4
<b>PD-L1</b>	Programmed death-ligand 1 (CD274)
<b>PI3K</b>	Phosphatidylinositol 4,5-bisphosphate 3-kinase
<b>PTPN</b>	Protein Tyrosine Phosphatase Non-Receptor Type

<b>PU.1</b>	Purine Rich Box Binding factor 1
<b>Puma</b>	p53 upregulated modulator of apoptosis
<b>P16</b>	Tumor suppressor (CDKN1A1)
<b>P53</b>	Tumor suppressor protein 53
<b>gC1qR</b>	Globular heads of the human C1q receptor
<b>MAGT</b>	Magnesium Transporter 1
<b>MAPK</b>	Mitogen-Activated Protein Kinase 1
<b>MLL</b>	Mixed Lymphocytic Leukemia
<b>Myc</b>	Myc Proto-oncogene
<b>MYCL</b>	Myc-like gene
<b>mTOR</b>	Mechanistic Target Of Rapamycin Kinase
<b>Ras</b>	Ras oncogene
<b>RASSF1A</b>	Ras Association Domain Family Member 1
<b>Rb</b>	Retinoblastoma Tumor suppressor gene
<b>RBPJ</b>	Recombination Signal Binding Protein For Immunoglobulin Kappa J Region
<b>RUNX1</b>	Runt-related transcription factor 1
<b>SIGLEC</b>	Sialic Acid Binding Ig Like Lectin 1
<b>SOCS2</b>	Suppressor Of Cytokine Signaling 2
<b>Src</b>	Rous sarcoma virus oncogene
<b>SREBP2</b>	Sterol regulatory element (SRE)-binding protein-2
<b>STING</b>	Stimulator of interferon <b>genes</b>
<b>TET</b>	Ten-Eleven Translocation (methylcytosine dioxygenase)
<b>TLR</b>	Toll-Like Receptor
<b>TNF</b>	Tumor Necrosis Factor
<b>TRAFF</b>	TNF receptor (TNFR) associated factor
<b>TRADD</b>	TNFRSF1A Associated Via Death Domain

## REFERENCES

1. Parkin DM. The global health burden of infection-associated cancers in the year 2002. *Int J Cancer*. (2006) 118:3030–44. doi: 10.1002/ijc.21731 [PubMed: 16404738]
2. de Martel C, Georges D, Bray F, Ferlay J, Clifford GM. Global burden of cancer attributable to infections in 2018: a worldwide incidence analysis. *Lancet Glob Health*. (2020) 8:e180–90. doi: 10.1016/S2214-109X(19)30488-7 [PubMed: 31862245]
3. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*. (2000) 100:57–70. doi: 10.1016/S0092-8674(00)81683-9 [PubMed: 10647931]
4. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. (2011) 144:646–74. doi: 10.1016/j.cell.2011.02.013 [PubMed: 21376230]
5. Mesri EA, Feitelson MA, Munger K. Human viral oncogenesis a cancer hallmarks analysis. *Cell Host Microbe*. (2014) 15:266–82. doi: 10.1016/j.chom.2014.02.011 [PubMed: 24629334]
6. Zur Hausen H. Cancers in humans: a lifelong search for contributions of infectious agents, autobiographic notes. *Annu Rev Virol*. (2019) 6:1–28. doi: 10.1146/annurev-virology-092818-015907 [PubMed: 31567062]
7. Pipas JM. DNA tumor viruses and their contributions to molecular biology. *J Virol*. (2019) 93:e01524–18. doi: 10.1128/JVI.01524-18 [PubMed: 30814278]
8. Lion T. Adenovirus persistence, reactivation, clinical management. *FEBS Lett*. (2019) 593:3571–82. doi: 10.1002/1873-3468.13576 [PubMed: 31411731]
9. Munger K, Scheffner M, Huibregtse JM, Howley PM. Interactions of HPV E6 and E7 oncoproteins with tumour suppressor gene products. *Cancer Surv*. (1992) 12:197–217 [PubMed: 1322242]
10. Rashid NN, Rothan HA, Yusoff MS. The association of mammalian DREAM complex and HPV16 E7 proteins. *Am J Cancer Res*. (2015) 5:3525–33. [PubMed: 26885443]
11. Hatterschide J, Bohidar AE, Grace M, Nulton TJ, Kim HW, Windle B, et al. . PTPN14 degradation by high-risk human papillomavirus E7 limits keratinocyte differentiation and contributes to HPV-mediated oncogenesis. *Proc Natl Acad Sci U S A*. (2019) 116:7033–42. doi: 10.1073/pnas.1819534116 [PubMed: 30894485]
12. Longworth MS, Laimins LA. The binding of histone deacetylases and the integrity of zinc finger-like motifs of the E7 protein are essential for the life cycle of human papillomavirus type 31. *J Virol*. (2004) 78:3533–41. doi: 10.1128/JVI.78.7.3533-3541.2004 [PubMed: 15016876]
13. Yang S, Chen T, Huang L, Xu S, Cao Z, Zhang S, et al. . High-risk human papillomavirus E7 maintains stemness *via* APH1B in cervical cancer stem-cell like cells. *Cancer Manag Res*. (2019) 11:9541–52. doi: 10.2147/CMAR.S194239 [PubMed: 31814758]
14. Panayiotou T, Michael S, Zaravinos A, Demirag E, Achilleos C, Strati K. Human papillomavirus E7 binds Oct4 and regulates its activity in HPV-associated cervical cancers. *PLoS Pathog*. (2020) 16:e1008468. doi: 10.1371/journal.ppat.1008468 [PubMed: 32298395]
15. Westrich JA, Warren CJ, Klausner MJ, Guo K, Liu CW, Santiago ML, et al. . Human papillomavirus 16 E7 stabilizes APOBEC3A protein by inhibiting cullin 2-dependent protein degradation. *J Virol*. (2018) 92:e01318–17. doi: 10.1128/JVI.01318-17 [PubMed: 29367246]
16. Liu X, Roberts J, Dakic A, Zhang Y, Schlegel R, HPV. E7 contributes to the telomerase activity of immortalized and tumorigenic cells and augments E6-induced hTERT promoter function. *Virology*. (2008) 375:611–23. doi: 10.1016/j.virol.2008.02.025 [PubMed: 18367227]
17. Katzenellenbogen RA. Activation of telomerase by HPVs. *Virus Res*. (2017) 231:50–5. doi: 10.1016/j.virusres.2016.11.003 [PubMed: 27863966]
18. Park DE, Cheng J, McGrath JP, Lim MY, Cushman C, Swanson SK, et al. . Merkel cell polyomavirus activates LSD1-mediated blockade of noncanonical BAF to regulate transformation and tumorigenesis. *Nat Cell Biol*. (2020) 22:603–15. doi: 10.1038/s41556-020-0503-2 [PubMed: 32284543]
19. Shuda M, Kwun HJ, Feng H, Chang Y, Moore PS. Human Merkel cell polyomavirus small T antigen is an oncoprotein targeting the 4E-BP1 translation regulator. *J Clin Invest*. (2011) 121:3623–34. doi: 10.1172/JCI46323 [PubMed: 21841310]



20. Kwun HJ, Shuda M, Camacho CJ, Gamper AM, Thant M, Chang Y, et al. Restricted protein phosphatase 2A targeting by Merkel cell polyomavirus small T antigen. *J Virol.* (2015) 89:4191–200. doi: 10.1128/JVI.00157-15 [PubMed: 25631078]
21. Nwogu N, Ortiz LE, Kwun HJ. Surface charge of Merkel cell polyomavirus small T antigen determines cell transformation through allosteric FBW7 WD40 domain targeting. *Oncogenesis.* (2020) 9:53. doi: 10.1038/s41389-020-0235-y [PubMed: 32427880]
22. Zhao J, Jia Y, Shen S, Kim J, Wang X, Lee E, et al. Merkel cell polyomavirus small T antigen activates noncanonical NFkappaB signaling to promote tumorigenesis. *Mol Cancer Res.* (2020) 18:1623–37. doi: 10.1158/1541-7786.MCR-20-0587 [PubMed: 32753470]
23. Tsang SH, Wang R, Nakamaru-Ogiso E, Knight SA, Buck CB, You J. The oncogenic small tumor antigen of Merkel cell polyomavirus is an iron-sulfur cluster protein that enhances viral DNA replication. *J Virol.* (2016) 90:1544–56. doi: 10.1128/JVI.02121-15 [PubMed: 26608318]
24. Thorley-Lawson DA. Epstein-Barr virus: exploiting the immune system. *Nat Rev Immunol.* (2001) 1:75–82. doi: 10.1038/35095584 [PubMed: 11905817]
25. Thorley-Lawson DA. EBV persistence-introducing the virus. *Curr Top Microbiol Immunol.* (2015) 390:151–209. doi: 10.1007/978-3-319-22822-8\_8 [PubMed: 26424647]
26. Wang LW, Jiang S, Gewurz BE. Epstein-Barr virus LMP1-Mediated oncogenicity. *J Virol.* (2017) 91:e01718–16. doi: 10.1128/JVI.01718-16 [PubMed: 28835489]
27. Rider MA, Cheerathodi MR, Hurwitz SN, Nkosi D, Howell LA, Tremblay DC, et al. The interactome of EBV LMP1 evaluated by proximity-based BioID approach. *Virology.* (2018) 516:55–70. doi: 10.1016/j.virol.2017.12.033 [PubMed: 29329079]
28. Longnecker R Epstein-Barr virus latency: LMP2, a regulator or means for Epstein-Barr virus persistence? *Adv Cancer Res.* (2000) 79:175–200. doi: 10.1016/S0065-230X(00)79006-3 [PubMed: 10818681]
29. Farrell PJ. Epstein-Barr virus and cancer. *Annu Rev Pathol.* (2019) 14:29–53. doi: 10.1146/annurev-pathmechdis-012418-013023 [PubMed: 30125149]
30. Pich D, Mrozek-Gorska P, Bouvet M, Sugimoto A, Akidil E, Grundhoff A, et al. First days in the life of naive human B lymphocytes infected with Epstein-Barr virus. *mBio.* (2019) 10. doi: 10.1128/mBio.01723-19
31. Ponnusamy R, Khatri R, Correia PB, Wood CD, Mancini EJ, Farrell PJ, et al. Increased association between Epstein-Barr virus EBNA2 from type 2 strains and the transcriptional repressor BS69 restricts EBNA2 activity. *PLoS Pathog.* (2019) 15:e1007458. doi: 10.1371/journal.ppat.1007458 [PubMed: 31283782]
32. Cesarman E, Damania B, Krown SE, Martin J, Bower M, Whitby D. Kaposi sarcoma. *Nat Rev Dis Primers.* (2019) 5:9. doi: 10.1038/s41572-019-0060-9 [PubMed: 30705286]
33. Lee N, Moss WN, Yario TA, Steitz JA. EBV noncoding RNA binds nascent RNA to drive host PAX5 to viral DNA. *Cell.* (2015) 160:607–18. doi: 10.1016/j.cell.2015.01.015 [PubMed: 25662012]
34. Houmani JL, Davis CI, Ruf IK. Growth-promoting properties of Epstein-Barr virus EBEB-1 RNA correlate with ribosomal protein L22 binding. *J Virol.* (2009) 83:9844–53. doi: 10.1128/JVI.01014-09 [PubMed: 19640998]
35. Li Z, Duan Y, Cheng S, Chen Y, Hu Y, Zhang L, et al. EBV-encoded RNA *via* TLR3 induces inflammation in nasopharyngeal carcinoma. *Oncotarget.* (2015) 6:24291–303. doi: 10.18632/oncotarget.4552 [PubMed: 26172457]
36. Baglio SR, van Eijndhoven MA, Koppers-Lalic D, Berenguer J, Lougheed SM, Gibbs S, et al. Sensing of latent EBV infection through exosomal transfer of 5'pppRNA. *Proc Natl Acad Sci U S A.* (2016) 113:E587–596. doi: 10.1073/pnas.1518130113 [PubMed: 26768848]
37. Hassani A, Khan G. Epstein-Barr virus and miRNAs: partners in crime in the pathogenesis of multiple sclerosis? *Front Immunol.* (2019) 10:695. doi: 10.3389/fimmu.2019.00695 [PubMed: 31001286]
38. Hussein HAM, Alfhili MA, Pakala P, Simon S, Hussain J, McCubrey JA, et al. miRNAs and their roles in KSHV pathogenesis. *Virus Res.* (2019) 266:15–24. doi: 10.1016/j.virusres.2019.03.024 [PubMed: 30951791]

39. Shinozaki-Ushiku A, Kunita A, Fukayama M. Update on Epstein-Barr virus and gastric cancer (review). *Int J Oncol.* (2015) 46:1421–34. doi: 10.3892/ijo.2015.2856 [PubMed: 25633561]
40. Zhang J, Li X, Hu J, Cao P, Yan Q, Zhang S, et al. Long noncoding RNAs involvement in Epstein-Barr virus infection and tumorigenesis. *Virology.* (2020) 17:51. doi: 10.1186/s12985-020-01308-y [PubMed: 32272952]
41. Liang C, Oh BH, Jung JU. Novel functions of viral anti-apoptotic factors. *Nat Rev Microbiol.* (2015) 13:7–12. doi: 10.1038/nrmicro3369 [PubMed: 25363821]
42. Banerjee S, Uppal T, Strahan R, Dabral P, Verma SC. The modulation of apoptotic pathways by gammaherpesviruses. *Front Microbiol.* (2016) 7:585. doi: 10.3389/fmicb.2016.00585 [PubMed: 27199919]
43. Kvensakul M. Viral infection and apoptosis. *Viruses.* (2017) 9:356. doi: 10.3390/v9120356
44. Altmann M, Hammerschmidt W. Epstein-Barr virus provides a new paradigm: a requirement for the immediate inhibition of apoptosis. *PLoS Biol.* (2005) 3:e404. doi: 10.1371/journal.pbio.0030404 [PubMed: 16277553]
45. Fitzsimmons L, Kelly GL. EBV and apoptosis: the viral master regulator of cell fate? *Viruses.* (2017) 9:339. doi: 10.3390/v9110339
46. Marquitz AR, Mathur A, Nam CS, Raab-Traub N. The Epstein-Barr virus BART microRNAs target the pro-apoptotic protein bim. *Virology.* (2011) 412:392–400. doi: 10.1016/j.viro.2011.01.028 [PubMed: 21333317]
47. Choy EY, Siu KL, Kok KH, Lung RW, Tsang CM, To KF, et al. An Epstein-Barr virus-encoded microRNA targets PUMA to promote host cell survival. *J Exp Med.* (2008) 205:2551–60. doi: 10.1084/jem.20072581 [PubMed: 18838543]
48. Gallo A, Lampe M, Gunther T, Brune W. The viral Bcl-2 homologs of Kaposi's sarcoma-associated herpesvirus and rhesus rhadinovirus share an essential role for viral replication. *J Virol.* (2017) 91:e01875–16. doi: 10.1128/JVI.01875-16 [PubMed: 28053098]
49. Arora R, Shuda M, Guastafierro A, Feng H, Toptan T, Tolstov Y, et al. Survivin is a therapeutic target in Merkel cell carcinoma. *Sci Transl Med.* (2012) 4:133ra156. doi: 10.1126/scitranslmed.3003713
50. Zhang TY, Chen HY, Cao JL, Xiong HL, Mo XB, Li TL, et al. Structural and functional analyses of hepatitis B virus X protein BH3-like domain and Bcl-xL interaction. *Nat Commun.* (2019) 10:3192. doi: 10.1038/s41467-019-11173-1 [PubMed: 31324803]
51. Jiang X, Kanda T, Wu S, Nakamoto S, Wakita T, Shirasawa H, et al. Hepatitis C virus nonstructural protein 5A inhibits thapsigargin-induced apoptosis. *PLoS ONE.* (2014) 9:e113499. doi: 10.1371/journal.pone.0113499 [PubMed: 25409163]
52. Muhleisen A, Giaisi M, Kohler R, Krammer PH, Li-Weber M. Tax contributes apoptosis resistance to HTLV-1-infected T cells *via* suppression of Bid and Bim expression. *Cell Death Dis.* (2014) 5:e1575. doi: 10.1038/cddis.2014.536 [PubMed: 25522269]
53. Pavlova NN, Thompson CB. The emerging hallmarks of cancer metabolism. *Cell Metab.* (2016) 23:27–47. doi: 10.1016/j.cmet.2015.12.006 [PubMed: 26771115]
54. Purdy JG, Luftig MA. Reprogramming of cellular metabolic pathways by human oncogenic viruses. *Curr Opin Virol.* (2019) 39:60–9. doi: 10.1016/j.coviro.2019.11.002 [PubMed: 31766001]
55. Fan R, Hou WJ, Zhao YJ, Liu SL, Qiu XS, Wang EH, et al. Overexpression of HPV16 E6/E7 mediated HIF-1 $\alpha$  upregulation of GLUT1 expression in lung cancer cells. *Tumour Biol.* (2016) 37:4655–63. doi: 10.1007/s13277-015-4221-5 [PubMed: 26508030]
56. Nakamura M, Bodily JM, Beglin M, Kyo S, Inoue M, Laimins LA. Hypoxia-specific stabilization of HIF-1 $\alpha$  by human papillomaviruses. *Virology.* (2009) 387:442–8. doi: 10.1016/j.viro.2009.02.036 [PubMed: 19321184]
57. Cuninghame S, Jackson R, Lees SJ, Zehbe I. Two common variants of human papillomavirus type 16 E6 differentially deregulate sugar metabolism and hypoxia signalling in permissive human keratinocytes. *J Gen Virol.* (2017) 98:2310–9. doi: 10.1099/jgv.0.000905 [PubMed: 28857035]
58. Zwerschke W, Mazurek S, Massimi P, Banks L, Eigenbrodt E, Jansen-Durr P. Modulation of type M2 pyruvate kinase activity by the human papillomavirus type 16 E7 oncoprotein. *Proc Natl Acad Sci U S A.* (1999) 96:1291–6. doi: 10.1073/pnas.96.4.1291 [PubMed: 9990017]

59. Berrios C, Padi M, Keibler MA, Park DE, Molla V, Cheng J, et al. Merkel cell polyomavirus small T antigen promotes pro-glycolytic metabolic perturbations required for transformation. *PLoS Pathog.* (2016) 12:e1006020. doi: 10.1371/journal.ppat.1006020 [PubMed: 27880818]
60. Hafez AY, Messinger JE, McFadden K, Fenyofalvi G, Shepard CN, Lenzi GM, et al. Limited nucleotide pools restrict Epstein-Barr virus-mediated B-cell immortalization. *Oncogenesis.* (2017) 6:e349. doi: 10.1038/oncis.2017.46 [PubMed: 28604764]
61. Wang LW, Shen H, Nobre L, Ersing I, Paulo JA, Trudeau S, et al. Epstein-Barr-virus-induced one-carbon metabolism drives B cell transformation. *Cell Metab.* (2019) 30:539–55.e511. doi: 10.1016/j.cmet.2019.06.003 [PubMed: 31257153]
62. Wang LW, Wang Z, Ersing I, Nobre L, Guo R, Jiang S, et al. Epstein-Barr virus subverts mevalonate and fatty acid pathways to promote infected B-cell proliferation and survival. *PLoS Pathog.* (2019) 15:e1008030. doi: 10.1371/journal.ppat.1008030 [PubMed: 31518366]
63. Yogev O, Lagos D, Enver T, Boshoff C. Kaposi's sarcoma herpesvirus microRNAs induce metabolic transformation of infected cells. *PLoS Pathog.* (2014) 10:e1004400. doi: 10.1371/journal.ppat.1004400 [PubMed: 25255370]
64. Bertout JA, Patel SA, Simon MC. The impact of O<sub>2</sub> availability on human cancer. *Nat Rev Cancer.* (2008) 8:967–75. doi: 10.1038/nrc2540 [PubMed: 18987634]
65. Hoppe-Seyler K, Mandl J, Adrian S, Kuhn BJ, Hoppe-Seyler F. Virus/Host cell crosstalk in hypoxic HPV-Positive cancer cells. *Viruses.* (2017) 9:174. doi: 10.3390/v9070174
66. Carroll PA, Kenerson HL, Yeung RS, Lagunoff M. Latent Kaposi's sarcoma-associated herpesvirus infection of endothelial cells activates hypoxia-induced factors. *J Virol.* (2006) 80:10802–12. doi: 10.1128/JVI.00673-06 [PubMed: 16956952]
67. Haque M, Wang V, Davis DA, Zheng ZM, Yarchoan R. Genetic organization and hypoxic activation of the Kaposi's sarcoma-associated herpesvirus ORF34–37 gene cluster. *J Virol.* (2006) 80:7037–51. doi: 10.1128/JVI.00553-06 [PubMed: 16809309]
68. Sodhi A, Montaner S, Patel V, Zohar M, Bais C, Mesri EA, et al. The Kaposi's sarcoma-associated herpes virus G protein-coupled receptor up-regulates vascular endothelial growth factor expression and secretion through mitogen-activated protein kinase and p38 pathways acting on hypoxia-inducible factor 1alpha. *Cancer Res.* (2000) 60:4873–80. [PubMed: 10987301]
69. Giffin L, Yan F, Ben Major M, Damania B. Modulation of Kaposi's sarcoma-associated herpesvirus interleukin-6 function by hypoxia-upregulated protein 1. *J Virol* (2014) 88:9429–41. doi: 10.1128/JVI.00511-14 [PubMed: 24920810]
70. Kumar V, Gabrilovich DI. Hypoxia-inducible factors in regulation of immune responses in tumour microenvironment. *Immunology.* (2014) 143:512–9. doi: 10.1111/imm.12380 [PubMed: 25196648]
71. Magalhaes I, Yogev O, Mattsson J, Schurich A. The metabolic profile of tumor and virally infected cells shapes their microenvironment counteracting T cell immunity. *Front Immunol.* (2019) 10:2309. doi: 10.3389/fimmu.2019.02309 [PubMed: 31636636]
72. Chang CH, Qiu J, O'Sullivan D, Buck MD, Noguchi T, Curtis JD, et al. Metabolic competition in the tumor microenvironment is a driver of cancer progression. *Cell.* (2015) 162:1229–41. doi: 10.1016/j.cell.2015.08.016 [PubMed: 26321679]
73. Bentz GL, Liu R, Hahn AM, Shackelford J, Pagano JS. Epstein-Barr virus BRLF1 inhibits transcription of IRF3 and IRF7 and suppresses induction of interferon-beta. *Virology.* (2010) 402:121–8. doi: 10.1016/j.virol.2010.03.014 [PubMed: 20381110]
74. Morrison TE, Mauser A, Wong A, Ting JP, Kenney SC. Inhibition of IFN-gamma signaling by an Epstein-Barr virus immediate-early protein. *Immunity.* (2001) 15:787–99. doi: 10.1016/S1074-7613(01)00226-6 [PubMed: 11728340]
75. Morrison TE, Mauser A, Klingelutz A, Kenney SC. Epstein-Barr virus immediate-early protein BZLF1 inhibits tumor necrosis factor alpha-induced signaling and apoptosis by downregulating tumor necrosis factor receptor 1. *J Virol.* (2004) 78:544–9. doi: 10.1128/JVI.78.1.544-549.2004 [PubMed: 14671137]
76. Michaud F, Coulombe F, Gaudreault E, Paquet-Bouchard C, Rola-Pleszczynski M, Gosselin J. Epstein-Barr virus interferes with the amplification of IFNalpha secretion by activating suppressor

- of cytokine signaling 3 in primary human monocytes. *PLoS ONE*. (2010) 5:e11908. doi: 10.1371/journal.pone.0011908 [PubMed: 20689596]
77. van Gent M, Griffin BD, Berkhoff EG, van Leeuwen D, Boer IG, Buisson M, et al. EBV lytic-phase protein BGLF5 contributes to TLR9 downregulation during productive infection. *J Immunol*. (2011) 186:1694–702. doi: 10.4049/jimmunol.0903120 [PubMed: 21191071]
78. McNamara RP, Chugh PE, Bailey A, Costantini LM, Ma Z, Bigi R, et al. Extracellular vesicles from Kaposi Sarcoma-associated herpesvirus lymphoma induce long-term endothelial cell reprogramming. *PLoS Pathog*. (2019) 15:e1007536. doi: 10.1371/journal.ppat.1007536 [PubMed: 30716130]
79. Hurwitz SN, Nkosi D, Conlon MM, York SB, Liu X, Tremblay DC, et al. CD63 regulates Epstein-Barr virus LMP1 exosomal packaging, enhancement of vesicle production, and noncanonical NF- $\kappa$ B signaling. *J Virol*. (2017) 91:e02251–16. doi: 10.1128/JVI.02251-16 [PubMed: 27974566]
80. Yogev O, Henderson S, Hayes MJ, Marelli SS, Ofir-Birin Y, Regev-Rudzi N, et al. Herpesviruses shape tumour microenvironment through exosomal transfer of viral microRNAs. *PLoS Pathog*. (2017) 13:e1006524. doi: 10.1371/journal.ppat.1006524 [PubMed: 28837697]
81. Mrizak D, Martin N, Barjon C, Jimenez-Pailhes AS, Mustapha R, Niki T, et al. Effect of nasopharyngeal carcinoma-derived exosomes on human regulatory T cells. *J Natl Cancer Inst*. (2015) 107:363. doi: 10.1093/jnci/dju363 [PubMed: 25505237]
82. Cao S, Wylie KM, Wyczalkowski MA, Karpova A, Ley J, Sun S, et al. Dynamic host immune response in virus-associated cancers. *Commun Biol*. (2019) 2:109. doi: 10.1038/s42003-019-0352-3 [PubMed: 30911684]
83. Anastasiadou E, Stroopinsky D, Alimperti S, Jiao AL, Pyzer AR, Cippitelli C, et al. Epstein-Barr virus-encoded EBNA2 alters immune checkpoint PD-L1 expression by downregulating miR-34a in B-cell lymphomas. *Leukemia*. (2019) 33:132–47. doi: 10.1038/s41375-018-0178-x [PubMed: 29946193]
84. Fang W, Zhang J, Hong S, Zhan J, Chen N, Qin T, et al. EBV-driven LMP1 and IFN- $\gamma$  up-regulate PD-L1 in nasopharyngeal carcinoma: implications for oncotargeted therapy. *Oncotarget*. (2014) 5:12189–202. doi: 10.18632/oncotarget.2608 [PubMed: 25361008]
85. Bi XW, Wang H, Zhang WW, Wang JH, Liu WJ, Xia ZJ, et al. PD-L1 is upregulated by EBV-driven LMP1 through NF- $\kappa$ B pathway and correlates with poor prognosis in natural killer/T-cell lymphoma. *J Hematol Oncol*. (2016) 9:109. doi: 10.1186/s13045-016-0341-7 [PubMed: 27737703]
86. Miyauchi S, Sanders PD, Guram K, Kim SS, Paolini F, Venuti A, et al. HPV16 E5 mediates resistance to PD-L1 blockade and can be targeted with rimantadine in head and neck cancer. *Cancer Res*. (2020) 80:732–46. doi: 10.1158/0008-5472.CAN-19-1771 [PubMed: 31848196]
87. Tsai TY, Huang MT, Sung PS, Peng CY, Tao MH, Yang HI, et al. SIGLEC-3 (CD33) serves as an immune checkpoint receptor for HBV infection. *J Clin Invest*. (2021) 131:e141965. doi: 10.1172/JCI141965
88. Osuch S, Metzner KJ, Caraballo Cortes K. Reversal of T cell exhaustion in chronic HCV infection. *Viruses*. (2020) 12:799. doi: 10.3390/v12080799
89. Lu F, Chen HS, Kossenkov AV, DeWisleare K, Won KJ, Lieberman PM. EBNA2 drives formation of new chromosome binding sites and target genes for B-Cell Master regulatory transcription factors RBP-jkappa and EBF1. *PLoS Pathog*. (2016) 12:e1005339. doi: 10.1371/journal.ppat.1005339 [PubMed: 26752713]
90. Liang J, Zhou H, Gerdt C, Tan M, Colson T, Kaye KM, et al. Epstein-Barr virus super-enhancer eRNAs are essential for MYC oncogene expression and lymphoblast proliferation. *Proc Natl Acad Sci U S A*. (2016) 113:14121–6. doi: 10.1073/pnas.1616697113 [PubMed: 27864512]
91. Jiang S, Zhou H, Liang J, Gerdt C, Wang C, Ke L, et al. The Epstein-Barr virus Regulome in Lymphoblastoid Cells. *Cell Host Microbe*. (2017) 22:561–73.e564. doi: 10.1016/j.chom.2017.09.001 [PubMed: 29024646]
92. Skalska L, White RE, Franz M, Ruhmann M, Allday MJ. Epigenetic repression of p16(INK4A) by latent Epstein-Barr virus requires the interaction of EBNA3A and EBNA3C with CtBP. *PLoS Pathog*. (2010) 6:e1000951. doi: 10.1371/journal.ppat.1000951 [PubMed: 20548956]

93. Paschos K, Parker GA, Watanatanasup E, White RE, Allday MJ. BIM promoter directly targeted by EBNA3C in polycomb-mediated repression by EBV. *Nucleic Acids Res.* (2012) 40:7233–46. doi: 10.1093/nar/gks391 [PubMed: 22584624]
94. Juillard F, Tan M, Li S, Kaye KM. Kaposi's sarcoma herpesvirus genome persistence. *Front Microbiol.* (2016) 7:1149. doi: 10.3389/fmicb.2016.01149 [PubMed: 27570517]
95. Ueda K KSHV genome replication and maintenance in latency. *Adv Exp Med Biol.* (2018) 1045:299–320. doi: 10.1007/978-981-10-7230-7\_14 [PubMed: 29896673]
96. Brulois K, Jung JU. Interplay between Kaposi's sarcoma-associated herpesvirus and the innate immune system. *Cytokine Growth Factor Rev.* (2014) 25:597–609. doi: 10.1016/j.cytogfr.2014.06.001 [PubMed: 25037686]
97. Giudice A, D'Arena G, Crispo A, Tecce MF, Nocerino F, Grimaldi M, et al. Role of viral miRNAs and epigenetic modifications in Epstein-Barr virus-associated gastric carcinogenesis. *Oxid Med Cell Longev.* (2016) 2016:6021934. doi: 10.1155/2016/6021934 [PubMed: 26977250]
98. Lou W, Liu J, Ding B, Chen D, Xu L, Ding J, et al. Identification of potential miRNA-mRNA regulatory network contributing to pathogenesis of HBV-related HCC. *J Transl Med.* (2019) 17:7. doi: 10.1186/s12967-018-1761-7 [PubMed: 30602391]
99. Zhao Q, Zhang Y, Zhang X, Sun Y, Lin Z. Mining of gene modules and identification of key genes in head and neck squamous cell carcinoma based on gene co-expression network analysis. *Medicine.* (2020) 99:e22655. doi: 10.1097/MD.00000000000022655 [PubMed: 33285674]
100. Kuss-Duerkop SK, Westrich JA, Pyeon D. DNA tumor virus regulation of host DNA methylation and its implications for immune evasion and oncogenesis. *Viruses.* (2018) 10:82. doi: 10.3390/v10020082
101. Matsusaka K, Funata S, Fukuyo M, Seto Y, Aburatani H, Fukayama M, et al. Epstein-Barr virus infection induces genome-wide *de novo* DNA methylation in non-neoplastic gastric epithelial cells. *J Pathol.* (2017) 242:391–9. doi: 10.1002/path.4909 [PubMed: 28418084]
102. Birdwell CE, Queen KJ, Kilgore PC, Rollyson P, Trutschl M, Cvek U, et al. Genome-wide DNA methylation as an epigenetic consequence of Epstein-Barr virus infection of immortalized keratinocytes. *J Virol.* (2014) 88:11442–58. doi: 10.1128/JVI.00972-14 [PubMed: 25056883]
103. Kaneda A, Matsusaka K, Aburatani H, Fukayama M. Epstein-Barr virus infection as an epigenetic driver of tumorigenesis. *Cancer Res.* (2012) 72:3445–50. doi: 10.1158/0008-5472.CAN-11-3919 [PubMed: 22761333]
104. Li L, Li C, Mao H, Du Z, Chan WY, Murray P, et al. Epigenetic inactivation of the CpG demethylase TET1 as a DNA methylation feedback loop in human cancers. *Sci Rep.* (2016) 6:26591. doi: 10.1038/srep34435 [PubMed: 27225590]
105. Namba-Fukuyo H, Funata S, Matsusaka K, Fukuyo M, Rahmutulla B, Mano Y, et al. TET2 functions as a resistance factor against DNA methylation acquisition during Epstein-Barr virus infection. *Oncotarget.* (2016) 7:81512–26. doi: 10.18632/oncotarget.13130 [PubMed: 27829228]
106. Tsai CN, Tsai CL, Tse KP, Chang HY, Chang YS. The Epstein-Barr virus oncogene product, latent membrane protein 1, induces the downregulation of E-cadherin gene expression *via* activation of DNA methyltransferases. *Proc Natl Acad Sci U S A.* (2002) 99:10084–9. doi: 10.1073/pnas.152059399 [PubMed: 12110730]
107. Tsai CL, Li HP, Lu YJ, Hsueh C, Liang Y, Chen CL, et al. Activation of DNA methyltransferase 1 by EBV LMP1 involves c-Jun NH(2)-terminal kinase signaling. *Cancer Res.* (2006) 66:11668–76. doi: 10.1158/0008-5472.CAN-06-2194 [PubMed: 17178861]
108. Hino R, Uozaki H, Murakami N, Ushiku T, Shinozaki A, Ishikawa S, et al. Activation of DNA methyltransferase 1 by EBV latent membrane protein 2A leads to promoter hypermethylation of PTEN gene in gastric carcinoma. *Cancer Res.* (2009) 69:2766–74. doi: 10.1158/0008-5472.CAN-08-3070 [PubMed: 19339266]
109. Chong JM, Sakuma K, Sudo M, Ushiku T, Uozaki H, Shibahara J, et al. Global and non-random CpG-island methylation in gastric carcinoma associated with Epstein-Barr virus. *Cancer Sci.* (2003) 94:76–80. doi: 10.1111/j.1349-7006.2003.tb01355.x [PubMed: 12708478]
110. Okabe A, Funata S, Matsusaka K, Namba H, Fukuyo M, Rahmutulla B, et al. Regulation of tumour related genes by dynamic epigenetic alteration at enhancer regions in gastric epithelial


- cells infected by Epstein-Barr virus. *Sci Rep.* (2017) 7:7924. doi: 10.1038/s41598-017-08370-7 [PubMed: 28801683]
111. Soto D, Song C, McLaughlin-Drubin ME. Epigenetic alterations in human papillomavirus-associated cancers. *Viruses.* (2017) 9:248. doi: 10.3390/v9090248
  112. Knipe DM. Nuclear sensing of viral DNA, epigenetic regulation of herpes simplex virus infection, innate immunity. *Virology.* (2015) 479–80:153–9. doi: 10.1016/j.virol.2015.02.009
  113. Knisbacher BA, Gerber D, Levanon EY, DNA editing by APOBECs: a genomic preserver and transformer. *Trends Genet.* (2016) 32:16–28. doi: 10.1016/j.tig.2015.10.005 [PubMed: 26608778]
  114. Zhou H, Schmidt SC, Jiang S, Willox B, Bernhardt K, Liang J, et al. Epstein-Barr virus oncoprotein super-enhancers control B cell growth. *Cell Host Microbe.* (2015) 17:205–16. doi: 10.1016/j.chom.2014.12.013 [PubMed: 25639793]
  115. Gunnell A, Webb HM, Wood CD, McClellan MJ, Wichaidit B, Kempkes B, et al. RUNX super-enhancer control through the Notch pathway by Epstein-Barr virus transcription factors regulates B cell growth. *Nucleic Acids Res.* (2016) 44:4636–50. doi: 10.1093/nar/gkw085 [PubMed: 26883634]
  116. Kim KD, Tanizawa H, De Leo A, Vladimirova O, Kossenkov A, Lu F, et al. Epigenetic specifications of host chromosome docking sites for latent Epstein-Barr virus. *Nat Commun.* (2020) 11:877. doi: 10.1038/s41467-019-14152-8 [PubMed: 32054837]
  117. Okabe A, Huang KK, Matsusaka K, Fukuyo M, Xing M, Ong X, et al. Cross-species chromatin interactions drive transcriptional rewiring in Epstein-Barr virus-positive gastric adenocarcinoma. *Nat Genet.* (2020) 52:919–30. doi: 10.1038/s41588-020-0665-7 [PubMed: 32719515]
  118. Moquin SA, Thomas S, Whalen S, Warburton A, Fernandez SG, McBride AA, et al. The Epstein-Barr virus episome maneuvers between nuclear chromatin compartments during reactivation. *J Virol.* (2018) 92. doi: 10.1128/JVI.01413-17
  119. Lieberman PM. Keeping it quiet: chromatin control of gammaherpesvirus latency. *Nat Rev Microbiol.* (2013) 11:863–75. doi: 10.1038/nrmicro3135 [PubMed: 24192651]
  120. Lieberman PM. Epigenetics and genetics of viral latency. *Cell Host Microbe.* (2016) 19:619–28. doi: 10.1016/j.chom.2016.04.008 [PubMed: 27173930]
  121. De Leo A, Calderon A, Lieberman PM. Control of viral latency by episome maintenance proteins. *Trends Microbiol.* (2020) 28:150–62. doi: 10.1016/j.tim.2019.09.002 [PubMed: 31624007]
  122. Watanabe T Adult T-cell leukemia: molecular basis for clonal expansion and transformation of HTLV-1-infected T cells. *Blood.* (2017) 129:1071–81. doi: 10.1182/blood-2016-09-692574 [PubMed: 28115366]
  123. Moore PS, Chang Y. Why do viruses cause cancer? Highlights of the first century of human tumour virology. *Nat Rev Cancer.* (2010) 10:878–89. doi: 10.1038/nrc2961 [PubMed: 21102637]
  124. Chen X, Kost J, Sulovari A, Wong N, Liang WS, Cao J, et al. A virome-wide clonal integration analysis platform for discovering cancer viral etiology. *Genome Res.* (2019) 29:819–30. doi: 10.1101/gr.242529.118 [PubMed: 30872350]
  125. Cao S, Wendl MC, Wyczalkowski MA, Wylie K, Ye K, Jayasinghe R, et al. Divergent viral presentation among human tumors and adjacent normal tissues. *Sci Rep.* (2016) 6:28294. doi: 10.1038/srep28294 [PubMed: 27339696]
  126. Romanczuk H, Howley PM. Disruption of either the E1 or the E2 regulatory gene of human papillomavirus type 16 increases viral immortalization capacity. *Proc Natl Acad Sci U S A.* (1992) 89:3159–63. doi: 10.1073/pnas.89.7.3159 [PubMed: 1313584]
  127. Cheung JL, Lo KW, Cheung TH, Tang JW, Chan PK. Viral load, E2 gene disruption status, and lineage of human papillomavirus type 16 infection in cervical neoplasia. *J Infect Dis.* (2006) 194:1706–12. doi: 10.1086/509622 [PubMed: 17109343]
  128. Shuda M, Feng H, Kwun HJ, Rosen ST, Gjoerup O, Moore PS, et al. T antigen mutations are a human tumor-specific signature for Merkel cell polyomavirus. *Proc Natl Acad Sci U S A.* (2008) 105:16272–7 doi: 10.1073/pnas.0806526105 [PubMed: 18812503]
  129. Naipauer J, Salyakina D, Journo G, Rosario S, Williams S, Abba M, et al. High-throughput sequencing analysis of a “hit and run” cell and animal model of KSHV tumorigenesis. *PLoS Pathog.* (2020) 16:e1008589. doi: 10.1371/journal.ppat.1008589 [PubMed: 32603362]

130. Cancian L, Hansen A, Boshoff C. Cellular origin of Kaposi's sarcoma and Kaposi's sarcoma-associated herpesvirus-induced cell reprogramming. *Trends Cell Biol.* (2013) 23:421–32. doi: 10.1016/j.tcb.2013.04.001 [PubMed: 23685018]
131. Iacovides D, Michael S, Achilleos C, Strati K. Shared mechanisms in stemness and carcinogenesis: lessons from oncogenic viruses. *Front Cell Infect Microbiol.* (2013) 3:66. doi: 10.3389/fcimb.2013.00066 [PubMed: 24400225]
132. Hibner U, Gregoire D. Viruses in cancer cell plasticity: the role of hepatitis C virus in hepatocellular carcinoma. *Contemp Oncol.* (2015) 19:A62–67. doi: 10.5114/wo.2014.47132
133. Xiang T, Lin YX, Ma W, Zhang HJ, Chen KM, He GP, et al. Vasculogenic mimicry formation in EBV-associated epithelial malignancies. *Nat Commun.* (2018) 9:5009. doi: 10.1038/s41467-018-07308-5 [PubMed: 30479336]
134. Reid P, Marcu LG, Olver I, Moghaddasi L, Staudacher AH, Bezak E. Diversity of cancer stem cells in head and neck carcinomas: the role of HPV in cancer stem cell heterogeneity, plasticity and treatment response. *Radiother Oncol.* (2019) 135:1–12. doi: 10.1016/j.radonc.2019.02.016 [PubMed: 31015153]
135. Shen S, Clairambault J. Cell plasticity in cancer cell populations. *F1000Res.* (2020) 9:F1000. doi: 10.12688/f1000research.24803.1
136. Nijman SMB. Perturbation-driven entropy as a source of cancer cell heterogeneity. *Trends Cancer.* (2020) 6:454–61. doi: 10.1016/j.trecan.2020.02.016 [PubMed: 32460001]
137. Chang Y, Moore PS, Weiss RA. Human oncogenic viruses: nature and discovery. *Philos Trans R Soc Lond B Biol Sci.* (2017) 372:20160264. doi: 10.1098/rstb.2016.0264 [PubMed: 28893931]
138. Dhawan A, Scott J, Sundaresan P, Veness M, Porceddu S, Hau E, et al. Role of gene signatures combined with pathology in classification of oropharynx head and neck cancer. *Sci Rep.* (2020) 10:10226. doi: 10.1038/s41598-020-66983-x [PubMed: 32576885]
139. Bhatia K, Goedert JJ, Modali R, Preiss L, Ayers LW. Merkel cell carcinoma subgroups by Merkel cell polyomavirus DNA relative abundance and oncogene expression. *Int J Cancer.* (2010) 126:2240–6. doi: 10.1002/ijc.24676 [PubMed: 19551862]
140. Ho DW, Tsui YM, Chan LK, Sze KM, Zhang X, Cheu JW, et al. Single-cell RNA sequencing shows the immunosuppressive landscape and tumor heterogeneity of HBV-associated hepatocellular carcinoma. *Nat Commun.* (2021) 12:3684. doi: 10.1038/s41467-021-24010-1 [PubMed: 34140495]
141. Krause J, von Felden J, Casar C, Frundt TW, Galaski J, Schmidt C, et al. Hepatocellular carcinoma: intratumoral EpCAM-positive cancer stem cell heterogeneity identifies high-risk tumor subtype. *BMC Cancer.* (2020) 20:1130. doi: 10.1186/s12885-020-07580-z [PubMed: 33225916]
142. Yamagishi M, Kubokawa M, Kuze Y, Suzuki A, Yokomizo A, Kobayashi S, et al. Chronological genome and single-cell transcriptome integration characterizes the evolutionary process of adult T cell leukemia-lymphoma. *Nat Commun.* (2021) 12:4821. doi: 10.1038/s41467-021-25101-9 [PubMed: 34376672]
143. Messinger JE, Dai J, Stanland LJ, Price AM, Luftig MA. Identification of host biomarkers of Epstein-Barr virus latency IIb and latency III. *MBio.* (2019) 10. doi: 10.1128/mBio.01006-19
144. Yi M, Cai J, Li J, Chen S, Zeng Z, Peng Q, et al. Rediscovery of NF-kappaB signaling in nasopharyngeal carcinoma: how genetic defects of NF-kappaB pathway interplay with EBV in driving oncogenesis? *J Cell Physiol.* (2018) 233:5537–49. doi: 10.1002/jcp.26410 [PubMed: 29266238]
145. Bailey C, Shoura MJ, Mischel PS, Swanton C. Extrachromosomal DNA-relieving heredity constraints, accelerating tumour evolution. *Ann Oncol.* (2020) 31:884–93. doi: 10.1016/j.annonc.2020.03.303 [PubMed: 32275948]
146. Manners O, Murphy JC, Coleman A, Hughes DJ, Whitehouse A. Contribution of the KSHV and EBV lytic cycles to tumourigenesis. *Curr Opin Virol.* (2018) 32:60–70. doi: 10.1016/j.coviro.2018.08.014 [PubMed: 30268927]
147. Raab-Traub N, Flynn K. The structure of the termini of the Epstein-Barr virus as a marker of clonal cellular proliferation. *Cell.* (1986) 47:883–9. doi: 10.1016/0092-8674(86)90803-2 [PubMed: 3022942]

148. Pathmanathan R, Prasad U, Sadler R, Flynn K, Raab-Traub N. Clonal proliferations of cells infected with Epstein-Barr virus in preinvasive lesions related to nasopharyngeal carcinoma. *N Engl J Med.* (1995) 333:693–8 doi: 10.1056/NEJM199509143331103 [PubMed: 7637746]
149. Feederle R, Neuhierl B, Bannert H, Geletneky K, Shannon-Lowe C, Delecluse HJ. Epstein-Barr virus B958 produced in 293 cells shows marked tropism for differentiated primary epithelial cells and reveals interindividual variation in susceptibility to viral infection. *Int J Cancer.* (2007) 121:588–94. doi: 10.1002/ijc.22727 [PubMed: 17417777]
150. Tsang CM, Yip YL, Lo KW, Deng W, To KF, Hau PM, et al. Cyclin D1 overexpression supports stable EBV infection in nasopharyngeal epithelial cells. *Proc Natl Acad Sci U S A.* (2012) 109:E3473–3482. doi: 10.1073/pnas.1202637109 [PubMed: 23161911]
151. Tsao SW, Tsang CM, Lo KW. Epstein-Barr virus infection and nasopharyngeal carcinoma. *Philos Trans R Soc Lond B Biol Sci.* (2017) 372. doi: 10.1098/rstb.2016.0270
152. Miller CA, McMichael J, Dang HX, Maher CA, Ding L, Ley TJ, et al. Visualizing tumor evolution with the fishplot package for R. *BMC Genomics.* (2016) 17:880. doi: 10.1186/s12864-016-3195-z [PubMed: 27821060]
153. Yu G, Hsu WL, Coghill AE, Yu KJ, Wang CP, Lou PJ, et al. Whole-Exome sequencing of nasopharyngeal carcinoma families reveals novel variants potentially involved in nasopharyngeal carcinoma. *Sci Rep.* (2019) 9:9916. doi: 10.1038/s41598-019-46137-4 [PubMed: 31289279]
154. Chaigne-Delalande B, Li FY, O'Connor GM, Lukacs MJ, Jiang P, Zheng L, et al. Mg<sup>2+</sup> regulates cytotoxic functions of NK and CD8T cells in chronic EBV infection through NKG2D. *Science.* (2013) 341:186–91. doi: 10.1126/science.1240094 [PubMed: 23846901]
155. Dhalla F, Murray S, Sadler R, Chaigne-Delalande B, Sadaoka T, Soilleux E, et al. Identification of a novel mutation in MAGT1 and progressive multifocal leucoencephalopathy in a 58-year-old man with XME disease. *J Clin Immunol.* (2015) 35:112–8. doi: 10.1007/s10875-014-0116-2 [PubMed: 25504528]
156. Nakhoul H, Lin Z, Wang X, Roberts C, Dong Y, Flemington E. High-Throughput sequence analysis of peripheral T-Cell lymphomas indicates subtype-specific viral gene expression patterns and immune cell microenvironments. *mSphere.* (2019) 4:e00248–19. doi: 10.1128/mSphere.00248-19 [PubMed: 31292228]
157. Chau SL, Tong JH, Chow C, Kwan JS, Lung RW, Chung LY, et al. Distinct molecular landscape of Epstein-Barr Virus associated pulmonary lymphoepithelioma-like carcinoma revealed by genomic sequencing. *Cancers.* (2020) 12:2065. doi: 10.3390/cancers12082065
158. Schiffman M, Wentzensen N. Human papillomavirus infection and the multistage carcinogenesis of cervical cancer. *Cancer Epidemiol Biomarkers Prev.* (2013) 22:553–60. doi: 10.1158/1055-9965.EPI-12-1406 [PubMed: 23549399]
159. Spurgeon ME, Liem A, Buehler D, Cheng J, DeCaprio JA, Lambert PF. The Merkel cell polyomavirus T antigens function as tumor promoters in murine skin. *Cancers.* (2021) 13:222. doi: 10.3390/cancers13020222
160. Izaki M, Yasunaga JI, Nosaka K, Sugata K, Utsunomiya H, Suehiro Y, et al. *In vivo* dynamics and adaptation of HTLV-1-infected clones under different clinical conditions. *PLoS Pathog.* (2021) 17:e1009271. doi: 10.1371/journal.ppat.1009271 [PubMed: 33524072]
161. Sartorius K, Swadling L, An P, Makarova J, Winkler C, Chuturgoon A, et al. The multiple roles of hepatitis b virus X Protein (HBx) dysregulated microRNA in hepatitis B virus-associated hepatocellular carcinoma (HBV-HCC) and immune pathways. *Viruses.* (2020) 12:746. doi: 10.3390/v12070746
162. Li Q, Wennborg A, Aurell E, Dekel E, Zou JZ, Xu Y, et al. Dynamics inside the cancer cell attractor reveal cell heterogeneity, limits of stability, and escape. *Proc Natl Acad Sci U S A.* (2016) 113:2672–7. doi: 10.1073/pnas.1519210113 [PubMed: 26929366]
163. Ladewig J, Koch P, Brustle O. Leveling Waddington: the emergence of direct programming and the loss of cell fate hierarchies. *Nat Rev Mol Cell Biol.* (2013) 14:225–36. doi: 10.1038/nrm3543
164. Waddington CH, Robertson E. Selection for developmental canalisation. *GenetRes.* (1966) 7:303–12. doi: 10.1017/S0016672300009769
165. Chen H, He X. The convergent cancer evolution toward a single cellular destination. *Mol Biol Evol.* (2016) 33:4–12. doi: 10.1093/molbev/msv212 [PubMed: 26464125]




166. Urban EA, Johnston RJ Jr. Buffering and amplifying transcriptional noise during cell fate specification. *Front Genet.* (2018) 9:591. doi:10.3389/fgene.2018.00591 [PubMed: 30555516]
167. Huang S, Ernberg I, Kauffman S. Cancer attractors: a systems view of tumors from a gene network dynamics and developmental perspective. *Semin Cell Dev Biol.* (2009) 20:869–76. doi: 10.1016/j.semcdb.2009.07.003 [PubMed: 19595782]
168. Scott J, Marusyk A. Somatic clonal evolution: a selection-centric perspective. *Biochim Biophys Acta Rev Cancer.* (2017) 1867:139–50. doi: 10.1016/j.bbcan.2017.01.006 [PubMed: 28161395]
169. Westrich JA, Warren CJ, Pyeon D. Evasion of host immune defenses by human papillomavirus. *Virus Res.* (2017) 231:21–33. doi: 10.1016/j.virusres.2016.11.023 [PubMed: 27890631]
170. Jenkinson G, Pujadas E, Goutsias J, Feinberg AP. Potential energy landscapes identify the information-theoretic nature of the epigenome. *Nat Genet.* (2017) 49:719–29. doi: 10.1038/ng.3811 [PubMed: 28346445]
171. Chan KCA, Woo JKS, King A, Zee BCY, Lam WKJ, Chan SL, et al. Analysis of plasma Epstein-Barr virus DNA to screen for nasopharyngeal cancer. *N Engl J Med.* (2017) 377:513–22. doi: 10.1056/NEJMoa1701717 [PubMed: 28792880]
172. Ji MF, Sheng W, Cheng WM, Ng MH, Wu BH, Yu X, et al. Incidence and mortality of nasopharyngeal carcinoma: interim analysis of a cluster randomized controlled screening trial (PRO-NPC-001) in southern China. *Ann Oncol.* (2019) 30:1630–7. doi: 10.1093/annonc/mdz231 [PubMed: 31373615]
173. Zhang Y, Li Y, Li T, Shen X, Zhu T, Tao Y, et al. Genetic load and potential mutational meltdown in cancer cell populations. *Mol Biol Evol.* (2019) 36:541–52. doi: 10.1093/molbev/msy231 [PubMed: 30649444]



**FIGURE 1 |**


Diversity of viral oncogenic mechanisms. Viral oncogenes have been shown to perturb multiple hallmarks of cancer. How each of these viral-host interactions contribute to carcinogenesis in a particular viral cancer is a challenge for identifying primary driving forces of viral cancers. Examples highlighted from EBV proteins EBNA-LP, EBNA1, EBNA2, EBNA3A, 3B, 3C, BNRF1, BZLF1; KSHV LANA, vGPCR, vCyclin, vIL6, ORF75; HPV E6, E7; MCPyV LT, ST; HBV HbX; HTLV-1 Tax, HBZ.

# Evolution of an EBV Cancer




**FIGURE 2 |**

Viral contributions to tumor evolution over time. The viral contributions to carcinogenesis may differ and change depending on the temporal stage of cancer cell evolution. The dynamic properties of viral genomes and infection enable greater plasticity than cellular genomes and gene expression mechanisms. Examples from EBV suggest that viral cancers evolve through changes in viral and cellular gene expression, including the loss of viral oncogenes, such as EBNA2 in BL and LMP1 in NPC, and compensatory oncogenic mutations in cellular genes, such as myc translocations and NFkB activation, at later stages of cancer cell development. In this way, viruses provide lower cost pathways to cellular oncogenesis.



**FIGURE 3 |** Heterogeneity of viral tumors. Hypothetical “fishplot” of EBV NPC tumors demonstrate highly heterogeneous patterns of viral and host gene expression in the different cells of an emerging tumor (151). Fishplots measure the clonal evolution of cells in a tumor microenvironment over time (152). Such fishplots reflect temporal, historical events in the cancer evolution process, and spatial, topological variations in the tumor microenvironment. Viruses contribute to the adaptability of tumor cells to these rapid changes in microenvironment.



**FIGURE 4 |**

Thermodynamic landscape of viral oncogenesis. A Waddington-like developmental landscape conceptualizes how viruses create alternative gene programs and biochemical pathways that facilitate the transition to an oncogenic state (163). The oncogenic state may be considered in terms of alternative attractor states (e.g., M1 and M2) with favorable thermodynamic properties and increased Darwinian fitness. The Waddington developmental landscape is related to the thermodynamic landscape for chemical reactions. Oncogenic viruses enable new attractors states by providing additional genetic and biochemical flexibility. This viral-borne adaptability may be considered an entropic driver of cancer cell evolution.