

Subcutaneous emphysema of the scrotum (pneumoscotum) due to traumatic pneumothorax: a case report

Vasileios Simaioforidis*, Stylianos Kontos, Ioannis Fokitis, Georgios Lefakis and Sotirios Koritsiadis

Address: Department of Urology, General Hospital of Nikea, 3 D. Mantouvalou St., Nikea, 18454, Piraeus, Greece

Email: Vasileios Simaioforidis* - vas_sim@hotmail.com; Stylianos Kontos - kontostylianos@gmail.com; Ioannis Fokitis - ifokitis@yahoo.gr; Georgios Lefakis - geolefakis@yahoo.gr; Sotirios Koritsiadis - skoritsiadis@nikaia-hosp.gr

* Corresponding author

Published: 1 November 2008

Received: 10 October 2008

Cases Journal 2008, 1:293 doi:10.1186/1757-1626-1-293

Accepted: 1 November 2008

This article is available from: <http://www.casesjournal.com/content/1/1/293>

© 2008 Simaioforidis et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Introduction: Subcutaneous emphysema of the scrotum due to traumatic pneumothorax is a rare medical situation and only a few cases are reported in the literature.

Case report: We present the case of a 22 year old man who was admitted to the emergency department after a motorcycle accident having a painless crepitant scrotum and chest excoriations. Further evaluation revealed subcutaneous emphysema of the scrotum caused by left pneumothorax.

Conclusion: In conclusion, subcutaneous emphysema of the scrotum (or pneumoscotum) due to traumatic pneumothorax is not an urgent condition and assessment should be supportive with intervention directed at the etiology, e.g. the pneumothorax.

Background

Subcutaneous emphysema of the scrotum (or pneumoscotum) is a rare condition and receives little or no discussion in standard texts of urology. This paper presents a case of pneumoscotum due to traumatic pneumothorax and discusses its clinical course, findings and treatment strategy with a review of the relevant literature.

Case presentation

A 22 year old man was admitted to the emergency department following a motorcycle accident. He had chest excoriations and a painless crepitant scrotum (Fig 1, 2.). Palpation of the scrotum was characteristic and gave the impression of palpating "snow". Testicles could not be palpated. During clinical examination, patient complained of incipient dyspnea. Auscultation of the lungs

placed the suspicion of pneumothorax. The CT imaging study demonstrated a left pneumothorax (Fig. 3) causing a subcutaneous emphysema that was expanded in the scrotum (Fig. 4). A left haemothorax and three rib fractures were also diagnosed. No lesions of the genitalia were diagnosed. A chest drainage tube was inserted and further assessment was supportive in collaboration with the thoracosurgery clinic. Patient was dismissed eight days after the accident. Pneumothorax was partially absorbed and further control was arranged as an outpatient. By one month, no air was found in the ultrasound imaging of the scrotum.

Discussion

Subcutaneous emphysema of the scrotum – especially of traumatic etiology – is not a common medical condition.



Figure 1
Oedematous scrotum (profile).

Causes that may result in the presence of air in the scrotum are:

1. Iatrogenic: such as various endoscopies (colonoscopy, polypectomy, laparoscopic procedures, tracheostomy, CPR, chest drain tube insertion, dental procedures) [1-3].



Figure 2
Oedematous scrotum.




Figure 3
CT demonstrates left pneumo-haemothorax with subcutaneous emphysema.

2. Gas producing infection (gangrene Fournier).
3. Pneumothorax [4,5].
4. Scrotal trauma.
5. Visceral perforation [6].

The true nature of the development of the subcutaneous emphysema of the scrotum has not been fully elucidated. Three possible mechanisms were proposed:

1. Rapture of the alveoli and expansion of the air through the mediastinum in the subcutaneous area [6].




Figure 4
CT demonstrates pneumoscrotum (red arrows).

2. Expansion of the emphysema along the Scarpa's Fascia [7].
3. Through the mediastinum air passes in the paranephric space and through the retroperitoneum and the inguinal canal is gathered in the scrotum [8-10].

In the case described above, spread of the subcutaneous emphysema towards scrotum is the most "desirable" potential, given the fact that if this expansion was towards neck it could obstruct the upper airways and lead to suffocation.

Conclusion

We can support that subcutaneous emphysema of the scrotum due to traumatic pneumothorax (blunt trauma), despite its impressive clinical presentation, is not really an urgent situation and assessment should be supportive with intervention directed at the etiology, that is the pneumothorax [11].

Abbreviations

CT: Computed Tomography; CPR: Cardio Pulmonary Resuscitation.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors have made substantial contribution to concept this case reports

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in Chief of this journal.

References

1. Stavem K, Hoivik B: **[Pneumosrotum after air leak from tension pneumothorax]**. *Tidsskr Nor Laegeforen* 116(10):1214-5. 1996 Apr 20
2. Humpheys F, Hewetson KA, Dellipianni AW: **Massive subcutaneous emphysema following colonoscopy**. *Endoscopy* 1984, 16(4):160-161.
3. Casey RG, Al-Dousari S, Murphy D, Power RE: **Chest drain insertion may result in the acute scrotum**. *Scand J Urol Nephrol* 2006, 40(1):78-9.
4. Wakabayashi Y, Bush WH Jr: **Pneumosrotum after blunt chest trauma**. *J Emerg Med* 1994, 12(5):603-5.
5. Firman R, Heiselman D, Lloyd T, Mardesich P: **Pneumosrotum**. *Ann Emerg Med* 1993, 22(8):1353-6.
6. Aslan Y, Sarihan H, Dinc H, Gedik Y, Aksoy A, Dereci S: **Gastric perforation presenting as bilateral scrotal pneumatocoles**. *Turk J pediatr* 1999, 41(2):267-71.
7. Millmond SH, Goldman SM: **Pneumosrotum after spontaneous pneumothorax with air leak**. *J Urol* 1991, 145(6):1271-1272.
8. Heimbach D, Hofmockel G, Wirth M, Frohmuller H: **Pneumosrotum: Case report and review of the literature**. *Urologe A* 1993, 32(6):503-506.
9. Casey RG, Al-Dousari S, Murphy D, Power RE: **Chest drain insertion may result in the acute scrotum**. *Scand J Urol Nephrol* 2006, 40(1):78-79.
10. Watson HS, Klugo RC, Coffield KS: **Pneumosrotum: report of two cases and review of the mechanisms or its development**. *Urology* 1992, 40(6):517-521.
11. Mateos Colino A, Golpe Gomez R, Gonzalez Rodriguez A, Sousa Escandon A, Gonzalez Uribarri C, Seirulo Salas M: **Neumoesrototo secundario a efisema sudcutaneo masivo tras el drenaje de un neumotorax espontaneo**. *Actas Urol Esp* 2004:606-609.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

