

MEETING ABSTRACT

Open Access

Environmental and occupational respiratory diseases – 1057. Correlation between air pollution and respiratory health of school children in Delhi

Jincy Mathew^{1*}, Radha Goyal², KK Taneja³, Naveen Arora¹

From 2nd WAO International Scientific Conference (WISC 2012)
Hyderabad, India. 6-9 December 2012

Background

Urban air pollution (indoor and outdoor) poses a major health hazard for city dwellers. Exposure to ambient air pollution may result in respiratory health disorders. The aim of this study was to correlate air pollutant levels to the respiratory health of school children in Delhi.

Methods

The questionnaire was designed based on the internationally valid questionnaires for respiratory illness. The respiratory health survey was approved by the Directorate of Education – Delhi and schools were selected based on land usage pattern i.e. commercial (Chandni Chowk), industrial (Mayapuri) and residential (Sarojini Nagar) areas. Approximately 1800 students (600 / zone) of age 10 – 14 years participated in the survey which included spirometry tests also. Indoor and outdoor levels of SO₂, NO₂ and PM were also measured in the school premises.

Results

The questionnaire data showed that the students having respiratory disorder symptoms was maximum in Chandni Chowk (66%) followed by Mayapuri (59%) and Sarojini Nagar (46%). Spirometry test results demonstrated that a significant population of subjects in Chandni Chowk (19%) had mild to severe pulmonary obstruction. However the percentage of subjects with such conditions was comparatively less in Mayapuri (17%) and Sarojini Nagar (14%) area. Indoor and outdoor PM₁₀ concentration at schools located in Chandni Chowk, a commercial zone, was observed to be 815±354.45 µg/m³ & 337±85 µg/m³

respectively, which is ten times above the permissible limits. The PM₁₀ concentration was lower in Mayapuri (694.6±322.9 µg/m³ & 274±78 µg/m³), an industrial zone, and was least in Sarojini Nagar (534.3±94.22µg/m³ & 197±48 µg/m³) which is a residential zone. However, levels of SO₂ and NO₂ were under the permissible limits in all three areas.

Conclusions

There is a correlation between pollutant levels and symptoms of respiratory illness in children. Commercial zones like Chandni Chowk with high traffic movement and human activities contribute more PM which affects the pulmonary health of children.

Author details

¹Allergy and Immunology Section, Institute of Genomics and Integrative Biology, Delhi, India. ²Air Pollution, National Environmental Engineering Research Institute, Delhi Zonal Lab, India. ³Lipid Biochemistry, Institute of Genomics and Integrative Biology, Delhi, India.

Published: 23 April 2013

doi:10.1186/1939-4551-6-S1-P55

Cite this article as: Mathew et al.: Environmental and occupational respiratory diseases – 1057. Correlation between air pollution and respiratory health of school children in Delhi. *World Allergy Organization Journal* 2013 **6**(Suppl 1):P55.

¹Allergy and Immunology Section, Institute of Genomics and Integrative Biology, Delhi, India

Full list of author information is available at the end of the article