


CASE REPORT

Atrial fibrillation, an epiphenomenon of acute Stanford type-A aortic dissection with suspected intimo-intimal intussusception

Michele M. Ciulla¹  | Patrizia Vivona² | Alessandro Lemos³ | Fabiola Sozzi² | Ugo Cioffi⁴ | Alberto Testori⁵

¹Laboratory of Clinical Informatics and Cardiovascular Imaging, Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy

²Cardiovascular Diseases Unit, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan, Italy

³Department of Radiology, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan, Italy

⁴Department of Surgery, University of Milan, Milan, Italy

⁵General and Thoracic Surgery, Humanitas Research Hospital, Milan, Italy

Correspondence: Michele M. Ciulla, Department of Clinical Sciences and Community Health, IRCCS Cà Granda Ospedale Maggiore Policlinico, University of Milan, Via Francesco Sforza 35, 20122, Milan, Italy (michele.ciulla@unimi.it).

Key Clinical Message

Supraventricular arrhythmias can sometimes be “only” epiphenomena appearing during acute hypoxia, pneumonia, pulmonary embolism, and thrombosis. Indeed, atrial fibrillation is not rare in acute aortic dissection as it is estimated in about one half of patients and may be secondary to a perfusion deficit of the sinoatrial node artery.

KEYWORDS

atrial fibrillation, intimo-intimal intussusception, Stanford type-A aortic dissection

1 | INTRODUCTION

Acute aortic dissection (AAD) is the most common catastrophic event affecting the aorta,¹ with an estimated annual incidence approximately 5-30 per million. Ascending AAD occurs most commonly between 50 and 60 years of age. In the 2010 guidelines, the American Heart Association and the American College of Cardiology introduced the aortic dissection detection risk score (ADDRS) as a simple and systematic bedside tool guiding the diagnostic approach to suspected AAD^{2,3} that was revised in 2017 including the D-dimer assay. Hereby, we present a 64-year-old patient with a typical atrial fibrillation (AF)—a condition actually nonincluded into the ADDRIS—masking an AAD Stanford type-A extended into

the right coronary artery (RCA) originating from the aortic false lumen, axillary artery aneurysm, and thrombosis. We finally present our considerations on the fact that supraventricular arrhythmias may be secondary to a perfusion deficit of the sinoatrial node artery that, in our case, was related to the AAD involving the RCA that was the supplier.

2 | CASE REPORT

The patient was admitted to our emergency department with right arm weakness and pain started 2 hours before admission. The ECG showed AF with rapid ventricular response and mild ST-segment depression in lower-lateral leads that




FIGURE 1 ECG Tracing Showing a Typical Atrial Fibrillation (AF) with Rapid Ventricular Response and Mild ST-Segment Depression in Lateral Leads that Could also be Interpreted as an Underlying Overload and/or Ischemia

could also be interpreted as an underlying overload and/or ischemia (Figure 1). The patient was currently followed at our hospital for essential arterial hypertension with a satisfactory control of blood pressure by lacidipine; from the clinical records, previous supraventricular arrhythmias were excluded.

On the examination, the right arm was cold and radial and humeral pulses were absent. Color-Doppler echocardiography of the right arm showed aneurysmal dilatation of the right axillary artery. Computed tomography (CT) scan demonstrated an AAD Stanford type-A dissection of the ascending aorta with a suspect intimo-intimal intussusception (Figure 2) and the patient was promptly transferred to the cardio-thoracic unit.

3 | DISCUSSION

As far as we know, this is the second case reported in the literature of AAD Stanford type-A presenting with AF,⁴ and the presence of this arrhythmia was not helpful, on the contrary, it has contributed to confusing, at least, the initial clinical picture, although the peculiar symptomatology has addressed correctly to the execution of a contrast-enhanced CT that was consistent for the final diagnosis. Contrast-enhanced CT scanning,⁵ magnetic resonance imaging, and Transesophageal

echocardiography are accurate techniques that are useful. In our case, the irregular appearance of the intimal flap at axial contrast-enhanced CT was firstly misinterpreted as an intimo-intimal intussusception; however, the absence of circumferential dissection and the so-called toe sign helped us in the differential diagnosis from a common intimal flap on volume rendering images as shown in the figure panels. Furthermore, AAD is characterized by a spectrum of clinical presentations that can be misleading; data from the International Registry of Acute Aortic Dissection¹ support that, among the electrocardiographic abnormalities, nonspecific ST segment change is the most common, with an incidence of acute myocardial infarction of 4.8%, while AF is not mentioned. In our case, AF is possibly an epiphenomenon of the acute occlusion of the artery that supplies the sinus node (SN), an event to be fully featured within the AF etiological factors. An ischemic damage of the SN, that in a large percentage of subjects is vascularized by the SN artery originated from the RC, is a well-known cause of cardiac arrhythmias.⁶ Indeed, not infrequently, the ischemic hypothesis⁷ of AF usually is not taken into great consideration, even if it could be an epiphenomenon appearing during acute hypoxia, pneumonia, pulmonary embolism.^{6,7} Finally, it should be said that even the acute thrombosis of the SN artery has been implied into sinus node dysfunction, in the context of coronary artery disease.⁸




FIGURE 2 Top Panels. Left: Volume Rendering 3D Reconstruction Shows Type A Dissection (arrows). Right: Axial contrast-enhancing CT scan with appropriate vascular window setting shows an AAD Stanford type-A dissection of the ascending aorta with suspected intimo-intimal intussusception. This suspect was ruled out by observing the absence of circumferential dissection and the so-called toe sign. Bottom Panels. Left: Axial contrast-enhancing CT scan shows no evidence of the origin of the right coronary artery, which was probably supplied by the false lumen. Right: Axial contrast-enhancing CT scan shows dilatation of the ascending aorta associated with dissection (type A, sec. Stanford). Patency of the left anterior descending artery (LAD). II, intimo-intimal intussusception; FL, false lumen; LAD, left anterior descending; LSA, left subclavian artery

4 | CONCLUSIONS

Aside the peculiar symptomatology, that in our case contributed, in a crucial way, to direct toward the solution of the clinical puzzle, it should be remembered that, even supraventricular AR, can sometimes be “only” epiphenomena of hypoxia. Indeed, AF is not rare in AAD,⁷ we suppose that the reduction of the oxygen supply to the sinoatrial nodal artery, that in 80% of the people origin from the right coronary,⁶ could induce arrhythmias. Finally, in order to reduce the time from diagnosis to surgery, that it is known to cut down the mortality of 50% in the first 48 hours,⁹ in our opinion, it is important to consider the supraventricular arrhythmias, alone with the D-dimer that has been recently included, to the ADDRS to assess the susceptibility to develop arrhythmias.¹⁰

INFORMED CONSENT

Written informed consent was obtained from the patient for publication of this manuscript and accompanying images. A copy of the written consent is available for review by the First Author of this report.

CONFLICT OF INTEREST

The authors have no conflict of interests to declare.

AUTHORSHIP

MMC, PV, and FS: planned the case, revised the literature, and drafted the manuscript. AL: contributed to the diagnosis by imaging. UC and AT: provided critical feedback on the surgical approach. All authors: discussed the results, contributed to the final manuscript, and approved the final version.

ORCID

Michele M. Ciulla  <http://orcid.org/0000-0002-6717-6065>

REFERENCES

1. Hagan PG, Nienaber CA, Isselbacher EM, et al. The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA*. 2000;283(7):897-903.
2. Nazerian P, Mueller C, Soeiro AM, et al. Diagnostic Accuracy of the aortic dissection detection risk score Plus D-Dimer for acute aortic syndromes: the ADvISED prospective multicenter study. *Circulation*. 2018;137(3):250-258.
3. Tang L, Hu XQ, Zhou SH. Acute stanford type A aortic dissection mimicking acute myocardial infarction: a hidden catastrophe which should prompt greater vigilance. *Acta Cardiol Sin*. 2014;30(5):493-496.
4. Lemos AA, Pezzullo JC, Fasani P, et al. Can the unenhanced phase be eliminated from dual-phase CT angiography for

- chest pain? Implications for diagnostic accuracy in acute aortic intramural hematoma. *Am J Roentgenol.* 2014;203(6):1171-1180.
5. Ciulla MM, Astuti M, Carugo S. The atherosclerosis of the sinus node artery is associated with an increased history of supra-ventricular arrhythmias: a retrospective study on 541 standard coronary angiograms. *PeerJ.* 2015;3:e1156.
 6. Ciulla MM, Astuti M, Vivona P, Gallazzi E, Meazza R, Lombardi F. The revascularization of the atrio-ventricular node artery in a case of right coronary artery occlusion promptly discontinue the atrio-ventricular block. *Hellenic J Cardiol.* 2017;58(1):77-79.
 7. Koracevic G, Djordjevic D, Glasnovic J. Is significance of atrial fibrillation in acute aortic dissection underestimated? *J Emerg Med.* 2009;37(2):168-171.
 8. Ando' G, Gaspardone A, Proietti I. Acute thrombosis of the sinus node artery: arrhythmological implications. *Heart.* 2003;89(2):E5.
 9. Pepper J. Differential aspects of the disease and treatment of Thoracic Acute Aortic Dissection (TAAD)-the European experience. *Ann Cardiothorac Surg.* 2016;5(4):360-367.
 10. Erbel R, Alfonso F, Boileau C, et al. Diagnosis and management of aortic dissection. *Eur Heart J.* 2001;22(18):1642-1681.

How to cite this article: Ciulla MM, Vivona P, Lemos A, Sozzi F, Cioffi U, Testori A. Atrial fibrillation, an epiphenomenon of acute Stanford type-A aortic dissection with suspected intimo-intimal intussusception. *Clin Case Rep.* 2018;6:1791–1794. <https://doi.org/10.1002/ccr3.1701>