

Research article

Open Access

Phenotypic alterations in breast cancer cells overexpressing the nuclear receptor co-activator AIB1

Sarah L Anzick^{1,2}, David O Azorsa³, S Stoney Simons Jr⁴ and Paul S Meltzer*¹

Address: ¹Cancer Genetics Branch, NHGRI, National Institutes of Health, Bethesda, MD, 20892, USA, ²Columbia School of Arts and Sciences, Program in Molecular and Cellular Oncology, George Washington University, Washington, D.C., 20037, USA, ³Translational Genomics Research Institute, Phoenix, AZ 85004, USA and ⁴Steroid Hormones Section, NIDDK, National Institutes of Health, Bethesda, MD 20892, USA

Email: Sarah L Anzick - sanzick@nhgri.nih.gov; David O Azorsa - dazorsa@tgen.org; S Stoney Simons - steroids@helix.gov; Paul S Meltzer* - pmeltzer@mail.nih.gov

* Corresponding author

Published: 10 September 2003

Received: 17 March 2003

BMC Cancer 2003, 3:22

Accepted: 10 September 2003

This article is available from: <http://www.biomedcentral.com/1471-2407/3/22>

© 2003 Anzick et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Estrogen signaling plays a critical role in a number of normal physiological processes and has important implications in the treatment of breast cancer. The p160 nuclear receptor coactivator, *AIB1* (amplified in breast cancer 1), is frequently amplified and overexpressed in human breast cancer and has been shown to enhance estrogen-dependent transactivation.

Methods: To better understand the molecular and physiological consequences of *AIB1* overexpression in breast cancer cells, an *AIB1* cDNA was transfected into the low *AIB1* expressing, estrogen-receptor (ER) negative breast cancer cell line, MDA-MB-436. The features of a derivative cell line, designated 436.1, which expresses high levels of *AIB1*, are described and compared with the parental cell line.

Results: A significant increase in the levels of CREB binding protein (CBP) was observed in 436.1 cells and immunofluorescent staining revealed altered *AIB1* and CBP staining patterns compared to the parental cells. Further, transient transfection assays demonstrated that the overall estrogen-dependent transactivation in 436.1 cells is approximately 20-fold higher than the parental cells and the estrogen dose-response curve is repositioned to the right. Finally, cDNA microarray analysis of approximately 7,100 cDNAs identified a number of differentially expressed genes in the 436.1 cells.

Conclusion: These observations lend insight into downstream signaling pathways that are influenced by *AIB1*.

Background

Estrogen hormone action is an essential regulator of physiologic responses ranging from reproductive development and function to bone formation and fat metabolism. Estrogens have also been considered one of the major adverse factors in the pathogenesis of breast cancer, the second leading cause of cancer-related deaths in American women [1,2].

Estrogen action is mediated through the estrogen receptors, ER- α and ER- β , members of the nuclear hormone receptor superfamily [3,4]. Also critical to estrogen signaling are the interactions of ER- α with transcriptional coactivators including p/CAF, CREB binding protein (CBP), p300, and the p160 family members (reviewed in [5]). It is thought that the coactivators activate transcription through the modification of histones and the bridging of

sequence specific transcription factors, including nuclear receptors, to the pre-initiation complex. Moreover, transcriptional activation through ER- α depends upon an ordered assembly of coactivator proteins on and off the promoter of the target gene in a cyclic fashion [6,7]. This selective modulation of cofactor assembly potentially contributes to the tightly regulated pattern of tissue-specific estrogen response.

The p160 coactivator, *AIB1*, was identified during a search for genes that are frequently amplified in human breast cancer [8]. The frequent gene amplification and overexpression of *AIB1* in ER-positive human breast cancers is hypothesized to confer a selective growth advantage through increased hormone signaling [8,9]. While it is likely that overexpression of *AIB1* contributes to tumorigenesis through this mechanism, overexpression of *AIB1* has also been observed in breast tumors that are ER-negative, suggesting increased expression of *AIB1* perturbs additional signaling pathways which may have important implications in the absence of ER [10].

In the present study, low *AIB1* expressing, ER-negative breast cancer cells, MDA-MB-436, were transfected with the human *AIB1* cDNA and selected for stable expression of *AIB1*. One of the clones expresses high levels of *AIB1* message and protein and is referred to as 436.1. These cells were used to investigate estrogen receptor activity in transient transfection assays and cDNA microarray analysis was performed to identify differentially expressed genes. Here, we report the characterization of the 436.1 cell line and suggest that it may be a useful tool for examining the effects of *AIB1* overexpression on a variety of signaling pathways.

Methods

The breast cancer cell line, MDA-MB-436, was obtained from the American Type Culture Collection (ATCC) and cultured in RPMI 1640 medium (Life Technologies) containing 10% fetal bovine serum. Stable transfections were performed using the standard calcium-phosphate technique using the CalPhos Maximizer System (Clontech) according to the manufacturer's recommended procedures. To generate stably transfected lines, MDA-MB-436 cells were seeded at 1.5×10^5 cells per well into 6-well culture dishes (20 mm²) and grown overnight. Cells were transfected with 2.0 μ g pcDNA3.1-*AIB1* (described in [8]) or empty pcDNA3.1 vector (Invitrogen) and refed the following day. After 48 hours, cells were transferred to 100 mm² dishes and selected in culture medium supplemented with 500 μ g/ml neomycin (G418) (Life Technologies). After approximately three weeks, neomycin-resistant colonies were isolated using cloning rings and transferred to 12-well dishes for expansion.

RNA was extracted from exponentially growing cells using the RNeasy kit (Qiagen). Following extraction, 10–15 μ g of total RNA was electrophoresed in a 1.0% formaldehyde agarose gel and transferred by standard methods [11]. An RNA ladder, 0.24–9.5 kb (Life Technologies), was used as a molecular size marker. Hybridization was performed using a ³²P-dCTP random-primed 2.2 kb Not/SacI fragment of pCMVSPORT-B11 as probe (described in [8]).

For Western blot analyses, cells were harvested and lysed in RIPA buffer (1X PBS, 1% NP40, 0.5% sodium deoxycholate, 0.1% SDS) containing complete protease inhibitor cocktail (Boehringer Mannheim). Cellular lysates were collected after centrifugation at 10,000 \times g for 15 minutes. Approximately 100 μ g of total protein was denatured by boiling for three minutes in sample loading buffer (Novex) and separated by SDS-polyacrylamide gel electrophoresis. To examine proteins from transient transfection assays, a total of 3 \times 10 μ l of cell extract from each triplicate was pooled for Western analysis. Resolved proteins were transferred onto nylon membranes and blocked overnight at 4°C in 5% milk/PBS. *AIB1* protein was detected by chemiluminescence using a mouse monoclonal antibody, AX15.2 [12]. Estrogen receptor and CBP were detected using the anti-ER- α polyclonal antibody HC-20 and the anti-CBP antibody A-11 (Santa Cruz Biotechnology), respectively. Secondary antibodies used were HRP-goat anti-mouse IgG Fc specific and HRP goat anti-rabbit IgG (Jackson ImmunoResearch) and detection was done using the ECL chemiluminescent substrate (Amersham) and exposing to autoradiography film (Hyperfilm-ECL, Amersham).

For fluorescent microscopy, adherent cells were grown on Falcon culture slides (Becton Dickinson) to 70% confluency. Cells were fixed with 2% paraformaldehyde, permeabilized with 1:1 methanol/acetone and blocked with 5% normal goat serum/1% casein/0.5% BSA/0.2% Triton X-100. Cells were incubated with 1:5 dilution of anti-*AIB1* mAb AAP.1 hybridoma culture supernatant [13] and anti-CBP for 1 hour at room temperature. After washing with PBS, the cells were incubated with 2 μ g/ml Rhodamine-conjugated goat anti-mouse IgG and 2 μ g/ml FITC-conjugated goat anti-rabbit IgG (Jackson ImmunoResearch) for 1 hour at room temperature. Following a final wash, DAPI mounting media and coverslips were added and slides were analyzed using a fluorescent microscope (Zeiss Axiophot).

To examine the effects of estrogen, cells were transiently transfected using Lipofectamine transfection reagent (Life Technologies). The transfections were performed with 1.0 μ g pGL3.luc.3ERE or pTK.luc.ERE reporter vectors, 0.4 or 10 ng of pHEGO-hyg (wild type estrogen receptor coding

sequence plus hygromycin resistance, ATCC), and 5.0 ng of pRL-tk (*Renilla* null luciferase reporter; Promega). The day following transfection, cells were incubated in the absence or presence of varying concentrations of 17 β -estradiol or 100 nM 4-hydroxytamoxifen. Cell lysates were harvested 48 hours post transfection and reporter activities were determined using the Dual Luciferase Reporter Assay system (Promega) and the supplier's recommended methods. The pGL3. luc.3ERE, which contains three tandem copies of the ERE upstream the Simian Virus 40 promoter driving the luciferase gene, was the kind gift of Fern Merdoch (University of Wisconsin). The tk. ERE. luc, which contains three ERE upstream the thymidine kinase promoter, was gift from Susan Fuqua (Baylor University). The activity for subsaturating concentrations of 17 β -estradiol was expressed as percent of maximal activity with saturating concentrations of 17 β -estradiol (10 nM). In cases in which saturation was reached at concentrations below 10 nM 17 β -estradiol, an average of the maximal activities was utilized to determine percent of maximal activity. The results are expressed in relative luciferase activity (Luciferase/*Renilla*).

cDNA microarray preparation and analyses were performed according to standard NHGRI protocol <http://www.nhgri.nih.gov/DIR/Microarray/main.html>. Microarrays consisted of 7,088 known human genes. Briefly, RNA was extracted from cells using TRIzol purification (Life Technologies) followed by purification with the RNeasy kit (Qiagen). Sixty to 80 μ g total RNA was fluorescently labeled with Cy5-dUTP or Cy3-dUTP (Amersham) using oligo-(dT)-primed reverse transcriptase (Superscript II, Life Technologies) and hybridized to PCR products immobilized on glass slides. Fluorescent images were acquired using a custom-designed confocal scanner equipped for dual laser excitation at the appropriate wavelengths. Pseudo-colored images were obtained by placing one fluorescent image into the green channel and the other into the red channel. A detection method was then employed to determine the actual target region based on the information from both red and green pixel values [14]. The ratios of the red intensity to the green intensity (R/G) for all targets were determined and normalized with 88 preselected internal control genes. The hybridization quality of each target area was based on fluorescence intensity background and signal to noise ratio, as described in detail at <http://www.nhgri.nih.gov/DIR/Microarray/main.html>.

Results

MDA-MB-436 cells were transfected with an expression vector containing the full-length human *AIB1* cDNA and cultured in the presence of neomycin for the selection of stable transfectants. While numerous resistant colonies

were selected for expansion, the majority gradually died following culture passage. The viable colonies were expanded and characterized. Additionally, five independently transfected populations were selected *en masse* and expanded for characterization.

To determine if the transfected cells expressed the exogenous *AIB1* cDNA, RNAs from viable clonal populations were analyzed by Northern blot. One of the clones, designated 436.1, showed significant expression of the predicted 5.7 kb transcript (Fig. 1A). This transcript was not detected in other transfected clones (e.g., 436.5) or in five *en masse* selected populations (data not shown), suggesting the exogenous *AIB1* cDNA was not highly expressed in these cell populations. Next, Western blot analysis of whole cell lysates from the parental MDA-MB-436 and the 436.1 cells was performed to ascertain if 436.1 cells displayed increased *AIB1* protein. Figure 1B shows high levels of *AIB1* protein in the 436.1 cells compared to the parental cells. Taken together, these results illustrate that the 436.1 cells integrated and significantly expressed the transfected *AIB1* cDNA.

Previous reports have demonstrated compensatory effects of p160 coactivator expression levels with the loss of one of the p160 coactivators [15,16]. In *SRC-1* knockout mice, for example, the expression of *TIF2* is increased in certain tissues. To examine if the increase in *AIB1* mRNA expression in 436.1 cells was paralleled by changes in *SRC-1* or *TIF2* coactivator levels, the Northern blot was stripped and hybridized with cDNAs of *SRC-1* and *TIF2*. Densitometric measurements indicated approximately 10% and 15% reduction in the expression levels of *SRC-1* and *TIF2*, respectively, in the 436.1 cells compared to the MDA-MB-436 parental cells or the *AIB1* transfected cell line, 436.5, which failed to express the 5.7 kb transcript (Fig. 1C).

To determine if the levels of the general co-integrator proteins, CBP or p300, had changed in the 436.1 cells, the Western blot from Figure 1B was stripped and probed with either anti-CBP or anti-p300 antibodies. Interestingly, a significant increase in the levels of CBP protein was observed in the 436.1 cells compared to the parental cells, whereas levels of p300 appeared slightly reduced (Fig. 1D).

The 436.1 cells are morphologically distinct from the parental MDA-MB-436 cells. While the MDA-MB-436 cells display a typical epithelial morphology, the most conspicuous feature of the 436.1 population was the appearance of giant cells with abundant cytoplasm, often exhibiting multiple nuclei or a multilobular nucleus (Fig. 2).




Figure 1

Analysis of coactivator RNA and protein levels in BT474 and wild-type and *AIB1*-transfected MDA-MB-436 cells. (A) RNAs from *AIB1* transfected 436.1 and 436.5 cells, and breast cancer cell lines, BT474 and MDA-MB-436, were analyzed by Northern blot. All cell lines expressed a full-length endogenous *AIB1* transcript of 9.0 kb, whereas a transcript of 5.7 kb derived from the exogenous *AIB1* cDNA was detected in an *AIB1* transfectant (436.1). The corresponding ethidium bromide-stained gel indicates equivalent loading. (B) Whole cell lysates from wild-type MDA-MB-436 and from *AIB1*-transfected 436.1 cells were analyzed by Western blot. The 160 kDa *AIB1* protein was detected in both cell lines, with increased amounts seen in *AIB1*-transfected 436.1 cells. Tubulin is included as a loading control. (C) The membrane used in A was stripped and hybridized with probes for *SRC-1* or *TIF2* cDNAs. (D) The membrane used in B was stripped and probed with anti-CBP and anti-p300 antibodies.

The cellular distribution of *AIB1* protein in MDA-MB-436 and 436.1 cells was examined by immunofluorescence (Fig. 3). *AIB1* staining of MDA-MB-436 cells produced a diffuse nuclear and faint cytoplasmic pattern (Fig. 3A). In addition to the nuclear staining, bright speckles were localized to the nucleoli in some cells. *AIB1* distribution in the majority of 436.1 cells was similar to that observed in the parental MDA-MB-436 line, but the staining




Figure 2

Comparison of the cellular characteristics of MDA-MB-436 and the transfected, 436.1 cells. MDA-MB-436 and 436.1 cell morphology was examined using a Nikon microscope equipped with 10× and 20× objectives. Images were captured using a 35 mm Nikon camera. Note the large, cells with multilobular nuclei in the 436.1 culture (bracket).

appeared brighter and more granular (Fig. 3B). The brighter speckles localized in the nucleoli were also apparent and often arranged at the nucleolar periphery.

The majority (~99%) of 436.1 cells showed bright diffuse nuclear and cytoplasmic *AIB1* staining, yet additional staining patterns of *AIB1* were also seen in 436.1 cells that were not apparent in the parental MDA-MB-436 cells. Some of the cells displayed bright, dot-like nuclear *AIB1* staining (Fig. 3C) or had reduced nuclear signal with profound cytoplasmic staining (Fig. 3D) that appeared to be concentrated in cytoplasmic organelles.

Since Western blot analysis revealed increased levels of CBP protein in the 436.1 cells, an anti-CBP antibody was used for immunofluorescent detection of endogenous CBP in each of the cell lines. Interestingly, CBP staining was faint and homogeneously diffuse throughout the nucleus in the MDA-MB-436 cells (Fig. 3A); while in the 436.1 cells, three or more bright, dot-like foci appeared in the nucleus, in addition to the diffuse nuclear staining (Fig. 3B,3C,3D). These results suggest upregulation of CBP leads to altered protein distribution within the cells. A clear correlation between the number of CBP dots and the different *AIB1* staining patterns could not be




Figure 3
Immunofluorescence of AIB1 and CBP in MDA-MB-436 (top row) and 436.1 (bracket) cells. Primary antibodies are as indicated. The cells were fixed at approximately 70% confluency and counterstained with DAPI (first panel). Red corresponds to AIB1 staining revealed with the Cy3-conjugated secondary goat anti-mouse antibody (second panel) and green corresponds to CBP staining with fluorescein isothiocyanate-conjugated goat anti-rabbit secondary antibody (third panel). The DAPI, AIB1, and CBP images from the same field of cells are merged in the last panel.

established, though in a few cells some CBP dots co-localized with AIB1 dots (data not shown).

The results described above demonstrate that the 436.1 cells have integrated and expressed exogenous *AIB1*, with small reductions in the expression of TIF2 and SRC-1, and a large increase in CBP levels. The effects of these alterations on estrogen receptor- α dependent transactivation in 436.1 cells were examined in transient transfection assays using an ER reporter and varying concentrations of an estrogen receptor- α expression plasmid. An estrogen-induced response was seen in both cell lines, but the 436.1 cells displayed a dramatic increase (approximately 20-fold) in the overall luciferase reporter gene activity compared with the parental line (Fig. 4). The increase in luciferase activity in 436.1 cells was observed in both the absence and presence of added ligand. Moreover, an increase in reporter activity was also observed with




Figure 4
AIB1 transfected 436.1 cells display increased reporter gene activity. Cells were transiently transfected with increasing amounts of estrogen receptor expression plasmid and 1.5 μ g of pGL3. ERE. luc reporter vector and incubated in the presence or absence of ethanol (EtOH), 10 nM 17 β -estradiol (E2), or 100 nM 4-hydroxytamoxifen (Tam). Results are representative of three experiments. Relative luciferase activity = Luciferase/Renilla.

increasing amounts of transfected receptor (50 ng to 500 ng). These results suggest increased expression of *AIB1* and CBP augments the capacity of 436.1 cells to assemble functional ER/ERE coregulator complexes.

The dose-response curve for a particular steroid predicts the biological response of the exposed cell to a given concentration of ligand. Since the levels of transactivation were profoundly higher in 436.1 cells compared to MDA-MB-436 cells, we asked if the estrogen dose-response curve was modified in the 436.1 line compared to MDA-MB-436 cells. For these studies, different concentrations of ER were used to ascertain that the receptor was limiting and that increased response is obtained over a range of steroid concentrations. Thus, transient transfections were performed in which 0.4 ng and 10 ng of ER- α plasmid were transfected into MDA-MB-436 and 436.1 cells and incubated with increasing concentrations of 17 β -estradiol. In 436.1 cells, increasing amounts of ER- α from 0.4 ng to 10 ng increased reporter gene expression at each concentration of steroid (Fig. 5A). To determine if the dose-response curve of estradiol has been altered by the addition of ER- α , the data from Figure 5A is replotted as a percent of the maximal activity induced by a saturating concentration of 17 β -estradiol (Fig. 5B). Significantly, concentrations. The the dose-response curve is shifted to a

lower EC_{50} with higher $ER-\alpha$ average left shift in the dose-response curve was 2.2 ± 0.33 fold (SEM, $n = 8$, $P = 0.0085$) with increasing $ER-\alpha$ plasmid. This effect was also seen in the MDA-MB-436 cells, but total gene expression reached maximal activity at a much lower ligand concentration. Figure 5C illustrates the dose-response curves of both MDA-MB-436 and 436.1 cells transfected with 0.4 ng $ER-\alpha$. The EC_{50} of 13 ± 4 pM (SEM, $n = 5$) in MDA-MB-436 cells was significantly right shifted in 436.1 cells to 118 ± 20 pM (SEM, $n = 8$, $P = 0.0016$ by Mann-Whitney test) 17 β -estradiol.

Next, we determined if the increased levels of AIB1 and CBP coactivators in 436.1 cells modified the levels of transfected $ER-\alpha$. Cellular lysates from MDA-MB-436 and 436.1 cells transfected with 10 ng and 100 ng of $ER-\alpha$ were analyzed by Western blot. The levels of full length $ER-\alpha$ protein were not significantly increased in the 436.1 cells relative to the parental cells, indicating the observed increase in ER-dependent transactivation in these cells is not the result of increased receptor levels (Fig. 5D).

Finally, cDNA microarray analysis was performed to examine endogenous gene expression [8] in the 436.1 cells compared to MDA-MB-436, and to identify potential downstream effects of AIB1 overexpression. Total RNAs from MDA-MB-436 and 436.1 cells were reverse-transcribed and labeled with fluorescent Cy3 and Cy5 dyes. Replicate experiments were also performed in which the fluorescent labels were reversed. Labeled RNAs were hybridized to a cDNA microarray containing 7,088 cDNAs arrayed on a glass slide. Out of 3,658 genes which met hybridization quality criteria (see methods), a total of 94 and 87 genes were upregulated and downregulated, respectively, by a factor of 3 or more in 436.1 cells compared to MDA-MB-436 cells. The 30 most significant outliers in each category are listed in Table 1 [see additional file 1].

Discussion

Vast epidemiological data supports a causative role for estrogens in the development of human breast cancer. Though significant progress has been made in the identification and characterization of a large number of proteins that participate in estrogen signaling, the precise molecular mechanisms that control specific cellular responses to endogenous and exogenous ligands of the ER signaling pathway are complex and not completely understood. However, increased understanding of hormone action suggests that the cell type and/or co-regulator levels dictate the ultimate response to ligand. The present study was initiated to examine the cellular and molecular consequences of AIB1 overexpression.

Our previous report demonstrated that the majority of ER-positive breast cancer cell lines have correspondingly high levels of endogenous AIB1 expression (four out of five cell lines analyzed); therefore, the number of suitable cell lines in which to generate stable overexpression of AIB1 are limiting [8]. Of interest, transfection of the AIB1 cDNA into the ER-positive, low AIB1-expressing cell line, T47-D, demonstrated a cell-lethal phenotype and no subclones with high levels of AIB1 expression were generated (data not shown).

Likewise, transfection of AIB1 into MDA-MB-436 breast cancer cells appears to be lethal since all but one of the surviving subclones failed to demonstrate expression of the transfected AIB1. The 436.1 line discussed in this study, however, has integrated the exogenous AIB1 cDNA and maintained high levels of AIB1 mRNA and protein expression. Most likely, 436.1 cells have acquired additional alterations which support its survival.

Though the parental MDA-MB-436 cells are $ER-\alpha$ negative, transient transfection of $ER\alpha$ was capable of inducing expression of an ERE-regulated reporter gene in the presence of estrogen, indicating these cells contain sufficient amounts of co-regulatory factors required for estrogen inducibility. Furthermore, since the levels of $ER-\alpha$ could be experimentally modulated in transient transfection studies, this derivative cell line provided a useful model to compare the effects of low-level, constitutive versus high-level expression of AIB1 in the presence or absence of varying concentrations of $ER-\alpha$.

Immunostaining patterns of AIB1 and CBP included diffuse fluorescent staining as well as large, bright, dot-like foci (Fig. 3). Although the punctate-staining pattern of AIB1 in 436.1 cells resembled CBP containing PML (promyelocytic leukemia protein) oncogenic domain (POD)-like structures [17,18], the AIB1 dots appeared in only a small percentage of the population (< 2%) and typically did not overlap with CBP. Analyses of populations derived from single cell isolates revealed that the punctate staining was not a clonal variant. Similarly, Baumann *et al* used GFP-fused to GRIP1 to investigate its intracellular distribution in HeLa cells and observed both diffuse and focal accumulation of GRIP1 [19]. CBP staining, on the other hand, was consistently diffuse throughout the nucleus in MDA-MB-436 cells but appeared to accumulate in PML-like bodies in the AIB1-transfected 436.1 cells. These results are consistent with observations made recently by Boisvert *et al* [20]; several cell lines do not accumulate CBP into PML bodies, but transfection of a GFP-CBP fusion protein into these cells leads to accumulation of CBP into distinct PML nuclear bodies. Taken together, these observations imply that increased amounts of AIB1 and CBP have caused the redistribution




Figure 5

Dose-dependent stimulation of luciferase activity by 17β-estradiol. MDA-MB-436 and 436.1 cells were transiently transfected with 0.4 or 10 ng of estrogen receptor plasmid (pHEGO-hyg) and incubated in the presence of ethanol or increasing amounts of 17β-estradiol. The total levels of luciferase seen with ethanol and varying levels 17β-estradiol in 436.1 cells are plotted in (A). The data from (A) was replotted in (B) as a percentage of the maximal activity with 10 nM 17β-estradiol, as described in the Materials and Methods section. To facilitate direct comparison of the position of the dose-response curves for the two cell lines, the data from (A) and that obtained with MDA-MB-436 cells with 0.4 ng ER (data not shown) is illustrated in panel (C). The data are representative of eight (436.1) and five (MDA-MB-436) experiments. (D) Whole cell lysates from ER-positive BT474 and MCF-7 breast cancer cells and transiently transfected MDA-MB-436 and 436.1 cells were analyzed by Western blot. The 58 kDa ER protein (arrow) was detected in ER-positive BT474 and MCF-7 cells, as well as both cell lines that were transfected with 10 and 100 ng of ER.

of these proteins into nuclear foci in 436.1 cells. Perhaps the levels of CBP and AIB1 are below the level required for recruitment to nuclear bodies in the non-transfected MDA-MB-436 cells. The mechanisms for upregulation of CBP in 436.1 cells remain elusive, but it will be interesting to determine if overexpression of AIB1 plays a direct role in the transcriptional regulation of CBP, or if levels of CBP are sensitized to and somehow adjusted with levels of AIB1, or vice versa.

While the overall level of reporter gene activity was dramatically increased in 436.1 cells, perhaps the most striking observation is in the right in the shift of the estrogen dose-response curve in 436.1 cells. The repositioning of the E2 dose-response curve to the right in 436.1 cells was somewhat unexpected since several studies with other receptors have demonstrated a left shift in the dose-response curve with the addition of p160 proteins, including TIF2 and SRC-1 [21–24]. However, there is no evidence that the total levels of transactivation and the position of the dose-response curve are linked. Studies with glucocorticoid [21,25–28] and progesterone [22,28] receptors show that the two phenomena can be totally disconnected. Increased total transactivation might be expected if the concentrations of two coactivators (AIB1 and CBP) are increased, but this does not predict the effects of these changes on the dose-response curve and EC₅₀.

One explanation for these observations is squelching, where increased levels of AIB1 compete for some limiting factor that is crucial for nuclear receptor transactivation, such as CBP. Because we have also demonstrated that the levels of CBP are significantly elevated in the 436.1 cells it is unlikely that the level of CBP is limiting. However, AIB1 and CBP could be competing for a common factor that is now limiting due to AIB1 and CBP abundance [24]. Also, cell specific differences, which have caused some coactivators and corepressors to be apparently inactive [21,22] or to have opposite activities [22,28], may be important.

We propose a hypothetical model to explain the increase in total transactivation and the overall right-shift in the dose-response curve, as schematized in Figure 6. In this model, the dose-response curve depends on the presence of a ternary complex composed of AIB1, CBP, and a third, unknown nuclear receptor complex participant, depicted as X. In MDA-MB-436 cells, the levels of AIB1, CBP, and factor X, are at concentrations sufficient to form ternary AIB1/CBP/X complexes that support the observed dose-response curve. The absence or dramatic reduction of such a ternary AIB1/CBP/X complex in 436.1 cells (due to increased levels of AIB1 and CBP and limiting levels of X) might result in a very different EC₅₀. In fact, by the law of

mass action, elevated levels of AIB1 and CBP would be expected to cause increased levels of the dimeric complexes (i.e., AIB1/X, AIB1/CBP, and CBP/X). Each of these may cause increased transactivation but not permit the same positioning of the dose-response curve as in MDA-MB-436 cells, resulting in a relative increase in the EC₅₀ and a right shift in the dose-response curve. The greatly decreased levels in 436.1 cells of the ternary AIB1/CBP/X complex, which has no effect on the level of total transactivation, would now have only minimal effects of the positioning of the dose-response curve.

Additional support for this model comes from several lines of evidence. First, convincing evidence for a limiting factor other than p160 coactivators has been demonstrated by Lopez *et al.* [23]. These investigators have shown that squelching between different receptors cannot be relieved by supplying excess amounts of CBP or p160 coactivator and suggest there is an unknown factor that is necessary for receptor coactivation. Similarly, Kino *et al.* reported that the amount of CBP that is required to see a decrease in glucocorticoid receptor (GR) transactivation is cell dependent [29]. The presence of a limiting complex component that participates in the positioning of the dose-response curve has been established by Chen *et al.* [24]. Here, the effects of modulators (GR, TIF2, or a cis-acting element in the reporter gene) were additive when all components were limiting, but no further effect was obtained if saturating amounts of any one modulator was present. Second, there is a well established physical interaction between CBP/p300 and p160 coactivators [13,30–32], and several groups have proposed that CBP and other coregulators, e.g., p160s, exist in the cytoplasm in preformed complexes [33,34].

Candidate genes whose altered expression might contribute to the phenotypic features of the 436.1 cells, such as cell survival, multinucleated morphology, and transcriptional properties were identified by cDNA microarray technology. Comparison of RNAs from MDA-MB-436 versus 436.1 revealed a number of profound changes in the expression of genes with a variety of functions, including cell adhesion, communication, maintenance of cell shape, apoptosis, and cell cycle control. For example, cyclin D3, an important regulator of endomitosis and polyploidization in megakaryocytes [35], was upregulated 3.75-fold in 436.1 cells, suggesting upregulation of cyclin D3 contributed to the altered morphological features observed in these cells. An 8.2-fold upregulation of the apoptosisinhibitor 2 gene may have facilitated the survival of these cells. Downregulation of *NFκB* and *C/EBPβ* cDNAs (ratios of 0.3 and 0.5, respectively) and a number of interleukins whose gene expression are regulated by these transcription factors (Table 1) possibly indicates a shift away from *NFκB* and *C/EBPβ* signaling to




Figure 6

A proposed model of how increased concentrations of AIB1 and CBP in 436.1 cells compete for a factor that is necessary for estrogen-induced transcription. As schematized on the left, transcriptional activation through the estrogen receptor relies on interaction with AIB1, CBP, and an unknown factor X. In MDA-MB-436 cells, none of these components are limiting. The 436.1 cells have limited amounts of factor X and excess quantities of AIB1 and CBP, which compete for binding to factor X. The absence or presence of a ternary complex of AIB1/CBP/X in 436.1 cells affects the EC₅₀ and transactivation, as schematized.

transcriptional activities mediated by other transcription factors. Finally, none of the general transcription factors, including TFIIA (ratio = 1.1), TFIIB (ratio = 1.0), TFIID (ratio=.53), TFIIIE (ratio = 1.25), TFIIF polypeptides 1 and 2 (ratio = 0.63, 0.94), and TFIIF (ratio = 1.16) showed increased expression in 436.1 cells, supporting the idea that AIB1 and CBP overexpression play a significant role in the observed increase in basal and E2-induced reporter gene activity in 436.1 cells. Future studies are necessary to confirm these observations and to determine which are genes regulated by transcriptional complexes containing AIB1.

In summary, we have characterized an *AIB1*-transfected cell line that displays high levels of *AIB1* expression. The

data presented here support a role for AIB1 in the estrogen-dose response and also suggest that, apart from ER-mediated pathways, AIB1 contributes to a number of signaling events which influence cell cycle, morphology, and survival. Future efforts directed at determining how overexpression of *AIB1* affects cell behavior in the presence or absence of specific hormone receptors should shed light onto specific signaling pathways that are altered as a consequence of *AIB1* amplification in breast cancer.

Competing interest

None declared.

Additional material

Additional File 1

The 30 most up- (ratio >3.0) or down- (ratio < 0.3) regulated genes in 436.1 cells compared to MDA-MB-436.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2407-3-22-S1.pdf>]

Acknowledgements

This work was done in partial fulfillment of the Ph.D. requirements of the Molecular and Cellular Oncology Program for S. L. Anzick. The authors wish to thank J. Lueders, B. Walker, K. Gayton, and H. Cunliffe for technical assistance and helpful discussions.

References

- McGuire WL, Chamness GC and Fuqua SA: **Estrogen receptor variants in clinical breast cancer.** *Mol Endocrinol* 1991, **5**:1571-1577.
- Dickson RB and Lippman ME: **Control of human breast cancer by estrogen, growth factors, and oncogenes.** *Cancer Treat Res* 1988, **40**:119-165.
- Evans RM: **The steroid and thyroid hormone receptor superfamily.** *Science* 1988, **240**:889-895.
- Dechering K, Boersma C and Mosselman S: **Estrogen receptors alpha and beta: two receptors of a kind?** *Curr Med Chem* 2000, **7**:561-576.
- Robyr D, Wolffe AP and Wahli W: **Nuclear hormone receptor coregulators in action: diversity for shared tasks.** *Mol Endocrinol* 2000, **14**:329-347.
- Kraus WL and Kadonaga JT: **p300 and estrogen receptor cooperatively activate transcription via differential enhancement of initiation and reinitiation.** *Genes Dev* 1998, **12**:331-342.
- Shang Y, Hu X, DiRenzo J, Lazar MA and Brown M: **Cofactor dynamics and sufficiency in estrogen receptor-regulated transcription.** *Cell* 2000, **103**:843-852.
- Anzick SL, Kononen J, Walker RL, Azorsa DO, Tanner MM, Guan XY, Sauter G, Kallioniemi OP, Trent JM and Meltzer PS: **AIB1, a steroid receptor coactivator amplified in breast and ovarian cancer.** *Science* 1997, **277**:965-968.
- Murphy LC, Simon SL, Parkes A, Leygue E, Dotzlaw H, Snell L, Troup S, Adeyinka A and Watson PH: **Altered expression of estrogen receptor coregulators during human breast tumorigenesis.** *Cancer Res* 2000, **60**:6266-6271.
- Bouras T, Southey MC and Venter DJ: **Overexpression of the steroid receptor coactivator AIB1 in breast cancer correlates with the absence of estrogen and progesterone receptors and positivity for p53 and HER2/neu.** *Cancer Res* 2001, **61**:903-907.
- Sambrook J, Fritsch EF and Maniatis T: **Molecular cloning: A laboratory manual.** Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N 1989.
- Azorsa DO and Meltzer PS: **Production and characterization of monoclonal antibodies to the steroid receptor coactivator AIB1.** *Hybridoma* 1999, **18**:281-287.
- Azorsa DO, Cunliffe HE and Meltzer PS: **Association of steroid receptor coactivator AIB1 with estrogen receptor-alpha in breast cancer cells.** *Breast Cancer Res Treat* 2001, **70**:89-101.
- Chen Y, Dougherty ER and Bittner ML: **Ratio-based decisions and the quantitative analysis of cDNA microarray images.** *J Biomed Optics* 1997, **2**:364-374.
- Wang Z, Rose DW, Hermanson O, Liu F, Herman T, Wu W, Szedo D, Gleiberman A, Kronen A, Pratt K, Rosenfeld R, Glass CK and Rosenfeld MG: **Regulation of somatic growth by the p160 coactivator p/CIP.** *Proc Natl Acad Sci U S A* 2000, **97**:13549-13554.
- Xu J, Qiu Y, DeMayo FJ, Tsai SY, Tsai MJ and O'Malley BW: **Partial hormone resistance in mice with disruption of the steroid receptor coactivator-1 (SRC-1) gene.** *Science* 1998, **279**:1922-1925.
- Doucas V, Tini M, Egan DA and Evans RM: **Modulation of CREB binding protein function by the promyelocytic (PML) oncoprotein suggests a role for nuclear bodies in hormone signaling.** *Proc Natl Acad Sci U S A* 1999, **96**:2627-2632.
- LaMorte VJ, Dyck JA, Ochs RL and Evans RM: **Localization of nascent RNA and CREB binding protein with the PML-containing nuclear body.** *Proc Natl Acad Sci U S A* 1998, **95**:4991-4996.
- Baumann CT, Ma H, Wolford R, Reyes JC, Maruvada P, Lim C, Yen PM, Stallcup MR and Hager GL: **The glucocorticoid receptor interacting protein 1 (GRIP1) localizes in discrete nuclear foci that associate with ND10 bodies and are enriched in components of the 26S proteasome.** *Mol Endocrinol* 2001, **15**:485-500.
- Boisvert FM, Kruhlak MJ, Box AK, Hendzel MJ and Bazett-Jones DP: **The transcription coactivator CBP is a dynamic component of the promyelocytic leukemia nuclear body.** *J Cell Biol* 2001, **152**:1099-1106.
- Szapary D, Huang Y and Simons SS Jr: **Opposing effects of corepressor and coactivators in determining the dose-response curve of agonists, and residual agonist activity of antagonists, for glucocorticoid receptor-regulated gene expression.** *Mol Endocrinol* 1999, **13**:2108-2121.
- Giannoukos G, Szapary D, Smith CL, Meeker JE and Simons SS Jr: **New anti-progestins with partial agonist activity: potential selective progesterone receptor modulators (SPRMs) and probes for receptor- and coregulator-induced changes in progesterone receptor induction properties.** *Mol Endocrinol* 2001, **15**:255-270.
- Lopez GN, Webb P, Shinsako JH, Baxter JD, Greene GL and Kushner PJ: **Titration by estrogen receptor activation function-2 of targets that are downstream from coactivators.** *Mol Endocrinol* 1999, **13**:897-909.
- Chen S, Sarlis NJ and Simons SS Jr: **Evidence for a common step in three different processes for modulating the kinetic properties of glucocorticoid receptor-induced gene transcription.** *J Biol Chem* 2000, **275**:30106-30117.
- Oshima H and Simons SS Jr: **Sequence-selective interactions of transcription factor elements with tandem glucocorticoid-responsive elements at physiological steroid concentrations.** *J Biol Chem* 1993, **268**:26858-26865.
- Szapary D, Xu M and Simons SS Jr: **Induction properties of a transiently transfected glucocorticoid-responsive gene vary with glucocorticoid receptor concentration.** *J Biol Chem* 1996, **271**:30576-30582.
- Zeng H, Plisov SY and Simons SS Jr: **Ability of the glucocorticoid modulatory element to modify glucocorticoid receptor transactivation indicates parallel pathways for the expression of glucocorticoid modulatory element and glucocorticoid response element activities.** *Mol Cell Endocrinol* 2000, **162**:221-234.
- Song LN, Huse B, Rusconi S and Simons SS Jr: **Transactivation specificity of glucocorticoid versus progesterone receptors. Role of functionally different interactions of transcription factors with amino- and carboxyl-terminal receptor domains.** *J Biol Chem* 2001, **276**:24806-24816.
- Kino T, Nordeen SK and Chrousos GP: **Conditional modulation of glucocorticoid receptor activities by CREB-binding protein (CBP) and p300.** *J Steroid Biochem Mol Biol* 1999, **70**:15-25.
- Torchia J, Rose DW, Inostroza J, Kamei Y, Westin S, Glass CK and Rosenfeld MG: **The transcriptional co-activator p/CIP binds CBP and mediates nuclear-receptor function.** *Nature* 1997, **387**:677-684.
- Chen H, Lin RJ, Schiltz RL, Chakravarti D, Nash A, Nagy L, Privalsky ML, Nakatani Y and Evans RM: **Nuclear receptor coactivator ACTR is a novel histone acetyltransferase and forms a multimeric activation complex with P/CAF and CBP/p300.** *Cell* 1997, **90**:569-580.
- Kamei Y, Xu L, Heinzel T, Torchia J, Kurokawa R, Glass B, Lin SC, Heyman RA, Rose DW, Glass CK and Rosenfeld MG: **A CBP integrator complex mediates transcriptional activation and AP-1 inhibition by nuclear receptors.** *Cell* 1996, **85**:403-414.
- McKenna NJ, Nawaz Z, Tsai SY, Tsai MJ and O'Malley BW: **Distinct steady-state nuclear receptor coregulator complexes exist in vivo.** *Proc Natl Acad Sci U S A* 1998, **95**:11697-11702.

34. Yao TP, Ku G, Zhou N, Scully R and Livingston DM: **The nuclear hormone receptor coactivator SRC-1 is a specific target of p300.** *Proc Natl Acad Sci U S A* 1996, **93**:10626-10631.
35. Zimmet JM, Ladd D, Jackson CW, Stenberg PE and Ravid K: **A role for cyclin D3 in the endomitotic cell cycle.** *Mol Cell Biol* 1997, **17**:7248-7259.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/3/22/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

