

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Methyl 2,6-diphenyl-1-*p*-tolyl-4-(*p*-tolyl-amino)-1,2,5,6-tetrahydropyridine-3-carboxylate

K. N. Venugopala,^{a*} Susanta K. Nayak^{b*} and Bharti Odhav^a

^aDepartment of Biotechnology and Food Technology, Durban University of Technology, Durban 4001, South Africa, and ^bCenter for Nano Science and Technology at PoliMi, Istituto Italiano di Tecnologia, Via Pascoli 70/3, 20133 Milan, Italy

Correspondence e-mail: katharigattav@dut.ac.za, nksusa@gmail.com


Received 28 June 2012; accepted 3 July 2012

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.044; wR factor = 0.112; data-to-parameter ratio = 15.4.

In the title compound, $\text{C}_{33}\text{H}_{32}\text{N}_2\text{O}_2$, the tetrahydropyridine ring adopts a boat conformation with the carbonyl group in an *s-cis* conformation with respect to the $\text{C}=\text{C}$ bond of the six-membered tetrahydropyridine ring. The molecular conformation is stabilized by intramolecular $\text{N}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions. Formation of centrosymmetric head-to-head dimers is observed through pairwise intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. Additional weak $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions stabilize the three-dimensional molecular assembly.

Related literature

For background to the applications of piperidines, see: Pearson *et al.* (2005); Sakai *et al.* (1986); Nayak *et al.* (2011); Mishra & Ghosh (2011). For ring-puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{33}\text{H}_{32}\text{N}_2\text{O}_2$
 $M_r = 488.61$
 Monoclinic, $P2_1/c$
 $a = 13.3701$ (11) Å
 $b = 6.1744$ (5) Å
 $c = 31.797$ (2) Å
 $\beta = 90.381$ (2)°
 $V = 2624.8$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹
 $T = 173$ K
 $0.29 \times 0.13 \times 0.05$ mm

Data collection

Bruker Kappa DUO APEXII diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2008b)
 $T_{\min} = 0.978$, $T_{\max} = 0.996$
 16243 measured reflections
 5175 independent reflections
 3513 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$
 Standard reflections: 0

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.112$
 $S = 1.03$
 5175 reflections
 337 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.18$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

C_{g1} , C_{g2} and C_{g3} are the centroids of the $C29-C34$, $C7-C12$ and $C22-C27$ rings, respectively.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{O1}$	0.88	2.11	2.7343 (17)	128
$\text{C19}-\text{H19}\cdots\text{O2}$	0.95	2.47	3.213 (3)	135
$\text{N2}-\text{H2}\cdots\text{O1}^i$	0.88	2.52	3.2471 (17)	140
$\text{C23}-\text{H23}\cdots\text{O1}^{ii}$	0.95	2.40	3.236 (2)	147
$\text{C24}-\text{H24}\cdots\text{O1}^{iii}$	0.95	2.58	3.363 (2)	140
$\text{C27}-\text{H27}\cdots\text{C}_{g1}$	0.95	2.83	3.476 (2)	126
$\text{C13}-\text{H13B}\cdots\text{C}_{g2}^{iv}$	0.98	2.81	3.679 (2)	148
$\text{C21}-\text{H21C}\cdots\text{C}_{g3}^v$	0.98	2.97	3.487 (2)	114

Symmetry codes: (i) $-x + 1, -y + 2, -z$; (ii) $x, y - 1, z$; (iii) $-x + 1, -y + 1, -z$; (iv) $-x + 1, y + \frac{5}{2}, -z + \frac{1}{2}$; (v) $x + 1, -y + \frac{5}{2}, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008a); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008a); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: PLATON (Spek, 2009) and PARST (Nardelli, 1995).

The authors thank Durban University of Technology for facilities. KNV thanks NRF South Africa for a DST/NRF Innovation Postdoctoral Fellowship 2012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2490).

References

- Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
 Mishra, S. & Ghosh, R. (2011). *Tetrahedron Lett.* **52**, 2857–2861.

- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Nayak, S. K., Venugopala, K. N., Chopra, D. & Guru Row, T. N. (2011). *CrystEngComm*, **13**, 591–605.
- Pearson, M. S. M., Mathe-Allainmat, M., Fargeas, V. & Lebreton, J. (2005). *Eur. J. Org. Chem.* **36**, 2159–2191.
- Sakai, R., Higa, T., Jefford, C. W., Bernardinelli, G. & Manzamine, A. (1986). *J. Am. Chem. Soc.* **108**, 6404–6405.
- Sheldrick, G. M. (2008a). *Acta Cryst. A* **64**, 112–122.
- Sheldrick, G. M. (2008b). *SADABS*. University of Göttingen, Germany.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2012). E68, o2392–o2393 [doi:10.1107/S1600536812030309]

Methyl 2,6-diphenyl-1-*p*-tolyl-4-(*p*-tolylamino)-1,2,5,6-tetrahydropyridine-3-carboxylate

K. N. Venugopala, Susanta K. Nayak and Bharti Odhav

Comment

Piperidine iminocyclitols have shown significant pharmacological results against HIV and in cancer therapy. Two deoxy-nojirimycin derivatives have already found clinical applications in treatment of type-II diabetes and Gaucher's disease (Pearson *et al.*, 2005). Alkaloids containing the piperidine nucleus exhibited a promising wide range of biological activities such as antimicrobial, antiparasitic, cytotoxicity, anti-inflammatory, pesticidal and anti-HIV-1 properties (Sakai *et al.*, 1986; Mishra & Ghosh, 2011). We have been investigating conformational and packing features of on tetrahydropyrimidine derivatives of compound (Nayak *et al.*, 2011). In the extension of our recent work, we have been focusing on the synthesis and biological properties of piperidine analogues. Here, we determined and analyzed the single-crystal structure of the title compound

In the title molecule, the tetrahydropyridine ring adopts a boat conformation. The Cremer & Pople (1975) parameters are $Q = 0.603$ (2) Å, $\theta = 98.16$ (15)° and $\varphi = 65.53$ (15)° respectively. The carbonyl group is in a *s-cis* conformation with respect to the C=C bond of the six-membered tetrahydropyridine ring. The molecular conformation is stabilized by intramolecular N—H⋯O, C—H⋯O and C—H⋯ π [C27—H27⋯Cg1 (centroid of C29—C34)] interactions (Fig.1, Table 1). Bifurcated N—H⋯O hydrogen bond patterns are observed with both intra- and intermolecular interactions. Pairwise intermolecular N—H⋯O interactions lead to formation of centrosymmetric head to head dimers (Fig. 2a). Further, weak C—H⋯O and C—H⋯ π [C13—H13B⋯Cg2 (centroid of C7—C12); C21—H21C⋯Cg3 (centroid of C22—C27)] interactions stabilize the 3-dimensional molecular assembly (Fig. 2b).

Experimental


To a mixture of *p*-toluidine (0.01 mol), methylacetoacetate (0.005 mmol), ZnCl₂ (0.01 mol) and ethanol (10 ml) in a 50 ml round bottom flask was added benzaldehyde (0.01 mol). The reaction mixture was stirred at room temperature for 16 h, as monitored by TLC and then cooled to room temperature. The solid precipitate obtained was filtered and washed with aqueous ethanol to obtain a crude product (Yield 60 %; m. p. 222 (2) °C). Suitable crystals for X-ray analysis were grown from a mixture of ACN:THF (V/V;1:1) solvent by slow evaporation at room temperature.

Refinement


All H atoms were positioned geometrically, C—H = 0.95 Å, 0.98 Å, 0.99 Å, 1.0 Å for aromatic, methyl, methylene and methine hydrogen respectively and N—H = 0.88 Å, and were refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{N})$ for aromatic, methylene, methine and amine hydrogens, and $1.5 U_{\text{eq}}(\text{C})$ for methyl H atoms respectively.

Computing details

Data collection: *APEX2* (Bruker, 2008); cell refinement: *S SAINT* (Bruker, 2008); data reduction: *S SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008a); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008a); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009) and *PARST* (Nardelli, 1995).

**Figure 1**

View of the molecular structure showing the atom labelling scheme with displacement ellipsoids for non-H atoms at the 30% probability level. Hydrogen atoms are shown as spheres of arbitrary radius. Dotted lines indicate intramolecular N—H...O, C—H...O, and C—H... π interactions.

**Figure 2**

(a) Bifurcated N—H···O hydrogen bonds with both intra- and intermolecular interactions, where the later interactions lead to formation of centrosymmetric head to head dimers, (b) Weak C—H··· π and C—H···O interactions further stabilize the 3-dimensional molecular assembly.

Methyl 2,6-diphenyl-1-*p*-tolyl-4-(*p*-tolylamino)-1,2,5,6-tetrahydropyridine-3-carboxylate

Crystal data

$C_{33}H_{32}N_2O_2$

$M_r = 488.61$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 13.3701$ (11) Å

$b = 6.1744$ (5) Å

$c = 31.797$ (2) Å

$\beta = 90.381$ (2)°

$V = 2624.8$ (4) Å³

$Z = 4$

$F(000) = 1040$

$D_x = 1.236$ Mg m⁻³

Melting point: 495(2) K

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 16734 reflections
 $\theta = 1.5\text{--}26.0^\circ$
 $\mu = 0.08\text{ mm}^{-1}$

$T = 173\text{ K}$
 Plate, yellow
 $0.29 \times 0.13 \times 0.05\text{ mm}$

Data collection

Bruker Kappa DUO APEXII
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 $0.5^\circ \varphi$ scans and ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 2008b)
 $T_{\min} = 0.978$, $T_{\max} = 0.996$

16243 measured reflections
 5175 independent reflections
 3513 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -15 \rightarrow 16$
 $k = -7 \rightarrow 7$
 $l = -39 \rightarrow 37$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.112$
 $S = 1.03$
 5175 reflections
 337 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0564P)^2 + 0.077P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.26\text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.39224 (9)	1.07628 (18)	0.01596 (3)	0.0325 (3)
O2	0.25859 (9)	1.23990 (17)	0.04498 (3)	0.0329 (3)
N1	0.26268 (10)	0.8892 (2)	0.15100 (4)	0.0282 (3)
N2	0.47196 (10)	0.7370 (2)	0.06014 (4)	0.0300 (3)
H2	0.4771	0.8083	0.0363	0.036*
C2	0.23238 (12)	0.9440 (3)	0.10784 (5)	0.0273 (4)
H2A	0.2120	1.0998	0.1082	0.033*
C3	0.31945 (12)	0.9271 (2)	0.07746 (5)	0.0263 (4)
C4	0.38851 (12)	0.7657 (2)	0.08314 (5)	0.0266 (4)
C5	0.36664 (13)	0.6094 (2)	0.11802 (5)	0.0290 (4)
H5A	0.4256	0.5155	0.1229	0.035*
H5B	0.3097	0.5159	0.1098	0.035*
C6	0.34127 (12)	0.7301 (2)	0.15878 (5)	0.0272 (4)
H6	0.3153	0.6218	0.1794	0.033*

C7	0.21331 (12)	0.9795 (3)	0.18521 (5)	0.0274 (4)
C8	0.14302 (14)	1.1459 (3)	0.18094 (5)	0.0408 (4)
H8	0.1251	1.1947	0.1536	0.049*
C9	0.09902 (15)	1.2410 (3)	0.21542 (6)	0.0479 (5)
H9	0.0531	1.3564	0.2111	0.057*
C10	0.11944 (14)	1.1743 (3)	0.25632 (5)	0.0410 (4)
C11	0.18535 (14)	1.0036 (3)	0.26032 (5)	0.0400 (4)
H11	0.1992	0.9487	0.2876	0.048*
C12	0.23208 (13)	0.9092 (3)	0.22634 (5)	0.0360 (4)
H12	0.2780	0.7940	0.2310	0.043*
C13	0.07216 (17)	1.2833 (4)	0.29375 (6)	0.0606 (6)
H13A	0.0836	1.1951	0.3190	0.091*
H13B	0.0001	1.2988	0.2888	0.091*
H13C	0.1021	1.4267	0.2978	0.091*
C14	0.14036 (12)	0.8157 (3)	0.09319 (5)	0.0299 (4)
C15	0.10575 (14)	0.6362 (3)	0.11428 (6)	0.0430 (5)
H15	0.1383	0.5919	0.1395	0.052*
C16	0.02464 (16)	0.5192 (3)	0.09954 (7)	0.0570 (6)
H16	0.0031	0.3944	0.1144	0.068*
C17	-0.02498 (16)	0.5812 (4)	0.06370 (6)	0.0561 (6)
H17	-0.0810	0.5011	0.0537	0.067*
C18	0.00758 (16)	0.7605 (4)	0.04252 (7)	0.0678 (7)
H18	-0.0262	0.8057	0.0177	0.081*
C19	0.08925 (15)	0.8762 (3)	0.05708 (6)	0.0541 (6)
H19	0.1108	1.0002	0.0419	0.065*
C20	0.32894 (12)	1.0818 (2)	0.04374 (5)	0.0267 (4)
C21	0.26050 (14)	1.3986 (3)	0.01188 (5)	0.0377 (4)
H21A	0.3238	1.4786	0.0131	0.057*
H21B	0.2046	1.4996	0.0154	0.057*
H21C	0.2544	1.3258	-0.0154	0.057*
C22	0.55320 (12)	0.5962 (3)	0.07236 (5)	0.0275 (4)
C23	0.56248 (13)	0.3923 (3)	0.05468 (5)	0.0316 (4)
H23	0.5148	0.3442	0.0344	0.038*
C24	0.64064 (13)	0.2587 (3)	0.06633 (5)	0.0352 (4)
H24	0.6460	0.1193	0.0539	0.042*
C25	0.71143 (13)	0.3237 (3)	0.09577 (5)	0.0348 (4)
C26	0.70173 (14)	0.5291 (3)	0.11301 (5)	0.0378 (4)
H26	0.7497	0.5776	0.1331	0.045*
C27	0.62366 (13)	0.6650 (3)	0.10161 (5)	0.0337 (4)
H27	0.6184	0.8049	0.1138	0.040*
C28	0.79502 (15)	0.1731 (3)	0.10946 (7)	0.0528 (5)
H28A	0.8358	0.2441	0.1312	0.079*
H28B	0.8370	0.1383	0.0852	0.079*
H28C	0.7664	0.0395	0.1209	0.079*
C29	0.43611 (12)	0.8289 (3)	0.17719 (5)	0.0277 (4)
C30	0.46725 (13)	1.0365 (3)	0.16685 (5)	0.0341 (4)
H30	0.4250	1.1271	0.1504	0.041*
C31	0.55940 (15)	1.1125 (3)	0.18032 (6)	0.0474 (5)
H31	0.5804	1.2539	0.1726	0.057*

C32	0.62073 (16)	0.9851 (4)	0.20476 (6)	0.0553 (6)
H32	0.6846	1.0367	0.2134	0.066*
C33	0.58895 (15)	0.7814 (4)	0.21667 (6)	0.0534 (6)
H33	0.6300	0.6947	0.2344	0.064*
C34	0.49762 (14)	0.7035 (3)	0.20290 (5)	0.0387 (4)
H34	0.4766	0.5627	0.2111	0.046*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0322 (7)	0.0414 (7)	0.0238 (6)	-0.0024 (5)	0.0044 (5)	0.0027 (5)
O2	0.0369 (7)	0.0333 (6)	0.0287 (6)	0.0033 (5)	0.0057 (5)	0.0072 (5)
N1	0.0286 (8)	0.0339 (7)	0.0220 (7)	0.0044 (6)	0.0020 (6)	0.0012 (6)
N2	0.0305 (8)	0.0362 (7)	0.0234 (7)	0.0005 (6)	0.0046 (6)	0.0017 (6)
C2	0.0296 (9)	0.0290 (8)	0.0232 (8)	0.0006 (7)	0.0018 (7)	0.0027 (7)
C3	0.0269 (9)	0.0295 (8)	0.0226 (8)	-0.0039 (7)	0.0014 (6)	-0.0016 (7)
C4	0.0297 (9)	0.0281 (8)	0.0218 (8)	-0.0054 (7)	0.0006 (7)	-0.0035 (7)
C5	0.0315 (10)	0.0259 (8)	0.0299 (9)	-0.0030 (7)	0.0044 (7)	0.0007 (7)
C6	0.0319 (9)	0.0251 (8)	0.0245 (8)	0.0011 (7)	0.0045 (7)	0.0042 (7)
C7	0.0243 (9)	0.0328 (8)	0.0252 (8)	-0.0034 (7)	0.0028 (6)	-0.0015 (7)
C8	0.0444 (12)	0.0486 (11)	0.0295 (9)	0.0122 (9)	0.0045 (8)	0.0043 (8)
C9	0.0477 (12)	0.0536 (12)	0.0424 (11)	0.0182 (10)	0.0075 (9)	-0.0021 (9)
C10	0.0340 (10)	0.0565 (11)	0.0325 (9)	0.0013 (9)	0.0060 (8)	-0.0097 (9)
C11	0.0349 (11)	0.0594 (11)	0.0258 (9)	0.0009 (9)	0.0020 (7)	-0.0012 (9)
C12	0.0351 (10)	0.0446 (10)	0.0284 (9)	0.0073 (8)	0.0025 (7)	0.0010 (8)
C13	0.0560 (14)	0.0836 (16)	0.0422 (11)	0.0148 (12)	0.0087 (10)	-0.0185 (11)
C14	0.0268 (9)	0.0357 (9)	0.0273 (8)	0.0029 (7)	0.0027 (7)	0.0051 (7)
C15	0.0372 (11)	0.0499 (11)	0.0416 (10)	-0.0104 (9)	-0.0085 (8)	0.0179 (9)
C16	0.0508 (14)	0.0558 (12)	0.0640 (14)	-0.0208 (10)	-0.0177 (11)	0.0249 (11)
C17	0.0398 (12)	0.0696 (14)	0.0586 (13)	-0.0189 (11)	-0.0161 (10)	0.0163 (11)
C18	0.0499 (14)	0.0917 (17)	0.0614 (14)	-0.0244 (13)	-0.0290 (11)	0.0386 (13)
C19	0.0430 (12)	0.0645 (13)	0.0547 (12)	-0.0165 (10)	-0.0146 (10)	0.0326 (11)
C20	0.0259 (9)	0.0311 (8)	0.0230 (8)	-0.0055 (7)	-0.0022 (7)	-0.0028 (7)
C21	0.0422 (11)	0.0378 (9)	0.0330 (9)	0.0016 (8)	0.0017 (8)	0.0115 (8)
C22	0.0271 (9)	0.0309 (8)	0.0245 (8)	-0.0020 (7)	0.0051 (7)	0.0014 (7)
C23	0.0358 (10)	0.0322 (9)	0.0269 (9)	-0.0057 (8)	0.0019 (7)	-0.0018 (7)
C24	0.0416 (11)	0.0285 (8)	0.0356 (9)	-0.0011 (8)	0.0065 (8)	-0.0003 (8)
C25	0.0309 (10)	0.0366 (9)	0.0369 (9)	-0.0015 (8)	0.0056 (8)	0.0093 (8)
C26	0.0352 (11)	0.0418 (10)	0.0365 (10)	-0.0067 (8)	-0.0057 (8)	0.0005 (8)
C27	0.0384 (11)	0.0323 (9)	0.0305 (9)	-0.0025 (8)	-0.0007 (8)	-0.0057 (7)
C28	0.0395 (12)	0.0491 (11)	0.0697 (14)	0.0027 (9)	-0.0011 (10)	0.0114 (11)
C29	0.0301 (9)	0.0328 (8)	0.0204 (8)	0.0037 (7)	0.0043 (7)	-0.0019 (7)
C30	0.0369 (10)	0.0326 (9)	0.0330 (9)	0.0000 (8)	0.0053 (8)	-0.0043 (8)
C31	0.0419 (12)	0.0509 (11)	0.0495 (12)	-0.0120 (10)	0.0110 (10)	-0.0199 (10)
C32	0.0337 (12)	0.0867 (16)	0.0457 (12)	-0.0020 (11)	0.0016 (9)	-0.0338 (12)
C33	0.0404 (12)	0.0858 (16)	0.0339 (10)	0.0226 (12)	-0.0071 (9)	-0.0114 (11)
C34	0.0423 (11)	0.0455 (10)	0.0282 (9)	0.0132 (9)	0.0058 (8)	0.0018 (8)

Geometric parameters (Å, °)

O1—C20	1.2277 (18)	C15—H15	0.9500
O2—C20	1.3562 (19)	C16—C17	1.369 (3)
O2—C21	1.4385 (18)	C16—H16	0.9500
N1—C7	1.393 (2)	C17—C18	1.368 (3)
N1—C6	1.458 (2)	C17—H17	0.9500
N1—C2	1.4679 (19)	C18—C19	1.382 (3)
N2—C4	1.3498 (19)	C18—H18	0.9500
N2—C22	1.443 (2)	C19—H19	0.9500
N2—H2	0.8800	C21—H21A	0.9800
C2—C3	1.521 (2)	C21—H21B	0.9800
C2—C14	1.533 (2)	C21—H21C	0.9800
C2—H2A	1.0000	C22—C23	1.384 (2)
C3—C4	1.370 (2)	C22—C27	1.386 (2)
C3—C20	1.442 (2)	C23—C24	1.380 (2)
C4—C5	1.500 (2)	C23—H23	0.9500
C5—C6	1.535 (2)	C24—C25	1.386 (2)
C5—H5A	0.9900	C24—H24	0.9500
C5—H5B	0.9900	C25—C26	1.388 (2)
C6—C29	1.521 (2)	C25—C28	1.516 (2)
C6—H6	1.0000	C26—C27	1.385 (2)
C7—C8	1.398 (2)	C26—H26	0.9500
C7—C12	1.399 (2)	C27—H27	0.9500
C8—C9	1.379 (2)	C28—H28A	0.9800
C8—H8	0.9500	C28—H28B	0.9800
C9—C10	1.389 (2)	C28—H28C	0.9800
C9—H9	0.9500	C29—C30	1.388 (2)
C10—C11	1.379 (3)	C29—C34	1.391 (2)
C10—C13	1.510 (2)	C30—C31	1.384 (3)
C11—C12	1.381 (2)	C30—H30	0.9500
C11—H11	0.9500	C31—C32	1.373 (3)
C12—H12	0.9500	C31—H31	0.9500
C13—H13A	0.9800	C32—C33	1.381 (3)
C13—H13B	0.9800	C32—H32	0.9500
C13—H13C	0.9800	C33—C34	1.381 (3)
C14—C15	1.377 (2)	C33—H33	0.9500
C14—C19	1.384 (2)	C34—H34	0.9500
C15—C16	1.382 (3)		
C20—O2—C21	116.96 (12)	C17—C16—H16	119.6
C7—N1—C6	118.83 (12)	C15—C16—H16	119.6
C7—N1—C2	120.57 (13)	C18—C17—C16	118.79 (19)
C6—N1—C2	120.52 (12)	C18—C17—H17	120.6
C4—N2—C22	123.83 (13)	C16—C17—H17	120.6
C4—N2—H2	118.1	C17—C18—C19	120.44 (18)
C22—N2—H2	118.1	C17—C18—H18	119.8
N1—C2—C3	111.74 (13)	C19—C18—H18	119.8
N1—C2—C14	112.39 (13)	C18—C19—C14	121.54 (18)
C3—C2—C14	112.80 (13)	C18—C19—H19	119.2

N1—C2—H2A	106.5	C14—C19—H19	119.2
C3—C2—H2A	106.5	O1—C20—O2	121.50 (14)
C14—C2—H2A	106.5	O1—C20—C3	125.50 (15)
C4—C3—C20	121.17 (14)	O2—C20—C3	113.00 (13)
C4—C3—C2	118.95 (13)	O2—C21—H21A	109.5
C20—C3—C2	119.88 (14)	O2—C21—H21B	109.5
N2—C4—C3	125.65 (14)	H21A—C21—H21B	109.5
N2—C4—C5	118.82 (14)	O2—C21—H21C	109.5
C3—C4—C5	115.53 (14)	H21A—C21—H21C	109.5
C4—C5—C6	110.94 (12)	H21B—C21—H21C	109.5
C4—C5—H5A	109.5	C23—C22—C27	119.24 (15)
C6—C5—H5A	109.5	C23—C22—N2	120.53 (14)
C4—C5—H5B	109.5	C27—C22—N2	120.23 (14)
C6—C5—H5B	109.5	C24—C23—C22	120.29 (16)
H5A—C5—H5B	108.0	C24—C23—H23	119.9
N1—C6—C29	113.15 (13)	C22—C23—H23	119.9
N1—C6—C5	110.29 (12)	C23—C24—C25	121.36 (16)
C29—C6—C5	109.36 (13)	C23—C24—H24	119.3
N1—C6—H6	108.0	C25—C24—H24	119.3
C29—C6—H6	108.0	C24—C25—C26	117.78 (16)
C5—C6—H6	108.0	C24—C25—C28	121.00 (16)
N1—C7—C8	122.69 (14)	C26—C25—C28	121.20 (17)
N1—C7—C12	121.50 (15)	C27—C26—C25	121.48 (16)
C8—C7—C12	115.80 (15)	C27—C26—H26	119.3
C9—C8—C7	121.71 (16)	C25—C26—H26	119.3
C9—C8—H8	119.1	C26—C27—C22	119.84 (16)
C7—C8—H8	119.1	C26—C27—H27	120.1
C8—C9—C10	122.38 (18)	C22—C27—H27	120.1
C8—C9—H9	118.8	C25—C28—H28A	109.5
C10—C9—H9	118.8	C25—C28—H28B	109.5
C11—C10—C9	115.75 (16)	H28A—C28—H28B	109.5
C11—C10—C13	122.59 (17)	C25—C28—H28C	109.5
C9—C10—C13	121.65 (18)	H28A—C28—H28C	109.5
C10—C11—C12	122.88 (17)	H28B—C28—H28C	109.5
C10—C11—H11	118.6	C30—C29—C34	118.43 (16)
C12—C11—H11	118.6	C30—C29—C6	122.00 (14)
C11—C12—C7	121.39 (16)	C34—C29—C6	119.41 (15)
C11—C12—H12	119.3	C31—C30—C29	120.52 (17)
C7—C12—H12	119.3	C31—C30—H30	119.7
C10—C13—H13A	109.5	C29—C30—H30	119.7
C10—C13—H13B	109.5	C32—C31—C30	120.53 (19)
H13A—C13—H13B	109.5	C32—C31—H31	119.7
C10—C13—H13C	109.5	C30—C31—H31	119.7
H13A—C13—H13C	109.5	C31—C32—C33	119.53 (19)
H13B—C13—H13C	109.5	C31—C32—H32	120.2
C15—C14—C19	117.11 (16)	C33—C32—H32	120.2
C15—C14—C2	122.64 (14)	C34—C33—C32	120.22 (19)
C19—C14—C2	120.24 (15)	C34—C33—H33	119.9
C14—C15—C16	121.39 (17)	C32—C33—H33	119.9

C14—C15—H15	119.3	C33—C34—C29	120.68 (18)
C16—C15—H15	119.3	C33—C34—H34	119.7
C17—C16—C15	120.71 (18)	C29—C34—H34	119.7
C7—N1—C2—C3	151.62 (14)	C3—C2—C14—C19	-65.7 (2)
C6—N1—C2—C3	-31.79 (19)	C19—C14—C15—C16	1.2 (3)
C7—N1—C2—C14	-80.38 (18)	C2—C14—C15—C16	-177.72 (19)
C6—N1—C2—C14	96.20 (17)	C14—C15—C16—C17	-1.2 (3)
N1—C2—C3—C4	36.66 (19)	C15—C16—C17—C18	0.4 (4)
C14—C2—C3—C4	-91.11 (17)	C16—C17—C18—C19	0.2 (4)
N1—C2—C3—C20	-142.40 (14)	C17—C18—C19—C14	-0.1 (4)
C14—C2—C3—C20	89.83 (17)	C15—C14—C19—C18	-0.6 (3)
C22—N2—C4—C3	166.10 (15)	C2—C14—C19—C18	178.4 (2)
C22—N2—C4—C5	-14.4 (2)	C21—O2—C20—O1	1.0 (2)
C20—C3—C4—N2	2.5 (2)	C21—O2—C20—C3	-178.39 (13)
C2—C3—C4—N2	-176.52 (14)	C4—C3—C20—O1	5.7 (2)
C20—C3—C4—C5	-176.98 (13)	C2—C3—C20—O1	-175.28 (15)
C2—C3—C4—C5	4.0 (2)	C4—C3—C20—O2	-174.94 (13)
N2—C4—C5—C6	131.03 (15)	C2—C3—C20—O2	4.1 (2)
C3—C4—C5—C6	-49.43 (18)	C4—N2—C22—C23	100.73 (18)
C7—N1—C6—C29	-71.72 (18)	C4—N2—C22—C27	-80.1 (2)
C2—N1—C6—C29	111.64 (15)	C27—C22—C23—C24	0.5 (2)
C7—N1—C6—C5	165.43 (13)	N2—C22—C23—C24	179.67 (15)
C2—N1—C6—C5	-11.21 (19)	C22—C23—C24—C25	0.0 (2)
C4—C5—C6—N1	52.06 (17)	C23—C24—C25—C26	-0.5 (2)
C4—C5—C6—C29	-72.98 (16)	C23—C24—C25—C28	177.92 (16)
C6—N1—C7—C8	174.53 (15)	C24—C25—C26—C27	0.5 (3)
C2—N1—C7—C8	-8.8 (2)	C28—C25—C26—C27	-177.91 (17)
C6—N1—C7—C12	-5.0 (2)	C25—C26—C27—C22	0.0 (3)
C2—N1—C7—C12	171.68 (15)	C23—C22—C27—C26	-0.5 (2)
N1—C7—C8—C9	-176.44 (17)	N2—C22—C27—C26	-179.67 (15)
C12—C7—C8—C9	3.1 (3)	N1—C6—C29—C30	-33.1 (2)
C7—C8—C9—C10	-1.8 (3)	C5—C6—C29—C30	90.31 (17)
C8—C9—C10—C11	-1.1 (3)	N1—C6—C29—C34	151.50 (14)
C8—C9—C10—C13	178.74 (19)	C5—C6—C29—C34	-85.13 (17)
C9—C10—C11—C12	2.7 (3)	C34—C29—C30—C31	2.9 (2)
C13—C10—C11—C12	-177.19 (19)	C6—C29—C30—C31	-172.61 (15)
C10—C11—C12—C7	-1.3 (3)	C29—C30—C31—C32	-1.2 (3)
N1—C7—C12—C11	177.96 (16)	C30—C31—C32—C33	-1.4 (3)
C8—C7—C12—C11	-1.6 (3)	C31—C32—C33—C34	2.2 (3)
N1—C2—C14—C15	-14.2 (2)	C32—C33—C34—C29	-0.5 (3)
C3—C2—C14—C15	113.24 (18)	C30—C29—C34—C33	-2.1 (2)
N1—C2—C14—C19	166.88 (16)	C6—C29—C34—C33	173.55 (15)

Hydrogen-bond geometry (Å, °)

Cg1, Cg2 and Cg3 are the centroids of the C29–C34, C7–C12 and C29–C34 rings, respectively. [Cg3 and Cg1 are the same ring (information taken from Comment section)?]

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2···O1	0.88	2.11	2.7343 (17)	128
C19—H19···O2	0.95	2.47	3.213 (3)	135
N2—H2···O1 ⁱ	0.88	2.52	3.2471 (17)	140
C23—H23···O1 ⁱⁱ	0.95	2.40	3.236 (2)	147
C24—H24···O1 ⁱⁱⁱ	0.95	2.58	3.363 (2)	140
C27—H27···Cg1	0.95	2.83	3.476 (2)	126
C13—H13B···Cg2 ^{iv}	0.98	2.81	3.679 (2)	148
C21—H21C···Cg3 ^v	0.98	2.97	3.487 (2)	114

Symmetry codes: (i) $-x+1, -y+2, -z$; (ii) $x, y-1, z$; (iii) $-x+1, -y+1, -z$; (iv) $-x+1, y+5/2, -z+1/2$; (v) $x+1, -y+5/2, z+1/2$.