



# Anterior Shoulder Instability and Open Procedures: History, Indications, and Clinical Outcomes

Mohamad Y. Fares, MD, Peter Boufadel, MD, Mohammad Daher, MD, Jonathan Koa, BSc, Akshay Khanna, BSc\*, Joseph A. Abboud, MD<sup>#</sup>

*Division of Shoulder and Elbow Surgery, Rothman Orthopaedic Institute, Philadelphia, PA,  
\*Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA, USA*

The shoulder, being the most mobile joint in the human body, is often susceptible to dislocations and subluxations more so than other joints. As such, shoulder instability constitutes a common complaint among patients worldwide, especially those who are young, participate in contact sports, and have increased innate flexibility in their joints. Management options in the setting of instability vary between conservative and surgical options that aim to mitigate symptoms and allow return of function. Surgical options can be arthroscopic and open, with a general shift among surgeons towards utilizing arthroscopic surgery in the past several decades. Nevertheless, open procedures still play a role in managing shoulder instability patients, especially those with significant bone loss, recurrent instability, coexisting shoulder pathologies, and high risk of failure with arthroscopic surgery. In these clinical settings, open procedures, like the Latarjet procedure, open Bankart repair, glenoid bone augmentation using iliac crest autograft or distal tibial allograft, and salvage options like glenohumeral arthrodesis and arthroplasty may show good clinical outcomes and low recurrence rates. Each of these open procedures possesses its own set of advantages and disadvantages and entails a specific set of indications based on published literature. It is important to cater treatment options to the individual patient in order to optimize outcomes and reduce the risk of complications. Future research on open shoulder stabilization procedures should focus on the long-term outcomes of recently utilized procedures, investigate different graft options for procedures involving bone augmentation, and conduct additional comparative analyses in order to establish concrete surgical management guidelines.

**Keywords:** *Latarjet, Bankart lesions, Glenoid bone loss, Contact athlete, Labrum*

The glenohumeral joint is the most mobile joint in the human body, and this mobility predisposes it to a higher risk of dislocations and instability.<sup>1)</sup> The natural stability of this joint is maintained through dynamic and static stabilizers that work together to ensure that the ball-and-socket joint remains in place.<sup>2)</sup> Dynamic stabilizers are composed

of the muscles and tendons that surround the shoulder, and these include the rotator cuff, which allows the active movement of the upper extremity in different directions.<sup>3,4)</sup> On the other hand, the static stabilizers include the joint's innate anatomy, the surrounding glenohumeral ligaments, and the glenoid labrum.<sup>2,5)</sup> The glenoid labrum is a thick tissue composed of fibrocartilage that surrounds the rim of the glenoid and helps stabilize the humeral head while it translates along the socket, preventing translocation of the humeral head beyond the glenoid.<sup>2,6)</sup> When a shoulder dislocation ensues, a tear in the labrum often results (termed a Bankart tear), along with a concomitant potential loss in glenoid bone, humeral head bone (termed a Hill-Sachs lesion), and injury to joint capsule.<sup>7,8)</sup> As such, joint instability rises and increases with every subsequent dislocation event.

Received January 16, 2023; Revised March 15, 2023;

Accepted April 16, 2023

Correspondence to: Joseph A. Abboud, MD

Division of Shoulder and Elbow Surgery, Rothman Orthopaedic Institute, Thomas Jefferson University, 925 Chestnut, Philadelphia, PA 19107, USA

Tel: +1-610-547-8351

E-mail: [abboudj@gmail.com](mailto:abboudj@gmail.com)

<sup>#</sup>Current affiliation: Division of Shoulder and Elbow Surgery, Rothman Orthopaedic Institute, Philadelphia, PA, USA

Shoulder instability is a very common complaint in the population, and it can present as an anterior (most common), posterior, or multidirectional instability depending on the type and severity of the dislocation event.<sup>9-11)</sup> Risk factors often include contact sports, young age, joint hypermobility, and prior dislocation events.<sup>7,12)</sup> It is estimated that the incidence of anterior shoulder instability is 0.08 per 1,000 person-years in the general population, with much higher reported rates for young high risk men at 3% per year.<sup>13,14)</sup> These high rates of anterior shoulder instability, along with the debilitating presenting symptoms—evident by pain, apprehensive mobility, and decreased function—necessitate careful consideration when deciding appropriate management plans.<sup>15)</sup> Treatment options for anterior instability vary between conservative and surgical measures; however, it is often advisable to undergo surgery following the initial dislocation event, especially in high risk young patients to prevent recurrent dislocations and concomitant bone loss.<sup>9)</sup> Surgical options include minimally invasive procedures like arthroscopic Bankart repair and more invasive open procedures like open Bankart repair, coracoid transfer procedure or Latarjet, and bone augmentation procedures using different autograft sources.<sup>7,9,16)</sup> Each of these procedures possesses its own set of advantages and disadvantages, and management decisions are often based on the severity of instability, evident through clinical and radiographic examination.<sup>9)</sup>

Arthroscopic Bankart repair is considered the most commonly used surgical procedure for shoulder instability, constituting 87% of shoulder stabilization techniques in 2006.<sup>17,18)</sup> Nevertheless, significantly high failure rates have been reported, mainly in patients with high risk of recurrent instability.<sup>19-22)</sup> In these cases, open techniques may be considered, as these can provide better and more sustainable outcomes.<sup>23)</sup> That being said, the purpose of this study

was to explore the history of open techniques for shoulder stabilization, describe the available open procedures, and highlight the relevant key surgical considerations based on reported clinical outcomes.

## HISTORY OF OPEN STABILIZATION PROCEDURES

It is important to describe the history of open shoulder stabilization procedures in order to reflect on current techniques and procedures available today. Prior to 1980, all shoulder instability cases were treated openly. As a matter of fact, the first anatomic repair of the labrum and the inferior glenohumeral ligament was described by Arthur Sydney Blundell Bankart in 1923, who by doing so, discovered the main lesion involved in anterior shoulder instability (Fig. 1).<sup>24)</sup> Since then, numerous open stabilization techniques evolved and developed in an attempt to achieve satisfactory patient outcomes. The Eden-Hybinette procedure, reportedly first described in 1932 as well, involved the utilization of an iliac crest autograft to augment the glenoid in shoulder instability cases (Fig. 1).<sup>25)</sup> The subscapularis transfer was described by Magnuson-Stack in 1943, followed by the subscapularis tendon and anterior capsular imbrication (Putti-Platt) described in 1948 (Fig. 1).<sup>26,27)</sup>

A major breakthrough occurred in 1954, when Michel Latarjet described an open coracoid transfer onto the anterior glenoid margin in order to help restore shoulder stability (Fig. 1).<sup>28)</sup> This was followed by the coracoid tip transfer described by Bristow-Helfet as a modification to Latarjet original procedure (Fig. 1).<sup>29)</sup> In 1980, Neer and Foster's open inferior capsular shift was described: an operation that involves tightening the shoulder capsule to achieve better joint stability before arthroscopic repair of the labrum started emerging later that year (Fig. 1).<sup>30)</sup>

History of open shoulder stabilization procedures




Fig. 1. Timeline of open procedures for shoulder stabilization procedures.

Modifications of these procedures were created and established as credible and beneficial treatment options for recurrent instability, but as times passed, open procedures decreased in frequency, and arthroscopic labral repair procedures gained notable popularity and prominence.<sup>17,18)</sup>

Indeed, arthroscopic options for labral repair witnessed prominent rise over the past decades. A study by Riff et al.<sup>17)</sup> explored the trends of shoulder stabilization techniques used in the United States and noted a 7.9% increase in the use of arthroscopic stabilization between 1994 and 2006, along with 9.1% decrease in the use of open Bankart procedures, and a 15.4% increase in the use of the Latarjet procedures. As a matter of fact, the breakdown of shoulder stabilization procedures in 2006 showed that 87% of shoulder stabilization procedures were arthroscopic Bankart repairs, 7% were open Bankart repairs, and 3.2% were Latarjet procedures.<sup>17)</sup> Considering that arthroscopic labral repairs were first described in 1980, it is impressive to see how popular its use has been in the United States.

It is certain that open procedures still play a role in shoulder stabilization, and their benefits are yet to be replaced by arthroscopic procedures. By adhering to the right indications, open stabilization procedures can provide superior clinical outcomes and achieve better satisfaction for the unstable patient.

## INDICATIONS FOR OPEN STABILIZATION PROCEDURES

Several indications support the use of open procedures for anterior shoulder stabilization, and these include critical glenoid bone loss, failed arthroscopic surgery, massive engaging Hill-Sachs lesions and concomitant shoulder pathologies, and high instability severity scores (ISSs). Open stabilization procedures for the shoulder are indicated in patients who have critical glenoid bone loss, and this can occur as a result of an osseous Bankart lesion, recurrent dislocations, or innate joint hypermobility and laxity.<sup>31)</sup> Glenoid bone loss is a risk factor that can lead to recurrent instability, and in these cases, using open stabilization techniques instead of an arthroscopic repair can result in better long-term outcomes and lower rates of recurrence.<sup>32,33)</sup> The definition of critical bone loss, which is considered highly unstable, has varied in the literature. Previous reports suggested that defects greater than 20%–25% are considered critical, but more recent ones consider defects as low as 13.5% to be critical and highly unstable.<sup>34-36)</sup> It has also been shown that with each dislocation event, 6.8% glenoid bone loss can occur.<sup>37)</sup> Accordingly,

an open Latarjet procedure can be indicated following three dislocation events necessitating formal reduction.<sup>7,38)</sup> Hence, with recurrent instability events, surgical options tend to lean more towards open procedures rather than arthroscopic.

Previously failed surgery for anterior glenohumeral instability is also an indication for an open revision stabilization procedure. Open procedures, such as the Latarjet, traditionally result in a lower recurrence rate for future anterior instability compared to other approaches, such as an arthroscopic revision.<sup>39,40)</sup> Although open stabilization procedures have low recurrence rates for future anterior instability, revision surgeries are inferior to primary stabilization surgeries and can lead to higher recurrence and lower outcome predictability.<sup>41)</sup>

Concomitant shoulder pathologies, like humeral head fractures, glenoid fractures, or humeral avulsion of anterior glenohumeral ligament, constitute an indication for open stabilization surgery. Similarly, large engaging Hill-Sachs lesions may constitute an indication for open procedures, mainly those that involve bone graft reconstruction or arthroplasty. It is important to determine the severity of Hill-Sachs lesions, as those that affect less than 20% of the humeral head may be treated non-surgically with physical therapy and careful observation. Larger lesions, however, especially those that constitute a defect greater than 40% of the humeral head, usually require open surgical management.<sup>7,8)</sup>

Finally, the ISS is a tool that predicts the patient's risk of experiencing recurrent shoulder instability after an arthroscopic Bankart repair. Multiple factors are included when calculating the ISS, such as age, the risks associated with the sport played by the patient, the presence of glenoid loss or a Hill-Sachs lesion, and hyperlaxity.<sup>42,43)</sup> Athletes involved in risky contact sports, especially younger athletes, have an increased chance of reinjury or re-dislocation.<sup>44)</sup> The ISS uses a point scoring guideline where differing amounts of points are assigned to the patient for the varying factors that are evaluated before the operation. Higher scores contraindicate an arthroscopic repair because an open procedure would provide more shoulder stability and better patient outcomes.<sup>42-44)</sup>

## OPEN STABILIZATION PROCEDURES

### Latarjet Procedure (Open Coracoid Transfer)

A Latarjet procedure, which involves augmentation of a defected and unstable glenoid using an open coracoid transfer, is indicated in the setting of patients who are at high risk for recurrent shoulder instability (primarily

anterior) (Fig. 2).<sup>45)</sup> The procedure is considered the gold standard of open stabilization procedures, and is especially popular among contact athletes under the age of 25, and patients with critical glenoid bone loss.<sup>7,45,46)</sup>


Patient outcomes following Latarjet procedures have been extensively explored in the literature. The Latarjet has shown low rates of recurrent shoulder instability and high rates of patient satisfaction in diverse patient populations.<sup>7,47)</sup> The procedure is especially popular in the sport setting due to its high return to sport rates. Yapp et al.<sup>48)</sup> and Elamo et al.<sup>49)</sup> explored the prognosis of athletes who had undergone a Latarjet procedure and found a 100% return to sport rate at 5- and 10-year postoperation. In another study, at 25-year postoperation, out of the 40 patients who underwent a Latarjet procedure, none reported redislocation.<sup>2,13)</sup> Another systematic review by Hurley et al.<sup>50)</sup> explored outcomes of Latarjet procedures in 2,134 athletes and reported an overall return to play rate of 88.8% at a mean time of 5.8 months, with 72.6% of athletes returning to their previous level of competition. The authors further stratified return to play and return to previous level of play by type of sport: rates of return to play were 88.2% and 90.3%, while rates of return to previous level of play were 69.5% and 80.6% among collision athletes and overhead athletes, respectively.<sup>50)</sup>

With regards to open stabilization procedures, the Latarjet remains the gold standard with superior outcomes when compared to other open procedures. Hovelius et al.<sup>51)</sup> performed a retrospective analysis of 185 shoulders with a mean follow-up duration of 17 years comparing open techniques: Bristow-Latarjet and open Bankart repair. Patients in the Bristow-Latarjet group had superior

shoulder function with significantly better patient-reported outcome scores than patients in the open Bankart group.<sup>51)</sup> Patients who underwent a Latarjet procedure also showed significantly lower rates of redislocation, subluxation, apprehension, and revisions than patients who underwent standard open Bankart repair.<sup>51,52)</sup> Comparisons have also been drawn between different Latarjet techniques. One study by Dumont et al.<sup>53)</sup> compared the computed tomography (CT) findings of patients treated with a traditional Latarjet to those treated with a congruent arc modified Latarjet. In a congruent arc Latarjet, the inferior surface of the coracoid is used to augment the glenoid rather than the lateral edge of the coracoid.<sup>54)</sup> In theory, this allows the reconstruction of a greater percentage of glenoid bone deficit, since the inferior portion of the coracoid is wider than its lateral edge.<sup>54)</sup> Nevertheless, Dumont et al.<sup>53)</sup> found greater bony contact with the glenoid (5.65 cm<sup>2</sup> vs. 3.64 cm<sup>2</sup>) and greater bone width on each side of the screws (7.1 mm vs. 4.1 mm) in the traditional technique compared to the modified technique.

The advantages of this procedure, implied by increased joint stability and lower recurrence rates, are challenged by its significant complication profile, which can range from hematomas to graft reunion failure.<sup>7,55,56)</sup> Regarding complications, reported rates showed a range of 7% to 30%.<sup>7,55,56)</sup> These include screw mispositioning, nerve injury, nonunion, bone overhang, and superior graft resorption.<sup>57)</sup> With regards to long-term complications, osteoarthritis appeared to be a concern as studies have shown that it can develop in 23% to 37% of cases.<sup>58,59)</sup> Domos et al.<sup>52)</sup> reported that postoperative osteoarthritis is usually present in 20% to 25% of open Latarjet procedures. In another long-term study on 822 patients who underwent a Latarjet procedure with an average follow-up of 16.6 years, 38.2% of patients reported degenerative changes, 35.7% reported residual shoulder pain, and 4.8% reported daily pain.<sup>60)</sup>

Differences in outcomes and prognostic parameters between arthroscopic stabilization procedures and Latarjet have been explored in the literature. One retrospective analysis of 364 shoulders performed by Zimmermann et al.<sup>61)</sup> compared open Latarjet procedure and arthroscopic Bankart repair and reported a significant improvement in Subjective Shoulder Value (SSV) score in both procedures.<sup>61)</sup> However, the open Latarjet repair group had a significantly greater increase in overall SSV than the arthroscopic repair group. This may be due to the fact that patients in the open Latarjet repair group scored lower in preoperative SSV and had a greater disability than patients in the arthroscopic repair group, and thus had a significantly greater subjective benefit.<sup>61)</sup> Zimmermann et



**Fig. 2.** The Latarjet procedure involves releasing and transferring the coracoid and conjoint tendon (A) and attaching it to the anterior glenoid using one or two screws to augment the anterior glenoid defect (B).

al.<sup>61)</sup> similarly recorded a significantly higher recurrence of redislocation, subluxation, apprehension, and revision surgery in the arthroscopic repair group compared to the open Latarjet repair group. In addition, the authors found that while the majority of patients were able to return to their previous sports activity, the Latarjet group had a significantly higher percentage than the arthroscopic Bankart repair group.<sup>61)</sup> With regards to arthroscopic Latarjet, outcomes have been similar to those of the open approach.<sup>62)</sup> One systematic review by Horner et al.<sup>62)</sup> examined all comparative studies exploring the two Latarjet approaches and showed that both produce similarly successful outcomes with no significant differences in complication rates and need for revision surgery. The authors<sup>62)</sup> did, however, report less early postoperative pain with arthroscopic Latarjet but increased required operative time.

### Open Bankart Repair

Indications for an open Bankart are usually similar to those of other open stabilization procedures.<sup>7,63,64)</sup> The popularity of this procedure has decreased over the recent decades due to its more invasive approach and subsequent higher rate of complications when compared to arthroscopic repair; nevertheless, it still plays a role in shoulder stabilization surgery.<sup>17,18)</sup>

Outcomes following an open Bankart repair have been explored extensively in previous literature. A systematic review by AlSomali et al.<sup>65)</sup> demonstrated that open Bankart repair is a reliable procedure that allows high-demand populations, such as athletes and people who work in heavy labor, to return to sports and work; in their study, 87% (358/410) of patients who underwent open Bankart repair returned to their previous sports and work. In another study done on 44 athletes who underwent this procedure, 88% of the athletes were able to return to their sport, with 66% reporting similar levels of performance at an average follow-up period of 14.2 months.<sup>66)</sup> Neviaser et al.<sup>67)</sup> explored 30 patients who underwent open Bankart revision stabilization and found no recurrence at 10 years of follow-up. The authors also noted that 22 out of 23 competitive athletes were able to return to sport.<sup>67)</sup> Several negative outcomes, however, have been reported with this procedure, and these often revolve around decreased range of motion in external and internal rotation, but this was typically experienced without concomitant loss of shoulder functionality.<sup>67-69)</sup>

When comparing open Bankart repair to arthroscopic repair, findings have been varied. While both procedures have been found to provide similar functional capacities to patients, differences mainly reside in post-

operative stability and range of motion.<sup>70)</sup> One study by Rhee et al.<sup>19)</sup> explored the outcomes of arthroscopic versus open Bankart repair on a group of 46 collision athletes and displayed augmented postoperative instability in those who underwent arthroscopy. A meta-analysis of 16 trials involving 827 shoulders by Chen et al.<sup>71)</sup> demonstrated that patients who underwent arthroscopic Bankart repair had better recovery of range of motion, but a significantly higher number of redislocations and revision operations than patients who underwent open Bankart repair. Hohmann et al.<sup>70)</sup> performed a meta-analysis of 22 studies separated into two groups by publication date (1995–2004 and 2005–2015) and evaluated recurrence rates between open and arthroscopic repair in the two decades. They found that there was a significant difference favoring open Bankart repair in the 1995–2004 decade, with arthroscopic Bankart repair having twice the risk of recurrence compared with open repair group (odds ratio, 1.96).<sup>70)</sup> Although no significant difference was recorded in the more recent 2005–2015 group as well as when all studies were pooled together, the recurrence rate in patients who underwent arthroscopic repair was found to only marginally decrease 2.5% between the two decades.<sup>70)</sup> These findings are disputed, however, as a study in patients older than 50 years of age found recurrence rates between the two procedures to be similar.<sup>7,72)</sup>

Open Bankart repair is a more invasive procedure than arthroscopic repair and places the patient at higher risk of complications. One of the main differences between the two procedures is that the subscapularis must be detached in open repair, which raises concerns about a potential compromise in function. However, several studies have found no significant differences between the two procedures with regards to subscapularis function. Hiemstra et al.<sup>73)</sup> performed a randomized control trial to assess strength deficits between open and arthroscopic repair and demonstrated that while both procedures resulted in deficits in internal and external rotation compared to the contralateral shoulder, there were no significant differences between the two. Degenerative changes to the glenohumeral joint is another concern regarding open repair.<sup>65)</sup> AlSomali et al.<sup>65)</sup> conducted a systematic review to explore patient outcomes following open Bankart repair and indicated osteoarthritic changes in 33% of patients (89/268), with mild to moderate changes in 81 shoulders and severe changes in 8 using the Samilson-Prieto classification at a mean follow-up of 11.5 years. On the other hand, Chen et al.<sup>71)</sup> conducted a systematic review of the clinical trials involving both open and arthroscopic Bankart repairs to assess their safety profile and reported no significant

differences between the two approaches with regards to complications, concluding that both procedures have the same level of safety. No statistical differences were found between the two surgical groups when comparing the number of patients with dysesthesia, persistent pain after surgery, wound infections, and delayed wound healing.<sup>71)</sup>

### Bone Augmentation Using Iliac Crest Graft


Bone augmentation procedures other than the Latarjet exist can be used in the setting of severe bone loss.<sup>74)</sup> Utilizing an iliac crest autograft can augment the glenoid and provide good outcomes in cases of significant anterior or posterior shoulder instability.<sup>74)</sup> An example of such cases is an epileptic patient with a history of seizures and numerous dislocations. In these cases, a Latarjet may not be suitable due to high risk of failure and complications.<sup>74)</sup> As such, glenoid augmentation using iliac crest autograft can be a suitable stabilization option that can provide more favorable outcomes.<sup>75)</sup>

Outcomes of iliac crest bone grafting for glenoid reconstruction have been generally favorable in the literature. One prospective study by Moroder et al.<sup>76)</sup> compared the outcomes between Latarjet procedure or open iliac crest bone graft transfer in 60 randomized anterior instability patients. Patients of both groups showed improvements and there were no clinical differences in patient-reported outcomes at any of the follow-up points (6, 12, and 24 months after procedure).<sup>76)</sup> A systematic review by Malahias et al.<sup>77)</sup> reported on 261 patients who underwent iliac crest bone block techniques for shoulder stabilization and reported an overall cause of reoperation rate of 6.1%, a rate of recurrent instability of 4.8%, a graft nonunion rate of 2.2%, and rates of graft fracture, infection, and osteolysis of 0.9%, 1.7%, and 0.4%, respectively, at a mean follow-up ranging between 20.6 and 42 months. The authors<sup>77)</sup> also reported a hardware-related complication rate of 3.9%. Another study by Ernstbrunner et al.<sup>78)</sup> explored 20 cases of iliac crest bone grafting conducted following failed Latarjet procedure. While 7 of the 20 patients reported recurrence of instability following the revision procedure, the remaining 13 reported significant improvements in Constant score (25 points increase), relative Constant score (25% increase), and SSV (54% increase).<sup>78)</sup>

Different techniques exist when conducting an iliac crest autograft bone blocking procedure. One interesting technique is the J-bone graft technique, which uses a bicortical iliac crest autologous bone graft that is sculpted into a J-shaped structure and inserted via a press-fit technique in order to reconstruct the glenoid (Fig. 3).<sup>74,79)</sup> The J-bone graft technique restores the anatomic integrity of

the defected glenoid and provides good strength, mobility, and functional capacity to the surgical patient.<sup>74,80)</sup> The procedure has been shown to produce visual analog scale scores that are comparable to the Latarjet procedure, as well as a similarly low rate of recurrent dislocations at around 3%.<sup>74,80,81)</sup> The J-bone technique may entail several advantages over the Latarjet procedure. Since it is inserted via a press-fit technique and does not involve insertion of metal hardware, the J-bone technique might prevent implant-related issues such as screw breakage or loosening, which frequently necessitate revision surgery.<sup>81)</sup> In addition, the J-bone grafts are often covered with soft-tissue that can potentially differentiate into hyaline and fibrous cartilage, and this may protect against dislocation arthropathy.<sup>82)</sup> Moreover, in contrast to the Latarjet procedure, extended immobilization is not necessary because stability is already present before the remodeling process begins.<sup>74,83)</sup>

Several complications can arise when attempting glenoid reconstruction using iliac bone graft and these include subluxations, nonunion, postoperative osteolysis, neurovascular injuries, and arthritis.<sup>74,81)</sup> One study showed that the rates of moderate to severe arthritis seen with this technique were comparable to the Latarjet at around 12%.<sup>74)</sup> Another major disadvantage is the requirement for a second iliac crest incision for the purpose of harvesting the bone block. This could increase the risk of problems like infection, poor wound healing, or hemorrhage at the donor site as well as the chance of injuring the lateral femoral cutaneous nerve, which could result in short-term or perhaps long-term pain.<sup>84,85)</sup> The high learning curve of the procedure when compared to the Latarjet technique, constitutes another limitation.<sup>82)</sup>



**Fig. 3.** Schematic representation of the J-bone graft procedure for shoulder stabilization.

An arthroscopic approach to this procedure has been described in the literature as well. Anderl et al.<sup>79)</sup> explored the outcomes of 14 patients treated with arthroscopic bone block using iliac crest autograft and reported adequate restoration of the glenoid anatomy, excellent clinical outcomes, and no signs of recurrent instability on follow-up. An interesting finding was that in the arthroscopic approach, no postoperative osteoarthritic progression was found, which may be explained by the maintenance of natural biomechanics.<sup>79)</sup> Nevertheless, a longer follow-up period is required to confirm this conclusion.<sup>79)</sup> When compared to the open technique, advantages of the arthroscopic technique include better cosmetic results, a reduced risk of infection, fewer scars, quicker recovery, the ability to effectively treat concurrent injuries, easier situation in case of revision surgery, and its muscle-sparing properties.<sup>79)</sup> Thus, the risk of decreased shoulder function can be reduced by limiting damage to the shoulder joint's musculotendinous unit, preserving its structural and biomechanical integrity, and further bolstering the anatomical approach of the described treatment.<sup>86)</sup>

### **Bone Augmentation Using Distal Tibial Allograft**

The use of fresh distal tibia allograft (DTA) for glenoid reconstruction has also been described in the literature.<sup>87-89)</sup> In this procedure, a fresh distal tibial allograft is prepared and fashioned to appropriately fit and augment the defect in the patient's glenoid. This specific type of implant was first used due to studies that showed, even among non-laterality-matched cadaveric specimens, a fairly similar radius of curvature between the lateral aspect of the distal tibia and that of the glenoid exists.<sup>88,90)</sup> It was shown that throughout the whole range of motion, the lateral third of the distal tibia's articular surface is congruent with the humeral head.<sup>88-91)</sup> A biomechanical analysis by Bhatia et al.<sup>92)</sup> compared this procedure with the Latarjet and showed a higher contact area and lower peak forces with the DTA graft when the shoulder was tested in 60° of abduction and in abducted external rotation, making this procedure viable for glenoid reconstruction.

Clinically, this technique showed excellent outcomes.<sup>89)</sup> A study by Provencher et al.<sup>89)</sup> explored 27 male patients who underwent glenoid reconstruction using DTA, with a minimum of 15% anterior glenoid bone loss prior to procedure. The authors noted significant improvements in patient-reported outcome scores and no cases of recurrent instability at an average follow-up of 45 months.<sup>89)</sup> The authors also reported an overall graft healing rate of 89% and a mean allograft lysis rate of 3% on CT scans performed at a median time of 1.4.<sup>89)</sup> Another study

by Robinson et al.<sup>93)</sup> retrospectively reviewed 12 patients, with an average glenoid bone loss of 33%, who underwent DTA glenoid reconstruction. The authors reported significant improvement in clinical patient-reported outcomes and range of motion at an average follow-up of 28 months.<sup>93)</sup>

Arthroscopic DTA glenoid reconstruction has also been described in the literature and has been gaining popularity worldwide. Wong et al.<sup>94)</sup> reported on 73 patients treated with arthroscopic glenoid reconstruction using DTA and reported significantly improved patient-reported outcome scores, no recurrent instability, and union rates around 100% at a mean follow-up of 4.7 years. Amar et al.<sup>95)</sup> explored this procedure and reported an excellent safety profile, no intraoperative issues such as neurovascular injury, adverse events, hemorrhage, or infections, and 100% graft healing on CT scan at a mean radiographic follow-up of 6.31 months in 31 out of 42 patients. As such, both open and arthroscopic DTA glenoid reconstructions are utilized in shoulder surgery with appropriate outcomes for both. Comparative studies exploring the two approaches should be conducted in order to assess which procedure has better therapeutic efficacy. In addition, studies with longer follow-up periods should be done to appropriately assess its efficacy and safety profile.

### **Salvage Procedures**

Patients with significant instability frequently pose diagnostic challenges and may continue to be resistant to surgical intervention. A tiny subgroup may experience multiple surgical failures, resulting in severe impairment from ongoing pain and instability. In these situations, salvage options, like glenohumeral arthrodesis or arthroplasty, may be utilized.

Stark et al.<sup>96)</sup> and Cofield et al.<sup>97)</sup> both reported improvements in pain, stability, and range of motion after arthrodesis for instability. Richards et al.,<sup>98)</sup> however, reported contradicting results, noting persistent postoperative pain and instability at a mean follow-up of 43 months in 4 out of 6 patients who underwent the procedure for multidirectional instability. Diaz et al.<sup>99)</sup> explored the outcomes of glenohumeral arthrodesis in 8 patients, with an average of 7 prior stabilization attempts at a mean follow up of 35 months. Despite some functional limitations and residual pain, the authors reported no postoperative instability complaints and good patient satisfaction postoperatively, with all patients stating that they would repeat the surgery under the same preoperative condition.<sup>99)</sup>

Shoulder arthroplasty procedures can also be considered salvage procedures for severe shoulder instabil-

ity.<sup>100,101</sup> This procedure can improve range of motion and reduce pain.<sup>101,102</sup> However, complications such as posterior instability and component loosening requiring revision may occur.<sup>101</sup> Having a shorter time between the injury and the arthroplasty can improve the postoperative outcomes.<sup>101</sup> Yet, a history of a previous surgery (which is often the case), the necessity of an anteromedial approach or the concurrent presence of rotator cuff repair can compromise the outcomes.<sup>101</sup>

As stated earlier, a critically large Hill-Sachs lesion may entail prominent instability and require open surgical management. In this setting, allograft bone reconstruction and humeral resurfacing are two of the main open surgical procedures available. Allograft humeral head reconstruction has been widely explored in the literature, with different reported techniques and good resultant outcomes. One study by Miniaci and Gish<sup>103</sup> reported on 18 cases of instability patients treated with humeral head structural osteoarticular allografts. Patients had failed previous stabilization procedures and had humeral head defects of at least 25%.<sup>103</sup> At a mean follow-up of 50 months, 89% of patients returned to work, average Constant score was 78.5, and no incidents of recurrent instability were reported.<sup>103</sup> With larger Hill-Sachs lesions, typically greater than 40%–45%, hemiarthroplasty can be used to treat instability symptoms.<sup>104,105</sup> Nevertheless, the use of this procedure in this setting has several limitations and the published available literature suggests mixed results.<sup>104,105</sup> While some studies report generally favorable outcomes, especially in older patients, others highlight the risk of prosthetic loosening and the need of additional retroversion to achieve proper stability.<sup>104,105</sup>

## RECOMMENDATIONS

Open stabilization procedures for the shoulder remain valid options for many presenting patients. It is important, however, to recognize when these procedures are more favorable over arthroscopic procedures. All patients are different, and hence, a holistic assessment of the history, presentation, and expected outcomes of the patients is always necessary before deciding on a management plan. In specific, factors like age, bone loss, type of athletic participation, coracoid size, and degree of laxity should factor in when choosing the appropriate surgical procedure. Generally, arthroscopic procedures are praised for their minimally invasive approach and retention of range of motion at the expense of higher recurrence rates and risk of failures. Open stabilization procedures, on the other hand, offer increased stability and lower recurrence rates

at the expense of a possible nonanatomic restoration of glenohumeral stability and a consequently higher complication profile. Hence, patient selection and education is key. While arthroscopic procedures are popular and offer significant benefits for the average patient, patients who are involved in contact sports and those with high risk of recurrence, severe glenoid bone loss, or concomitant glenohumeral pathologies may be more suited for open stabilization procedures.

Research and innovation targeting both open and arthroscopic stabilization procedures have produced several new surgical procedures with much promise. Quite recently, the addition of soft-tissue repair to Bankart procedures, such as the case of remplissage, has been garnering a lot of attention.<sup>106-111</sup> Some studies in the literature show superior results and very low rates of revision or recurrence at long-term follow-up when compared to other procedures, while other studies challenge any superior benefit from this technique.<sup>106-111</sup> With continuous medical and surgical advancements and innovations, it is possible that the addition of soft-tissue repair to arthroscopic stabilization procedures may replace the need for some of the more robust and invasive open procedures. That being said, conducting additional research that compares different stabilization procedures (open and arthroscopic) is necessary for the establishment of proper treatment guidelines and strategies. In addition, exploring new graft options for bone-block procedures can be beneficial in light of the cost and supply restraints of available cadaveric options.

## CONCLUSION

Shoulder instability is a very common pathology that affects many patients worldwide. While arthroscopic management of this pathology has been the standard treatment for the average presenting patient, many cases involve much more complex presentations that necessitate the use of an open approach. Open stabilization procedures are usually indicated for patients with recurrent instability, severe glenoid bone loss, concomitant shoulder pathologies, and high risk for failure with arthroscopic management. Some of these procedures include the Latarjet procedure, open Bankart repair, bone-blocking procedures using iliac crest autograft or DTA, and salvage procedures like shoulder replacement or glenohumeral arthrodesis. Each of these procedures entail different advantages and disadvantages, and accordingly, it is of pivotal importance to cater management to the individual patient. Additional research should be conducted to determine the best treat-



ment approach for different patient presentations, report on long-term outcomes of some of the newer procedures, and explore additional graft options for bone-blocking procedures.

### CONFLICT OF INTEREST

JAA would like to disclose royalties from DJO Global, Zimmer-Biomet, Smith and Nephew, Stryker, Globus Medical, INC.; research support as a PI from Lima Corporation - Italy, Orthofix, Arthrex, OREF; royalties, financial or material support from Wolters Kluwer; and Board Member/Committee appointments for American Shoulder and Elbow Society, Pacira.

### ACKNOWLEDGEMENTS

BioRender.com was used to create some of the figures in this manuscript.

### ORCID

Mohamad Y Fares <https://orcid.org/0000-0001-8228-3953>  
 Peter Boufadel <https://orcid.org/0000-0001-7078-9148>  
 Mohammad Daher <https://orcid.org/0000-0002-9256-9952>  
 Jonathan Koa <https://orcid.org/0000-0002-3922-359X>  
 Joseph A. Abboud <https://orcid.org/0000-0002-3845-7220>

### REFERENCES

- Chang LR, Anand P, Varacallo M. Anatomy, shoulder and upper limb, glenohumeral joint [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 [cited 2023 Apr 1]. Available from: <https://pubmed.ncbi.nlm.nih.gov/30725703/>.
- Wilk KE, Arrigo CA, Andrews JR. Current concepts: the stabilizing structures of the glenohumeral joint. *J Orthop Sports Phys Ther.* 1997;25(6):364-79.
- Halder AM, Halder CG, Zhao KD, O'Driscoll SW, Morrey BF, An KN. Dynamic inferior stabilizers of the shoulder joint. *Clin Biomech (Bristol, Avon).* 2001;16(2):138-43.
- Halder AM, Zhao KD, Odriscoll SW, Morrey BF, An KN. Dynamic contributions to superior shoulder stability. *J Orthop Res.* 2001;19(2):206-12.
- Balvanyossy P. Static stabilizers of the shoulder joint. *Unfallchirurg.* 1990;93(1):27-31.
- Beltran J, Jbara M, Maimon R. Shoulder: labrum and bicipital tendon. *Top Magn Reson Imaging.* 2003;14(1):35-49.
- Haratian A, Yensen K, Bell JA, et al. Open stabilization procedures of the shoulder in the athlete: indications, techniques, and outcomes. *Open Access J Sports Med.* 2021;12:159-69.
- Provencher MT, Frank RM, Leclere LE, et al. The Hill-Sachs lesion: diagnosis, classification, and management. *J Am Acad Orthop Surg.* 2012;20(4):242-52.
- Provencher MT, Midtgaard KS, Owens BD, Tokish JM. Diagnosis and management of traumatic anterior shoulder instability. *J Am Acad Orthop Surg.* 2021;29(2):e51-61.
- Millett PJ, Clavert P, Hatch GF 3rd, Warner JJ. Recurrent posterior shoulder instability. *J Am Acad Orthop Surg.* 2006;14(8):464-76.
- Gaskill TR, Taylor DC, Millett PJ. Management of multidirectional instability of the shoulder. *J Am Acad Orthop Surg.* 2011;19(12):758-67.
- Owens BD, Campbell SE, Cameron KL. Risk factors for anterior glenohumeral instability. *Am J Sports Med.* 2014;42(11):2591-6.
- Waterman B, Owens BD, Tokish JM. Anterior shoulder instability in the military athlete. *Sports Health.* 2016;8(6):514-9.
- Owens BD, Duffey ML, Nelson BJ, DeBerardino TM, Taylor DC, Mountcastle SB. The incidence and characteristics of shoulder instability at the United States Military Academy. *Am J Sports Med.* 2007;35(7):1168-73.
- DeFroda SF, Goyal D, Patel N, Gupta N, Mulcahey MK. Shoulder instability in the overhead athlete. *Curr Sports Med Rep.* 2018;17(9):308-14.
- Castricini R, Castioni D, De Benedetto M, et al. Arthroscopic Latarjet for primary shoulder instability with off-track lesions or revision surgery yields satisfactory clinical results and reliable return to sport and work at minimum 3-year follow-up. *Arthroscopy.* 2022;38(10):2809-18.
- Riff AJ, Frank RM, Sumner S, et al. Trends in shoulder stabilization techniques used in the United States based on a large private-payer database. *Orthop J Sports Med.* 2017;5(12):2325967117745511.
- Frank RM, Chalmers PN, Moric M, Leroux T, Provencher MT, Romeo AA. Incidence and changing trends of shoulder stabilization in the United States. *Arthroscopy.* 2018;34(3):784-92.
- Rhee YG, Ha JH, Cho NS. Anterior shoulder stabilization in collision athletes: arthroscopic versus open Bankart repair.

- Am J Sports Med. 2006;34(6):979-85.
20. Tokish JM, Lafosse L, Giacomo GD, Arciero R. Patients in whom arthroscopic Bankart repair is not enough: evaluation and management of complex anterior glenohumeral instability. *Instr Course Lect.* 2017;66:79-89.
  21. Carlson Strother CR, McLaughlin RJ, Krych AJ, Sanchez-Sotelo J, Camp CL. Open shoulder stabilization for instability: anterior labral repair with capsular shift. *Arthrosc Tech.* 2019;8(7):e749-54.
  22. Chan AG, Kilcoyne KG, Chan S, Dickens JF, Waterman BR. Evaluation of the Instability Severity Index score in predicting failure following arthroscopic Bankart surgery in an active military population. *J Shoulder Elbow Surg.* 2019;28(5):e156-63.
  23. Bliven KC, Parr GP. Outcomes of the Latarjet procedure compared with Bankart repair for recurrent traumatic anterior shoulder instability. *J Athl Train.* 2018;53(2):181-83.
  24. Bankart AS. Recurrent or habitual dislocation of the shoulder-joint. *Br Med J.* 1923;2(3285):1132-3.
  25. Hybbinette SR. De la transplantation d'un fragment osseux pour remédier aux luxations récidivantes de l'épaule: constatations et résultats opératoires. *Acta Chir Scand.* 1932;71(411-445):26.
  26. Magnuson PB, Stack JK. Recurrent dislocation of the shoulder. 1943. *Clin Orthop Relat Res.* 1991;(269):4-8.
  27. Osmond-Clarke H. Habitual dislocation of the shoulder: the Putti-Platt operation. *J Bone Joint Surg Br.* 1948;30(1):19-25.
  28. Latarjet M. Treatment of recurrent dislocation of the shoulder. *Lyon Chir.* 1954;49(8):994-7.
  29. Helfet AJ. Coracoid transplantation for recurring dislocation of the shoulder. *J Bone Joint Surg Br.* 1958;40(2):198-202.
  30. Neer CS 2nd, Foster CR. Inferior capsular shift for involuntary inferior and multidirectional instability of the shoulder. A preliminary report. *J Bone Joint Surg Am.* 1980;62(6):897-908.
  31. Skupinski J, Piechota MZ, Wawrzynek W, Maczuch J, Babiniska A. The bony Bankart lesion: how to measure the glenoid bone loss. *Pol J Radiol.* 2017;82:58-63.
  32. Hurley ET, Davey MS, Montgomery C, et al. Arthroscopic Bankart repair versus open Latarjet for recurrent shoulder instability in athletes. *Orthop J Sports Med.* 2021;9:23259671211023801.
  33. Wolf JM, Cameron KL, Owens BD. Impact of joint laxity and hypermobility on the musculoskeletal system. *J Am Acad Orthop Surg.* 2011;19(8):463-71.
  34. Streubel PN, Krych AJ, Simone JP, et al. Anterior glenohumeral instability: a pathology-based surgical treatment strategy. *J Am Acad Orthop Surg.* 2014;22(5):283-94.
  35. Di Giacomo G, Itoi E, Burkhart SS. Evolving concept of bipolar bone loss and the Hill-Sachs lesion: from "engaging/non-engaging" lesion to "on-track/off-track" lesion. *Arthroscopy.* 2014;30(1):90-8.
  36. James M, Kwong CA, More KD, LeBlanc J, Lo IK, Bois AJ. Bony apprehension test for identifying bone loss in patients with traumatic anterior shoulder instability: a validation study. *Am J Sports Med.* 2022;50(6):1520-8.
  37. Bois AJ, Mayer MJ, Fening SD, Jones MH, Miniaci A. Management of bone loss in recurrent traumatic anterior shoulder instability: a survey of North American surgeons. *JSES Int.* 2020;4(3):574-83.
  38. Fabricant PD, Taylor SA, McCarthy MM, et al. Open and arthroscopic anterior shoulder stabilization. *JBJS Rev.* 2015;3(2):e4.
  39. Jeon YS, Jeong HY, Lee DK, Rhee YG. Borderline glenoid bone defect in anterior shoulder instability: Latarjet procedure versus Bankart repair. *Am J Sports Med.* 2018;46(9):2170-6.
  40. Rollick NC, Ono Y, Kurji HM, et al. Long-term outcomes of the Bankart and Latarjet repairs: a systematic review. *Open Access J Sports Med.* 2017;8:97-105.
  41. Levine WN, Arroyo JS, Pollock RG, Flatow EL, Bigliani LU. Open revision stabilization surgery for recurrent anterior glenohumeral instability. *Am J Sports Med.* 2000;28(2):156-60.
  42. Rosenberg SI, Padanilam SJ, Pagni BA, Tjong VK, Sheth U. A lower Instability Severity Index score threshold may better predict recurrent anterior shoulder instability after arthroscopic Bankart repair: a systematic review. *J ISAKOS.* 2021;6(5):295-301.
  43. Hurley ET, Davey MS, Montgomery C, et al. Arthroscopic Bankart repair versus open Latarjet for first-time dislocators in athletes. *Orthop J Sports Med.* 2021;9(8):23259671211023803.
  44. Monk AP, Crua E, Gatenby GC, et al. Clinical outcomes following open anterior shoulder stabilization for glenohumeral instability in the young collision athlete. *J Shoulder Elbow Surg.* 2022;31(7):1474-8.
  45. DeFroda SF, Perry AK, Bodendorfer BM, Verma NN. Evolving concepts in the management of shoulder instability. *Indian J Orthop.* 2021;55(2):285-98.
  46. Boe B, Stoen RO, Blich I, Moatshe G, Ludvigsen TC. Learning curve for arthroscopic shoulder Latarjet procedure shows shorter operating time and fewer complications with experience. *Arthroscopy.* 2022;38(8):2391-8.

47. Kowalski TJ, Khan AZ, Cohen JR, et al. Open shoulder stabilization: current trends and 1-year postoperative complications. *JSES Open Access*. 2017;1(2):72-8.
48. Yapp LZ, Nicholson JA, McCallum C, Macdonald DJ, Robinson CM. Latarjet as a primary and revision procedure for anterior shoulder instability: a comparative study of survivorship, complications and functional outcomes in the medium to long-term. *Shoulder Elbow*. 2020;12(5):338-48.
49. Elamo S, Selanne L, Lehtimäki K, et al. Bankart versus Latarjet operation as a revision procedure after a failed arthroscopic Bankart repair. *JSES Int*. 2020;4(2):292-6.
50. Hurley ET, Montgomery C, Jamal MS, et al. Return to play after the Latarjet procedure for anterior shoulder instability: a systematic review. *Am J Sports Med*. 2019;47(12):3002-8.
51. Hovelius L, Vikersjö O, Olofsson A, Svensson O, Rahme H. Bristow-Latarjet and Bankart: a comparative study of shoulder stabilization in 185 shoulders during a seventeen-year follow-up. *J Shoulder Elbow Surg*. 2011;20(7):1095-101.
52. Doms P, Lunini E, Walch G. Contraindications and complications of the Latarjet procedure. *Shoulder Elbow*. 2018;10(1):15-24.
53. Dumont GD, Vopat BG, Parada S, et al. Traditional versus congruent arc Latarjet technique: effect on surface area for union and bone width surrounding screws. *Arthroscopy*. 2017;33(5):946-52.
54. Rossi LA, Tanoira I, Gorodischer T, Pasqualini I, Muscolo DL, Ranalletta M. Are the classic and the congruent arc Latarjet procedures equally effective for the treatment of recurrent shoulder instability in athletes? *Am J Sports Med*. 2020;48(9):2081-9.
55. Scanlon JP, Hurley ET, Davey MS, et al. 90-Day complication rate after the Latarjet procedure in a high-volume center. *Am J Sports Med*. 2020;48(14):3467-71.
56. Willemot L, De Boey S, Van Tongel A, Declercq G, De Wilde L, Verborgt O. Analysis of failures after the Bristow-Latarjet procedure for recurrent shoulder instability. *Int Orthop*. 2019;43(8):1899-907.
57. Alkaduhimi H, Willigenburg NW, Wessel RN, et al. Ninety-day complication rate based on 532 Latarjet procedures in Dutch hospitals with different operation volumes. *J Shoulder Elbow Surg*. 2023;32(6):1207-13.
58. Merrill CA, Arciero R. Open Bankart repair: a reproducible technique. *Oper Tech Sports Med*. 2019;27(1):42-8.
59. Coughlin RP, Crapser A, Coughlin K, Coughlin LP. Open Bankart revisited. *Arthrosc Tech*. 2017;6(1):e233-7.
60. Hurley ET, Jamal MS, Ali ZS, Montgomery C, Pauzenberger L, Mullett H. Long-term outcomes of the Latarjet procedure for anterior shoulder instability: a systematic review of studies at 10-year follow-up. *J Shoulder Elbow Surg*. 2019;28(2):e33-9.
61. Zimmermann SM, Scheyerer MJ, Farshad M, Catanzaro S, Rahm S, Gerber C. Long-term restoration of anterior shoulder stability: a retrospective analysis of arthroscopic Bankart repair versus open Latarjet procedure. *J Bone Joint Surg Am*. 2016;98(23):1954-61.
62. Horner NS, Moroz PA, Bhullar R, et al. Open versus arthroscopic Latarjet procedures for the treatment of shoulder instability: a systematic review of comparative studies. *BMC Musculoskelet Disord*. 2018;19(1):255.
63. Eberbach H, Jaeger M, Bode L, et al. Arthroscopic Bankart repair with an individualized capsular shift restores physiological capsular volume in patients with anterior shoulder instability. *Knee Surg Sports Traumatol Arthrosc*. 2021;29(1):230-9.
64. Prada C, Al-Mohrej OA, Patel A, Flood B, Leroux T, Khan M. Managing bone loss in shoulder instability-techniques and outcomes: a scoping review. *Curr Rev Musculoskelet Med*. 2021;14(6):447-61.
65. AlSomali K, Kholinne E, Van Nguyen T, et al. Outcomes and return to sport and work after open Bankart repair for recurrent shoulder instability: a systematic review. *Orthop J Sports Med*. 2021;9(10):232596712111026907.
66. Pavlik A, Csepai D, Hidas P, Banoczy A. Sports ability after Bankart procedure in professional athletes. *Knee Surg Sports Traumatol Arthrosc*. 1996;4(2):116-20.
67. Neviasser AS, Benke MT, Neviasser RJ. Open Bankart repair for revision of failed prior stabilization: outcome analysis at a mean of more than 10 years. *J Shoulder Elbow Surg*. 2015;24(6):897-901.
68. Hatch MD, Hennrikus WL. The open Bankart repair for traumatic anterior shoulder instability in teenage athletes. *J Pediatr Orthop*. 2018;38(1):27-31.
69. Steinbeck J, Jerosch J. Open Bankart repair using suture anchors in posttraumatic shoulder instability: 2 to 5-year results. *Unfallchirurg*. 1997;100(12):938-42.
70. Hohmann E, Tetsworth K, Glatt V. Open versus arthroscopic surgical treatment for anterior shoulder dislocation: a comparative systematic review and meta-analysis over the past 20 years. *J Shoulder Elbow Surg*. 2017;26(10):1873-80.
71. Chen L, Xu Z, Peng J, Xing F, Wang H, Xiang Z. Effectiveness and safety of arthroscopic versus open Bankart repair for recurrent anterior shoulder dislocation: a meta-analysis of clinical trial data. *Arch Orthop Trauma Surg*. 2015;135(4):529-38.
72. Sperling JW, Duncan SF, Torchia ME, O'Driscoll SW, Cofield RH. Bankart repair in patients aged fifty years or

- greater: results of arthroscopic and open repairs. *J Shoulder Elbow Surg.* 2005;14(2):111-3.
73. Hiemstra LA, Sasyniuk TM, Mohtadi NG, Fick GH. Shoulder strength after open versus arthroscopic stabilization. *Am J Sports Med.* 2008;36(5):861-7.
  74. Boesmueller S, Berchtold M, Lorenz G, et al. Implant-free iliac crest bone graft procedure shows anatomic remodelling without redislocation in recurrent anterior shoulder instability after short-term follow-up. *Arch Orthop Trauma Surg.* 2022;142(6):1047-54.
  75. Falbo R, Moore A, Singleton A, Steffenson A, Levine J, Miller R. Glenoid bone augmentation: a contemporary and comprehensive systematic review of open procedures. *Orthop Rev (Pavia).* 2022;14(3):37834.
  76. Moroder P, Schulz E, Wierer G, et al. Neer Award 2019: Latarjet procedure vs. iliac crest bone graft transfer for treatment of anterior shoulder instability with glenoid bone loss. A prospective randomized trial. *J Shoulder Elbow Surg.* 2019;28(7):1298-307.
  77. Malahias MA, Chytas D, Raoulis V, Chronopoulos E, Brilakis E, Antonogiannakis E. Iliac crest bone grafting for the management of anterior shoulder instability in patients with glenoid bone loss: a systematic review of contemporary literature. *Sports Med Open.* 2020;6(1):12.
  78. Ernstbrunner L, Pastor T, Waltenspul M, Gerber C, Wieser K. Salvage iliac crest bone grafting for a failed Latarjet procedure: analysis of failed and successful procedures. *Am J Sports Med.* 2021;49(13):3620-7.
  79. Anderl W, Pauzenberger L, Laky B, Kriegleder B, Heuberger PR. Arthroscopic implant-free bone grafting for shoulder instability with glenoid bone loss: clinical and radiological outcome at a minimum 2-year follow-up. *Am J Sports Med.* 2016;44(5):1137-45.
  80. Moroder P, Plachel F, Becker J, et al. Clinical and radiological long-term results after implant-free, autologous, iliac crest bone graft procedure for the treatment of anterior shoulder instability. *Am J Sports Med.* 2018;46(12):2975-80.
  81. Griesser MJ, Harris JD, McCoy BW, et al. Complications and re-operations after Bristow-Latarjet shoulder stabilization: a systematic review. *J Shoulder Elbow Surg.* 2013;22(2):286-92.
  82. Auffarth A, Resch H, Matis N, et al. Cartilage morphological and histological findings after reconstruction of the glenoid with an iliac crest bone graft. *Am J Sports Med.* 2018;46(5):1039-45.
  83. Pauzenberger L, Dyrna F, Obopilwe E, et al. Biomechanical evaluation of glenoid reconstruction with an implant-free j-bone graft for anterior glenoid bone loss. *Am J Sports Med.* 2017;45(12):2849-57.
  84. Rausch V, Konigshausen M, Gebmann J, Schildhauer TA, Seybold D. Bony Bankart lesions and glenoid defects: from refixation techniques to bony augmentation. *Unfallchirurg.* 2018;121(2):117-25.
  85. Auffarth A, Kralinger F, Resch H. Anatomical glenoid reconstruction via a J-bone graft for recurrent posttraumatic anterior shoulder dislocation. *Oper Orthop Traumatol.* 2011;23(5):453-61.
  86. Liem D, Kleeschulte K, Dedy N, Schulte TL, Steinbeck J, Marquardt B. Subscapularis function after transosseous repair in shoulder arthroplasty: transosseous subscapularis repair in shoulder arthroplasty. *J Shoulder Elbow Surg.* 2012; 21(10):1322-7.
  87. Frank RM, Romeo AA, Provencher MT. Glenoid reconstruction with distal tibia allograft for recurrent anterior shoulder instability. *Orthopedics.* 2017;40(1):e199-205.
  88. Provencher MT, Ghodadra N, LeClere L, Solomon DJ, Romeo AA. Anatomic osteochondral glenoid reconstruction for recurrent glenohumeral instability with glenoid deficiency using a distal tibia allograft. *Arthroscopy.* 2009;25(4):446-52.
  89. Provencher MT, Frank RM, Golijanin P, et al. Distal tibia allograft glenoid reconstruction in recurrent anterior shoulder instability: clinical and radiographic outcomes. *Arthroscopy.* 2017;33(5):891-7.
  90. Provencher MT, LeClere LE, Ghodadra N, Solomon DJ. Postsurgical glenohumeral anchor arthropathy treated with a fresh distal tibia allograft to the glenoid and a fresh allograft to the humeral head. *J Shoulder Elbow Surg.* 2010; 19(6):e6-11.
  91. Frank RM, Romeo AA, Richardson C, et al. Outcomes of Latarjet versus distal tibia allograft for anterior shoulder instability repair: a matched cohort analysis. *Am J Sports Med.* 2018;46(5):1030-8.
  92. Bhatia S, Van Thiel GS, Gupta D, et al. Comparison of glenohumeral contact pressures and contact areas after glenoid reconstruction with latarjet or distal tibial osteochondral allografts. *Am J Sports Med.* 2013;41(8):1900-8.
  93. Robinson SP, Patel V, Rangarajan R, Lee BK, Blout C, Itamura JM. Distal tibia allograft glenoid reconstruction for shoulder instability: outcomes after lesser tuberosity osteotomy. *JSES Int.* 2020;5(1):60-5.
  94. Wong I, John R, Ma J, Coady CM. Arthroscopic anatomic glenoid reconstruction using distal tibial allograft for recurrent anterior shoulder instability: clinical and radiographic outcomes. *Am J Sports Med.* 2020;48(13):3316-21.
  95. Amar E, Konstantinidis G, Coady C, Wong IH. Arthroscopic treatment of shoulder instability with glenoid bone loss using distal tibial allograft augmentation: safety profile and

- short-term radiological outcomes. *Orthop J Sports Med.* 2018;6(5):2325967118774507.
96. Stark DM, Bennett JB, Tullos HS. Rigid internal fixation for shoulder arthrodesis. *Orthopedics.* 1991;14(8):849-55.
97. Cofield RH, Briggs BT. Glenohumeral arthrodesis: operative and long-term functional results. *J Bone Joint Surg Am.* 1979;61(5):668-77.
98. Richards RR, Beaton D, Hudson AR. Shoulder arthrodesis with plate fixation: functional outcome analysis. *J Shoulder Elbow Surg.* 1993;2(5):225-39.
99. Diaz JA, Cohen SB, Warren RF, Craig EV, Allen AA. Arthrodesis as a salvage procedure for recurrent instability of the shoulder. *J Shoulder Elbow Surg.* 2003;12(3):237-41.
100. Matsoukis J, Tabib W, Guiffault P, et al. Primary unconstrained shoulder arthroplasty in patients with a fixed anterior glenohumeral dislocation. *J Bone Joint Surg Am.* 2006;88(3):547-52.
101. Wooten C, Klika B, Schleck CD, Harmsen WS, Sperling JW, Cofield RH. Anatomic shoulder arthroplasty as treatment for locked posterior dislocation of the shoulder. *J Bone Joint Surg Am.* 2014;96(3):e19.
102. Gavriilidis I, Magosch P, Lichtenberg S, Habermeyer P, Kircher J. Chronic locked posterior shoulder dislocation with severe head involvement. *Int Orthop.* 2010;34(1):79-84.
103. Miniaci A, Gish MW. Management of anterior glenohumeral instability associated with large Hill–Sachs defects. *Tech Shoulder Elb Surg.* 2004;5(3):170-5.
104. Kropf EJ, Sekiya JK. Osteoarticular allograft transplantation for large humeral head defects in glenohumeral instability. *Arthroscopy.* 2007;23(3):322.
105. Armitage MS, Faber KJ, Drosdoweck DS, Litchfield RB, Athwal GS. Humeral head bone defects: remplissage, allograft, and arthroplasty. *Orthop Clin North Am.* 2010;41(3):417-25.
106. Paul RW, Reddy MP, Onor G, et al. Bankart repair with or without concomitant remplissage results in similar shoulder motion and postoperative outcomes in the treatment of shoulder instability. *Arthrosc Sports Med Rehabil.* 2022;5(1):e171-8.
107. Wu D, Zhou Z, Song W, et al. Arthroscopic autologous iliac crest grafting results in similar outcomes and low recurrence compared to remplissage plus Bankart repair for anterior shoulder instability with bipolar bone defects. *Arthroscopy.* 2023;39(7):1600-7.
108. Gouveia K, Harbour E, Athwal GS, Khan M. Return to sport after arthroscopic Bankart repair with remplissage: a systematic review. *Arthroscopy.* 2023;39(4):1046-59.
109. Davis WH, DiPasquale JA, Patel RK, et al. Arthroscopic remplissage combined with Bankart repair results in a higher rate of return to sport in athletes compared with Bankart repair alone or the Latarjet procedure: a systematic review and meta-analysis. *Am J Sports Med.* 2023;3635465221138559.
110. Horinek JL, Menendez ME, Narbona P, Ladermann A, Barth J, Denard PJ. Arthroscopic Bankart repair with remplissage as an alternative to Latarjet for anterior glenohumeral instability with more than 15% glenoid bone loss. *Orthop J Sports Med.* 2022;10(12):23259671221142257.
111. Paul RW, Reddy MP, Sonnier JH, et al. Increased rates of subjective shoulder instability after Bankart repair with remplissage compared to Latarjet surgery. *J Shoulder Elbow Surg.* 2023;32(5):939-46.