

Consensus on clinical management of tumor-induced osteomalacia

Yan Jiang¹, Xiang Li¹, Li Huo², Yong Liu³, Wei Lyu⁴, Lian Zhou⁵, Wei Yu⁶, Huan-Wen Wu⁷, Xiao-Ping Xing¹, Mei Li¹, Ou Wang¹, Yue Chi¹, Rui-Zhi Jiajue¹, Yu Pei⁸, Jian-Min Liu⁹, Jian-Ming Ba⁸, Qiao Zhang¹⁰, Zhi-Feng Sheng¹¹, Zhen-Lin Zhang¹², Jia-Jun Zhao¹³, Salvatore Minisola¹⁴, Wei-Bo Xia¹; on behalf of Chinese Society of Osteoporosis and Bone Mineral Research and Chinese Society of Endocrinology

¹Department of Endocrinology, Key Laboratory of Endocrinology, State Key Laboratory of Complex Severe and Rare Diseases, NHC, Peking Union Medical College Hospital, Chinese Academy of Medical Science and Peking Union Medical College, Beijing 100730, China;

²Department of Nuclear Medicine, Peking Union Medical College Hospital, Chinese Academy of Medical Science, Beijing 100730, China;

³Department of Orthopaedic Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Science, Beijing 100730, China;

⁴Department of ENT, Peking Union Medical College Hospital, Chinese Academy of Medical Science, Beijing 100730, China;

⁵Department of Stomatology, Peking Union Medical College Hospital, Chinese Academy of Medical Science, Beijing 100730, China;

⁶Department of Radiology, Peking Union Medical College Hospital, Chinese Academy of Medical Science, Beijing 100730, China;

⁷Department of Pathology, Peking Union Medical College Hospital, Chinese Academy of Medical Science, Beijing 100730, China;

⁸Department of Endocrinology, Chinese PLA General Hospital, Beijing 100853, China;

⁹Department of Endocrine and Metabolic Diseases, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai Institute of Endocrine and Metabolic Diseases, Shanghai Clinical Center for Endocrine and Metabolic Diseases, Shanghai 200025, China;

¹⁰Department of Endocrinology and Metabolism, Affiliated Hospital of Guiyang Medical University, Guiyang, Guizhou 550004, China;

¹¹Department of Metabolism and Endocrinology, The Second Xiangya Hospital, Central South University, Changsha, Hunan 410011, China;

¹²Department of Osteoporosis and Bone Diseases, Shanghai Clinical Research Center of Bone Disease, Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai 200233, China;

¹³Department of Endocrinology and Metabolism, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, Shandong 250021, China;

¹⁴Department of Internal Medicine and Medical Disciplines, Sapienza University of Rome, Rome, Italy.

Tumor-induced osteomalacia (TIO) is a rare paraneoplastic syndrome caused by excessive fibroblast growth factor 23 (FGF23) production by a tumor, which often arises from a mesenchymal origin.^[1-3] Most clinical symptoms of TIO are the consequences of prolonged FGF23-mediated hypophosphatemia as muscle weakness, bone pain, impaired mobility, and fractures.^[4] Clinical diagnosis and management of TIO are challenging because knowledge about this condition is still restricted to a few specialized centers, leading to delay in diagnosis and appropriate treatment. The scope of the present consensus is to provide up-to-date guidance on the assessment and treatment of TIO.

The writing committee consists of experts representing endocrinology, pathology, radiology, nuclear medicine, orthopedics, stomatology, and rhinology departments. From the evidence, especially high-quality evidence is limited or even non-existent for this rare disease; we provide recommendations based on expert's review on the limited data, as well as their experiences and opinions

when data are unavailable. This process may be less systematic than the grade methodological framework; however, it is unrealistic to gather more reliable evidence without an international consensus to promote standard management of TIO.

All participants signed a conflict of interest declaration, and the consensus was strictly supported by funding from academic or professional societies only, with no sponsorship from the pharmaceutical industry. The details of consensus on clinical management of TIO are shown in Supplementary Material, <http://links.lww.com/CM9/A506>.

Yan Jiang and Xiang Li contributed equally to this work.

Correspondence to: Wei-Bo Xia, Department of Endocrinology, Key Laboratory of Endocrinology, State Key Laboratory of Complex Severe and Rare Diseases, NHC, Peking Union Medical College Hospital, Chinese Academy of Medical Science and Peking Union Medical College, Beijing 100730, China

E-Mail: xiaweibo8301@163.com;

Salvatore Minisola, Department of Internal Medicine and Medical Disciplines, Sapienza University of Rome, Rome 00161, Italy

E-Mail: salvatore.minisola@uniroma1.it;

Jia-Jun Zhao, Department of Endocrinology, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, Shandong 250021, China

E-mail: jjzhao@sdu.edu.cn;

Zhen-Lin Zhang, Department of Osteoporosis and Bone Diseases, Shanghai Clinical Research Center of Bone Disease, Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai 200233, China

E-mail: zhangzl@sjtu.edu.cn.

Copyright © 2021 The Chinese Medical Association, produced by Wolters Kluwer, Inc. under the CC-BY-NC-ND license. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Chinese Medical Journal 2021;134(11)

Received: 11-01-2021 Edited by: Li-Shao Guo

Access this article online	
Quick Response Code:	Website: www.cmj.org
	DOI: 10.1097/CM9.0000000000001448

Diagnosis

The diagnosis of TIO is based on the association of clinical manifestations, biochemical findings, and the identification of the tumor (most importantly). Patients with symptoms, such as bone pain, weakness, and/or radiological signs of osteomalacia like obscure bone structure, concave changes of vertebrae, inward bending of the pelvic sidewall, and pseudofracture (Looser zone), should be suspected as TIO.^[4-6] Suspected patients should have further tests. Decreased serum phosphate, tubular maximum reabsorption of phosphate/glomerular filtration rate, low or inappropriately normal 1,25-dihydroxyvitamin D (1,25(OH)₂D) levels, and elevated alkaline phosphatase (ALP) accompanied with elevated FGF23 are typical laboratory findings. Besides, inherited disorders or other acquired causes of hypophosphatemic rickets/osteomalacia should be excluded at the very beginning of the diagnosis.^[2]

If clinical features, laboratory findings, and radiological results hint a diagnose of TIO together, a stepwise approach to locate the causative tumor is widely recommended since tumors are usually small, slow-growing with unexpected locations over the whole body.^[1,2] The first step is to screen the whole body for suspected lesions. This step comprises a thorough inquiry and physical examination, as well as functional imaging. The somatostatin receptor imaging method is recommended as a first-line imaging investigation, which comprises ⁶⁸Ga-DOTA-conjugated-somatostatin-receptor-targeting-peptides positron emission tomography/ computed tomography (CT) scan and octreoscan with single-photon emission computed tomography/CT. Once the TIO tumors are suspected by functional imaging or physical examinations, the next step is to confirm the lesions by anatomical imaging. Based on different sites of suspected masses, techniques including radiography, magnetic resonance imaging (MRI), CT, or ultrasound may be used. When accessible, MRI and CT are recommended because of their advantage in high resolution. Selective venous sampling may be useful to confirm causative tumors in patients with multiple suspicious regions. For patients who fail to locate the tumor, we recommend to start medical therapy and repeat imaging studies within 1 to 2 years.

Pathology

The overwhelming majority of TIO-associated tumors are of mesenchymal origin. Despite its polymorphous histology, it is widely accepted that almost all TIO-associated mesenchymal tumors are caused by a single morphologically and genetically distinct neoplasm, known as phosphaturic mesenchymal tumor (PMT).^[7] Most PMTs present as non-specific soft tissue or bone masses and typically produce a characteristic “smudgy” matrix that calcifies in a peculiar “grungy” or flocculent fashion.

Management and Treatment

Surgery

We recommend that surgical treatment is the first-line therapy when feasible. From the surgical perspective, the optimal treatment for TIO involves the complete removal of

the disease-causing tumor.^[8] In most cases, this procedure can correct biochemical abnormalities and accelerate the process of bone remineralization. However, even a small amount of tumor tissue remains, the patient’s symptoms continue to present or relapse easily. We recommend that non-remission and recurrent patients should repeat the diagnostic work-up to exclude other diseases and re-located causative tumor and receive reoperations when possible.^[9]

Medical treatment

We recommend that inoperable patients should initiate medical therapy to alleviate symptoms, and biochemical tests should be monitored every 3 to 6 months. Burosumab or phosphate with active vitamin D should be the first consideration when accessible.

The therapeutic goal of conventional medical treatment is to alleviate clinical symptoms, increase the serum phosphate levels, normalize ALP, and maintain parathyroid hormone (PTH) in the normal range. Complete normalization of serum phosphate usually represents an overdose. We recommend a dose of 20 to 40 mg/kg daily (1–3 g/d for adults) of element phosphate and a dose of 20 to 30 ng/kg daily (0.5–1.5 μg/d for adults) of calcitriol. The equivalent dosage of alphacalcidol is 1.5 to 2 times that of calcitriol. Phosphate supplements should be divided into 4 to 6 doses/d and titrated to the target dose over several days to weeks. Renal function, serum calcium, phosphate, ALP, PTH, and 24-h urinary calcium should be monitored every 3 to 6 months to adjust the medical dosage to prevent complications including secondary or tertiary hyperparathyroidism, nephrolithiasis, and reduced renal function.

Burosumab, a fully human monoclonal antibody against FGF23, is the most promising drug in the near future. The recommended initial dosage of burosumab for TIO is 0.5 mg/kg once every 4 weeks; round dose to the nearest 10 mg; and maximum dosage 2 mg/kg (not to exceed 180 mg) for every 2 weeks. Dosage adjustment based on serum phosphate: evaluate fasting serum phosphate monthly, measured 2 weeks post-dose, for the first 3 months of treatment and as clinically necessary thereafter.

Radiofrequency ablation and cryoablation can be tried in patients with TIO who are neither willing nor qualified to undergo complete excision surgeries on tumors. The effectiveness of other treatments, such as FGF receptor inhibitor or cinacalcet, is still doubtful because the evidence is insufficient.

Monitoring

Upon the removal of the tumor completely, the serum phosphate levels are normalized in 5 days (2–10 days).^[4] The serum 1,25(OH)₂D levels rise synchronously with the decrease of FGF23. For patients with complete tumor removal, biochemical parameters, especially serum phosphate, should be measured initially every 6 months and then at yearly intervals with a dual-energy X-ray absorptiometry examination. For patients who fail to locate the tumor and adopt a long-term medical treatment, the interval examinations for biochemical parameters




Figure 1: Diagnostic and management diagram of TIO. 1,25(OH)₂D: 1,25-dihydroxyvitamin D; ALP: Alkaline phosphatase; ⁶⁸Ga-DOTA-SST: ⁶⁸Ga-DOTA-conjugated-somatostatin-receptor-targeting-peptides; GFR: Glomerular filtration rate; PTH: Parathyroid hormone; TmP: Tubular maximum reabsorption of phosphate; TIO: Tumor-induced osteomalacia.

should be shortened to every 3 to 6 months to adjust the drug doses and prevent the side effects.^[10] The diagnostic and management diagram of TIO is shown in Figure 1.

Future Prospects

In terms of diagnosis, we need to propose some specific and easily-obtained criteria to help making suspicious diagnoses quickly in primary health care institutions. The mechanisms of tumorigenesis and FGF23 overproduction, detailed models and risk markers for stratifying severity, and refractory of TIO need to be further explored which would accurately benefit clinical management markedly. Prospective evidence of the effectiveness of novel drugs including burosumab, FGFR1 inhibitors would greatly expand treatment options of TIO in the future.

Acknowledgements

The authors deeply appreciate Prof. Seiji Fukumoto for his meticulous and insightful opinions in the process of revising the consensus. The authors thank Dr. Chen-Xi Jin, Dian-Yin Shi, Guo-Zhu Hou, Shu-Zhong Liu, Xiao-Lin Ni, and Zhen-Zhen Zhu for their indispensable contributions in developing the consensus.

Funding

This work was supported by grants from the CAMS Innovation Fund for Medical Science (No. 2016-I2M-3-003) and the National Natural Science Foundation of China (No. 81970757 and No. 81670714).

Conflicts of interest

None.

References

1. Minisola S, Peacock M, Fukumoto S, Cipriani C, Pepe J, Tella SH, *et al.* Tumour-induced osteomalacia. *Nat Rev Dis Primers* 2017;3:17044. doi: 10.1038/nrdp.2017.44.
2. Florenzano P, Hartley IR, Jimenez M, Roszko K, Gafni RI, Collins MT. Tumor-induced osteomalacia. *Calcif Tissue Int* 2020;108:128–142. doi: 10.1007/s00223-020-00691-6.
3. Shimada T, Mizutani S, Muto T, Yoneya T, Hino R, Takeda S, *et al.* Cloning and characterization of FGF23 as a causative factor of tumor-induced osteomalacia. *Proc Natl Acad Sci U S A* 2001;98:6500–6505. doi: 10.1073/pnas.101545198.
4. Jiang Y, Xia WB, Xing XP, Silva BC, Li M, Wang O, *et al.* Tumor-induced osteomalacia: an important cause of adult-onset hypophosphatemic osteomalacia in China: report of 39 cases and review of the literature. *J Bone Miner Res* 2012;27:1967–1975. doi: 10.1002/jbmr.1642.
5. Chong WH, Andreopoulou P, Chen CC, Reynolds J, Guthrie L, Kelly M, *et al.* Tumor localization and biochemical response to cure in tumor-induced osteomalacia. *J Bone Miner Res* 2013;28:1386–1398. doi: 10.1002/jbmr.1881.
6. Yu WJ, He JW, Fu WZ, Wang C, Zhang ZL. Reports of 17 Chinese patients with tumor-induced osteomalacia. *J Bone Miner Metab* 2017;35:298–307. doi: 10.1007/s00774-016-0756-9.
7. Folpe AL, Fanburg-Smith JC, Billings SD, Bisceglia M, Bertoni F, Cho JY, *et al.* Most osteomalacia-associated mesenchymal tumors are a single histopathologic entity: an analysis of 32 cases and a comprehensive review of the literature. *Am J Surg Pathol* 2004;28:1–30. doi: 10.1097/00000478-200401000-00001.
8. Sun ZJ, Jin J, Qiu GX, Gao P, Liu Y. Surgical treatment of tumor-induced osteomalacia: a retrospective review of 40 cases with extremity tumors. *BMC Musculoskelet Disord* 2015;16:43. doi: 10.1186/s12891-015-0496-3.
9. Li X, Jiang Y, Huo L, Wu H, Liu Y, Jin J, *et al.* Nonremission and recurrent tumor-induced osteomalacia: a retrospective study. *J Bone Miner Res* 2020;35:469–477. doi: 10.1002/jbmr.3903.
10. Chong WH, Molinolo AA, Chen CC, Collins MT. Tumor-induced osteomalacia. *Endocr Relat Cancer* 2011;18:R53–77. doi: 10.1530/ERC-11-0006.

How to cite this article: Jiang Y, Li X, Huo L, Liu Y, Lyu W, Zhou L, Yu W, Wu HW, Xing XP, Li M, Wang O, Chi Y, Jiajue RZ, Pei Y, Liu JM, Ba JM, Zhang Q, Sheng ZF, Zhang ZL, Zhao JJ, Minisola S, Xia WB; on behalf of Chinese Society of Osteoporosis and Bone Mineral Research and Chinese Society of Endocrinology. Consensus on clinical management of tumor-induced osteomalacia. *Chin Med J* 2021;134:1264–1266. doi: 10.1097/CM9.0000000000001448