

ORAL PRESENTATION

Open Access

Neurocognitive impairment screening in Romanian HCV infected patients

Ioana-Catrinel Cercel^{1*}, Şerban Polli¹, Oana Streinu-Cercel^{1,2}, Anca Streinu-Cercel^{1,2}, Adrian Marinescu¹, Adrian Streinu-Cercel^{1,2}

From The 9th Edition of the Scientific Days of the National Institute for Infectious Diseases Prof Dr Matei Bals Bucharest, Romania. 23-25 October 2013

Background

Neurocognitive impairment can occur in HCV infected patients. A series of studies have shown that patients with HCV may experience diminished attention and concentration, or suffer from severe cognitive deficits, such as disorientation and fluctuating consciousness. We performed a study to determine if there are neurocognitive differences or dissimilarities in the quality of life between a control group and a group of HCV infected patients.

Methods

We assessed patients by applying the EQ-5D-5L questionnaire and the Montreal Cognitive Assessment. EQ-5D-3L comprises the following 5 dimensions: mobility, self-care, usual activities, pain/discomfort and anxiety/depression. Montreal Cognitive Assessment indicates if patients have neurocognitive impairment. All subjects were evaluated with these questionnaires during an appointment with the clinical psychologist.

Results

We assessed 11 HCV infected patients and 11 controls. The HCV group had ages between 25-58 with a median of 40 ± 12 (2 retired, 1 doctor, 1 economist, 2 engineers, 1 architect, 1 editor, 1 clerk, 1 financial analyst) and the control group had ages between 26-38, median 30 ± 4 (7 doctors, 3 nurses and 1 secretary). EQ-5D-5L scores were 85 ± 10 percent (min: 65, max: 95) in the HCV group and 95 ± 7 percent (min: 85, max: 100) in the control group. The median MOCA scores were 28 ± 4 (min: 19, max: 31) in the HCV group and 30 ± 3 (min: 22, max: 31) in the control group.

* Correspondence: catycercel@yahoo.com

¹National Institute for Infectious Diseases "Prof. Dr. Matei Balş", Bucharest, Romania

Full list of author information is available at the end of the article

Conclusion

At a first look, we identified apparent neurocognitive differences between the two groups but the number of subjects was too low to offer a statistical significance. We intend to evaluate larger groups of patients, and reapply the tests periodically to determine if there are longitudinal changes in the neurocognitive status.

Authors' details

¹National Institute for Infectious Diseases "Prof. Dr. Matei Balş", Bucharest, Romania. ²Carol Davila University of Medicine and Pharmacy, Bucharest, Romania.

Published: 16 December 2013

doi:10.1186/1471-2334-13-S1-O29

Cite this article as: Cercel et al.: Neurocognitive impairment screening in Romanian HCV infected patients. *BMC Infectious Diseases* 2013 13(Suppl 1):O29.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

