



OPEN

Identifying optimal reference genes for gene expression studies in Eurasian spruce bark beetle, *Ips typographus* (Coleoptera: Curculionidae: Scolytinae)

Gothandapani Sellamuthu¹, Jan Bílý¹, Mallikarjuna Reddy Joga¹, Jiří Synek¹ & Amit Roy^{1,2}✉

Eurasian spruce bark beetle (*Ips typographus* [L.]) causes substantial damage to spruce forests worldwide. Undoubtedly, more aggressive measures are necessary to restrict the enduring loss. Finishing genome sequencing is a landmark achievement for deploying molecular techniques (i.e., RNA interference) to manage this pest. Gene expression studies assist in understanding insect physiology and deployment of molecular approaches for pest management. RT-qPCR is a valuable technique for such studies. However, accuracy and reliability depend on suitable reference genes. With the genome sequence available and the growing requirement of molecular tools for aggressive forest pest management, it is crucial to find suitable reference genes in *Ips typographus* under different experimental conditions. Hence, we evaluated the stability of twelve candidate reference genes under diverse experimental conditions such as biotic (developmental, sex and tissues) and abiotic factors (i.e., temperature and juvenile hormone treatment) to identify the reference genes. Our results revealed that *ribosomal protein 3a* (*RPS3-a*) was the best reference gene across all the experimental conditions, with minor exceptions. However, the stability of the reference gene can differ based on experiments. Nevertheless, present study provides a comprehensive list of reference genes under different experimental conditions for *Ips typographus* and contributes to “future genomic and functional genomic research”.

The Eurasian spruce bark beetle, *Ips typographus* (Coleoptera: Curculionidae: Scolytinae), is the most devastating pest in Eurasian coniferous forests of Norway spruce (*Picea abies* (L.) Karst) and has caused significant ecological and economic damage in recent years under the influence of climate change. It primarily affected European countries^{1–4} and was recently found in Kent and East Sussex (England) (Forestry Commission England, 2021, <https://www.gov.uk/government/news/forestry-commission-acts-on-bark-beetle-tree-pest>). Some recent publications even indicated the risk of *Ips typographus* and its associated pathogen invasion to Canada and America based on climatic and host suitability^{5,6}. In Europe, tree mortality exceeded 2.9 million m³ per year from 1950 to 2000 and has increased further in the last decades^{7–9}. In the Czech Republic, the loss of vegetation was estimated to be 14.5 million m³ during recent years¹⁰. *Ips typographus* thrives on dying trees when their populations are at the endemic stage, and mass attack on healthy trees happens only at epidemic levels^{1,11,12}. However, climate change often rises the occurrence and severity of outbreaks^{9,13–15}. The management of *I. typographus* has been primarily relied on mass-trapped using pheromone, sanitary cutting, trap trees, salvage logging, and insecticides¹⁶. Protecting the forest health and environment from toxic chemical insecticides has led to searching for superior alternatives for forest pest insect management strategies such as RNAi, as reviewed¹⁷. Therefore, molecular studies are essential for facilitating new pest management strategies using available genomic resources.

Gene expression analyses have become immensely important for revealing gene function and molecular regulation under the different environmental responses during the bark beetle life cycle or beetle-host interaction. Reverse transcription-quantitative polymerase chain reaction (RT-qPCR) has become the most extensively accepted methodology for detecting and quantifying target gene expression with higher accuracy, sensitivity,

¹Excellent Team for Mitigation (ETM), Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic. ²EVA 4.0 Unit, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic. ✉email: roy@fld.czu.cz

and reproducibility compared with other traditional methods molecular techniques^{18–23}. It is especially useful to detect low-abundance mRNAs in limited samples^{24,25}. Nevertheless, RT-qPCR data are influenced by many factors such as initial RNA sample quantity and quality, the efficiency of cDNA synthesis, mRNA recovery, primer and PCR efficiency^{26–31}. Furthermore, the reliability of RT-qPCR data is highly dependent on the appropriate reference genes as internal controls from the same samples across various biotic and abiotic stresses and treatments²⁵. A literature search indicates that reference genes that are constitutively expressed under different environmental factors and maintain the essential cellular functions have been used extensively as internal controls for expression normalization of target genes^{32,33}. It is pretty clear now that a single reference gene is not appropriate for the more comprehensive experimental conditions, and it can generate an error in the gene expression estimations causing nonoptimal interpretation of the data^{25,34,35}. For mitigation, it is often recommended to use multiple reference genes to minimize variations by RT-qPCR normalization^{23,36,37}. Several research studies revealed that most reference gene expression depends on samples/experimental conditions, suggesting no universal reference gene is available for all experimental conditions³⁸. Hence, for accurate gene expression normalization, it is essential to evaluate the stability of reference genes for different environmental conditions, life stages, sex-specific and tissue-specific stages for each insect^{25,31,32,39–43}.

In the present study, we analyzed the expression level of 12 commonly occurring reference genes in different coleopteran insects based on published articles^{38,43,44}. These genes mostly perform conserved cellular functions in the coleopteran insects, hence expected to be constitutively and stably expressed in all tissues and cells under different experimental conditions³⁸. Our objective is to find suitable reference genes for future gene expressions studies in *Ips typographus*. Candidate reference genes were evaluated across different experimental conditions are *elongation factor 1 α* (*EF-1 α*), *ribosomal protein L13a* (*RPL13a3*), *arginine kinase isoform XI* (*ArgK*), *ribosomal protein L7* (*RPL7*), *glyceraldehyde-3-phosphate dehydrogenase* (*GAPDH*), *β -actin* (*Actin*), *ribosomal protein 3a* (*RPS3-a*), *Tubulin beta-1 chain* (*β -Tubulin*), *ubiquitin C variant* (*UbiQ*), *V-type proton ATPase catalytic subunit A* (*V-ATPase-A*), *ribosomal protein S7* (*RPS7*), *ribosomal protein L6* (*RPL6*). Reference genes were tested across developmental stages (first, second and third instar larvae, pupa, adult male and female) and tissues (head, mid-gut, and fat body). Additionally, adult beetles were exposed to juvenile hormone treatment (JHIII) and varied temperatures to identify reference genes for those conditions. Wild and lab-reared beetles were also evaluated for the same intent. Nevertheless, the present study tried to identify reference genes suitable for various tested experimental conditions for *I. typographus*.

Material and methods

Beetles. *Ips typographus* were obtained from Kostelec nad Černými lesy (50° 00' 07.2" N 14° 50' 56.3" E, under School Forest Enterprise) in summer 2021. School Forest Enterprise in Kostelec nad Černými lesy (Czech University of Life Sciences Prague) is located in the Central Bohemia region, 40 km southwest from Prague. It is a relatively warmer and dryer area, mean annual temperature of 7–7.5 °C, the mean annual sum of precipitation 600 mm, length of vegetation season 150 days. Range of elevation 350–520 m a.s.l. Natural vegetation was composed mainly of beech and oak mixed forest with fir. However, the current species composition is dominated by spruce (50%) and pine (18%)⁴⁵. Due to the extreme drought in 2018, the whole area was under tremendous pressure from the bark beetle outbreak, mainly *I. typographus*⁴⁶. Collected beetles from the forest were maintained in an insect rearing chamber with fresh Norway spruce logs (*Picea abies*) at 27 ± 1 °C under 70 ± 5% humidity and a 16:8-h light/dark (L:D) photoperiod at Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Prague. The wild beetle population was supplied with fresh spruce logs for the next generation and maintained for 35–40 days.

Experimental conditions. *Biotic factors (life stages and tissue types).* Developmental stages such as three larval stages, pupae, callow male and female, and fed adult male and female were collected from *I. typographus*. Tissues, including head, fat body, and gut, were dissected from the callow and fed adult males and females (Fig. 1). The samples details are summarized in Table 1. Each replicate was derived by pooling tissues from ten beetles for tissue-specific comparisons. Four biological replicates were used for each bark beetle developmental stages and tissues.

Abiotic factors (temperature, JHIII treatment and wild beetles vs lab-reared beetles). To examine the effect of temperature treatment, the freshly emerged adults were exposed to 4, 27, and 37 °C for 72 h. For the juvenile hormone treatment (JHIII), fed adult male and female beetles were treated topically on the ventral surface of the abdomen with 10 µg JHIII (Sigma-Aldrich, St. Louis, MO, USA) and acetone as control^{47,48}. Beetles were maintained in similar conditions as mentioned above for 72 h. Wild beetles (adults) were collected from the infested Norway spruce stands from Kostelec nad Černými lesy, Czech Republic, to examine the climatic effect. Similarly, laboratory-reared adults emerged from eggs produced by the wild- beetles. The lab-reared beetles were collected after three generations. Four independent biological replications for each experimental condition were used. The experimental design is summarized in Fig. 1.

Selection of candidate reference genes for evaluation. The reference genes selected for this study were reference genes in other Coleoptera species and showed stable expression in our in-house *I. typographus* genome data⁴⁴ (Data not shown). The primers for those selected genes were designed via IDT (Integrated DNA Technologies), and the primer efficiencies (E) and correlation coefficients (R2) were also calculated (Table 2).

Total RNA extraction and RT-qPCR analysis. Total RNA was extracted from each tissue using TRIzol® (Invitrogen, Carlsbad, CA) following the manufacturer's protocol. Isolated RNA was further treated with




Figure 1. Schematic representation of experimental plan for finding reference genes in *Ips typographus*.

| Sample category | Sample name | Abbreviation |
|---------------------------|------------------|------------------|
| Developmental stages | Larval 1 | L1 |
| | Larval 2 | L2 |
| | Larval 3 | L3 |
| | Pupae | P |
| | Callow male | CM |
| | Callow female | CF |
| | Fed adult male | AMF |
| | Fed adult female | AFF |
| | Tissues | Callow male head |
| Callow male gut | | CMG |
| Callow male fat body | | CMFB |
| Callow female head | | CFH |
| Callow female gut | | CFG |
| Callow female fat body | | CFFB |
| Fed adult male head | | AMFH |
| Fed adult male gut | | AMFG |
| Fed adult male fat body | | AMFFB |
| Fed adult female head | | AFFH |
| Fed adult female gut | | AFFG |
| Fed adult female fat body | | AFFFB |

Table 1. Detailed sample list of *I. typographus* life stages and tissue types used for reference gene experiments.

DNases using a TURBO DNAase Kit (Ambion, USA). cDNA was synthesized from 1 µg RNA using the High-Capacity cDNA Reverse Transcription kits (Applied Biosystems-Life Technologies) following the manufacturer's recommendations and stored at -20°C . cDNA samples were diluted ten times before RT-qPCR reaction. Four independent biological replicates from each treatment were included in each RT-qPCR experiment. RT-qPCR reaction mix contained 5.0 µL of SYBR[®] Green PCR Master Mix (Applied Biosystems), 1.0 µL of cDNA, 1.0 µL optimized concentrations of primers (Table 2), and RNase-free water (Invitrogen) to a total volume of 10.0 µL. Amplification conditions were as follows: initial denaturation at 95°C for 10 min, followed by 40 cycles of 95°C for 15 s, 60°C for 1 min. The reactions were performed in an Applied Biosystems™ StepOne™ Real-Time PCR System (Applied Biosystems). To confirm the primer specificity, melting curve analysis was used to confirm gene-specific amplification by a steady increase in temperature from 60 to 95°C . All RT-qPCR assays were carried out in four biological replicates, including three technical replicates.

Data analyses. Four different algorithms, namely geNorm²⁵, NormFinder⁴⁹, BestKeeper³⁹ and ΔCT method⁵⁰, were used for measuring reference gene expression stability. geNorm computes the expression stability value (M) and pairwise variation comparison. NormFinder ranks the set of reference genes based on their expression stability within a given set of samples. BestKeeper is also a freely available algorithm that considers the Cq (quantification cycle) value of all reference genes to evaluate standard deviation and correlation coef-

| Gene symbol | Gene name | BioProject accession numbers PRJNA671615* | Primer sequences (5'-3') | Amplicon length (bp) | PCR efficiency % | Regression coefficient |
|--------------------|--|---|-----------------------------------|----------------------|------------------|------------------------|
| <i>EF1-a</i> | Elongation factor 1-alpha | Ityp13440 | FP: GCCATTCTTCCACCA TCTCGTC | 210 | 98.99 | 0.989 |
| | | | RP: CAGGAGCAAACA CCACTACCATAAC | | | |
| <i>RPL13a</i> | 60S ribosomal protein L13a | Ityp07303 | FP: CTGGAGCACTTAGGG TTGTC | 215 | 97.2 | 0.972 |
| | | | RP: TGAATGGCTTACCT GCTTG | | | |
| <i>ArgK</i> | Arginine kinase | Ityp12981 | FP: GTCTTCGCTGACTTG TTCGACC | 104 | 119.24 | 0.985 |
| | | | RP: GCCAAATACGCTCAC ATCACCC | | | |
| <i>RPL7</i> | 60S ribosomal protein L7 | Ityp01351 | FP: GCTGAACACCCCAAC TGGAG | 100 | 97.6 | 0.976 |
| | | | RP: CTGTTGCCAAAGTCA CCACC | | | |
| <i>GAPDH</i> | Glyceraldehyde-3-phosphate dehydrogenase | Ityp02436 | FP: GGCAAATTGTGGAGA GAGGGAC | 200 | 96.9 | 0.969 |
| | | | RP: GCGGGCTTCTTGAT CTTGG | | | |
| <i>Actin</i> | β -actin | Ityp12788 | FP: ACGAAAGATTCCGTT GCCCC | 170 | 114.79 | 0.954 |
| | | | RP: GCGTACAAGTCCTTA CGAATGTCC | | | |
| <i>RPS3-a</i> | 40S ribosomal protein S3-A | Ityp04549 | FP: GCCCCTTCAATGTT GCCAC | 119 | 96.2 | 0.972 |
| | | | RP: AAGTCCGCCAAACTA ACCTCA | | | |
| β -Tubulin | Tubulin beta-1 chain | Ityp08900 | FP: TGATGACGAGTACGA AGCGG | 138 | 114.67 | 0.984 |
| | | | RP: CAAAGCAAGGCA CTCTTGGTC | | | |
| <i>UbiQ</i> | Ubiquitin C variant | Ityp20676 | FP: CGCACACAAATCATG GGTGG | 120 | 94.1 | 0.944 |
| | | | RP: GTCAAAGGCACGTCT GATTTTC | | | |
| <i>V-ATPase-A</i> | V-type proton ATPase catalytic subunit A | Ityp14486 | FP: GTCGCCTCCTACAGG GAATG | 120 | 92.2 | 0.922 |
| | | | RP: CCAGAGACCGAA GGCAAACA | | | |
| <i>RPS7</i> | 40S ribosomal protein S7 | Ityp08836 | FP: CTGGTTAGGGAGTTG GAGA | 140 | 118.48 | 0.969 |
| | | | RP: GCATCATATACAGCC GTCAG | | | |
| <i>RPL6</i> | 60S ribosomal protein L6 | Ityp04817 | FP: CCACCAGCGAACATT GAG | 148 | 115.94 | 0.971 |
| | | | RP: TTCCGTAGGGACCTG TAAC | | | |
| Target Gene | | | | | | |
| <i>CYP03903</i> | Cytochrome P450 | Ityp03903 | FP: GTATTGCGCTGCTCA TTCC | 114 | 102.44 | 0.951 |
| | | | RP: CCGGTCTACTGGATC TGTT | | | |

Table 2. Primer sequence, amplicon length and RT-qPCR analysis of candidate reference genes and a target gene. *Powell et al. 2021.

efficient. Δ CT method directly evaluates the relative expression of 'gene pairs' within each sample. The mean Cq values of each reference gene from each experiment are offered as input data and subsequently processed using the web-based tool RefFinder (<https://www.heartcure.com.au/reffinder/>), which delivers a comprehensive stability index that ultimately ranks each reference gene. Pairwise variation (V), estimated by geNorm, was used to determine the optimal number of reference genes for precise RT-qPCR normalization. The V_n/V_{n+1} value exhibited the pairwise variation between two sequential normalization factors²⁵.

Validation of selected reference genes. In insects, cytochrome P450 monooxygenases (P450s) are key enzymes that detoxify a broad spectrum of xenobiotics such as plant allelochemicals and synthetic insecticides. To validate the selected reference gene, we analysed relative expression levels of cytochrome P450 (*CYP03903*;




Figure 2. Expression range of Ct (Cycle threshold) values of candidate reference genes under different experimental conditions in *I. typographus*. The different conditions included in biotic conditions: developmental stages (i.e., larvae, pupae, and adult stages), sex-specific (male and female), tissue types (head, midgut, fat body) and abiotic conditions: different treatments (such as temperature; Juvenile hormone III; wild-collected vs long-term laboratory-reared beetles).

Table 2) in life stages and various tissue types of the beetle according to the $2 - \Delta\Delta CT$ method for gene normalization using most and least stable reference genes⁵¹. Target genes mRNA expression level was analysed using one-way ANOVA using GraphPad Prism software. P -value < 0.05 was deployed as the cut-off to indicate significant differences between tested samples.

Results

Candidate reference gene selection and PCR efficiency. Total twelve candidate reference genes, *EF-1α*, *RPL13a*, *ArgK*, *RPL7*, *GAPDH*, *Actin*, *RPS3-a*, *β-Tubulin*, *UbiQ*, *V-ATPase-A*, *RPS7*, and *RPL6* were selected for identifying suitable reference genes from *I. typographus*. Each reference gene was produced a single amplicon, as deduced by agarose gel electrophoresis (Figure S1) and melting curve analyses (Figure S2). The amplification efficiency of each primer pair fluctuated from 92.2 to 119.24%, and the correlation coefficients (R^2) were larger than 0.94 (Table 2). The Ct values of the twelve candidate reference genes ranged from 19.87 to 34.78 and covered all experimental conditions (Fig. 2). While most Ct values ranged from 19 to 27, *Actin*, *eEF2*, *β-Tubulin*, and *RPS3* were the most abundant transcripts under almost all experimental conditions. The least frequently expressed reference genes were *NADH*, *RPL17*, and *HSP83*. The five remaining reference genes were expressed at moderate levels.

Stability of candidate reference genes. Four different algorithms (geNorm, NormFinder, BestKeeper, and delta-CT) were used to find stable reference genes. Gene expression stability was assessed under different experimental conditions such as biotic factors (Table 1; “Section Biotic factors (life stages and tissue types)”) and abiotic factors (“Section Abiotic factors (temperature, JHIII treatment and wild beetles vs lab-reared beetles)”) using the web-based tool RefFinder.

Biotic conditions. For different developmental stages that include three larval stages (from the first instar to third instar; L1–L3), one pupal stage (P), and two adults (callow and fed), the top three most stable candidates were *RPS3-a*, *EF-1α*, and *RPS7* based on Ct method, Bestkeeper, RefFinder and Normfinder (Table 3). The stability ranking order of the first three most stable reference genes that was obtained from four programs, was inconsistent (Table 3). The geNorm ranking of the top most stable reference genes was *RPS3-a*, *RPL6*, *RPS7*, and *RPL13a* (Fig. 3A). Integrating the results of all programs identified the consensus top three candidates, *RPS3-a*, *EF-1α*, and *RPS7*, as the most stable reference genes across the developmental stages. Alternatively, *GAPDH*, *UbiQ*, and *RPL7* were the least stable genes (Table 3 and Fig. 3A).

| Conditions | Genes | ΔCt Method | | Best keeper | | Reffinder | | NormFinder | | Recommended Genes |
|-------------------|-------------------|------------|------|-------------|-------|-----------|-------|------------|------|--|
| | | Stability | Rank | Stability | Rank | Stability | Rank | Stability | Rank | |
| Life stage | <i>RPS3-a</i> | 1.25 | 1 | 1.55 | 5 | 1.5 | 1 | 0.238 | 2 | <i>RPS3-a</i> , <i>EF-1α</i> and <i>RPS7</i> |
| | <i>EF-1α</i> | 1.31 | 2 | 1.64 | 7 | 3.44 | 3 | 0.245 | 3 | |
| | <i>RPS7</i> | 1.32 | 3 | 1.48 | 4 | 3.22 | 2 | 0.274 | 6 | |
| | <i>RPL13a</i> | 1.32 | 4 | 1.63 | 6 | 4.43 | 7 | 0.894 | 12 | |
| | <i>RPL6</i> | 1.35 | 5 | 1.75 | 9 | 4.05 | 4 | 0.224 | 1 | |
| | <i>V-ATPase-A</i> | 1.38 | 6 | 1.25 | 2 | 4.36 | 6 | 0.309 | 7 | |
| | <i>ArgK</i> | 1.58 | 7 | 0.93 | 1 | 4.3 | 5 | 0.264 | 5 | |
| | <i>RPL7</i> | 1.77 | 8 | 1.74 | 8 | 8 | 9 | 0.382 | 10 | |
| | <i>β-Tubulin</i> | 1.88 | 9 | 2.14 | 11 | 9.46 | 10 | 0.316 | 8 | |
| | <i>Actin</i> | 2.02 | 10 | 1.4 | 3 | 7.4 | 8 | 0.258 | 4 | |
| | <i>UbiQ</i> | 2.04 | 11 | 2.28 | 12 | 11.24 | 11 | 0.326 | 9 | |
| <i>GAPDH</i> | 2.07 | 12 | 1.85 | 10 | 11.47 | 12 | 0.403 | 11 | | |
| Male | <i>RPS3-a</i> | 1.2 | 1 | 1.12 | 6 | 1.57 | 1 | 0.246 | 1 | <i>RPS3-a</i> , <i>RPL7</i> and <i>EF-1α</i> |
| | <i>RPL7</i> | 1.29 | 2 | 0.91 | 2 | 2.38 | 2 | 0.295 | 8 | |
| | <i>EF-1α</i> | 1.38 | 3 | 1.22 | 8 | 4.56 | 4 | 0.316 | 11 | |
| | <i>RPL6</i> | 1.4 | 4 | 1.5 | 10 | 5.62 | 6 | 0.248 | 2 | |
| | <i>GAPDH</i> | 1.42 | 5 | 1.47 | 9 | 5.96 | 8 | 0.271 | 5 | |
| | <i>RPL13a</i> | 1.43 | 6 | 1.1 | 4 | 3.46 | 3 | 0.323 | 12 | |
| | <i>RPS7</i> | 1.47 | 7 | 1.21 | 7 | 5.86 | 7 | 0.250 | 3 | |
| | <i>β-Tubulin</i> | 1.51 | 8 | 0.83 | 1 | 4.6 | 5 | 0.308 | 9 | |
| | <i>UbiQ</i> | 1.72 | 9 | 1.07 | 3 | 6.84 | 9 | 0.314 | 10 | |
| | <i>ArgK</i> | 1.77 | 10 | 1.53 | 11 | 10.24 | 11 | 0.294 | 7 | |
| | <i>Actin</i> | 2 | 11 | 2.09 | 12 | 11.24 | 12 | 0.258 | 4 | |
| <i>V-ATPase-A</i> | 2.08 | 12 | 1.11 | 5 | 9.64 | 10 | 0.285 | 6 | | |
| Female | <i>RPS3-a</i> | 1.16 | 1 | 1.52 | 8 | 1.68 | 1 | 0.378 | 9 | <i>β-Tubulin</i> , <i>RPS3-a</i> and <i>RPL6</i> |
| | <i>RPL7</i> | 1.34 | 6 | 1.58 | 9 | 5.8 | 6 | 0.378 | 10 | |
| | <i>EF-1α</i> | 1.29 | 5 | 1.4 | 6 | 5.69 | 5 | 0.314 | 4 | |
| | <i>RPL6</i> | 1.18 | 2 | 1.71 | 10 | 2.78 | 2 | 0.318 | 5 | |
| | <i>GAPDH</i> | 1.34 | 7 | 1.73 | 11 | 7.8 | 10 | 0.248 | 1 | |
| | <i>RPL13a</i> | 1.46 | 10 | 1.11 | 3 | 7.21 | 9 | 0.342 | 7 | |
| | <i>RPS7</i> | 1.19 | 3 | 1.41 | 7 | 3.6 | 4 | 0.382 | 11 | |
| | <i>β-Tubulin</i> | 1.23 | 4 | 0.89 | 2 | 3.56 | 3 | 0.330 | 6 | |
| | <i>UbiQ</i> | 1.35 | 8 | 1.33 | 4 | 6.26 | 8 | 0.289 | 3 | |
| | <i>ArgK</i> | 1.45 | 9 | 1.38 | 5 | 7.98 | 11 | 0.263 | 2 | |
| | <i>Actin</i> | 1.87 | 12 | 2.37 | 12 | 12 | 12 | 0.364 | 8 | |
| <i>V-ATPase-A</i> | 1.7 | 11 | 0.76 | 1 | 6.04 | 7 | 0.393 | 12 | | |

Table 3. Ranking of the candidate reference genes based on stability values performed by Delta Ct, BestKeeper, Reffinder, and NormFinder, in life stages (or developmental stages) and sex-specific conditions.

For sex-specific tissue comparisons (head, fat body, gut tissues of male and female beetles), the consensus top three candidates were calculated separately for male and female [callow male (CM), female (CF), fed adult male (AMF) and female (AFF)]. The top three most stable genes in male tissues were *RPS3-a*, *EF-1α*, and *RPS7*. The least stable genes defined by four programs were *V-ATPase-A*, *Actin*, and *ArgK* (Table 3). However, based on Reffinder, the most stable reference genes were *EF-1α*, *GAPDH*, *ArgK*, and *RPL13a*, and the least were *Actin*, *V-ATPase-A*, and *RPL7* (Fig. 3B). In contrast, *β-Tubulin*, *RPS3-a*, and *RPL6* were the most stable genes in females. The least stable genes were *V-ATPase-A*, *Actin*, and *RPL7* (Table 3). The stability ranking for most stable reference genes was constant for females, i.e., *RPS7* and *RPL6* in the Reffinder (Fig. 3C).

Tissue-specific expression stability of candidate reference genes was calculated for various bark beetle tissues (head, gut, and fat body). In head tissues, the most stable genes were *RPL7*, *RPL6*, and *RPS3-a* (Table 4). *RPS3-a* and *RPL6* were the best combinations of reference genes based on Reffinder (Fig. 4A). Similarly, *RPS7*, *GAPDH*, *ArgK* and *EF-1α*, *RPS3-a*, *RPL6* were the most stable genes within the fat body and gut tissues, respectively (Table 4). In contrast, the best combination of the genes was *RPS3-a/RPL6* and *V-ATPase-A/RPL6*, respectively, as per Reffinder (Fig. 4B,C). The least stable genes defined by four programs were *RPL13a*, *RPL7*, and *UbiQ* in the head, fat body, and gut tissues (Table 4). One interesting unanimous observation was that *RPS3-a* was the most stable reference gene except for the fat body under biotic conditions.




Figure 3. The average expression stability values (M) of twelve reference genes under different conditions (developmental stage and sex-specific conditions) were calculated by geNorm displaying the least stable (left) to the most stable (right). (A) Developmental stages (or life stages), (B) callow and fed adult male, (C) callow and fed adult female.

Abiotic conditions. Abiotic conditions such as temperature, juvenile hormone III (JHIII) treatment, and laboratory-reared vs wild beetles were considered to evaluate the suitable reference genes using the aforementioned

| Conditions | Genes | ΔCt Method | | Best keeper | | RefFinder | | NormFinder | | Recommended Genes |
|---------------|-------------------|------------|------|-------------|-------|-----------|-------|------------|------|-------------------------------|
| | | Stability | Rank | Stability | Rank | Stability | Rank | Stability | Rank | |
| Head | <i>RPL7</i> | 1.02 | 1 | 0.63 | 2 | 1.68 | 1 | 0.242 | 3 | <i>RPL7, RPL6 and RPs3-a</i> |
| | <i>RPL6</i> | 1.03 | 2 | 0.74 | 3 | 1.86 | 2 | 0.300 | 11 | |
| | <i>Actin</i> | 1.09 | 3 | 1 | 9 | 3.95 | 5 | 0.258 | 4 | |
| | <i>RPS3-a</i> | 1.12 | 4 | 0.79 | 4 | 2.83 | 3 | 0.288 | 9 | |
| | <i>β-Tubulin</i> | 1.16 | 5 | 0.5 | 1 | 3.87 | 4 | 0.284 | 8 | |
| | <i>UbiQ</i> | 1.18 | 6 | 0.89 | 8 | 6.93 | 6 | 0.279 | 7 | |
| | <i>ArgK</i> | 1.23 | 7 | 1.07 | 10 | 7.65 | 7 | 0.261 | 5 | |
| | <i>EF-1α</i> | 1.27 | 8 | 1.16 | 11 | 7.7 | 8 | 0.314 | 12 | |
| | <i>GAPDH</i> | 1.31 | 9 | 1.43 | 12 | 8.97 | 10 | 0.186 | 1 | |
| | <i>V-ATPase-A</i> | 1.35 | 10 | 0.83 | 6 | 8.57 | 9 | 0.228 | 2 | |
| | <i>RPS7</i> | 1.44 | 11 | 0.8 | 5 | 9.03 | 11 | 0.270 | 6 | |
| <i>RPL13a</i> | 1.71 | 12 | 0.83 | 7 | 10.49 | 12 | 0.293 | 10 | | |
| Fat body | <i>RPL7</i> | 1.49 | 12 | 1.31 | 8 | 10.84 | 12 | 0.464 | 11 | <i>RPS7, GAPDH and ArgK</i> |
| | <i>RPL6</i> | 1.13 | 7 | 1.43 | 12 | 8.76 | 9 | 0.356 | 5 | |
| | <i>Actin</i> | 1.47 | 11 | 1.29 | 7 | 9.82 | 11 | 0.478 | 12 | |
| | <i>RPS3-a</i> | 1.07 | 4 | 1.29 | 6 | 5.42 | 8 | 0.387 | 8 | |
| | <i>β-Tubulin</i> | 1.18 | 9 | 0.86 | 1 | 4.74 | 6 | 0.405 | 9 | |
| | <i>UbiQ</i> | 1.18 | 10 | 1.37 | 10 | 9.46 | 10 | 0.296 | 2 | |
| | <i>ArgK</i> | 1.08 | 5 | 0.99 | 3 | 3.08 | 3 | 0.336 | 4 | |
| | <i>EF-1α</i> | 1.07 | 3 | 1.39 | 11 | 4.46 | 5 | 0.368 | 7 | |
| | <i>GAPDH</i> | 1.03 | 1 | 1.32 | 9 | 2.71 | 2 | 0.252 | 1 | |
| | <i>V-ATPase-A</i> | 1.16 | 8 | 1.07 | 4 | 4.12 | 4 | 0.428 | 10 | |
| | <i>RPS7</i> | 1.04 | 2 | 0.9 | 2 | 2.21 | 1 | 0.364 | 6 | |
| <i>RPL13a</i> | 1.11 | 6 | 1.09 | 5 | 5.23 | 7 | 0.314 | 3 | | |
| Gut | <i>RPL7</i> | 1.02 | 4 | 1.19 | 8 | 5.26 | 5 | 0.314 | 6 | <i>EF-1α, RPS3-a and RPL6</i> |
| | <i>RPL6</i> | 1.05 | 5 | 1.16 | 7 | 5.14 | 4 | 0.283 | 4 | |
| | <i>Actin</i> | 1.32 | 12 | 1.69 | 12 | 12 | 12 | 0.248 | 2 | |
| | <i>RPS3-a</i> | 0.9 | 1 | 1.06 | 4 | 1.41 | 1 | 0.329 | 7 | |
| | <i>β-Tubulin</i> | 1.09 | 7 | 0.94 | 3 | 5.66 | 7 | 0.249 | 3 | |
| | <i>UbiQ</i> | 1.21 | 9 | 1.36 | 11 | 9.46 | 11 | 0.364 | 10 | |
| | <i>ArgK</i> | 1.31 | 11 | 0.68 | 2 | 7.18 | 9 | 0.297 | 5 | |
| | <i>EF-1α</i> | 0.95 | 2 | 1.08 | 5 | 2.78 | 2 | 0.228 | 1 | |
| | <i>GAPDH</i> | 0.99 | 3 | 1.24 | 10 | 3.08 | 3 | 0.336 | 8 | |
| | <i>V-ATPase-A</i> | 1.29 | 10 | 0.46 | 1 | 5.62 | 6 | 0.352 | 9 | |
| | <i>RPS7</i> | 1.06 | 6 | 1.23 | 9 | 6.34 | 8 | 0.372 | 11 | |
| <i>RPL13a</i> | 1.1 | 8 | 1.09 | 6 | 7.44 | 10 | 0.412 | 12 | | |

Table 4. Ranking of the candidate reference genes based on stability values performed by Delta Ct, BestKeeper, RefFinder, and NormFinder in tissue-specific stages.

algorithms. After different temperatures incubations (4, 27, and 37 °C), *RPS3-a*, *RPS7*, and *V-ATPase-A* were documented as the most stable reference gene from four algorithms (Table 5). The best combination genes with the lowest expression stability value (M) or highest expression stability for different temperature exposure were *RPS3-a* and *RPS7* (Fig. 5A). The top three stable genes for juvenile hormone III treatment were *EF-1α*, *RPS7*, and *UbiQ* (Table 5). Based on RefFinder comprehensive ranking, *Actin*, *RPL6* and *V-ATPase-A* (Fig. 5B) were the most stable reference genes. The most stable reference gene expressions between wild and laboratory-reared beetles were *EF-1α*, *RPS3-a*, and *RPL13a* (Table 5), whereas the top three most stable reference genes via RefFinder were *RPL13a*, *RPL7* and *RPS3-a* (Fig. 5C).

Selection of optimal reference genes for normalization. To determine the consistent and more accurate results of the optimal number of reference genes in each experimental condition, pairwise variation (V) was assessed using geNorm²⁵. The optimal number of reference genes by calculating pairing variable value $V_{n/n+1}$ with a cut-off value of 0.15, where the n + 1 reference gene is unnecessary with values under the threshold. Alternatively, the first references gene is sufficient to normalize the target gene expression in those cases (Fig. 6). For biotic conditions minimum of two genes is required for normalization as per V-value above 0.15 in $V_{2/3}$.




Figure 4. The average expression stability values (M) of twelve reference genes under different tissues calculated by geNorm demonstrating the least stable (left) to the most stable (right). (A) head, (B) fat body, (C) gut.

Validation of reference gene selection. To validate the best reference gene, the expression of *CYP450* in the developmental stages and tissue types were normalized with single reference genes and gene combinations recommended by geNorm (Fig. 6). The relative expression pattern of *CYP450* was normalized with the expression level of most stable (*RPS3-a*, *RPS3-a/RPL6*) and the least stable reference genes (*UbiQ* for gut, *RPL13a* for head tissues). The results showed that *CYP450* expression in third instar larva, pupa, and female tissues increased after normalization with the most stable reference gene alone or in combination (Fig. 7A). In compari-

| Conditions | Genes | ΔCt Method | | Best keeper | | RefFinder | | NormFinder | | Recommended Genes |
|--------------|-------------------|------------|------|-------------|-------|-----------|------|------------|------|---|
| | | Stability | Rank | Stability | Rank | Stability | Rank | Stability | Rank | |
| Temp | <i>RPS3-a</i> | 0.56 | 1 | 0.97 | 6 | 1.57 | 1 | 0.15 | 7 | <i>V-ATPase-A</i> , <i>RPS3-a</i> and <i>RPS7</i> |
| | <i>RPS7</i> | 0.6 | 2 | 1.01 | 7 | 2.74 | 2 | 0.07 | 11 | |
| | <i>UbiQ</i> | 0.61 | 3 | 1.07 | 8 | 3.83 | 4 | 0.24 | 9 | |
| | <i>V-ATPase-A</i> | 0.61 | 4 | 0.82 | 2 | 2.99 | 3 | 0.15 | 10 | |
| | <i>EF-1α</i> | 0.63 | 5 | 1.25 | 11 | 5.76 | 6 | 0.28 | 1 | |
| | <i>GAPDH</i> | 0.67 | 6 | 1.12 | 9 | 6.64 | 7 | 0.08 | 5 | |
| | <i>ArgK</i> | 0.72 | 7 | 0.57 | 1 | 4.3 | 5 | 0.14 | 3 | |
| | <i>β-Tubulin</i> | 0.73 | 8 | 1.2 | 10 | 9.16 | 11 | 0.12 | 8 | |
| | <i>RPL13a</i> | 0.76 | 9 | 0.97 | 5 | 8.19 | 10 | 0.19 | 2 | |
| | <i>Actin</i> | 0.77 | 10 | 0.82 | 3 | 7.02 | 8 | 0.19 | 6 | |
| | <i>RPL7</i> | 0.78 | 11 | 0.89 | 4 | 7.89 | 9 | 0.20 | 4 | |
| <i>RPL6</i> | 0.8 | 12 | 1.37 | 12 | 12 | 12 | 0.21 | 12 | | |
| JHIII | <i>EF-1α</i> | 0.86 | 1 | 0.76 | 6 | 2.21 | 1 | 0.37 | 1 | <i>EF-1α</i> , <i>RPS7</i> and <i>UbiQ</i> |
| | <i>RPS3-a</i> | 0.87 | 2 | 0.84 | 7 | 3.03 | 4 | 0.23 | 7 | |
| | <i>UbiQ</i> | 0.9 | 3 | 0.73 | 4 | 3.66 | 5 | 0.27 | 9 | |
| | <i>RPS7</i> | 0.91 | 4 | 0.69 | 2 | 2.38 | 2 | 0.14 | 11 | |
| | <i>RPL6</i> | 0.92 | 5 | 0.71 | 3 | 2.94 | 3 | 0.18 | 12 | |
| | <i>V-ATPase-A</i> | 0.93 | 6 | 0.97 | 9 | 6.64 | 8 | 0.22 | 10 | |
| | <i>GAPDH</i> | 0.99 | 7 | 0.73 | 5 | 6.44 | 7 | 0.33 | 5 | |
| | <i>β-Tubulin</i> | 1.08 | 8 | 0.97 | 10 | 8.71 | 9 | 0.34 | 8 | |
| | <i>ArgK</i> | 1.09 | 9 | 0.63 | 1 | 5.05 | 6 | 0.32 | 3 | |
| | <i>RPL7</i> | 1.22 | 10 | 1.02 | 11 | 10.24 | 10 | 0.37 | 4 | |
| | <i>RPL13a</i> | 1.74 | 11 | 1.23 | 12 | 11.24 | 12 | 0.48 | 2 | |
| | <i>Actin</i> | 1.78 | 12 | 0.96 | 8 | 10.84 | 11 | 0.26 | 6 | |
| Lab vs. Wild | <i>RPS3-a</i> | 0.77 | 1 | 0.89 | 7 | 2.14 | 2 | 0.12 | 7 | <i>EF-1α</i> , <i>RPS3-a</i> and <i>RPL13a</i> |
| | <i>EF-1α</i> | 0.79 | 2 | 0.81 | 4 | 2 | 1 | 0.22 | 1 | |
| | <i>RPL13a</i> | 0.81 | 3 | 0.98 | 9 | 4.82 | 5 | 0.19 | 2 | |
| | <i>UbiQ</i> | 0.83 | 4 | 0.96 | 8 | 4.43 | 4 | 0.24 | 9 | |
| | <i>V-ATPase-A</i> | 0.86 | 5 | 0.73 | 2 | 2.99 | 3 | 0.13 | 10 | |
| | <i>RPL7</i> | 0.9 | 6 | 1.06 | 10 | 6.77 | 8 | 0.16 | 4 | |
| | <i>RPS7</i> | 0.98 | 7 | 1.18 | 11 | 8.1 | 9 | 0.26 | 11 | |
| | <i>RPL6</i> | 1.02 | 8 | 1.29 | 12 | 8.49 | 10 | 0.22 | 12 | |
| | <i>GAPDH</i> | 1.05 | 9 | 0.74 | 3 | 6.18 | 7 | 0.23 | 5 | |
| | <i>β-Tubulin</i> | 1.33 | 10 | 0.68 | 1 | 5.62 | 6 | 0.17 | 8 | |
| | <i>Actin</i> | 1.41 | 11 | 0.83 | 5 | 9.03 | 11 | 0.23 | 6 | |
| <i>ArgK</i> | 1.55 | 12 | 0.85 | 6 | 10.09 | 12 | 0.27 | 3 | | |

Table 5. Under the influence of abiotic factors, the candidate reference genes were ranked based on stability values performed by Delta Ct, BestKeeper, RefFinder, and NormFinder. The abiotic factors were temperature (temp), JHIII treatment, lab-reared vs wild-collected beetles.

son, when the least stable *UbiQ* was used as a reference gene, the *CYP450* expression patterns were inconsistent across different stages (Fig. 7A). Similarly, the expression patterns of *CYP450* after normalization based on the single (*RPS3-a*) and combined (*RPS3-a/RPL6*) genes expression in gut and head tissues were showed a similar pattern (Fig. 7B,C). On the contrary, *UbiQ* expression-based normalization reduced the expression of *CYP450* from 2.9 to 0.2-fold in the gut, whereas *RPL13a* based normalization increased 4.5-fold in head tissue. Hence, our result demonstrates the importance of selecting and validating reference genes to avoid misinterpretation of gene expression results.

Discussion

RT-qPCR is one of the effective and reliable techniques for quantifying the expression of mRNA levels under different experimental conditions. However, multiple factors, such as RNA extraction, cDNA synthesis, primers, and materials handling, impact the RT-qPCR results. The reliable reference gene can overcome confounding variations in the data. Therefore, the stability of the reference gene must be evaluated for each experimental condition to accurate and reliable data interpretation^{37,42,49,52,53}. Ideally, the reference gene should have a steady and unaltered expression level during the entire experimental conditions²⁵. Hence, screening the stable reference genes in different experimental conditions before examining the target gene expression is necessary for working on any insect system.




Figure 5. The average expression stability values (M) of twelve reference genes under different conditions calculated by geNorm where the least stable (left) to the most stable (right). (A) Temperature, (B) Juvenile hormone III, (C) Laboratory-reared vs wild beetles.

Although *I. typographus* is one of the most destructive pests of Norway spruce (*Picea abies*)^{1,3,4,54}, environmental stress such as drought significantly impacts its host colonization. It can provoke transitions from endemic to an epidemic bark beetle population, suitable for the mass attack to healthy trees^{55,56}. Unravelling the




Figure 6. The optimal number of reference genes for the normalization of *I. typographus* under different experimental conditions [developmental stages, sex-specific (male and female), tissue-specific (head, midgut, fat body, temperature; Juvenile hormone III; wild-collected vs long-term laboratory-reared beetles)]. Based on geNorm analysis, average pairwise variations were calculated between the normalization factors NFn and NFn + 1. Values < 0.15 imply that n + 1 genes were unnecessary to normalise gene expression.

mechanisms underlying *I. typographus* and host interaction was a daunting task until a decade ago. However, recent advances in genomic and transcriptomic studies on *I. typographus*⁴⁴ have managed to demonstrate the power of gathering valuable genetic information in delineating the beetle-host interaction dynamics. However, it is just the beginning, and without reference genes, there is no way forward for gene expression and RNAi based functional genomics studies. Our study delivers a catalogue of reference genes for the impending genomic and functional studies on *I. typographus*.

In this study, twelve candidate reference genes were tested for the first time in the bark beetle *I. typographus* across various experimental conditions, including developmental stage, sex-specific, tissues specific, and exposure to abiotic conditions (Tables 3, 4, 5; Figs. 3, 4, 5). The results showed that all reference gene primers acquire good amplification efficiency (92.2% to 119.24%) and regression coefficient (0.944 to 0.998). Among the twelve reference genes in this study, we found that *RPS3-a*, *EF-1 α* , *RPL6*, *RPS7*, and *RPL7* were the most stable reference genes for *I. typographus* under various experimental conditions (Fig. 8).

Ribosomal proteins (RPs: *RPL*- Ribosomal protein large subunit and *RPS*- Ribosomal protein small subunit) are a large group of proteins that play a crucial role in the cellular process such as protein synthesis, cell growth, and development^{57–59}. Lee et al.⁶⁰ demonstrated that ribosomal mutation controls the cellular processes but not direct consequences of ribosome depletion. Several reports indicated that ribosomal protein was widely used as a reference gene in insect functional genomics^{31,38,61}. Not surprisingly, ribosomal protein (*RPL* and *RPS*) was consistently expressed throughout most of the experimental conditions of the insect species. For example, *RPL10* exhibited the most stable expression in different tissues, different diets conditions, and populations of *Spodoptera litura*, whereas *RPS3* was expressed most stably in larvae after starving⁶². *RPS13* and *RPS7* exhibited the most stable expression under larval-crowding conditions in *Mythimna separata*⁶³. Similarly, different developmental stages (*RPL32*, *RPS18*) and different tissues (*RPS18*) of *Spodoptera frugiperda* showed the most stable expression⁶⁴, whereas *RPS26* and *RPL32* genes showed the same in *Thermobia domestica*⁶⁵; *RPS11*, *RPL28*, and *RPL10* genes showed *Tuta absoluta*⁵⁸; *RPS18* and *RPL13* genes showed *Rhopalosiphum padi* tissues⁶⁶. Sellamuthu et al.⁴³ reported that *RPS3* is stably expressed in the developmental stage, sex-specific and tissue-specific conditions in pine-feeding bark beetle, *I. sexdentatus*. Furthermore, *RPL36* displayed stable expression in the developmental stage, sex-specific and tissue-specific conditions of *Glenea cantor*⁶⁷. Developmental stage and tissue-specific expression of *RPS18* and *RPS11* were highly stable in *Cicindela campestris*³⁶. Similarly, *RP49* expression was the most stable in tissue and sex-specific conditions in *Harmonia axyridis*³³. Wang et al.⁶⁸ reported that *RPL22e* was stably expressed in sex-specific conditions of *Mytilus cichorii*. *RPS23* and *RPL12* were identified as reference genes in *Anthonomus eugenii*²². Similarly, *RPL32* gene was found as the most stable gene in adult (male and female) tissues of *Anoplophora glabripennis*⁶⁹. It was reported that *RPL13A* ribosomal protein was more stable expression under low-temperature treatments in *Thitarodes armoricanus*⁷⁰ and *Paederus fuscipes*⁷¹. Other ribosomal proteins such as *RPL27* and *RPS7*, *RPL7* of *S. frugiperda*, were the most stable reference genes under low-temperature and high-temperature⁶⁴. Tao et al.⁷² showed that *RPL-33* and *RPS-26* are the most stable




Figure 7. The relative expression levels of the target gene *CYP450* were studied under developmental and various tissues after normalizing with the least and most suitable reference genes separately or in combination. The most stable reference gene *RPS3-a* and *RPL6* for developmental stages, head, and gut tissue. The least stable reference gene *UbiQ* was used for both developmental and gut. For head tissue, *RPL13a* was used as the least stable reference gene. (A) Developmental stage, (B) Gut. (C) Head. Data represent mean values \pm SD of four biological replicates. Asterisks indicate significant differences in the expression of the target gene normalized separately by different reference genes (**** $P < 0.0001$ *** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$, ns indicate no significant difference).

reference genes from microarray data and *RPS-2* and *RPS-4* from RNA-seq of *Caenorhabditis elegans* out of thirteen ribosomal proteins. Moreover, *RPS20* was detected as the least stably expressed gene for analyzing *Plutella xylostella* under different conditions⁷³. Our results also demonstrated stable ribosomal gene expression in biotic and abiotic conditions. According to developmental, sex-specific, and tissue-specific stages, *RPS3-a*, *RPL7*, and *RPS6* showed higher expression stability. Furthermore, *RPS3-a* and *RPS7* were stably expressed as second top rank in various abiotic conditions. These results also suggest that no single reference gene for different experimental conditions is observed in the present study.

Elongation factor 1 (EF-1) plays a central role that promotes the delivery of aminoacyl-tRNA to the acceptor site of the ribosome during protein biosynthesis^{32,74}. *EF-1* has been a stable reference gene for many years and




Figure 8. Recommended reference genes under different experimental conditions for *I. typographus*. 1. Developmental stages (or life stages). 2. Temperature treatment. 3. JHIII treatment. 4. Wild /lab-reared beetles/ pheromone traps. 5. Gut tissues. 6. Fat body tissues 7. Head tissues. 8. Female tissues. 9. Male tissues.

is widely used for reference genes in many insects^{61–63,70}. Our results exhibited that *elongation factor 1* was the second most stable expression in biotic and abiotic conditions. Similarly, Su et al.⁶⁷ reported that *EF-1* was the most appropriate reference gene for all samples of *Glenaea cantor*. Teng et al.⁷⁵ reported *elongation factor2* (*eEF2*) as the most stably expressed gene in different developmental stages of *Plutella xylostella*. The expression stability of *eEF2* in developmental stages and tissues was also documented in *Agrilus planipennis*⁷⁶; *I. sexdentatus*⁴³; *Sogatella furcifera*⁷⁷; *Mythimna separata*^{63,78}; *Tuta absoluta*⁶¹; *Hippodamia variegata*⁴²; *Spodoptera frugiperda*⁶⁴. However, Ponton et al.³² reported that the elongation factor was unsuitable for normalization of relative expression due to the different expression variability in different experimental conditions in *Drosophila melanogaster*. Nevertheless, our results confirmed that *EF-1α* had stable expression between JHIII treated vs control bark beetles, wild vs lab-reared beetles, and within gut tissues, male tissues, and developmental stages.

To further validate our findings and confirm the stable reference gene, we analyzed the expression of cytochrome P450 (*CYP03903*) in the developmental stages, head, and gut tissues of the adult beetle. P450s is a large class of enzymes that often plays an essential role in detoxifying xenobiotics⁷⁹. It evolves during the insect-host interaction to metabolize a wide range of plant allelochemicals⁸⁰. Our results demonstrated the expression trends in developmental and tissue-specific conditions using single reference gene or gene combinations. The *CYP03903* transcript was increased in pupa and female tissue compared to other stages. However, the same expression profile of *CYP03903* significantly differs when normalized with the least stable reference gene showing the importance of having a suitable reference gene. For instance, the expression of *CYP450* demonstrated just the opposite pattern based on the most or least stable gene was used for normalization. Nevertheless, the *CYP03903* expression profile between the gut and head tissues is different, suggesting different functions in the respective tissues^{79,81}.

To summarise the present study, twelve candidate reference genes of *I. typographus* were selected and systematically evaluated for their expression stability using four widely used software programs under different experimental conditions to obtain the best reference genes for each condition. The results showed that the most suitable candidate reference genes were *RPS3-a*, *EF-1α*, *RPS7*, *RPL7*, and *RPL6* under different experimental

conditions. Based on our comprehensive analysis, we recommended a list of reference genes and combinations of reference genes be used to normalize gene expression in *I. typographus* subjected to different experimental conditions. This is a much-awaited reference gene validation work on *I. typographus*, setting the foundation for future molecular (i.e., gene expression) and functional genomics studies (i.e., RNAi). The same genes can be further evaluated to identify suitable reference genes for other *Ips* species (Coleoptera: Curculionidae: Scolytinae).

Data availability

The raw data supporting the conclusions of this manuscript will be made available by the authors.

Received: 30 September 2021; Accepted: 7 March 2022

Published online: 18 March 2022

References

1. Wermelinger, B. Ecology and management of the spruce bark beetle *Ips typographus*: A review of recent research. *For. Ecol. Manag.* **202**(1–3), 67–82. <https://doi.org/10.1016/j.foreco.2004.07.018> (2004).
2. Sun, X. L., Yang, Q. Y., Sweeney, J. D. & Gao, C. G. A review: Chemical ecology of *Ips typographus* (Coleoptera, Scolytidae). *J. For. Res.* **17**, 65–70. <https://doi.org/10.1007/s11676-006-0016-2> (2006).
3. Raffa, K. F., Grégoire, J. C. & Lindgren, B. S. Natural history and ecology of bark beetles. In *Bark Beetles: Biology and Ecology of Native and Invasive Species* (eds Vega, F. E. & Hofstetter, R. W.) 1–40 (Academic Press, 2015). <https://doi.org/10.1016/B978-0-12-417156-5.00001-0>.
4. Biedermann, P. et al. Bark beetle population dynamics in the anthropocene: Challenges and solutions. *Trends Ecol. Evol.* **34**, 914–924. <https://doi.org/10.1016/j.tree.2019.06.002> (2019).
5. Tanin, S. M., Kandasamy, D. & Krokene, P. Fungal interactions and host tree preferences in the spruce bark beetle *Ips typographus*. *Front. Microbiol.* **12**, 695167. <https://doi.org/10.3389/fmicb.2021.695167> (2021).
6. Sambaraju, K. R. & Côté, C. Are climates in Canada and the United States suitable for the European Spruce Bark Beetle, *Ips typographus*, and Its Fungal Associate, *Endoconidiophora polonica*? *Forests* **12**(12), 1725. <https://doi.org/10.3390/f12121725> (2021).
7. Schelhaas, M. J., Nabuurs, G. J. & Schuck, A. Natural disturbances in the European forests in the 19th and 20th centuries. *Glob. Change Biol.* **9**, 1620–1633 (2003).
8. Seidl, R., Rammer, W. & Lexer, M. J. Adaptation options to reduce climate change vulnerability of sustainable forest management in the Austrian Alps. *Can. J. For. Res.* **41**, 694–706 (2011).
9. Bentz, B. J. et al. *Ips typographus* and *Dendroctonus ponderosae* models project thermal suitability for intra- and inter-continental establishment in a changing climate. *Front. For. Glob.* **2**, 1 (2019).
10. Knížek, M. Historie a současnost kůrovcových kalamit ve střední Evropě. Zpravodaj ochrany lesa, Škodliví činitelé v lesích Česka, Svazek, 23 (2020). https://www.vulhm.cz/files/uploads/2020/10/ZOL_23_2020.pdf.
11. Müller, J., Bußler, H., Goßner, M., Rettelbach, T. & Duelli, P. The European spruce bark beetle *Ips typographus* in a national park: From pest to keystone species. *Biodivers. Conserv.* **17**, 2979. <https://doi.org/10.1007/s10531-008-9409-1> (2008).
12. Hroščo, B. et al. Drivers of Spruce Bark Beetle (*Ips typographus*) Infestations on downed trees after severe windthrow. *Forests* **11**, 1290. <https://doi.org/10.3390/f11121290> (2020).
13. Seidl, R., Rammer, W., Jäger, D. & Lexer, M. J. Impact of bark beetle (*Ips typographus* L.) disturbance on timber production and carbon sequestration in different management strategies under climate change. *For. Ecol. Manag.* **256**, 209–220. <https://doi.org/10.1016/j.foreco.2008.04.002> (2008).
14. Jönsson, A. M., Appelberg, G., Harding, S. & Bähring, L. Spatio-temporal impact of climate change on the activity and voltinism of the spruce bark beetle, *Ips typographus*. *Glob. Change Biol.* **15**, 486–499. <https://doi.org/10.1111/j.1365-2486.2008.01742.x> (2009).
15. Ghimire, R. P. et al. Effect of bark beetle (*Ips typographus* L.) attack on bark VOC emissions of Norway spruce (*Picea abies* Karst.) trees. *Atmos. Environ.* **126**, 145–152. <https://doi.org/10.1016/j.atmosenv.2015.11.049> (2016).
16. Grodzki, W. et al. Effects of intensive versus no management strategies during an outbreak of the bark beetle *Ips typographus* (L.) (Col.: Curculionidae, Scolytinae) in the Tatra Mts. in Poland and Slovakia. *Ann. For. Sci.* **63**, 55–61 (2006).
17. Joga, M. R., Mogilicherla, K., Smaghe, G. & Roy, A. RNA interference-based forest protection products (FPPs) against wood-boring coleopterans: Hope or hype? *Front. Plant Sci.* **12**, 733608. <https://doi.org/10.3389/fpls.2021.733608> (2021).
18. Bustin, S. A. et al. The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.* **55**(4), 611–622. <https://doi.org/10.1373/clinchem.2008.112797> (2009).
19. Ginzinger, D. G. Gene quantification using real-time quantitative PCR: An emerging technology hits the mainstream. *Exp. Hematol.* **30**(6), 503–512. [https://doi.org/10.1016/s0301-472x\(02\)00806-8](https://doi.org/10.1016/s0301-472x(02)00806-8) (2002).
20. Bustin, S. A. Quantification of mRNA using real-time reverse transcription PCR (RT-PCR): Trends and problems. *J. Mol. Endocrinol.* **29**, 23–39. <https://doi.org/10.1677/jme.0.0290023> (2002).
21. Bustin, S. A. & Nolan, T. Pitfalls of quantitative real-time reverse-transcription polymerase chain reaction. *J. Biomol. Tech.* **15**, 155–166 (2004).
22. Pinheiro, D. H. & Siegfried, B. D. Selection of reference genes for normalization of RT-qPCR data in gene expression studies in *Anthonomus eugenii* Cano (Coleoptera: Curculionidae). *Sci. Rep.* **10**, 5070. <https://doi.org/10.1038/s41598-020-61739-z> (2020).
23. Wang, Z. et al. Identification and evaluation of reference genes for normalization of gene expression in developmental stages, sexes, and tissues of *Diaphania caesalis* (Lepidoptera, Pyralidae). *J. Insect Sci.* **20**, 1. <https://doi.org/10.1093/jisesa/iez130> (2020).
24. Higuchi, R., Fockler, C., Dollinger, G. & Watson, R. Kinetic PCR analysis: Real-time monitoring of DNA amplification reactions. *Biotechnology (N. Y.)* **11**, 1026–1030. <https://doi.org/10.1038/nbt0993-1026> (1993).
25. Vandesompele, J. et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* **3**, 1–11. <https://doi.org/10.1186/gb-2002-3-7-research0034> (2002).
26. Bustin, S. A. et al. MIQE précis: Practical implementation of minimum standard guidelines for fluorescence-based quantitative real-time PCR experiments. *BMC Mol. Biol.* **11**, 74. <https://doi.org/10.1186/1471-2199-11-74> (2010).
27. Bustin, S. A. et al. Primer sequence disclosure: A clarification of the MIQE guidelines. *Clin. Chem.* **57**(6), 919–921. <https://doi.org/10.1373/clinchem.2011.162958> (2011).
28. Yeung, N., Botvinick, M. M. & Cohen, J. D. The neural basis of error detection: Conflict monitoring and the error-related negativity. *Psychol. Rev.* **111**(4), 931–959. <https://doi.org/10.1037/0033-295x.111.4.939> (2004).
29. Bustin, S. A., Benes, V., Nolan, T. & Pfaffl, M. W. Quantitative real-time RT-PCR—A perspective. *J. Mol. Endocrinol.* **34**, 597–601. <https://doi.org/10.1677/jme.1.01755> (2005).
30. Strube, C., Buschbaum, S., Wolken, S. & Schnieder, T. Evaluation of reference genes for quantitative real-time PCR to investigate protein disulfide isomerase transcription pattern in the bovine lungworm *Dictyocaulus viviparus*. *Gene* **425**, 36–43. <https://doi.org/10.1016/j.gene.2008.08.001> (2008).
31. Gao, P., Wang, J. & Wen, J. Selection of reference genes for tissue/organ samples of adults of *Eucryptorrhynchus scrobiculatus*. *PLoS ONE* **15**(2), e0228308. <https://doi.org/10.1371/journal.pone.0228308> (2020).

32. Ponton, F., Chapuis, M. P., Pernice, M., Sword, G. A. & Simpson, S. J. Evaluation of potential reference genes for reverse transcription-qPCR studies of physiological responses in *Drosophila melanogaster*. *J. Insect Physiol.* **57**(6), 840–850. <https://doi.org/10.1016/j.jinsphys.2011.03.014> (2011).
33. Yang, X. W., Pan, H. P., Yuan, L. & Zhou, X. Reference gene selection for RT-qPCR analysis in *Harmonia axyridis*, a global invasive lady beetle. *Sci. Rep.* **8**, 2689. <https://doi.org/10.1038/s41598-018-20612-w> (2018).
34. Koramutla, M. K., Aminedi, R. & Bhattacharya, R. Comprehensive evaluation of candidate reference genes for qRT-PCR studies of gene expression in mustard aphid, *Lipaphis erysimi* (Kalt). *Sci. Rep.* **6**, 25883. <https://doi.org/10.1038/srep25883> (2016).
35. Arya, S. K. *et al.* Reference genes validation in *Phenacoccus solenopsis* under various biotic and abiotic stress conditions. *Sci. Rep.* **7**, 13520. <https://doi.org/10.1038/s41598-017-13925-9> (2017).
36. García-Reina, A., Rodríguez-García, M. J. & Galián, J. Validation of reference genes for quantitative real-time PCR in tiger beetles across sexes, body parts, sexual maturity and immune challenge. *Sci. Rep.* **8**, 10743. <https://doi.org/10.1038/s41598-018-28978-7> (2018).
37. Shakeel, M., Rodríguez, A., Tahir, U. B. & Jin, F. Gene expression studies of reference genes for quantitative real-time PCR: An overview in insects. *Biotechnol. Lett.* **40**, 227–236. <https://doi.org/10.1007/s10529-017-2465-4> (2018).
38. Lu, J., Yang, C., Zhang, Y. & Pan, H. Selection of reference genes for the normalization of RT-qPCR data in gene expression studies in insects: A systematic review. *Front. Physiol.* **9**, 1560. <https://doi.org/10.3389/fphys.2018.01560> (2018).
39. Pfaffl, M. W., Tichopad, A., Prgomet, C. & Neuvians, T. P. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper-Excel-based tool using pairwise correlations. *Biotechnol. Lett.* **26**, 509–515. <https://doi.org/10.1023/b:bile.0000019559.84305.47> (2004).
40. Qu, C. *et al.* Selection and evaluation of reference genes for expression analysis using quantitative real-time PCR in the Asian Ladybird *Harmonia axyridis* (Coleoptera: Coccinellidae). *PLoS ONE* **13**(6), e0192521. <https://doi.org/10.1371/journal.pone.0192521> (2018).
41. Basu, S. *et al.* Evaluation of reference genes for real-time quantitative PCR analysis in southern corn rootworm, *Diabrotica undecimpunctata howardi* (Barber). *Sci. Rep.* **9**, 10703. <https://doi.org/10.1038/s41598-019-47020-y> (2019).
42. Xie, J. *et al.* Identification and evaluation of suitable reference genes for RT-qPCR analysis in *Hippodamia variegata* (Coleoptera: Coccinellidae) under different biotic and abiotic conditions. *Front. Physiol.* **12**, 669510. <https://doi.org/10.3389/fphys.2021.669510> (2021).
43. Sellamuthu, G. *et al.* Reference gene selection for normalizing gene expression in *Ips sexdentatus* (Coleoptera: Curculionidae: Scolytinae) under different experimental conditions. *Front. Physiol.* <https://doi.org/10.3389/fphys.2021.752768> (2021).
44. Powell, D. *et al.* A highly-contiguous genome assembly of the Eurasian spruce bark beetle, *Ips typographus*, provides insight into a major forest pest. *Commun. Biol.* **4**, 1059. <https://doi.org/10.1038/s42003-021-02602-3> (2021).
45. Remes, J. & Podrázský, V. Fertilization of spruce monocultures on the School Training Forest territory in Kostelec nad Černými lesy. *J. For. Sci.* **52**(1), 73–78 (2006).
46. Hlásný, T. *et al.* Devastating outbreak of bark beetles in the Czech Republic: Drivers, impacts, and management implications. *For. Ecol. Manag.* **490**, 119075 (2021).
47. Hall, G. M. *et al.* Midgut tissue of male pine engraver, *Ips pini*, synthesizes monoterpene component ipsdienol de novo. *Naturwissenschaften* **89**, 79–83 (2002).
48. Sun, Y. *et al.* Function of mevalonate pathway genes in the synthesis of frontalin in Chinese white pine beetle, *Dendroctonus armandi* (curculionidae: Scolytinae). *Arch. Insect Biochem. Physiol.* **107**(4), 1–13. <https://doi.org/10.1002/arch.21828> (2021).
49. Andersen, C. L., Jensen, J. L. & Orntoft, T. F. Normalization of real-time quantitative reverse transcription-PCR data: A model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res.* **64**, 5245–5250. <https://doi.org/10.1158/0008-5472.can-04-0496> (2004).
50. Silver, N., Best, S., Jiang, J. & Thein, S. L. Selection of housekeeping genes for gene expression studies in human reticulocytes using real-time PCR. *BMC Mol. Biol.* **6**, 33. <https://doi.org/10.1186/1471-2199-7-33> (2006).
51. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔC_T} method. *Methods* **25**, 402–408. <https://doi.org/10.1006/meth.2001.1262> (2001).
52. Tellin, O. *et al.* Housekeeping genes as internal standards: Use and limits. *J. Biotechnol.* **75**, 291–295. [https://doi.org/10.1016/S0168-1656\(99\)00163-7](https://doi.org/10.1016/S0168-1656(99)00163-7) (1999).
53. Huggett, J., Dheda, K., Bustin, S. & Zumla, A. Real-time RT-PCR normalisation; strategies and considerations. *Genes Immun.* **6**, 279–284. <https://doi.org/10.1038/sj.gene.6364190> (2005).
54. Ryan, M. G., Sapes, G., Sala, A. & Hood, S. M. Tree physiology and bark beetles. *New Phytol.* **205**, 955–957 (2015).
55. Kausrud, K. *et al.* Population dynamics in changing environments: The case of an eruptive forest pest species. *Biol. Rev.* **87**, 34–51. <https://doi.org/10.1111/j.1469185X.2011.00183.x> (2012).
56. Fettig C. J., *et al.* Bark Beetle Management, Ecology, and Climate Change (ed. Gandhi K. J. K & Hofstetter, R. W). 345–394, Academic Press (2021)
57. Zhou, X., Liao, W. J., Liao, J. M., Liao, P. & Lu, H. Ribosomal proteins: Functions beyond the ribosome. *J. Mol. Cell Biol.* **7**(2), 92–104. <https://doi.org/10.1093/jmcb/mjv014> (2015).
58. Saha, A. *et al.* Genome-wide identification and comprehensive expression profiling of ribosomal protein small subunit (RPS) genes and their comparative analysis with the large subunit (RPL) genes in rice. *Front. Plant Sci.* **8**, 1553. <https://doi.org/10.3389/fpls.2017.01553> (2017).
59. Petibon, C., Malik, G. M., Catala, M. & Abou, E. S. Regulation of ribosomal protein genes: An ordered anarchy. *Wiley Interdiscip. Rev. RNA* **12**(3), 1632. <https://doi.org/10.1002/wrna.1632> (2021).
60. Lee, C. H. *et al.* A regulatory response to ribosomal protein mutations controls translation, growth, and cell competition. *Develop. Cell* **46**(4), 456–469. <https://doi.org/10.1016/j.devcel.2018.07.003> (2018).
61. Yan, X., Zhang, Y., Xu, K., Wang, Y. & Yang, W. Selection and validation of reference genes for gene expression analysis in *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae). *Insects* **12**, 589. <https://doi.org/10.3390/insects12070589> (2021).
62. Lu, Y. *et al.* Identification and validation of reference genes for gene expression analysis using quantitative PCR in *Spodoptera litura* (Lepidoptera: Noctuidae). *PLoS ONE* **8**, e68059. <https://doi.org/10.1371/journal.pone.0068059> (2013).
63. Li, H. B. *et al.* Screening potential reference genes for quantitative real-time PCR analysis in the oriental armyworm, *Mythimna separata*. *PLoS ONE* **13**(4), e0195096. <https://doi.org/10.1371/journal.pone.0195096> (2018).
64. Zhou, L., Meng, J. Y., Ruan, H. Y., Yang, C. L. & Zhang, C. Y. Expression stability of candidate RT-qPCR housekeeping genes in *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Arch. Insect Biochem. Physiol.* **108**, e21831 (2021).
65. Bai, Y. *et al.* Selection of reference genes for normalization of gene expression in *Thermobia domestica* (Insecta: Zygentoma: Lepismatidae). *Genes* **12**(1), 21. <https://doi.org/10.3390/genes12010021> (2021).
66. Li, M. *et al.* Selection and validation of reference genes for qRT-PCR analysis of *Rhopalosiphum padi* (Hemiptera: Aphididae). *Front. Physiol.* **12**, 663338. <https://doi.org/10.3389/fphys.2021.663338> (2021).
67. Su, R. R. *et al.* Evaluation of reference genes in *Glenea cantor* (Fabricius) by using qRT-PCR. *Genes* **12**, 1984. <https://doi.org/10.3390/genes12121984> (2021).
68. Wang, Y., Wang, Z. K., Huang, Y., Liao, Y. F. & Yin, Y. P. Identification of suitable reference genes for gene expression studies by qRT-PCR in the blister beetle *Mylabris cichorii*. *J. Insect Sci.* **14**, 94. <https://doi.org/10.1093/jis/14.1.94> (2014).

69. Rodrigues, T. B., Dhandapani, R. K., Duan, J. J. & Palli, S. R. RNA interference in the Asian Longhorned Beetle: Identification of key RNAi genes and reference genes for RT-qPCR. *Sci. Rep.* **7**(1), 8913. <https://doi.org/10.1038/s41598-017-08813-1> (2017).
70. Liu, G. *et al.* Evaluation of reference genes for reverse transcription quantitative PCR studies of physiological responses in the ghost moth, *Thitarodes armoricanus* (Lepidoptera, Hepialidae). *PLoS ONE* **11**(7), e0159060. <https://doi.org/10.1371/journal.pone.0159060> (2016).
71. Khan, M. M. *et al.* Screening and validation of reference genes using in RT-qPCR for gene expression studies in *Paederus fuscipes*, a medically and agriculturally important insect. *J. King Saud. Univ. Sci.* **1**, 101654. <https://doi.org/10.1016/j.jksus.2021.101654> (2022).
72. Tao, J. *et al.* Systematic identification of housekeeping genes possibly used as references in *Caenorhabditis elegans* by large-scale data integration. *Cells* **9**(3), 786. <https://doi.org/10.3390/cells9030786> (2020).
73. Fu, W. *et al.* Exploring valid reference genes for quantitative real-time PCR analysis in *Plutella xylostella* (Lepidoptera: Plutellidae). *Int. J. Biol. Sci.* **9**(8), 792–802. <https://doi.org/10.7150/ijbs.5862> (2013).
74. Negrutskii, B. S. & El'skaya, A. V. Eukaryotic translation elongation factor 1 alpha: Structure, expression, functions, and possible role in aminoacyl-tRNA channeling. *Prog. Nucleic Acid Res. Mol. Biol.* **60**, 47–78. [https://doi.org/10.1016/s0079-6603\(08\)60889-2](https://doi.org/10.1016/s0079-6603(08)60889-2) (1998).
75. Teng, X., Zhang, Z., He, G., Yang, L. & Li, F. Validation of reference genes for quantitative expression analysis by real-time RT-PCR in four Lepidopteran insects. *J. Insect Sci.* **12**, 1–17. <https://doi.org/10.1673/031.012.6001> (2012).
76. Rajarapu, S. P., Mamidala, P. & Mittapalli, O. Validation of reference genes for gene expression studies in the emerald ash borer (*Agrilus planipennis*). *Insect Sci.* **19**, 41–46. <https://doi.org/10.1111/j.1744-7917.2011.01447.x> (2012).
77. An, X. K., Hou, M. L. & Liu, Y. D. Reference gene selection and evaluation for gene expression studies using qRT-PCR in the white-backed Planthopper, *Sogatella furcifera* (Hemiptera: Delphacidae). *J. Econ. Entomol.* **109**, 879. <https://doi.org/10.1093/jee/tov333> (2016).
78. Li, K., Xu, N., Yang, Y. J., Zhang, J. H. & Yin, H. Identification and validation of reference genes for RT-qPCR normalization in *Mythimna separata* (Lepidoptera: Noctuidae). *Biomed. Res. Int.* <https://doi.org/10.1155/2018/1828253> (2018).
79. Zhang, H., Zhao, M., Liu, Y., Zhou, Z. & Guo, J. Identification of cytochrome P450 monooxygenase genes and their expression in response to high temperature in the alligator weed flea beetle *Agasicles hygrophila* (Coleoptera: Chrysomelidae). *Sci. Rep.* **8**, 17847 (2018).
80. Peng, L. *et al.* Functional study of cytochrome P450 enzymes from the brown planthopper (*Nilaparvata lugens* Stål) to analyze its adaptation to BPH-resistant rice. *Front. Physiol.* **8**, 972. <https://doi.org/10.3389/fphys.2017.00972> (2017).
81. Chung, H. *et al.* Characterization of *Drosophila melanogaster* cytochrome P450 genes. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 5731–5736. <https://doi.org/10.1073/pnas.0812141106> (2009).

Acknowledgements

The project is funded by the "EXTEMIT-K" and Internal Grant Agency (IGA) from the Faculty of Forestry and Wood Sciences, Czech University of Life sciences. Infrastructural support and salary for GS, JB, MRJ and AR are obtained from grant "EXTEMIT-K," No. CZ.02.1.01/0.0/0.0/15_003/0000433 financed by OP RDE. Additional financial support for AR, JS is provided by grant "EVA 4.0," No. CZ.02.1.01/0.0/0.0/16_019/0000803 financed by OP RDE. We acknowledge Dr Roman Modlinger, FLD, CZU, for his input about the bark beetle sample collection site in Czechia. Last but not least, the authors would also acknowledge the anonymous peer reviewers who gave valuable comments on the earlier version of the manuscript.

Author contributions

G.S., J.B., and A.R. conceived and designed the research. J.S. collected, sorted, and dissected beetles. G.S., J.B. and M.R.J. conducted real-time experiments. G.S. and M.R.J. analysed the data. G.S. wrote the first draft. G.S., A.R. prepared the figures and final manuscript. All authors have read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-08434-3>.

Correspondence and requests for materials should be addressed to A.R.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022