

Research Article

Evaluation of Traumatic Subdural Hematoma Volume by Using Image Segmentation Assessment Based on Deep Learning

Dan Chen ¹, Lin Bian ², Hao-Yuan He ¹, Ya-Dong Li ¹, Chao Ma ¹,
and Lian-Gang Mao ³

¹Department of Neurosurgery, The Third People's Hospital of Hefei, Hefei 230022, China

²Department of Neurosurgery, HwaMei Hospital, University of Chinese Academy of Sciences, Ningbo Institute of Life and Health Industry, University of Chinese Academy of Sciences, Ningbo 315010, China

³Department of Clinical Laboratory, HwaMei Hospital, University of Chinese Academy of Sciences, Ningbo 315010, China

Correspondence should be addressed to Lin Bian; anthony_bian@126.com and Lian-Gang Mao; bianlin1995@126.com

Received 19 April 2022; Revised 31 May 2022; Accepted 9 June 2022; Published 28 June 2022

Academic Editor: Yao Chen

Copyright © 2022 Dan Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rapid and accurate evaluations of hematoma volume can guide the treatment of traumatic subdural hematoma. We aim to explore the consistency between the measurement results of traumatic subdural hematoma (TSDH) using a deep learn-based image segmentation algorithm. A retrospective study was conducted on 90 CT images of patients diagnosed with TSDH in our hospital from January 2019 to January 2022. All image data were measured by manual segmentation, convolutional neural networks (CNN) algorithm segmentation, and $ABC/2$ volume formula. With manual segmentation as the “golden standard,” a consistency test was carried out with CNN algorithm segmentation and $ABC/2$ volume formula, respectively. The percentage error of CNN algorithm segmentation is less than $ABC/2$ volume formula. There is no significant difference between CNN algorithm segmentation and manual segmentation ($P > 0.05$). The area under curve of the $ABC/2$ volume formula, manual segmentation, and CNN algorithm segmentation is 0.811 (95% CI: 0.717–0.905), 0.840 (95% CI: 0.753–0.928), and 0.832 (95% CI: 0.742–0.922), respectively. From our results, the algorithm based on CNN has a good efficiency in segmentation and accurate calculation of TSDH hematoma volume.

1. Introduction

Traumatic subdural hematoma (TSDH) is a common disease in neurosurgery [1]. The pathogenesis of this disease is mainly that the patient's head is violently hit, which causes the dura mater of his brain tissue to shift. Eventually, the bridging vein between the cerebral cortex and venous sinus was broken, which led to bleeding [2, 3]. At present, patients with this disease are mainly clinically treated by surgery or conservative treatment [4, 5]. After the onset of TSDH, patients will have symptoms such as increased intracranial pressure, coma, cerebral hernia, and hemiplegia. Some patients will have dilated pupils, a severe threat to their physical and mental health and life safety [6]. Therefore, accurate calculation of hematoma volume for patients has important guiding significance for later treatment and prognosis.

Previous studies have found that for TSDH, drainage of only 20% of hemorrhage can effectively help hematoma absorption [7]. There are many methods for measuring and calculating different types of intracranial hemorrhage. The traditional manual segmentation method is the “gold standard” for calculating hematoma volume, but it is time-consuming and laborious [8, 9]. Another conventional method is Tada formula, namely, $A \times B \times C/2$ volume formula [10]. Although this method is suitable for subdural hematoma, the shape of subdural hematoma is not all symmetrical crescent, especially since many chronic subdural hematomas are comma-like [11]. The accuracy of the calculation results is controversial [12, 13].

With the development of computer technology such as deep learning, more and more scholars began to try to calculate the hematoma volume by using related technologies

[14–16]. Yu et al. [17] constructed a deep learning algorithm covering all types of intracerebral hemorrhage hematoma volume. This algorithm has high accuracy and consistency with manual segmentation estimation of cerebral parenchymal hemorrhage hematoma volume. However, there is no specific algorithm for subdural hematoma at present. This study is aimed at proposing a segmentation method based on the CNN algorithm and compare it with Tada formula and manual segmentation to explore an accurate and convenient measurement method for subdural hematoma volume.

2. Methods

2.1. General Information. A total of 90 patients with TSDH diagnosed by CT in our hospital from January 2019 to January 2022 were collected retrospectively. There were 67 males and 23 females, aged from 23 to 80 years, with an average of (56.5 ± 13.9) years.

Inclusion criteria are (1) select CT images of the head of patients with subdural hematoma. (2) All of them were obtained before or without surgery. (3) All of them are in standard DICOM format, and the scanning parameters and machine brands are not limited. (4) Age > 18 years old. Exclusion criteria were patients with bilateral subdural hematoma.

2.2. Manual Segmentation. DICOM-formatted CT scan images of each patient were imported into Insight Toolkit SNAP open source software. Using manual segmentation, two radiologists manually sketched a three-dimensional region of interest (ROI) layer by layer along the hematoma boundary on the cross-sectional CT plain scan image and output the hematoma volume.

2.3. Tada Volume Formula. Tada volume formula measurement regards cerebral hemorrhage focus as an ellipsoid. A physician calculated the blood loss with 3 years of experiences in neuroimaging diagnosis. The specific calculation formula is

$$V(\text{bleeding}) = A \times B \times C \times \frac{1}{2}. \quad (1)$$

Tada volume formula can be expressed by $ABC/2$ volume formula [18]. The detailed calculation process is that A is the largest length (anterior to posterior) of the SDH, B is the maximum width (lateral to midline) 90° to A , and C is the maximum height (coronal plane or multiplication of slices) of the hematoma.

2.4. Deep Learning Segmentation

2.4.1. Construction Model. The Deep Red AI Toolbox generates the deep learning model in this research. To reduce the overfitting risk of the segmentation result of the construction algorithm and improve the robustness, the CT images of the included cases were verified by 5-fold crossover method [19]. According to the ratio of training set to test set of 4:1, 90 patients were divided into training set ($n = 72$) and test set ($n = 18$). Four groups were selected as the training set, the




FIGURE 1: Convolution neural network modeling flow chart.

remaining one group was selected as the verification set, and five training times were conducted. Summarize the results of five models in their respective verification sets to evaluate the performance of algorithm models (Figure 1).

2.4.2. Radiomics Feature Extraction. The convolutional neural network (CNN) has good feature extraction performance, so this paper chooses a convolutional neural network to extract image features to get the best image fusion result. Firstly, the CT image is preprocessed by fusion, and the hematoma area on the image is automatically detected and segmented by CNN algorithm (Figure 2). The software automatically measures the bleeding volume and records the measurement results.

The image is deaveraged when the CNN algorithm analyzes the image, as shown in the following formula.

$$a^* = a - \lambda. \quad (2)$$

The λ is the mean value of the image set. a is the characteristic value.

A complete CNN is a multilayer structure, including input layer, pooling layer, full connection layer, and convolution layer. Convolution layer is the most critical part, which contains many neural network nodes and can extract the features of image fusion. The operation of convolution layer can be described as the following formula.

$$a_j^l == e \left(\sum_i \in N_j a_j^{l-1} \times h_{ij}^l + d_j^l \right). \quad (3)$$

Because of the large number of image features, it is necessary to reduce the dimension of image fusion features. This step is mainly realized through the pooling layer, and the optimal image fusion features are identified through the maximum pooling function, which can be described as the following formula.




FIGURE 2: Region of interest intercepted by convolutional neural network model. (a) Origin image. (b) Image clipping. (c) Extraction of eigenvalues of the ROI. (d) Extracted hematoma.




FIGURE 3: Image segmentation flow chart.

$$a_j^l = \text{down}(a_j^{l-1}). \quad (4)$$

l represents the number of feature layers. a_j^{l-1} represents the j th feature of the $l-1$ layer.

In the training process, the segmentation accuracy on the loss function and verification set is checked in real time, the convergence speed and trend are judged, and the network parameters are adjusted. Satisfactory model is obtained in training set, and algorithm segmentation result is obtained in test set.

2.5. Measurement Error of Segmentation Method. The CT image data included in this study used manual image segmentation method, ABC/2 volume formula measurement method, and the CNN algorithm image segmentation method to measure hematoma volume. The results measured by CT manual segmentation method are “gold standard” [20]. The specific process is shown in Figure 3. In

addition, the absolute percentage error (APE) is used to represent the measurement error, and the specific calculation formula is

$$\text{APE} = \frac{|V_m - V_r|}{V_r} \times 100\%. \quad (5)$$

V_m and V_r represent the measured and actual value of bleeding volume, respectively.

2.6. Statistical Analysis. SPSS 23.0 statistical software was used for data processing and analysis. Measurement data is expressed by mean \pm standard deviation. Group comparison was performed by independent sample t test. Measurement data with nonnormal distribution are expressed by median and quartile interval $[M(P_{25}, P_{75})]$, and the Mann-Whitney U test is used. $P \leq 0.05$ indicates that the difference is statistically significant.

TABLE 1: Volume and percentage error of subdural hematoma measured by different image segmentation methods [$M(P_{25}, P_{75})$] (ml).

Methods	Volume	Minimum	Maximum	Percentage error (%)
Manual segmentation	26.15	7.43	44.46	—
CNN segmentation	21.38	4.31	38.44	19.48 (11.45, 52.43)
ABC/2 volume formula	38.90	13.52	63.64	24.53 (14.25, 43.85)

Note: “—”: not available.




FIGURE 4: Comparison of the mean value of hematoma volume among three image segmentation methods. $*P < 0.05$.

3. Results

3.1. Volume of Subdural Hematoma Measured by Different Image Segmentation Methods. After examination and calculation, the percentage error of hematoma volume obtained by the CNN algorithm segmentation, ABC/2 volume formula segmentation, and the manual segmentation is shown in Table 1. Taking the manual segmentation result as the “golden standard,” the percentage error of CNN algorithm segmentation is less than ABC/2 volume formula. In addition, there was significant difference between ABC/2 volume formula segmentation and manual segmentation ($P > 0.05$). However, there was no significant difference between CNN algorithm segmentation and manual segmentation ($P < 0.05$) (Figure 4).

3.2. Evaluation of Model Prediction Efficiency. The area under curve (AUC) of the ABC/2 volume formula is 0.811 (95% CI: 0.717~0.905). The AUC of the manual segmentation is 0.840 (95% CI: 0.753~0.928). The AUC of the CNN algorithm segmentation is 0.832 (95% CI: 0.742~0.922). It is suggested that the segmentation result of CNN algorithm is closer to the manual segmentation than the ABC/2 volume formula (Figure 5).

4. Discussion

TSDH is one of the common secondary injuries of traumatic brain injury, accounting for about 40% of traumatic intracranial hematoma. It is still one of the main causes of traumatic brain injury death [21]. With the increase of traffic accidents and falling accidents, the incidence of this disease is increasing year by year [1]. The brain function of patients with TSDH is seriously damaged, and there are many kinds




FIGURE 5: Receiver operating characteristic curve analysis of different segmentation methods.

of dysfunction, which is a serious threat to patients’ health and life safety [22]. Therefore, it is of great significance to make a clear diagnosis of the disease and determine the exact volume of hematoma to guide clinical treatment.

The calculation method of Tada formula is simplified from the ellipsoid volume formula, which is widely used in the measurement of cerebral parenchymal hemorrhage and hematoma as a simple and fast estimation tool for hematoma volume beside the bed [23, 24]. Previous studies have shown that the correlation between ABC/2 volume formula and manual segmentation algorithm in acute subdural hematoma is high, but lower than parenchymal hemorrhage [25]. However, the percentage error of Tada formula in our study was larger than that in previous studies. The reason may be that the error of Tada formula increased with the increase of hematoma volume.

In order to make up for the deficiency of traditional algorithms, more and more researches have applied CNN correlation algorithms for hematoma segmentation in recent years [14, 19]. In this study, an automatic segmentation algorithm of intracranial hematoma based on CNN is adopted, automatically identifying and segment subdural hematoma displayed by head CT plain scan and calculating the hematoma volume. Taking the manual segmentation method as the “gold standard” [26], the consistency of CNN algorithm segmentation and ABC/2 volume formula is tested, respectively. The results show that the percentage error of CNN algorithm segmentation is less than ABC/2 volume formula. This shows that the hematoma volume

segmented by CNN algorithm is closer to manual segmentation. In addition, there was no significant difference between CNN algorithm segmentation and manual segmentation. However, in this study, the 5-fold crossover test was adopted, and the data set was randomly divided into five groups in turn as the training set and test set. The final results showed the average capability of the models [20].

To better grasp the disease condition and answer clinical questions more accurately, especially for the inaccurate calculation of subdural hematoma by traditional multifield formula, it is a challenge faced by clinicians. In addition, there is a big error in the multifield formula, and manual segmentation is time-consuming and laborious, limiting its application in the study of the relationship between hematoma volume and prognosis [27]. With the in-depth application of artificial intelligence technology in clinical medicine, it is possible to accurately and conveniently segment intracranial hematoma and explore hematoma volume as an indication of operation [28]. In our study, the AUC of the CNN algorithm segmentation is 0.832 (95% CI: 0.742~0.922). At the same time, that of the manual segmentation is 0.840 (95% CI: 0.753~0.928). It shows that the CNN algorithm segmentation method is closer to the manual segmentation method. More algorithms can be applied to evaluate the subdural hematoma in the future [29].

5. Conclusions

In the TSDH patients, compared with the traditional ABC/2 volume formula, the CNN algorithm used to calculate the volume of hematoma in head CT plain scan images is in good agreement with the manual segmentation results. However, the cases included in this study were single centers, and the sample size was small. There was some error in the algorithm segmentation. Therefore, the clinical application needs further exploration.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest to report regarding the present study.

Acknowledgments

This project is supported by the key project of Hefei Third People's Hospital (SYKZ202101).

References

- [1] E. Léveillé, S. Schur, A. AlAzri, C. Couturier, M. Maleki, and J. Marcoux, "Clinical characterization of traumatic acute interhemispheric subdural hematoma," *The Canadian Journal of Neurological Sciences*, vol. 47, no. 4, pp. 504–510, 2020.
- [2] K. Kobayashi, S. Imagama, K. Ando, Y. Nishida, and N. Ishiguro, "Acute non-traumatic idiopathic spinal subdural hematoma: radiographic findings and surgical results with a literature review," *European Spine Journal*, vol. 26, no. 11, pp. 2739–2743, 2017.
- [3] K. Rumalla, A. Y. Reddy, and M. K. Mittal, "Traumatic subdural hematoma: is there a weekend effect?," *Clinical Neurology and Neurosurgery*, vol. 154, pp. 67–73, 2017.
- [4] J. Huang, C. Gao, J. Dong, J. Zhang, and R. Jiang, "Drug treatment of chronic subdural hematoma," *Expert Opinion on Pharmacotherapy*, vol. 21, no. 4, pp. 435–444, 2020.
- [5] J. Feghali, W. Yang, and J. Huang, "Updates in chronic subdural hematoma: epidemiology, etiology, pathogenesis, treatment, and outcome," *World Neurosurgery*, vol. 141, pp. 339–345, 2020.
- [6] J. M. Ahn, K. S. Lee, J. H. Shim, J. S. Oh, J. J. Shim, and S. M. Yoon, "Clinical features of interhemispheric subdural hematomas," *Korean Journal of Neurotrauma*, vol. 13, no. 2, pp. 103–107, 2017.
- [7] R. Sahyouni, K. Goshtasbi, A. Mahmoodi, D. K. Tran, and J. W. Chen, "Chronic subdural hematoma: a historical and clinical perspective," *World Neurosurgery*, vol. 108, pp. 948–953, 2017.
- [8] V. N. Nguyen, D. Wallace, S. Ajmera et al., "Management of subdural hematomas in abusive head trauma," *Neurosurgery*, vol. 86, no. 2, pp. 281–287, 2020.
- [9] M. R. Liechti, U. J. Muehlethaler, A. F. Schneider et al., "Manual prostate cancer segmentation in mri: interreader agreement and volumetric correlation with transperineal template core needle biopsy," *European Radiology*, vol. 30, no. 9, pp. 4806–4815, 2020.
- [10] S. Y. Won, A. Zagorcic, D. Dubinski et al., "Excellent accuracy of abc/2 volume formula compared to computer-assisted volumetric analysis of subdural hematomas," *PLoS One*, vol. 13, no. 6, article e0199809, 2018.
- [11] A. Kashanian, H. Sparks, T. Kaprelian, and N. Pouratian, "Assessing the volume of large cerebral arteriovenous malformations: can the ABC/2 formula reliably predict true volume?," *Journal of Clinical Neuroscience*, vol. 65, pp. 1–5, 2019.
- [12] C. Roark, V. Vadlamudi, N. Chaudhary et al., "ABC/2 method does not accurately predict cerebral arteriovenous malformation volume," *Neurosurgery*, vol. 82, no. 2, pp. 220–225, 2018.
- [13] D. Xiao, J. Liu, T. Hu et al., "Simple ways to estimate meningioma volume: can abc- and sh-derived methods be used in clinical practice reliably?," *Journal of Oncology*, vol. 2021, Article ID 9712287, 11 pages, 2021.
- [14] X. Zhao, K. Chen, G. Wu et al., "Deep learning shows good reliability for automatic segmentation and volume measurement of brain hemorrhage, intraventricular extension, and peripheral edema," *European Radiology*, vol. 31, no. 7, pp. 5012–5020, 2021.
- [15] D. Dreizin, Y. Zhou, Y. Zhang, N. Tirada, and A. L. Yuille, "Performance of a deep learning algorithm for automated segmentation and quantification of traumatic pelvic hematomas on ct," *Journal of Digital Imaging*, vol. 33, no. 1, pp. 243–251, 2020.
- [16] Y. Liu, Q. Fang, A. Jiang, Q. Meng, G. Pang, and X. Deng, "Texture analysis based on u-net neural network for intracranial hemorrhage identification predicts early enlargement," *Computer Methods and Programs in Biomedicine*, vol. 206, article 106140, 2021.
- [17] N. Yu, H. Yu, H. Li, N. Ma, C. Hu, and J. Wang, "A robust deep learning segmentation method for hematoma volumetric

- detection in intracerebral hemorrhage,” *Stroke*, vol. 53, no. 1, pp. 167–176, 2022.
- [18] L. Rehman, A. Afzal, H. F. Aziz, S. Akbar, A. Abbas, and R. Rizvi, “Radiological parameters to predict hemorrhagic progression of traumatic contusional brain injury,” *Journal of Neurosciences in Rural Practice*, vol. 10, no. 2, pp. 212–217, 2019.
- [19] T. Dissanayake, T. Fernando, S. Denman, S. Sridharan, H. Ghaemmaghami, and C. Fookes, “A robust interpretable deep learning classifier for heart anomaly detection without segmentation,” *IEEE Journal of Biomedical and Health Informatics*, vol. 25, no. 6, pp. 2162–2171, 2021.
- [20] J. Liu and H. Zhao, “Application of convolution neural network in medical image processing,” *Technology and Health Care*, vol. 29, no. 2, pp. 407–417, 2021.
- [21] W. Yang and J. Huang, “Chronic subdural hematoma: epidemiology and natural history,” *Neurosurgery Clinics of North America*, vol. 28, no. 2, pp. 205–210, 2017.
- [22] D. C. Holl, V. Volovici, C. M. F. Dirven et al., “Pathophysiology and nonsurgical treatment of chronic subdural hematoma: from past to present to future,” *World Neurosurgery*, vol. 116, pp. 402–411.e2, 2018.
- [23] F. Cuce, G. Tulum, O. Dandin, T. Ergin, O. Karadas, and O. Osman, “A new practical intracerebral hematoma volume calculation method and comparison to simple abc/2,” *Turkish Neurosurgery*, vol. 30, no. 4, pp. 520–526, 2020.
- [24] W. Boisseau, C. Dargazanli, S. Smajda et al., “Use of the abc/2 method to select patients for thrombectomy after 6 hours of symptom onset,” *Neurology*, vol. 96, no. 1, pp. e10–e18, 2021.
- [25] O. P. Leary, L. H. Merck, S. D. Yeatts et al., “Computer-assisted measurement of traumatic brain hemorrhage volume is more predictive of functional outcome and mortality than standard abc/2 method: an analysis of computed tomography imaging data from the progesterone for traumatic brain injury experimental clinical treatment phase-iii trial,” *Journal of Neurotrauma*, vol. 38, no. 5, pp. 604–615, 2021.
- [26] H. S. Nguyen, M. Patel, L. Li, S. Kurpad, and W. Mueller, “Quantitative estimation of a ratio of intracranial cerebrospinal fluid volume to brain volume based on segmentation of ct images in patients with extra-axial hematoma,” *The Neuro-radiology Journal*, vol. 30, no. 1, pp. 10–14, 2017.
- [27] D. D. Oge, E. M. Arsava, M. Y. Pektezel, R. Gocmen, and M. A. Topcuoglu, “Intracerebral hemorrhage volume estimation: is modification of the abc/2 formula necessary according to the hematoma shape?,” *Clinical Neurology and Neurosurgery*, vol. 207, article 106779, 2021.
- [28] R. T. Kellogg, J. Vargas, G. Barros et al., “Segmentation of chronic subdural hematomas using 3D convolutional neural networks,” *World Neurosurgery*, vol. 148, pp. e58–e65, 2021.
- [29] J. Li, S. Fong, R. K. Wong, R. Millham, and K. K. L. Wong, “Elitist binary wolf search algorithm for heuristic feature selection in high-dimensional bioinformatics datasets,” *Scientific Reports*, vol. 7, no. 1, p. 4354, 2017.