

Heart rate response and functional capacity in patients with chronic heart failure with preserved ejection fraction

Eloy Domínguez¹, Patricia Palau¹, Eduardo Núñez², José María Ramón², Laura López³, Joana Melero¹, Alejandro Bellver¹, Enrique Santas², Francisco J. Chorro² and Julio Núñez^{2*}

¹Cardiology Department, Hospital General Universitari de Castelló, Universitat Jaume I, Castellón, Spain; ²Cardiology Department, Hospital Clínico Universitario de Valencia, INCLIVA, Universitat de València, Spain; ³Facultat de Fisioteràpia, Universitat de València, Valencia, Spain

Abstract

Aims The mechanisms of exercise intolerance in heart failure with preserved ejection fraction (HFpEF) are not yet elucidated. Chronotropic incompetence has emerged as a potential mechanism. We aimed to evaluate whether heart rate (HR) response to exercise is associated to functional capacity in patients with symptomatic HFpEF.

Methods and results We prospectively studied 74 HFpEF patients [35.1% New York Heart Association Class III, 53% female, age (mean \pm standard deviation) 72.5 \pm 9.1 years, and 59.5% atrial fibrillation]. Functional performance was assessed by peak oxygen consumption (peak VO₂). The mean (standard deviation) peak VO₂ was 10 \pm 2.8 mL/min/kg. The following chronotropic parameters were calculated: Delta-HR (HR at peak exercise – HR at rest), chronotropic index (CI) = (HR at peak exercise – resting HR)/[(220 – age) – resting HR], and CI according to the equation developed by Keteyian *et al.* (CIK) (HR at peak exercise – HR at rest)/[119 + (HR at rest/2) – (age/2) – 5 – HR at rest]. In a bivariate setting, peak VO₂ was positively and significantly correlated with Delta-HR ($r = 0.35$, $P = 0.003$), CI ($r = 0.27$, $P = 0.022$), CIK ($r = 0.28$, $P = 0.018$), and borderline with HR at peak exercise ($r = 0.22$, $P = 0.055$). In a multivariable linear regression analysis that included clinical, analytical, echocardiographic, and functional capacity covariates, the chronotropic parameters were positively associated with peak VO₂. We found a linear relationship between Delta-HR and peak VO₂ (β coefficient of 0.03; 95% confidence interval: 0.004–0.05; $P = 0.030$); conversely, the association among CIs and peak VO₂ was exponentially shaped.

Conclusions In patients with chronic HFpEF, the HR response to exercise was positively associated to patient's functional capacity.

Keywords Heart failure with preserved ejection fraction; Functional capacity; Chronotropic response

Received: 13 January 2017; Accepted: 5 February 2018

*Correspondence to: Julio Núñez, Cardiology Department, Hospital Clínico Universitario de Valencia, Universitat de València, Avda. Blasco Ibáñez 17, Valencia CP 46010, Spain. Email: yulnunez@gmail.com

Introduction

Heart failure (HF) with preserved ejection fraction (HFpEF) has become the most common form of HF in elderly patients.^{1,2} Exercise intolerance and reduced functional capacity are the main clinical manifestations in HFpEF and are associated to decreased quality of life and poor prognosis.^{3,4} Despite great efforts, the pathophysiological mechanisms of impaired exercise capacity in these patients are not yet fully elucidated.⁵ Although diastolic dysfunction

has been the focus of attention of the researchers, different pathophysiological mechanisms have also been proposed.^{6,7} Among them, some evidence endorsed abnormal heart rate (HR) response to exercise as a potential mechanism linked to exercise intolerance in these patients;^{8–10} however, the evidence is scarce, and its contribution in patients with chronic HFpEF remains unclear. In this work, we aimed to evaluate the relationship between HR response to exercise and exercise capacity patients with stable chronic HFpEF.

Methods

Study population

Between June 2012 and May 2016, we prospectively included 74 outpatients with symptomatic HFpEF followed in the outpatient HF clinic of a single academic centre. The HFpEF diagnosis was performed by trained cardiologists according to the definition proposed by the European Society of Cardiology guidelines.¹¹ Patients fulfilled all the following inclusion criteria: (a) New York Heart Association (NYHA) Functional Class \geq II, (b) previous admission for acute HF, (c) clinical stability during the last 3 months, (d) left ventricular ejection fraction $>50\%$, and (e) left ventricle hypertrophy and/or left atrial enlargement and/or diastolic dysfunction estimated by two-dimensional echocardiography. Exclusion criteria were (a) inability to perform a valid exercise test; (b) significant primary moderate to severe valve disease; (c) acute coronary syndrome, cardiac surgery, or revascularization within the previous 3 months; and (d) significant primary pulmonary disease, including pulmonary arterial hypertension, chronic thromboembolic pulmonary disease, or chronic obstructive pulmonary disease.

This study was approved by an institutional review committee conforming to the ethical guidelines of the 1975 Declaration of Helsinki, and all patients gave informed consent. Clinical, electrocardiographic, and treatment characteristics were recorded in electronic forms. Patients enrolled underwent cardiopulmonary exercise testing, two-dimensional transthoracic echocardiography, 6 minute walk test, and a blood laboratory test.

Cardiopulmonary exercise testing

Maximal functional capacity was evaluated with an incremental and symptom-limited cardiopulmonary exercise testing (CORTEX Metamax 3B) on a bicycle ergometer, beginning with a workload of 10 W and increasing stepwise at 10 W increments every 1 min. During exercise, patients were continuously monitored with 12-lead electrocardiogram and blood pressure measurements every 2 min. Gas exchange data and cardiopulmonary variables were averaged every 10 s values. Peak oxygen consumption (peak VO_2) was considered the highest value of VO_2 during the last 20 s of exercise. The ventilatory efficiency (VE/VCO_2 slope), defined as the slope of the linear relationship between minute ventilation (VE) and carbon dioxide production (VCO_2), was determined by measuring the slope across the entire course of exercise. HR was assessed by continuous electrocardiogram. Resting HR was calculated previous to testing as an average HR during 3 minutes of seated rest; and peak HR was defined as the highest HR achieved during exercise.

Endpoints

Cardiopulmonary exercise testing results were used to establish the chronotropic response to exercise effort. The primary endpoint was to determine the relationship between Delta-HR and chronotropic index (CI) with peak VO_2 . Delta-HR was defined as the absolute difference in HR (HR at peak exercise – HR at rest) and CI defined as $\text{CI} = (\text{HR at peak exercise} - \text{HR at rest}) / [(220 - \text{age}) - \text{HR at rest}]$.^{12,13} We defined chronotropic incompetence based on established criteria^{14–16}: CI <0.8 for those not taking beta-blockers and <0.62 for those taking beta-blockers. CI according to the equation developed by Keteyian *et al.*¹⁷ (CIK) defined as $(\text{HR at peak exercise} - \text{HR at rest}) / [119 + (\text{HR at rest}/2) - (\text{age}/2) - 5 - \text{HR at rest}]$ was selected as a secondary endpoint. The CIK has been considered suitable to predict maximum HR in patients with HF taking beta-blockers.^{17,18} In a sensitivity analyses, we explored whether the effect of Delta-HR on peak VO_2 differed according age (>70 vs. ≤ 70 years), gender (male vs. female), rhythm [sinus rhythm (SR) vs. atrial fibrillation (AF)], body mass index (>30 vs. ≤ 30 kg/m^2), N-terminal pro-BNP (NT-pro-BNP) (>1000 vs. ≤ 1000 pg/mL), and treatment with beta-blockers (yes vs. no).

Statistical analysis

The bivariate correlations of Delta-HR, CI, and CIK with peak VO_2 were assessed with Pearson/Spearman correlation coefficient as appropriate. In addition, each of these three exposures was tested as independent predictors of peak VO_2 using multivariable linear regression analysis. As covariates, most of the variables presented in *Table 1* were initially included in each of the three multivariable models. From there, a reduced model was achieved through a backward selection procedure; for each continuous predictor variable, the linearity assumption with peak VO_2 tested and the variable transformed to the appropriate fractional polynomial if necessary. To eliminate the potential bias due to ‘regression to the mean’, HR at baseline was included as an additional covariate in the regression models for Delta-HR (analysis of covariance). The discriminative ability of the multivariate models was evaluated with the adjusted R^2 . A two-sided P -value of <0.05 was considered to be statistically significant for all analyses. All analyses were performed using STATA 14.1.

Results

The mean (standard deviation) of age was 72.5 ± 9.1 ; 53% of patients were female, 35.1% displayed NYHA III/IV, 59.5% showed AF, and 79.7% were under beta-blocker treatment.

Table 1 Baseline characteristics of study population

Variables	Included patients (n = 74)
Demographic, medical history and vital signs	
Age, years	72.5 ± 9.1
Female, n (%)	39 (53)
Weight, kg	82 ± 15.1
Body surface area, m ²	1.9 ± 0.2
Body mass index, kg/m ²	31.2 (28.2–35)
Hypertension, n (%)	67 (90.5)
Dyslipidaemia, n (%)	61 (81)
Diabetes mellitus, n (%)	37 (50)
Ischaemic heart disease, n (%)	27 (36.5)
Baseline NYHA Class III/IV, n (%)	26 (35.1)
Previous smoker, n (%)	26 (35.1)
Atrial fibrillation, n (%)	44 (59.5)
Systolic blood pressure, mmHg	129 ± 16
HR, b.p.m.	69 ± 14
HR at peak exercise, b.p.m.	99 ± 21
Delta-HR, b.p.m.	30 ± 16
CI	0.39 ± 0.22
CIK	0.70 ± 0.38
Laboratory	
Haemoglobin, g/dL	12.8 ± 1.5
Ferritin, ng/mL	91 (48–175)
GFR, mL/min/m ²	58.4 ± 24.3
NT-pro-BNP, pg/mL	1021 (363–2109)
Echocardiography	
LVEF, %	68.4 ± 9.8
TAPSE, mm	21.8 ± 4.2
LAVI, mL/m ²	46 (39–58)
LVMI g/m ²	125 (104–125)
E/e' ratio	16.9 (13.4–23.4)
PASP, ^a mmHg	46 (39–53)
Exercise performance	
Peak VO ₂ , mL/min/kg	10 ± 2.8
VE/VCO ₂ slope	35 ± 7.6
RER	1.04 ± 0.1
METs	2.4 ± 0.9
6-MWT, m	262 (198–350)
Treatment	
Beta-blockers, %	59 (79.7)
ACEI, %	14 (18.9)
ARB, %	36 (48.6)
Anti-aldosterone, %	21 (28.3)
Loop diuretics, %	58 (78.4)
Thiazides, %	22 (29.7)
Digoxin, %	5 (6.7)

ACEI, angiotensin-converting enzyme inhibitor; ARB, angiotensin II receptor antagonist; CI, chronotropic index based on the Astrand formula (220 – age); CIK, chronotropic index according to the equation of Keteyian *et al.*; Delta-HR, absolute difference between heart rate at peak exercise and heart rate at baseline; GFR, estimated glomerular filtration rate using the Modification of Diet in Renal Disease formula; HR, heart rate; LAVI, left atrial volume index; LVEF, left ventricular ejection fraction; LVMI, left ventricular mass index; METs, metabolic equivalents; NT-pro-BNP, NT-pro-BNP; NYHA, New York Heart Association; PASP, pulmonary artery systolic pressure; peak VO₂, peak oxygen consumption; RER, respiratory exchange ratio; 6-MWT, distance walked in 6 minutes; TAPSE, tricuspid annular plane systolic excursion; TSAT, transferrin saturation index; VE/VCO₂ slope, relationship between minute ventilation and the rate of CO₂ elimination. Continuous variables are presented as mean ± standard deviation or median (interquartile range) as appropriate; categorical variables as percentages.

^aData available in 59 patients.

The means (standard deviation) of resting HR and systolic blood pressures were 69 ± 14 b.p.m. and 129 ± 16 mmHg, respectively. The mean peak VO₂, VE/VCO₂ slope, respiratory exchange ratio (RER), and peak HR were 10 ± 2.8 mL/min/kg, 35 ± 7.6, 1.04 ± 0.1, and 99 ± 21 b.p.m., respectively. Following the criteria of chronotropic incompetence defined above, we found that 66 patients (89.2%) displayed chronotropic incompetence. Baseline characteristics of the study population are summarized in *Table 1*.

There were no differences in the medians (interquartile range) of Delta-HR across age (>70 vs. ≤70 years), gender (male vs. female), rhythm (SR vs. AF), body mass index (>30 vs. ≤30 kg/m²), NT-pro-BNP (>1000 vs. ≤1000 pg/mL), and treatment with beta-blockers (yes vs. no) as shown in *Figure 1*.

In a bivariate setting, peak VO₂ was negatively correlated with age ($r = -0.39$, $P = 0.001$), NT-pro-BNP ($r = -0.26$, $P = 0.028$), E/e' septal ($r = -0.26$, $P = 0.029$), and pulmonary artery systolic pressure ($r = -0.32$, $P = 0.015$) while positively correlated with haemoglobin ($r = 0.41$, $P < 0.001$), transferrin saturation index ($r = 0.39$, $P = 0.001$), serum ferritin ($r = 0.28$, $P = 0.022$), Delta-HR ($r = 0.35$, $P = 0.003$), CI ($r = 0.27$, $P = 0.022$), CIK ($r = 0.28$, $P = 0.018$), and RER ($r = 0.53$, $P < 0.001$). HR at peak exercise ($r = 0.22$, $P = 0.055$) and right ventricular function (tricuspid annular plane systolic excursion: $r = 0.20$, $P = 0.088$) were borderline correlated. Left atrial volume index ($r = -0.16$, $P = 0.179$), glomerular filtration rate (GFR: $r = 0.15$, $P = 0.210$), and HR at rest ($r = -0.06$, $P = 0.633$) were not correlated with peak VO₂.

We performed multivariable linear regression analyses for each of the three exposures; Delta-HR (Model 1), CI (Model 2), and CIK (Model 3) included the following covariates: age, gender, body surface area, AF rhythm, NYHA ≥3, GFR, haemoglobin, NT-pro-BNP, E/e' septal, and RER. HR at baseline was included as additional covariate in the regression models for Delta-HR. The variables treatment with digoxin and beta-blockers did not achieve statistical significance, and thus, they were excluded. Each of the three exposures was positively associated with peak VO₂.

In Model 1, Delta-HR was linear and positively associated with peak VO₂ [β coefficient of 0.03; 95% confidence interval (0.004–0.05); $P = 0.030$]. *Figure 2A* depicts this association, after centring Delta-HR at its median value (27 b.p.m.). The R^2 estimated for this model was 0.68.

In Model 2, CI showed a positive and exponential association with peak VO₂ as depicted in *Figure 2B*. For instance, using as a reference point the median of CI (0.35) when CI is 0.41, 0.6, 0.81, and 1.02, the predicted mean change in peak VO₂ is 0.06, 0.41, 1.16, and 2.41 mL/min/kg, respectively. The R^2 estimated for this model was 0.70.

Likewise, in Model 3, the adjusted association of CIK with peak VO₂ was also significant, positive, and exponentially shaped as shown in *Figure 2C*. For instance, using as reference point the median of CI (0.62) when CIK is 0.91, 1.24, 1.5, and 1.76, the predicted mean change in peak VO₂ is

Figure 1 Delta heart rate across subgroups of age, gender, type of rhythm, body mass index, NT-pro-BNP, and beta-blocker treatment. AF, atrial fibrillation; BB, beta-blocker treatment; BMI, body mass index; NT-pro-BNP, N-terminal pro-BNP; SR, sinus rhythm.





Figure 2 Functional form of the adjusted association among different chronotropic indexes and predicted change in peak VO_2 . Variables modelled with fractional polynomials. CI, chronotropic index, a measure of chronotropic response based on the Astrand formula; CIK, chronotropic index according to the equation of Keteyian *et al.*; Delta heart rate, absolute difference between heart rate at peak exercise and heart rate at baseline; peak VO_2 , peak oxygen consumption.



0.22, 0.72, 1.35, and 2.25 mL/min/kg, respectively. The R^2 estimated for this model was 0.70.

In a sensitivity analyses, we found that the effect of Delta-HR on peak VO_2 did not significantly differ according the pre-specified subgroups: age (>70 vs. ≤70 years: P -value for the interaction = 0.672), gender (male vs. female: P -value

for the interaction = 0.606), rhythm (SR vs. AF: P -value for the interaction = 0.160), body mass index (>30 vs. ≤30 kg/m²: P -value for the interaction = 0.631), NT-pro-BNP (>1000 vs. ≤1000 pg/mL: P -value for the interaction = 0.268), and treatment with beta-blockers (yes vs. no: P -value for the interaction = 0.986).

Discussion

In this study, we found that patients with symptomatic HFpEF displayed a high prevalence of blunted HR response to exercise and it was independently associated to patient's functional capacity impairment evaluated by peak VO_2 .

Prevalence and pathophysiology of chronotropic incompetence

The prevalence of chronotropic incompetence reported among previous studies is widely variable, ranging from 20% to 75%.^{8–10,19–21} The criteria used to define chronotropic incompetence, as well as patient characteristics (age, disease severity, heart rhythm, and drug therapy), could explain these differences. In the present study, we postulate that the greater severity and the proportion of beta-blocker treatment (79.7%) of the patients may explain the higher prevalence of chronotropic incompetence here reported.

An appropriate increase in peak VO_2 during exercise depends on adequate HR response according to the Fick equation²²: $\text{VO}_2 = [\text{HR} \times \text{stroke volume}] \times [\text{arterial-venous oxygen content difference}]$. In this way, chronotropic incompetence, defined as the inability of the heart to increase adequately during physical exertion, has been proposed as one of the key mechanisms linked to exercise intolerance in HFpEF in some studies.^{8–10,19} The pathophysiology of impaired HR response is complex and has been associated to autonomic imbalances such as (a) the decrease in beta-receptor density;²³ (b) the desensitization of beta-receptors despite the presence of increased circulating catecholamine levels;^{24–26} and (c) the remodelling of sinus node function.²⁷ However, the causative role of chronotropic incompetence in the pathophysiology of this syndrome is difficult to establish. Blunted HR response may just be a consequence of decreased exercise capacity in which other mechanisms such as premature cessation of exercise due to dyspnoea or muscle fatigue might be playing a pivotal role.^{6,28,29} However, the high prevalence of chronotropic incompetence in the presence of adequate effort and the robust multivariate associations found in this study lead us to speculate abnormal HR response to exercise may play a causative role.

Clinical implications

Lack of effective medical treatments for patients with HFpEF has driven an empiric medical management.³⁰ In agreement with present findings, previous studies exploring the effect of medical therapies such as beta-blockers^{31–35} to reduce HR response have shown neutral or negative results on functional capacity³⁵ and adverse clinical endpoints³³ in patients with HFpEF. In this regard, studies of the effect of ivabradine treatment in terms of improvement of functional capacity in

HFpEF patients have shown contradictory results.^{36,37} The main reasons underlying these controversial findings are unclear but may reflect several differences in the study designs, patient's characteristics, and the doses of ivabradine. Overall, we believe these inconsistent results reflect the complex and heterogeneous pathophysiology and different clinical phenotypes of this syndrome. In light of these findings, we postulate that the more advanced the disease, especially in elderly patients, the greater the importance of chronotropic incompetence as a crucial pathophysiological mechanism in patients with HFpEF.

Conversely, in the setting of HF with reduced ejection fraction (HFrEF), recently, Jamil *et al.*³⁸ examined the relationship between exercise HR rise and exercise capacity in patients with clinical stability and previous pacemaker implantation. The negative results obtained in this study contribute to our understanding of the role of HR response in patients with HFrEF, but the controversy in patients with HFpEF remains alive.

In our opinion, this line of research has important therapeutic implications. For instance, and based on this and other results, we may envision a potential clinical benefit of HR-lowering drug withdrawal in patients with HFpEF and chronotropic incompetence. Along this line, the RAPID-HF trial (NCT02145351) is currently testing the hypothesis that improved HR responsiveness via rate-adaptive atrial pacing will improve exercise capacity in patients with HFpEF. In addition, despite current evidence not strongly supporting beta-blocker therapy in HFpEF, these agents are frequently prescribed in daily clinical practice.³⁹ The results of this study called into question the use of beta-blockers in patients with HFpEF and chronotropic incompetence.

Study limitations

There are some limitations that need to be acknowledged. First, the main limitations of this study are the lack of a control group of meaningful similarities and differences and the limited potential for generalizing our results to other populations with HF due to the small sample size. Second, although we did not find any differential effect of chronotropic parameters across the type of rhythm, we cannot exclude a Type II error. Further larger studies evaluating separately patients with SR and AF are warranted. Finally, with the present data, we cannot unravel the pathophysiological mechanism endorsing this association.

Conclusions

In patients with chronic HFpEF, the HR response to exercise is positively associated to patient's functional capacity evaluated by peak VO_2 . Further studies are warranted to

explore the pathophysiological role of chronotropic incompetence in HFpEF.

Conflict of interest

None declared.

Funding

This work was supported in part by grants from Sociedad Española de Cardiología: Investigación Clínica en Cardiología, Grant SEC 2015, CIBER CV 16/11/00420, 16/11/00403, FEDER, and PIE15/00013.

References

- Lam CS, Donal E, Kraigher-Krainer E, Vasan RS. Epidemiology and clinical course of heart failure with preserved ejection fraction. *Eur J Heart Fail* 2011; **13**: 18–28.
- Abbate A, Arena R, Abouzaki N, Van Tassel BW, Canada J, Shah K, Biondi-Zoccai G, Voelkel NF. Heart failure with preserved ejection fraction: refocusing on diastole. *Int J Cardiol* 2015; **179**: 430–440.
- Kitzman DW, Little WH, Brubaker P, Anderson RT, Hundley WG. Pathophysiological characterization of isolated diastolic heart failure in comparison to systolic heart failure. *JAMA* 2002; **288**: 2144–2150.
- Fonarow GC, Stough WG, Abraham WT, Albert NM, Gheorghiade M, Greenberg BH, O'Connor CM, Sun JL, Yancy CW, Young JB. Characteristics, treatments, and outcomes of patients with preserved systolic function hospitalized for heart failure: a report from the OPTIMIZE-HF Registry. *J Am Coll Cardiol* 2007; **50**: 768–777.
- Borlaug BA. Mechanisms of exercise intolerance in heart failure with preserved ejection fraction. *Circ J* 2014; **78**: 20–32.
- Upadhyaya B, Taffet GE, Cheng CP, Kitman DW. Heart failure with preserved ejection fraction in the elderly: scope of the problem. *J Mol Cell Cardiol* 2015; **83**: 73–87.
- Paulus WJ, Tschoepe C. A novel paradigm for heart failure with preserved ejection fraction: comorbidities drive myocardial dysfunction and remodeling through coronary microvascular endothelial inflammation. *J Am Coll Cardiol* 2013; **62**: 263–271.
- Brubaker PH, Joo KC, Stewart KP, Fray B, Moore B, Kitman DW. Chronotropic incompetence and its contribution to exercise intolerance in older heart failure patients. *J Cardiopulm Rehabil* 2006; **26**: 86–89.
- Phan TT, Shivu GN, Abozguia K, Davies C, Nassimzadeh M, Jimenez D, Weaver R, Ahmed I, Frenneaux M. Impaired heart rate recovery and chronotropic incompetence in patients with heart failure with preserved ejection fraction. *Circ Heart Fail* 2010; **3**: 29–34.
- Borlaug BA, Melenovsky V, Russell SD, Kessler K, Pacak K, Becker LC, Kass DA. Impaired chronotropic and vasodilator reserves limit exercise capacity in patients with heart failure and a preserved ejection fraction. *Circulation* 2006; **114**: 2138–2147.
- Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, Falk V, González-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GMC, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P, ESC Scientific Document Group. 2016 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2016; **37**: 2129–2200.
- Astrand I. Aerobic work capacity in men and women with special reference to age. *Acta Physiol Scand Suppl* 1960; **49**: 1–92.
- Brubaker PH, Kitman DW. Chronotropic incompetence: causes, consequences and management. *Circulation* 2011; **123**: 1010–1020.
- Wilkoff BL, Miller RE. Exercise testing for chronotropic assessment. *Cardiol Clin* 1992; **10**: 705–717.
- Lauer MS, Francis GS, Okin PM, Pashkow FJ, Snader CE, Marwick TH. Impaired chronotropic response to exercise stress testing as a predictor of mortality. *JAMA* 1999; **281**: 524–529.
- Khan ME, Pothier CE, Lauer MS. Chronotropic incompetence as a predictor of death among patients with normal electrocardiograms taking beta blockers. *Am J Cardiol* 2005; **96**: 1328–1333.
- Keteyian SJ, Kitman D, Zannad F, Landzberg J, Arnold JM, Brubaker P, Brawner CA, Bensimhon D, Hellkamp AS, Ewald G. Predicting maximal heart rate in heart failure patients on beta-blockade therapy. *Med Sci Sports Exerc* 2012; **44**: 371–376.
- Dobre D, Zannad F, Keteyian SJ, Stevens SR, Rossignol P, Kitman DW, Landzberg J, Howlett J, Kraus WE, Ellis SJ. Association between resting heart, chronotropic index, and long-term outcomes in patients with heart failure receiving β -blocker therapy: data from the HF-ACTION trial. *Eur Heart J* 2013; **34**: 2271–2280.
- Wang J, Fang F, Yip GW, Sanderson JE, Feng W, Xie JM, Luo XX, Lee AP, Lam YY. Importance of chronotropic response and left ventricular long-axis function for exercise performance in patients with heart failure and preserved ejection fraction. *Int J Cardiol* 2016; **202**: 339–343.
- Borlaug BA, Olson TP, Lam CS, Flood KS, Lerman A, Johnson BD, Redfield MM. Global cardiovascular reserve dysfunction in heart failure with preserved ejection fraction. *J Am Coll Cardiol* 2010; **56**: 845–854.
- Klein DA, Katz DH, Beussink-Nelson L, Sanchez CL, Strzelczyk TA, Shah SJ. Association of chronic kidney disease with chronotropic incompetence in heart failure with preserved ejection fraction. *Am J Cardiol* 2015; **116**: 1093–1100.
- Fick A. Über die Messung des Blutquantums in den Herzventrikeln. Sitzungsberichte der Physiologisch-Medizinischen Gesellschaft zu Würzburg 1870; **2**: 16.
- Bristow MR, Hershberger RE, Port JD. Beta-adrenergic pathways in non-failing and failing human ventricular myocardium. *Circulation* 1990; **82**: 12–25.
- Colucci WS, Ribeiro JP, Rocco MB, Quigg RJ, Creager MA, Marsh JD, Gauthier DF, Hartley LH. Impaired chronotropic response to exercise in patients with congestive heart failure. *Circulation* 1989; **80**: 314–323.
- Samejima H, Omiya K, Uno M, Inoue K, Tamura M, Itoh K, Suzuki K, Akashi Y, Seki A, Suzuki N, Osada N, Tanabe K, Miyake F, Itoh H. Relationship between impaired chronotropic response, cardiac output during exercise, exercise tolerance in patients with chronic heart failure. *Jpn Heart J* 2003; **44**: 515–525.
- Messias LR, Messias AC, de Miranda SM, Wiefels CC, Ferreria AG, Santos LM, Teixeira JA, Marostica E, Mesquita CT. Abnormal adrenergic activation is the major determinant of reduced functional capacity in heart failure with preserved ejection fraction. *Int J Cardiol* 2015; **203**: 900–902.

27. Sanders P, Kistler PM, Morton JB, Spence SJ, Kalman JM. Remodeling of sinus node function in patients with congestive heart failure. *Circulation* 2004; **110**: 897–903.
28. Haykowsky M, Brubaker P, Kitzman D. Role of physical training in heart failure with preserved ejection fraction. *Curr Heart Fail Rep* 2012; **9**: 101–106.
29. Abudiab MM, Redfield MM, Melenosvsky V, Olson TP, Kass DA, Johnson BD, Borlaug BA. Cardiac output response to exercise in relation to metabolic demand in heart failure with preserved ejection fraction. *Eur J Heart Fail* 2013; **15**: 776–785.
30. Senni M, Paulus WJ, Gavazzi A, Fraser AG, Díez J, Solomon SD, Smiseth OA, Guazzi M, Lam CSP, Maggioni AP, Tschöpe C, Metra M, Hummel SC, Edelmann F, Ambrosio G, Stewart AJ, Filippatos GS, Gheorghiadu M, Anker SD, Levy D, Pfeffer MA, Stough WG, Piesk BM. New strategies for heart failure with preserved ejection fraction: the importance of targeted therapies for heart failure phenotypes. *Eur Heart J* 2014; **35**: 2797–2815.
31. Massie BM, Nelson JJ, Lukas MA, Greenberg B, Fowler MB, Gilbert EM, Abraham WT, Lottes SR, Franciosa JA, COHERE Participant Physicians. Comparison of outcomes and usefulness of carvedilol across a spectrum of left ventricular ejection fractions in patients with heart failure in clinical practice. *Am J Cardiol* 2007; **99**: 1263–1268.
32. Hernandez AF, Hammill BG, O'Connor CM, Schulman KA, Curtis LH, Fonarow GC. Clinical effectiveness of β -blockers in heart failure: findings from the OPTIMIZE-HF (Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure) registry. *J Am Coll Cardiol* 2009; **53**: 184–192.
33. Yamamoto K, Origasa H, Hori M, Investigators JD. Effects of carvedilol on heart failure with preserved ejection fraction: the Japanese Diastolic Heart Failure Study (J-HFS). *Eur J Heart Fail* 2013; **15**: 110–118.
34. Lund LH, Benson L, Dahlström U, Edner M, Friberg L. Association between use of β -blockers and outcomes in patients with heart failure and preserved ejection fraction. *JAMA* 2014; **312**: 2008–2018.
35. Conraads VM, Metra M, Kamp O, Keulenaer GW, Pieske B, Zamorano J, Vardas PE, Böhm M, Cas LD. Effects of the long-term administration of nebivolol on the clinical symptoms, exercise capacity, and left ventricular function of patients with diastolic dysfunction: results of the ELANDD study. *Eur J Heart Fail* 2012; **14**: 219–225.
36. Kosmala W, Holland DJ, Rojek A, Wright L, Przewlocka-Kosmala M, Marwick TH. Effect of If-channel inhibition on hemodynamic status and exercise tolerance in heart failure with preserved ejection fraction. *J Am Coll Cardiol* 2013; **62**: 1330–1338.
37. Pal N, Sivaswamy N, Mahmood M, Yavari A, Rudd A, Singh S, Dawson DK, Francis JM, Dwight JS, Watkins H, Neubauer S, Frenneaux M, Ashrafian H. Effect of selective heart rate slowing in heart failure with preserved ejection fraction. *Circulation* 2015; **132**: 1719–1725.
38. Jamil HA, Gierula J, Paton MF, Byrom R, Lowry JE, Cubbon RM, Cairns DA, Kearney MT, Witte KK. Chronotropic incompetence does not limit exercise capacity in chronic heart failure. *J Am Coll Cardiol* 2016; **67**: 1885–1896.
39. Santas E, Valero E, Mollar A, García-Blas S, Palau P, Miñana G, Núñez E, Sanchis J, Chorro FJ, Núñez J. Burden of recurrent hospitalizations following an admission for acute heart failure: preserved versus reduced ejection fraction. *Rev Esp Cardiol* 2016; **70**: 239–246.