

POSTER PRESENTATION

Open Access

Zero-lag synchronization in cortical motifs

Leonardo L Gollo^{1,2*}, Claudio Mirasso¹, Olaf Sporns³, Michael Breakspear^{2,4,5}

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

Zero-lag synchronization between distant cortical areas has been observed in diverse experimental settings, and between many different regions of the brain [1,2]. Several mechanisms have been proposed to account for such isochronous synchronization in the presence of long conduction delays: Of these, the phenomena of "dynamical relaying" - a mechanism that relies on a specific network motif (M9) - has proven to be the most robust with respect to parameter and system noise [2-4]. Surprisingly, despite a prevailing belief in the community, the common driving motif (M3) is an unreliable means of establishing zero-lag synchrony. Although dynamical relaying has been validated in empirical and computational studies [4], the deeper dynamical mechanisms and comparison to dynamics on other motifs is lacking (see Figure 1). Given the presence of different network motifs in cortical

systems, such deeper insights are of high priority. By systematically comparing synchronization on a variety of small motifs, we establish that the presence of a single reciprocally connected pair - a "resonance pair" - plays a crucial role in disambiguating those motifs that foster zero-lag synchrony in the presence of conduction delays (such as dynamical relaying, M9) from those that do not (such as the common driving triad, M3). Remarkably, minor structural changes to M3 that incorporate a reciprocal pair (hence M6, M9, M3+1) recovers robust zero-lag synchrony. The findings are observed in computational models of spiking neurons, populations of spiking neurons and mean field neural models, and arise whether the systems are periodic, chaotic, noise-free or driven by stochastic inputs. The influence of the resonant pair is also robust to parameter mismatch and asymmetrical time delays




Figure 1 Motifs and zero-lag cross-correlations between nodes 1 and 3 in mean field neural models for varying coupling strength (left panel) and delay (right panel).

* Correspondence: leonardo.gollo@qimr.edu.au

¹IFISC, Instituto de Física Interdisciplinar y Sistemas Complejos (CSIC-UIB), Palma de Mallorca, Spain

amongst the elements of the motif. The synchronization of the commonly driven nodes is optimal when the driver node is part of a resonant pair, or is under the influence of another resonant pair (since its effect can propagate in the network). We call this manner of facilitating zero-lag synchrony *resonance-induced synchronization* and propose that it may be a critical feature of zero-lag synchrony in the brain.

Author details

¹IFISC, Instituto de Física Interdisciplinar y Sistemas Complejos (CSIC-UIB), Palma de Mallorca, Spain. ²Program of Mental Health Research, Queensland Institute of Medical Research, Brisbane, QLD, Australia. ³Department of Psychological and Brain Sciences, Indiana University, Bloomington, Indiana, USA. ⁴School of Psychiatry, University of New South Wales and The Black Dog Institute, Sydney, NSW, Australia. ⁵The Royal Brisbane and Woman's Hospital, Brisbane, QLD, Australia.

Published: 8 July 2013

References

1. Roelfsema PR, Engel AK, Konig P, Singer W: Visuomotor integration is associated with zero time-lag synchronization among cortical areas. *Nature* 1997, **385**:157-161.
2. Vicente R, Gollo LL, Mirasso CR, Fischer I, Pipa G: Dynamical relaying can yield zero time lag neuronal synchrony despite long conduction delays. *Proc Natl Acad Sci* 2008, **105**(44):17157-17162.
3. Gollo LL, Mirasso CR, Villa AE: Dynamic control for synchronization of separated cortical areas through thalamic relay. *Neuroimage* 2010, **52**(3):947-55.
4. Gollo LL, Mirasso CR, Atienza M, Crespo-Garcia M, Cantero JL: Theta band zero-lag long-range cortical synchronization via hippocampal dynamical relaying. *PLoS ONE* 2011, **6**(3):e17756.

doi:10.1186/1471-2202-14-S1-P37

Cite this article as: Gollo et al.: Zero-lag synchronization in cortical motifs. *BMC Neuroscience* 2013 **14**(Suppl 1):P37.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

