

DNA sequence comparisons of *Ophiostoma* spp., including *Ophiostoma aurorae* sp. nov., associated with pine bark beetles in South Africa

Xudong Zhou^{1*}, Z. Wilhelm de Beer² and Michael J. Wingfield¹

¹Tree Protection Co-operative Programme (TPCP), Forestry and Agricultural Biotechnology Institute (FABI); ²Department of Microbiology and Plant Pathology, University of Pretoria, Pretoria, 0002, Republic of South Africa

*Correspondence: Xudong Zhou, Xu.Zhou@fabi.up.ac.za

Abstract: Bark beetles (Coleoptera: Scolytinae) are well-recognized vectors of *Ophiostoma* species. Three non-native bark beetle species infest various *Pinus* species in South Africa, and they are known to carry at least 12 different species of ophiostomatoid fungi. Some of these fungi have not been identified to species level. The aim of this study was to determine or confirm the identities of *Ophiostoma* species associated with bark beetles in South Africa using comparisons of DNA sequence data. Identities of *Ophiostoma ips*, *O. floccosum*, *O. pluriannulatum*, *O. quercus* and *O. stenoceras* were confirmed. *Ophiostoma abietinum*, *O. piliferum* and *Pesotum fragrans* are recognised for the first time and the new species, *O. aurorae* sp. nov., is described from pine-infesting bark beetles in South Africa.

Taxonomic novelty: *Ophiostoma aurorae* X.D. Zhou & M.J. Wingf. sp. nov.

Key words: Bark beetle, *Ophiostoma*, phylogeny, taxonomy.

INTRODUCTION

Conifer-infesting bark beetles (Coleoptera: Scolytinae) are economically important forest insects. They include many primary pest species, which can attack healthy living trees and have caused significant economic losses to the global forestry industry (Wood & Bright 1992). In South Africa, three non-native bark beetle species, *Hylastes angustatus*, *Hylurgus ligniperda*, and *Orthotomicus erosus* infest various *Pinus* spp. (Tribe 1992). They are generally considered as secondary pests, although *H. angustatus* may undergo maturation feeding on healthy living seedlings causing significant losses during plantation establishment (Tribe 1992).

Bark beetles are well-known vectors of fungi, especially *Ophiostoma* species (Six 2003, Kirisits 2004, Harrington 2005). The ophiostomatoid fungi are a polyphyletic group of morphologically similar fungi, adapted for insect dispersal. Several ophiostomatoid fungi are important pathogens of conifers (Harrington & Cobb 1988, Wingfield et al. 1993b, Jacobs & Wingfield 2001), while many others can cause sapstain on logs and freshly cut wood (Wingfield et al. 1993b). The group includes the genera *Ceratocystis* Ellis & Halst., *Gondwanamyces* G.J. Marais & M.J. Wingf., *Sphaeronaemella* P. Karst. and *Cornuvesica* C.D. Viljoen, M.J. Wingf. & K. Jacobs and their anamorphs in the *Microascales* (Spatafora & Blackwell 1994, Hausner et al. 2000), and *Ophiostoma* Syd. & P. Syd., *Grosmannia* Goid. and *Ceratocystiopsis* H.P. Upadhyay & W.B. Kendr., with their *Pesotum* J.L. Crane & Schokn., *Leptographium* Lagerb. & Melin, *Sporothrix* Hektoen & C.F. Perkins and *Hyalorhinocladiella* H.P. Upadhyay & W.B. Kendr. anamorphs in the *Ophiostomatales* (Zipfel et al. 2006).

More than 30 ophiostomatoid fungi have been reported from South Africa (Table 1), of which at

least 12 are associated with the three exotic pine-infesting bark beetle species in the country (Zhou et al. 2001). These fungi have been isolated from the insects or their galleries and identified based on their morphological characteristics (Zhou et al. 2001). Eight of these species belong to the genus *Ophiostoma* (*sensu* Zipfel et al. 2006) or its anamorphs. However especially those of which only the anamorphs were observed remained to be identified to species level (Zhou et al. 2001). The aim of this study was to use DNA sequence comparisons to confirm the identities of the *Ophiostoma* spp. (Zipfel et al. 2006) from South African pine bark beetles, previously identified based only on morphology (Zhou et al. 2001).

MATERIALS AND METHODS

Fungal isolates

Twelve isolates (Table 2) used in this study originated from a previous investigation of ophiostomatoid fungi associated with the three pine-infesting bark beetle species in South Africa (Zhou et al. 2001). All cultures are maintained in the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa. A relevant sub-set of cultures has been deposited with the Centraalbureau voor Schimmelcultures (CBS), Utrecht, Netherlands.

DNA sequencing and phylogenetic analyses

Single hyphal-tip cultures from the 12 isolates were grown on 2 % MEA (20 g Biolab malt extract, 20 g Biolab agar, and 1000 mL deionised water). DNA was extracted using PrepMan Ultra Sample reagent (Applied Biosystems) as described by Aghayeva et al. (2004). The ITS (internal transcribed spacer) region of

the ribosomal RNA operon was amplified using primers ITS1-F (Gardes & Bruns 1993) and ITS4 (White *et al.* 1990). PCR products were sequenced with the same primers. Conditions for PCR amplification and sequencing reactions were as described by Zhou *et al.* (2004b). For comparisons, ITS sequences of closely related taxa (Table 3) were obtained from GenBank.

All sequences were aligned using MAFFT v. 5.667 (Katoh *et al.* 2002). Phylogenetic relationships among the isolates were determined using distance analyses in MEGA3 (<http://www.megasoftware.net/>). Trees were constructed using the Neighbour-joining tree-building algorithm (Saitou & Nei 1987) and rooted using GenBank sequences of *Leptographium guttulatum* M. J. Wingf. & K. Jacobs (AY649782 and AY649783). Bootstrap analyses (1000 replicates) were run to determine confidence levels of the branching points (Felsenstein 1985).

Three of the 12 isolates (CMW 19362, CMW 19363, and CMW 19364) grouped in a clade separate from the other isolates, all of which grouped with known taxa. For these three isolates, part of the β -tubulin gene was amplified using primers Bt2a and Bt2b (Glass & Donaldson 1995). For each of the two regions, phylogenetic analyses were done separately, followed by a distance analysis of the combined data set. A partition homogeneity test was performed in PAUP v. 4.0b8 (Phylogenetic Analyses Using Parsimony) (Swofford 2002) to determine the congruence of the two data sets.

Morphology

Isolates (CMW 19362, CMW 19363, and CMW 19364) that resided in a defined phylogenetic clade of unknown identity were grown on 2 % WA (20 g Biolab agar and 1000 mL deionised water) with sterilised pine twigs,




Fig. 1. Neighbour-joining tree of *Ophiostoma* species associated with bark beetles in South Africa based on ITS sequences (ITS1 and ITS2 regions, as well as 5.8S rRNA gene). Isolates sequenced in this study are printed in bold. Bar = total nucleotide differences between taxa. Bootstrap values (1000 replicates) are indicated above the branches.

and on 1.5 % oatmeal agar (15 g oats powder, 20 g Biolab agar and 1000 mL deionised water) to induce production of perithecia. Perithecia with ascospores were formed in two isolates (CMW 19362 and CMW 19363) on oatmeal agar. Thirty measurements were made for each structure, and the ranges and averages were computed. Anamorph structures were observed on 7-d-old slide cultures (Riddell 1950), mounted in lactophenol.

RESULTS

DNA Sequence analyses

PCR of the ITS regions delivered products ranging from about 530 to 610 bp in size. Comparison of the ITS sequences with GenBank sequences confirmed the identities of seven *Ophiostoma* spp. (Fig. 1). These included *O. stenoceras* (Robak) Nannf., *O. abietinum* Marm. & Butin, *O. piliferum* (Fr.) Syd. & P. Syd., *O. pluriannulatum* (Hedg.) Syd. & P. Syd., *O. quercus* (Georgev.) Nannf., *O. floccosum* Math.-Käärik, and *Pesotum fragrans* (Math.-Käärik) G. Okada & Seifert. The identity of *O. ips* (Rumbold) Nannf. (also included in the study) had previously been confirmed based on DNA sequence comparisons (Zhou *et al.* 2004a).

Fragments 541 bp in size from the ITS region, and 345 from the partial β -tubulin gene were amplified for the three unidentified isolates (CMW 19362, CMW 19363, and CMW 19364). The β -tubulin region included intron 5, but no intron 4 was present. This corresponds with species in the *O. stenoceras* -complex (Zipfel *et al.* 2006). Sequences of isolates representing the majority of species in this complex were thus selected for further phylogenetic analyses, with *O. nigrocarpum* as outgroup. *Ophiostoma* spp. from outside the *O. stenoceras* -complex were not included in these

analyses because of the presence of intron 4, but no intron 5 (Zipfel *et al.* 2006). The partition homogeneity test ($P = 0.003$) confirmed that the ITS and β -tubulin data sets were congruent. Distance analyses for the combined data set showed that the three unidentified isolates grouped together with a bootstrap support of 100 % (Fig. 2), and that they either represented an undescribed species or a species for which no sequence data are available.

Morphology

The three isolates (CMW 19362, CMW 19363, and CMW 19364) are morphologically similar to each other and different from any other described *Ophiostoma* species. They produced a typical *Sporothrix* anamorph in culture with swollen clavate conidia. Two of the isolates produced ascomata with allantoid rounded ascospores.

TAXONOMY

Based on combined sequence comparisons of the ITS regions and partial β -tubulin gene, as well as morphology, we conclude the three isolates from *H. angustatus* infesting pines in South Africa represent an undescribed taxon. This is described as follows:

***Ophiostoma aurorae* X.D. Zhou & M.J. Wingf., sp. nov.** MycoBank MB500888. Figs 3A–3F.

Anamorph: *Sporothrix* (Fig. 3D–F).

Etymology: The type locality of this species is in Mpumalanga Province, South Africa. In siSwati, the name of the province means “the place where the sun rises”. Aurora was the Roman (Latin) goddess of dawn, so the specific epithet is an oblique reference to the type locality.




Fig. 2. Neighbour-joining tree of the *Ophiostoma stenoceras* - *Sporothrix schenckii* complex of species, including *O. aurorae* based on the combined ITS and β -tubulin sequences. Isolates sequenced in this study are printed in bold. Bar = total nucleotide differences between taxa. Bootstrap values (1000 replicates) are indicated above the branches.

Table 1. Ophiostomatoid fungi, including species with affinities to both the *Microascales* and *Ophiostomatales*, reported from South Africa. Currently accepted species names are listed first, with the name used in the original report in square brackets. Species reported as associates of bark or ambrosia beetles are printed in blue.

Year	Species	Host	References
1927	<i>Sporothrix schenckii</i> Hektoen & C.F. Perkins [= <i>Sporotrichum beurmanii</i> Matr. & Ramond]	Human	Doidge (1950)
1931	<i>Thielaviopsis basicola</i> (Berk. & Broome) Ferraris	<i>Nicotiana tabacum</i>	Gorter (1977)
1937	^a <i>Ophiostoma piliferum</i> (Fr.) Syd. & P. Syd. [= <i>Ceratostomella pilifera</i> (Fr.) G. Winter]	Logs of <i>Pinus radiata</i>	Laughton (1937)
1937	<i>Thielaviopsis paradoxa</i> (De Seynes) Höhn.	<i>Saccharum officinarum</i>	Doidge (1950)
1947	^b <i>Graphium</i> sp. associated with <i>Sporotrichum</i> sp., producing perithecia	Timber and air	Brown et al. (1947)
1956	<i>Ceratocystis adiposa</i> (E.J. Butler) C. Moreau	Shoots of <i>Pinus</i> sp.	Talbot (1956)
1965	<i>Chalara terrestris</i> Agnihothr. & K.C. Barna	<i>Eucalyptus saligna</i>	Marasas et al. (1966)
1970	<i>Raffaelea albimanens</i> D.B. Scott & J.W. du Toit	<i>Platypus extenedentatus ex Ficus sycamorus</i>	Scott & Du Toit (1970)
1970	<i>R. hennebertii</i> D.B. Scott & J.W. du Toit	<i>P. extenedentatus ex F. sycamorus</i>	Scott & Du Toit (1970)
1970	<i>R. arxii</i> D.B. Scott & J.W. du Toit	<i>Xyleborus torquatus ex Cussonia umbellifera</i>	Scott & Du Toit (1970)
1974	<i>Graphium putredinis</i> (Corda) S. Hughes	Soil	Eicker (1974)
1977	<i>Ceratocystis fimbriata</i> Ellis & Halst.	<i>Protea gigantea</i>	Gorter (1977)
1978	<i>Leptographium reconditum</i> Jooste	<i>Triticum</i> rhizosphere	Jooste (1978)
1980	<i>Ophiostoma ips</i> (Rumbold) Nannf. [= <i>Ceratocystis ips</i> (Rumbold) C. Moreau]	<i>Orthotomicus erosus ex Pinus</i> spp.	Wingfield & Marasas (1980a), Zhou et al. (2001, 2004a)
1980	^c <i>Grosmannia serpens</i> Goid. [= <i>Verticildiella alacris</i> M.J. Wingf. & Marasas]	Roots of <i>Pinus pinaster</i> and <i>P. radiata</i>	Wingfield & Marasas (1980b), Zhou et al. (2001)
1983	<i>Leptographium truncatum</i> (M.J. Wingf. & Marasas) M.J. Wingf. [= <i>Verticildiella truncata</i> M.J. Wingf. & Marasas]	Roots of <i>Pinus taeda</i>	Wingfield & Marasas (1983)
1988	<i>Gondwanamyces proteae</i> (M.J. Wingf., P.S. van Wyk & Marasas) Marais & M.J. Wingf. [= <i>Ceratocystiopsis proteae</i> M.J. Wingf., P.S. van Wyk & Marasas]	<i>Protea repens</i>	Wingfield et al. (1988a)
1993	^d <i>Quambalaria eucalypti</i> (M.J. Wingf., Crous & W.J. Swart) J.A. Simpson [= <i>Sporothrix eucalypti</i> M.J. Wingf., Crous & W.J. Swart]	<i>Eucalyptus grandis</i>	Wingfield et al. (1993a)
1993	<i>Gondwanamyces capensis</i> (M.J. Wingf. & P.S. van Wyk) Marais & M.J. Wingf. [= <i>Ophiostoma capense</i> M.J. Wingf. & P.S. van Wyk]	<i>Protea</i> spp.	Wingfield & Van Wyk (1993)
1994	<i>Ophiostoma splendens</i> G.J. Marais & M.J. Wingf.	<i>Protea</i> spp.	Marais & Wingfield (1994)
1994	^e <i>Graphium pseudormiticum</i> M. Mouton & M.J. Wingf.	<i>Orthotomicus erosus</i>	Wingfield et al. (1988b), Mouton et al. (1994b)
1995	<i>Ophiostoma quercus</i> (Georgev.) Nannf.	<i>Olinia</i> sp., <i>Eucalyptus grandis</i> , <i>Quercus robur</i>	De Beer et al. (1995, 2003b)
1996	<i>Ceratocystis albifundus</i> M.J. Wingf., De Beer & M.J. Morris	<i>Protea</i> sp.	Wingfield et al. (1996)
1997	<i>Ophiostoma protearum</i> G.J. Marais & M.J. Wingf.	<i>Protea</i> sp.	Marais & Wingfield (1997)
1999	<i>Ophiostoma stenoceras</i> (Robak) Nannf.	<i>Eucalyptus</i> spp., <i>Acacia mearnsii</i> , <i>Malus</i> sp.	De Beer et al. (1999, 2003a), Zhou et al. (2001)
1999	<i>Ceratocystis radicicola</i> (Bliss) C. Moreau	<i>Phoenix dactylifera</i>	Linde & Smit (1999)
2000	<i>Ophiostoma galeiforme</i> (B.K. Bakshi) Math.-Käärik	<i>P. pinaster</i> , <i>Hylurgus ligniperda</i> ex <i>Pinus elliottii</i>	Zhou et al. (2001, 2004b)
2000	<i>Leptographium procerum</i> (W.B. Kendrick) M.J. Wingf.	<i>Hylastes angustatus</i> ex <i>Pinus</i> sp.	Zhou et al. (2001)
2001	<i>Ophiostoma africanum</i> G.J. Marais & M.J. Wingf.	<i>Protea gaguedi</i>	Marais & Wingfield (2001)
2001	<i>Ceratocystiopsis minuta</i> (Siemaszko) H.P. Upadhyay & W.B. Kendr.	<i>O. erosus</i> , <i>Hylastes angustatus</i> , <i>H. ligniperda</i> ex <i>Pinus</i> spp.	Zhou et al. (2001)
2001	^f <i>Ophiostoma piceae</i> (Münch) Syd. & P. Syd.	<i>H. ligniperda</i>	Zhou et al. (2001)
2001	<i>Ophiostoma pluriannulatum</i> (Hedg.) Syd. & P. Syd.	<i>O. erosus</i> , <i>H. ligniperda</i> , <i>H. angustatus</i> ex <i>Pinus</i> spp.	Zhou et al. (2001)
2001	^g <i>Leptographium lundbergii</i> Lagerb. & Melin	Roots of <i>Pinus taeda</i>	Zhou et al. (2001)
2001	<i>Pesotum</i> spp.	<i>O. erosus</i> , <i>Hylastes angustatus</i> , <i>H. ligniperda</i> ex <i>Pinus</i> spp.	Zhou et al. (2001)
	<i>Sporothrix</i> sp.		
	<i>Hyalorhinocladiella</i> sp.		

Table 1. (Continued).

Year	Species	Host	References
2003	<i>Ophiostoma floccosum</i> Math.-Käärik	<i>Pinus elliottii</i>	De Beer <i>et al.</i> (2003b)
2003	<i>Ceratocystis moniliformis</i> (Hedg.) C. Moreau	<i>Erythrina</i> sp.	Barnes <i>et al.</i> (2003)
2004	<i>Chalara hughesii</i> Nag Raj & W.B. Kendr.	<i>Elegia capensis</i>	Lee <i>et al.</i> (2004)
2004	^b <i>Graphium calicoides</i> (Fr.) Cooke & Massee	<i>Leucadendron</i> sp.	Lee <i>et al.</i> (2004)
2004	<i>Ceratocystis pililliformis</i> I. Barnes & M.J. Wingf.	<i>Eucalyptus grandis</i>	Roux <i>et al.</i> (2004)

^aDoubtful report based on perithecia without ascospores (Laughton 1937). The present study provides the first substantiated report of *O. piliferum* in South Africa.

^bProbably a *Pesotum* sp. with *Sporothrix* synanamorph.

^cOnly the *Leptographium* anamorph of this species has been observed in South Africa.

^dInitially reported as a *Sporothrix* sp., *Q. eucalypti* has now been shown to be a smut in the *Exobasidiomycetes* order *Microstromatales* (De Beer *et al.* 2006).

^eIn 1988 reported as a *Graphium* sp. (Wingfield *et al.* 1998b).

^fThis identification of *O. piceae* was based on morphology and host specificity (Zhou *et al.* 2001). Sequence data in the present study show that this identification was incorrect and that the fungus was *O. quercus*.

^g*Leptographium truncatum* was treated a synonym of *L. lundbergii* (Jacobs & Wingfield 2001); however, Jacobs *et al.* (2005) showed that South African isolates identified as *L. lundbergii* by Zhou *et al.* (2001), have sequences distinct from *L. lundbergii*, and should be treated as *L. truncatum*.

^h*Graphium calicoides* has been placed in the *Chaetothyriales* (black yeasts) (Okada *et al.* 1998).

Coloniae in agar 1.5 % avenae in medio 45 mm diam aetate duarum hebdromadum in 25 °C, laete hyalinae vel albae. Mycelium aerum adest. Ascomata superficialia vel subimmersa in agar 1.5 % avenae. Bases perithiorum globosae, obscurae, 130–220(–350) µm diam, hyphis laete griseis 65–150(–280) µm longis, 1.5–2.0 (–2.5) µm latis ornatae. Colla perithiorum brunnea vel nigra, laevia, 340–800 (1415) µm longa, ad basim 35–42(–58) µm, ad apicem 12–15(–27) µm lata. Hyphae ostiolares adsunt. Ascospores hyalinae, non septatae, allantoidae, in sectione transversali rotundae, 2–3(–3.5) x 1–1.5(–2) µm.

Cellulae conidiogenae micronematae, mononematae, hyalinae, 12–60(–85) x 1.5–2(–2.5) µm, ad apicem incrassatum denticulos acres perferentes; conidia hyalina, unicellularia, clavata vel guttuliformia, 3–4.5(–8) x 1–1.5(–2.5) µm.

Ascomata with globose bases, dark, 130–220(–350) µm diam (Fig. 3A), ornamented with light grey hyphae, 65–150(–280) µm long, 1.5–2(–2.5) µm wide. **Perithecial necks** brown to black, smooth, 340–800 (1415) µm long, 35–42(–58) µm wide at the base, 12–15(–27) µm at the apex (Fig. 3A, B). **Ostolar hyphae** present (Fig. 3B). **Ascospores** hyaline, aseptate, allantoid, round in side view, 2–3(–3.5) x 1–1.5(–2) µm (Fig. 3C).

Conidiogenous cells (Fig. 3D–E), micronematous, mononematous, hyaline, 12–60(–85) x 1.5–2(–2.5) µm, sharp denticles present in the swollen apical part. **Conidia** (Fig. 3F) hyaline, single 1–celled, clavate to guttuliform, 3–4.5(–8) x 1–1.5(–2.5) µm.

Cultural characteristics: Colonies on 1.5 % oatmeal agar reaching on average 45 mm diam in two weeks at 25 °C. Colonies light hyaline to cotton-white. Aerial mycelium present. Perithecia produced superficially on or partially immersed in 1.5 % oatmeal agar.

Substrates: *Hylastes angustatus* and infested bark of *Pinus patula*.

Distribution: Mpumalanga Province, South Africa.

Specimens examined: **South Africa**, Mpumalanga Province, *Hylastes angustatus*, Sep. 1999, X.D. Zhou, **holotype** PREM 58886, culture ex-type CBS 118837 = CMW 19362; paratype PREM 58887,

culture ex-paratype = CMW 19363; paratype PREM 58888, culture ex-paratype CBS 118827 = CMW 19364.

DISCUSSION

Results of this study have confirmed the identities of five *Ophiostoma* spp. associated with the non-native pine-infesting bark beetles *H. angustatus*, *H. ligniperda*, and *O. erosus* in South Africa. These fungi are *O. ips*, *O. floccosum*, *O. plurianulatum*, *O. quercus* and *O. stenoceras*. In addition, *O. abietinum*, *O. piliferum* and *P. fragrans* are recognised for the first time from South Africa. One of the fungi associated with these bark beetles represents an undescribed taxon, for which the name *O. aurorae* has been provided.

The three fungal species *O. abietinum*, *O. piliferum* and *P. fragrans* reported from South Africa for the first time, are well-known associates of conifer timber. *Ophiostoma abietinum* was first described from *Abies vejari* attacked by a *Pseudohylesinus* sp. in Mexico (Marmolejo & Butin 1990), and was considered as an intermediate between *O. stenoceras* and *O. nigrocarpum* (R. Davidson) De Hoog (De Beer *et al.* 2003a). *Ophiostoma piliferum* is considered economically important to the forestry industry, and a colourless mutant of this species has been marketed as biocontrol agent against sapstaining fungi (Farrell *et al.* 1993). *Pesotum fragrans* was described from galleries of *Ips sexdentatus* infesting *Pinus sylvestris* in Sweden (Mathiesen-Käärik 1953), and the species has been reported from Australia, California, Canada, and New Zealand (Harrington *et al.* 2001, Jacobs *et al.* 2003).

Ophiostoma aurorae described in this study is morphologically similar to species in the *O. stenoceras*-complex (De Beer *et al.* 2003a, Aghayeva *et al.* 2004, 2005). Species in the complex have typical orange-

Table 2. Fungal isolates from pine bark beetles in South Africa used in this study.

Species	Isolate Number	GenBank no.	Host	Insect vector	Area
	ITS	β-tubulin			
<i>Pesotum fragrans</i>	^a CMW 19357	DQ396790	<i>Pinus patula</i>	<i>Hylastes angustatus</i>	Mpumalanga
<i>Ophiostoma abietinum</i>	CMW 397	DQ396788	—	<i>Orthotomicus erosus</i>	Western Cape
<i>O. floccosum</i>	CMW 19358	DQ396791	<i>P. elliotii</i>	<i>Hyllurgus ligniperda</i>	Kwazulu-Natal
	CMW 19359	DQ396792	<i>P. patula</i>	<i>O. erosus</i>	Mpumalanga
<i>O. plurianulatum</i>	CMW 19360	DQ396793	<i>P. elliotii</i>	<i>H. ligniperda</i>	Kwazulu-Natal
<i>O. piliferum</i>	CMW 554	DQ396789	—	<i>O. erosus</i>	Western Cape
<i>O. quercus</i>	CMW 19361	DQ396794	<i>P. patula</i>	<i>H. angustatus</i>	Mpumalanga
	CMW 19365	DQ396795	<i>P. elliotii</i>	<i>H. ligniperda</i>	Kwazulu-Natal
<i>O. stenoceras</i>	CMW 544	DQ396799	—	<i>H. angustatus</i>	Western Cape
<i>O. aurorae</i>	CMW 19362	DQ396796	DQ396800	<i>P. elliotii</i>	Mpumalanga
	CMW 19363	DQ396797	DQ396801	<i>P. elliotii</i>	Mpumalanga
	CMW 19364	DQ396798	DQ396802	<i>P. elliotii</i>	Mpumalanga

^aCulture Collection of the Forestry and Agricultural Biotechnology Institute, University of Pretoria, Pretoria, South Africa.**Table 3.** Isolates of selected species of *Ophiostoma* used for comparative purpose in this study.

Species	Strain No.	GenBank no.	Collector / supplier	Origin	Host / insect
	ITS	β-tubulin			
<i>Leptographium guttulatum</i>	^a CMW 1310	AY649782	J.N. Gibbs	England	<i>Pinus / Tomicus piniperda</i>
	^b CMW 742	AY649783	M. Morelet	France	<i>P. sylvestris / Tomicus sp.</i>
<i>Pesotum fragrans</i>	^{b,c} CBS 279.54	AF198248	A. Mathiesen-Käärik	Sweden	<i>P. sylvestris / ips sexdentatus</i>
<i>Ophiostoma abietinum</i>	^b CBS 125.89	AF484453	J.G. Marmolejo	Mexico	<i>Abies vejarii / Pseudohylesinus sp.</i>
<i>O. dentifundum</i>	^b CBS 115790	AY495434	C. Delatour	Hungary	<i>Quercus wood</i>
	CBS 115865	AY495435	T. Kowalski	Poland	<i>Quercus robur</i>
<i>O. floccosum</i>	^b CBS 799.73	AF198231	A. Käärik	Sweden	<i>Picea or Pinus</i>
<i>O. fusiforme</i>	^b CBS 112912	AY280481	D.N. Aghayeva	Azerbaijan	<i>Populus nigra</i>
	CBS 112909	AY280482	AY280462	D.N. Aghayeva	<i>Castanea sativa</i>
<i>O. ips</i>	^b CBS 137.36	AY546704	C.T. Rumbold	U.S.A.	<i>Ips integer</i>
	CMW 6418	AY546702	X.D. Zhou	South Africa	<i>Pinus elliotii / Orthotomicus erosus</i>

Table 3. (Continued).

Species	Strain No.	GenBank no.		Collector / supplier	Origin	Host / insect
		ITS	β -tubulin			
<i>O. lunatum</i>	^b CBS 112927	AY280485	AY280466	T. Kirisits	Austria	<i>Carpinus betulus</i>
	CBS 112928	AY280486	AY280467	T. Kirisits	Austria	<i>Larix decidua</i>
	CBS 357.77			R.W. Davidson	U.S.A.	<i>Pinus</i> sp.
<i>O. narcissi</i>	^c C 1648	AF484451	—	D.P. Limber	U.K.	<i>Narcissus</i> sp.
	^b CBS 138.50	AY194510		R.W. Davidson	Netherlands	<i>Narcissus</i> sp.
<i>O. nigrocarpum</i>	^b CBS 637.66	AY280489	AY280479	R.W. Davidson	U.S.A.	<i>Abies</i> sp.
	CBS 638.66	AY280490	AY280480	R.W. Davidson	U.S.A.	<i>Pseudotsuga menziesii</i>
<i>O. piceae</i>	^b CBS 108.21	AF198226		E. Münch	Germany	<i>Abies</i> or <i>Picea</i>
	CMW 7648	AF493249		D.B. Redfern, J.F. Webber	U.K.	<i>Picea sitchensis</i>
<i>O. piliferum</i>	CBS 129.32	AF221070		H. Diddens	—	<i>Pinus sylvestris</i>
<i>O. plurianulatum</i>	CMW 75			R.W. Davidson	U.S.A.	—
<i>O. pulvinisporum</i>	^b CMW 9020	AY546713		X.D. Zhou	Mexico	<i>P. pseudostrobos</i> / <i>Dendroctonus mexicanus</i>
	CMW 9026	AY546715		X.D. Zhou	Mexico	<i>P. pseudostrobos</i> / <i>Ips calligraphus</i>
<i>O. quercus</i>	^b CMW 2467	AY466626		M. Morelet	France	<i>Quercus</i> sp.
	CMW 7645	AF493246		T. Kirisits, E. Halmschlager	Austria	<i>Q. robur</i>
<i>O. stenoceras</i>	^b CBS 237.32	AY484462	AY280471	H. Robak	Norway	<i>Pinus</i> pulp
	CMW 11193	AY280493	AY280475	R. Farrell	New Zealand	wood
<i>Sporothrix infilata</i>	^b CBS 239.68	AY495426	AY495437	W. Gams	Germany	wheatfield soil
	CBS 841.73	AY495431	AY495442	J. Grinbergs	Chile	soil
<i>S. schenckii</i>	CMW 7612	AY280494	AY280476	H.F. Vismér	South Africa	human sporotrichosis
	CMW 7614	AY280495	AY280477	H.F. Vismér	South Africa	human sporotrichosis
	CMW 7615	AY280496	AY280478	H.F. Vismér	South Africa	human sporotrichosis

aCMW = Culture Collection of the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, South Africa.

bEx-type culture or authentic strain.

cCBS = Culture collection of the Centraalbureau voor Schimmelmilities, Utrecht, Netherlands.
dC = Culture collection of T.C. Harrington, Department of Plant Pathology, Iowa State University, U.S.A.




Fig. 3. *Ophiostoma aurorae* (CMW 19362) on 1.5 % oatmeal agar. A. Perithecium with long neck. (Scale bar = 190 µm). B. Apex of the neck with ostiolar hyphae. (Scale bar = 15 µm). C. Allantoid round ascospores. (Scale bar = 1.5 µm). D. Conidiophore. (Scale bar = 3.5 µm). E. Conidiogenous cell. (Scale bar = 1.5 µm). F. Clavate conidia. (Scale bar = 1.5 µm).

section-shaped ascospores and *Sporothrix* anamorphs. *Ophiostoma aurorae* can be distinguished from other species in the complex by its very obviously rounded ascospores and swollen clavate conidia. Its association with the root feeding scolytid bark beetle *H. angustatus* also appears to be a useful characteristic that might be applied in identification. In addition to its morphologically unique nature, analyses of ITS and partial β-tubulin gene sequences confirmed that *O. aurorae* resides in a phylogenetic clade, distinct from all morphologically similar *Ophiostoma* spp. for which sequence data are available.

Results of this study emphasise that a surprisingly large number of *Ophiostoma* spp. are associated with the three non-native conifer-infesting bark beetles accidentally introduced into South Africa. They also highlight the fact that the introduction of what might initially appear to be a single organism (plant, insect, fungus) is often considerably more complex. It seems likely that most of the fungi treated in this study are specifically associated with the insects in their areas of origin and like their insect vectors, they are also introduced exotics.

Species such as *O. quercus* that have a wide distribution on many woody substrates in South Africa could have invaded the bark beetle niche. It would be interesting to understand the long-term changes in such vector/fungus relationships, as has recently been found

with *Tomicus piniperda* (Linnaeus) and *Leptographium wingfieldii* M. Morelet in the United States (Jacobs et al. 2004). Clearly, the bark beetle/ fungal association represents a complex and dynamic environment that deserves further study.

ACKNOWLEDGEMENTS

We thank the National Research Foundation (NRF), members of Tree Protection Co-operative Programme (TPCP) and the THRIP initiative of the Department of Trade and Industry, South Africa for financial support. We also acknowledge Sappi for a fellowship awarded to the first author and Dr Hugh F. Glen for providing the Latin diagnosis.

REFERENCES

- Aghayeva DN, Wingfield MJ, De Beer ZW, Kirisits T (2004). Two new *Ophiostoma* species with *Sporothrix* anamorphs from Austria and Azerbaijan. *Mycologia* **96**: 866–878.
- Aghayeva DN, Wingfield MJ, Kirisits T, Wingfield BD (2005). *Ophiostoma dentifundum* sp. nov. from oak in Europe, characterized using molecular phylogenetic data and morphology. *Mycological Research* **109**: 1127–1136.
- Barnes I, Roux J, Wingfield BD, Dudzinski MJ, Old KM, Wingfield MJ (2003). *Ceratocystis pirilliformis*, a new species from *Eucalyptus nitens* in Australia. *Mycologia* **95**: 865–871.
- Brown R, Weinrib D, Simpson MW (1947). Timber as a source of sporotrichosis infection. In: *Sporotrichosis infection on Mines of the Witwatersrand*. A symposium. Proceedings of the Transvaal Mine Medical Officers' Association, eds). The Transvaal Chamber of Mines, Johannesburg, South Africa: 5–33.
- De Beer ZW, Begerow D, Bauer R, Pegg GS, Crous PW, Wingfield MJ (2006). Phylogeny of *Quambalariaeaceae* fam. nov., including important *Eucalyptus* pathogens from South Africa and Australia. *Studies in Mycology* **55**: 293–302.
- De Beer ZW, Harrington TC, Vismer HF, Wingfield BD, Wingfield MJ (2003a). Phylogeny of the *Ophiostoma stenoceras*–*Sporothrix schenckii* complex. *Mycologia* **95**: 434–441.
- De Beer ZW, Wingfield BD, Wingfield MJ (2003b). The *Ophiostoma piceae*-complex in the Southern Hemisphere: a phylogenetic study. *Mycological Research* **107**: 469–476.
- De Beer ZW, Wingfield MJ, Kemp GHJ (1995). First report of *Ophiostoma querci* in South Africa. 32nd Annual Congress of the South African Society for Plant Pathology, 23–26 January 1994, Christiana, South Africa. *South African Journal of Science* **91**: vi.
- De Beer ZW, Witthuhn RC, Britz H, Wingfield MJ, Wingfield BD (1999). 18S rDNA sequences place *Ophiostoma* isolates from hardwoods in the southern hemisphere in the species *Ophiostoma stenoceras*. Proceedings of the 37th Annual Congress of the Southern African Society for Plant Pathology, 17–20 January 1999, Pietermaritzburg, South Africa. *South African Journal of Science* **95**: vi.
- Dodge EM (1950). The South African fungi and lichens to the end of 1945. *Bothalia* **5**: 1–1094.
- Eicker A (1974). The mycoflora of an alkaline soil of the open-savannah of the Transvaal. *Transactions British Mycological Society* **63**: 281–288.
- Farrell RL, Blanchette RA, Brush TS, Hadar Y, Iverson S, Krisa K, Wendler PA, Zimmerman W (1993). Cartapip™: a biopulping product for control of pitch and resin acid problems in pulp mills. *Journal of Biotechnology* **30**: 115–122.
- Felsenstein J (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**: 783–791.
- Gardes M, Bruns TD (1993). ITS primers with enhanced specificity for *Basidiomycetes*—application to the identification of mycorrhiza and rusts. *Molecular Ecology* **2**: 113–118.
- Glass NL, Donaldson GC (1995). Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous Ascomycetes. *Applied Environmental Microbiology* **61**: 1323–1330.
- Gorter GMJA (1977). Index of plant pathogens and the diseases they

- cause in cultivated plants in South Africa. *PPRI Science Bulletin* **392**: 177 pp.
- Harrington TC (2005). Ecology and evolution of mycophagous bark beetles and their fungal partners. In: *Insect-Fungal Associations: Ecology and Evolution* (Vega FE, Blackwell M, eds). Oxford University Press, New York: 1–22.
- Harrington TC, Cobb FW (1988). *Leptographium root diseases on conifers*. American Phytopathological Press, St. Paul, Minnesota, U.S.A.
- Harrington TC, McNew D, Steimel J, Hofstra D, Farrell R (2001). Phylogeny and taxonomy of the *Ophiostoma piceae* complex and the Dutch elm disease fungi. *Mycologia* **93**: 111–136.
- Hausner G, Reid J, Klassen GR (2000). On the phylogeny of members of *Ceratocystis* s.s. and *Ophiostoma* that possess different anamorphic states, with emphasis on the anamorph genus *Leptographium*, based on partial ribosomal DNA sequences. *Canadian Journal of Botany* **78**: 903–916.
- Jacobs K, Bergdahl DR, Wingfield MJ, Halk S, Seifert KA, Bright DE, Wingfield BD (2004). *Leptographium wingfieldii* introduced into North America and found associated with exotic *Tomicus piniperda* and native bark beetles. *Mycological Research* **108**: 411–418.
- Jacobs K, Seifert KA, Harrison KJ, Kirisits T (2003). Identity and phylogenetic relationships of ophiostomatoid fungi associated with invasive and native *Tetropium* species (Coleoptera: Cerambycidae) in Atlantic Canada. *Canadian Journal of Botany* **81**: 316–329.
- Jacobs K, Solheim H, Wingfield BD, Wingfield MJ (2005). Taxonomic re-evaluation of *Leptographium lundbergii* based on DNA sequence comparisons and morphology. *Mycological Research* **109**: 1149–1161.
- Jacobs K, Wingfield MJ (2001). *Leptographium species – tree pathogens, insect associates and agents of blue-stain*. American Phytopathological Press, St. Paul, Minnesota, U.S.A.
- Jooste WJ (1978). *Leptographium reconditum* sp. nov. and observations on conidiogenesis in *Verticiladiella*. *Transactions of the British Mycological Society* **70**: 152–155.
- Katoh K, Misawa K, Kuma K, Miyata T (2002). MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**: 3059–3066.
- Kirisits T (2004). Fungal associates of European bark beetles with special emphasis on the ophiostomatoid fungi. In: *Bark and Wood boring insects in living trees in Europe, a synthesis*. (Lieutier F, Day KR, Battisti A, Grégoire JC, Evans HF, eds). Kluwer Academic Press, The Netherlands: 181–235.
- Laughton EM (1937). The incidence of fungal disease on timber trees in South Africa. *South African Journal of Science* **33**: 377–382.
- Lee S, Mel'nik V, Taylor JE, Crous PW (2004). Diversity of saprobic hyphomycetes on *Proteaceae* and *Restionaceae* from South Africa. *Fungal Diversity* **17**: 91–114.
- Linde C, Smit WA (1999). First report of rhizosporosis caused by *Ceratocystis radicicola* on date palms in South Africa. *Plant Disease* **83**: 880.
- Marais GJ, Wingfield MJ (1994). Fungi associated with infructescences of *Protea* species in South Africa, including a new species of *Ophiostoma*. *Mycological Research* **98**: 369–374.
- Marais GJ, Wingfield MJ (1997). *Ophiostoma protearum* sp. nov. associated with *Protea caffra* infructescences. *Canadian Journal of Botany* **75**: 362–367.
- Marais GJ, Wingfield MJ (2001). *Ophiostoma africanum* sp. nov., and a key to ophiostomatoid species from *Protea* infructescences. *Mycological Research* **105**: 240–246.
- Marasas WFO, Van der Westhuizen GCA, Van Warmelo KT, Papendorf MC (1966). New and interesting records of South African fungi. Part V. *Bothalia* **9**: 229–243.
- Marmolejo JG, Butin H (1990). New conifer-inhabiting species of *Ophiostoma* and *Ceratocystiopsis* (Ascomycetes, Microascales) from Mexico. *Sydowia* **42**: 193–199.
- Mathiesen-Käärik A (1953). Eine Übersicht über die gewöhnlichsten mit Borkenkäfern assoziierten Bläuepilze in Schweden und einige für Schweden neue Bläuepilze. *Meddelanden från Statens Skogsforskningsinstitut* **43**: 1–74.
- Mouton M, Wingfield MJ, Van Wyk PWJ (1994). *Graphium pseudomititicum* sp. nov.: a new hyphomycete with unusual conidiogenesis. *Mycological Research* **98**: 1272–1276.
- Okada G, Seifert KA, Takematsu A, Yamaoka Y, Miyazaki S, Tubaki K (1998). A molecular phylogenetic reappraisal of the *Graphium* complex based on 18S rDNA sequences. *Canadian Journal of Botany* **76**: 1495–1506.
- Riddell RW (1950). Permanent stained mycological preparations obtained by slide culture. *Mycologia* **42**: 265–270.
- Roux J, van Wyk M, Hatting H, Wingfield MJ (2004). *Ceratocystis* species infecting stem wounds on *Eucalyptus grandis* in South Africa. *Plant Pathology* **53**: 414–421.
- Saitou N, Nei M (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology Evolution* **4**: 406–425.
- Scott DB, Du Toit JW (1970). Three new *Raffaelea* species. *Transactions of the British Mycological Society* **55**: 181–186.
- Six DL (2003). Bark beetle-fungus symbiosis. In: *Insect Symbiosis* (Bourtzis K, Miller TA, eds). CRC Press, New York: 97–114.
- Spatafora JW, Blackwell M (1994). The polyphyletic origins of ophiostomatoid fungi. *Mycological Research* **98**: 1–9.
- Swofford DL (2002). PAUP*. Phylogenetic Analysis Using Parsimony (*and other methods). Version 4.0b10. Sinauer Associates, Sunderland, Massachusetts.
- Talbot PHB (1956). New and interesting records of South African fungi. Part II. *Bothalia* **6**: 489–500.
- Tribe GD (1992). Colonization sites on *Pinus radiata* logs of the bark beetles, *Orthotomicus erosus*, *Hylastes angustatus* and *Hylurgus ligniperda* (Coleoptera: Scolytidae). *Journal of the Entomological Society of Southern Africa* **1**: 77–84.
- White TJ, Bruns T, Lee S, Taylor J (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR protocols: genes for phylogenetics* (Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds). Academic Press, San Diego, California, USA: 315–322.
- Wingfield MJ, Crous PW, Swart WJ (1993a). *Sporothrix eucalypti* (sp. nov.), a shoot and leaf pathogen of *Eucalyptus* in South Africa. *Mycopathologia* **123**: 159–164.
- Wingfield MJ, De Beer C, Visser C, Wingfield BD (1996). A new *Ceratocystis* species defined using morphological and ribosomal DNA sequence comparisons. *Systematic and Applied Microbiology* **19**: 191–202.
- Wingfield MJ, Marasas WFO (1980a). *Ceratocystis ips* associated with *Orthotomicus erosus* (Coleoptera: Scolytidae) on *Pinus* spp. in the Cape Province of South Africa. *Phytophylactica* **12**: 65–69.
- Wingfield MJ, Marasas WFO (1980b). *Verticiladiella alacris* sp. nov., associated with a root disease of pines in the South Africa. *Transactions of the British Mycological Society* **75**: 21–28.
- Wingfield MJ, Marasas WFO (1983). Some *Verticiladiella* species, including *V. truncata* sp. nov., associated with root diseases of pine in New Zealand and South Africa. *Transactions of the British Mycological Society* **80**: 231–236.
- Wingfield MJ, Seifert KA, Webber JF (1993b). *Ceratocystis* and *Ophiostoma*. *Taxonomy, Ecology and Pathogenicity*. American Phytopathological Society Press, St. Paul, Minnesota, USA.
- Wingfield MJ, Van Wyk PS (1993). A new species of *Ophiostoma* from *Protea* infructescences in South Africa. *Mycological Research* **97**: 709–716.
- Wingfield MJ, Van Wyk PS, Marasas WFO (1988a). *Ceratocystiopsis proteae* sp. nov., with a new anamorph genus. *Mycologia* **80**: 23–30.
- Wingfield MJ, Van Wyk PS, Marasas WFO (1988b). A new *Graphium* species with unusual conidium development. Annual meeting of the South African Society for Plant Pathology. *Phytophylactica* **20**: 103.
- Wood SL, Bright DE (1992). A catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic index. *Great Basin Naturalist Memoirs A* **13**: 1–6.
- Zhou XD, De Beer ZW, Cibrián D, Wingfield BD, Wingfield MJ (2004a). Characterisation of *Ophiostoma* species associated with pine bark beetles from Mexico, including *O. pulvinisporum* sp. nov. *Mycological Research* **108**: 690–698.
- Zhou XD, De Beer ZW, Harrington TC, McNew D, Kirisits T, Wingfield BD, Wingfield MJ (2004b). Epitypification of *Ophiostoma galeiforme* and phylogeny of species in the *O. galeiforme* complex. *Mycologia* **96**: 1306–1315.
- Zhou XD, De Beer ZW, Wingfield BD, Wingfield MJ (2001). Ophiostomatoid fungi associated with three pine-infesting bark beetles in South Africa. *Sydowia* **53**: 290–300.
- Zipfel RD, De Beer ZW, Jacobs K, Wingfield BD, Wingfield MJ (2006). Multi-gene phylogenies define *Ceratocystiopsis* and *Grosmannia* distinct from *Ophiostoma*. *Studies in Mycology* **55**: 77–99.