



Published in final edited form as:

Nat Med. 2010 October ; 16(10): 1152–1156. doi:10.1038/nm.2214.

Cryptochrome Mediates Circadian Regulation of cAMP Signaling and Hepatic Gluconeogenesis

Eric E. Zhang^{1,2,*}, Yi Liu^{3,5,*}, Renaud Dentin³, Pagkapol Y. Pongsawakul¹, Andrew C. Liu^{1,2,6}, Tsuyoshi Hirota^{1,2}, Dmitri A. Nusinow¹, Xiujie Sun⁵, Severine Landais³, Yuzo Kodama⁴, David A. Brenner⁴, Marc Montminy^{3,†}, and Steve A. Kay^{1,†}

¹ Section of Cell & Developmental Biology, Division of Biological Sciences, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0130, USA

² Genomics Institute of the Novartis Research Foundation, 10675 John Jay Hopkins Drive, San Diego, CA 92121, USA

³ The Salk Institute for Biological Studies, 10010 North Torrey Pines Road, La Jolla, CA 92037, USA

⁴ School of Medicine, University of California San Diego, 9500 Gilman Drive, La Jolla, CA 92093, USA

⁵ Key Laboratory of Nutrition and Metabolism, Institute for Nutritional Sciences, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, 294 Taiyuan Road, Shanghai 200031, P.R. China

⁶ Department of Biological Sciences, The University of Memphis, 3774 Walker Avenue, Memphis, TN 38152, USA

Abstract

During fasting, mammals maintain glucose homeostasis by stimulating hepatic gluconeogenesis¹. Elevations in circulating glucagon (GLU) and epinephrine trigger the cAMP mediated phosphorylation of Creb and dephosphorylation of the Creb coactivator Crtc2². Although the underlying mechanism is unclear, hepatic gluconeogenesis is also regulated by the circadian clock, which coordinates glucose metabolism with changes in the external environment^{3–6}. Here we show that Creb activity during fasting is modulated by Cryptochromes (Cry1 and Cry2), core components of the clock that are rhythmically expressed in the liver. Cry was elevated during the night/day transition, when it reduced fasting gluconeogenic gene expression by blocking GLU-

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence and requests for materials should be addressed to SAK (skay@ucsd.edu) or MM (montminy@salk.edu).

*These authors contributed equally to this work.

Author Contributions

EEZ and SAK conceived the project. EEZ, YL, MM and SAK designed the research. EEZ, YL, RD, PYP, ACL, TH, DAN, XS, SL and YK performed the experiments. EEZ and YL analyzed the data. EEZ, YL, DAB, MM and SAK wrote the paper.

Author Information

All mouse works were conducted under regulations of Institutional Animal Care and Use Committee at Salk, GNF and CAS. The authors declare that there is no competing financial interest with this work. Reprints and permissions information is available at <http://npg.nature.com/reprintsandpermissions>.

mediated increases in intracellular cAMP concentrations and in the PKA-mediated phosphorylation of Creb. In biochemical reconstitution studies, we found that Cry inhibited accumulation of cAMP in response to G protein coupled receptor (GPCR) activation but not to forskolin, a direct activator of adenylyl cyclase. Cry appeared to modulate GPCR activity directly through interaction with G α . As hepatic over-expression of Cry lowered blood glucose concentrations and improved insulin sensitivity in insulin resistant *db/db* mice, our results suggest that compounds which enhance Cry activity may provide therapeutic benefit to individuals with type II diabetes.

We used an adenovirally encoded CRE:Luciferase reporter to evaluate the rhythmicity of hepatic Creb activity under fasting conditions⁷. Intraperitoneal GLU administration (GLU-IP) increased Ad-CRE:Luc activity 40-fold better in mice fasted at *Zeitgeber* time 10–13 (ZT10–13, the day-to-night transition) than at ZT22–1, corresponding to the night-to-day transition (Fig. 1a and Supplementary Fig. 1). Consistent with a rhythmic oscillation in Creb activity, GLU re-injection to the same mice fasted at ZT6–9, near the day-to-night transition, caused a robust increase in hepatic CRE:Luciferase reporter activity. These results suggest that the amplitude of hepatic Creb activation by fasting is gated rhythmically.

Based on the changes in fasting CRE:Luc reporter activity, we hypothesized that the circadian clock modulates Creb and *Crtc2* effects on the gluconeogenic program. Supporting this idea, GLU administration increased gluconeogenic gene expression (*G6Pase*, *Pck1*) to a greater extent in mice fasted at ZT10–13 (Fig. 1b). Consistently, GLU-IP administration increased hepatic Creb phosphorylation and *Crtc2* de-phosphorylation robustly in ZT10–13 fasted mice, but less in ZT22–1 fasted mice (Fig. 1c). Together, these results indicate that the clock modulates fasting gluconeogenesis in part by regulating Creb-*Crtc2* activities. These observations are consistent with recent reports showing that the phosphorylation of Creb is rhythmic *in vivo*⁸.

Owing to the rhythmic Creb phosphorylation, we considered that the liver clock may exert global effects on the expression of cAMP inducible genes. In line with this idea, we identified 194 Creb targets among a set of 652 clock-controlled genes (CCGs) in published microarray databases of mouse liver^{9–11}. We plotted the sum of the Creb targeted CCGs, along with their ratio compared to the total CCGs at each time-point, versus circadian time (CT) based on their peak expression. Both the number and ratio curves show a clear rhythmic pattern with depletion of Creb targets at CT22 (Supplementary Fig. 2), suggesting that Creb targets are generally rhythmic in the liver with the trough phase coinciding with that of *Per2*.

We evaluated the extent to which rhythmic gating of Creb activity is cell autonomous. Exposure to vasoactive intestinal peptide (VIP) increased CRE:Luc activity in mouse adult fibroblasts harboring the receptor *Vipr2*¹² (Fig. 2). After synchronizing the cells by serum shock, exposure to VIP at specific times resulted in CRE:Luc reporter activation that showed a rhythmic pattern, with peak (24, 48 h) and trough (36–39 h) activities coinciding with those for a *Per2*:Luciferase reporter (Fig. 2a). These results show the regulation of Creb pathway induction by cell autonomous circadian clock, and are reminiscent of the circadian

gating seen in the light signaling induction of a PER2::LUC reporter in melanopsin-containing fibroblasts¹³.

Because CRE:Luc activity in liver and in synchronized fibroblasts is lowest when Per-Cry activity is highest (ZT22), we tested the potential role of Cry in regulating Creb signaling. Over-expression of *Cry1* or *Cry2* in *Vipr2*/CRE:Luc fibroblasts or HEK293T cells attenuated CRE:Luc activity in cells exposed to VIP (Fig. 2b–d), while *Per1* did not show significant effect (Fig. 2c). The repression was dosage-dependent (Fig. 2d). Consistent with the absence of E-boxes within the CRE:Luc reporter, Clock-Bmal1 did not activate CRE:Luc expression (Supplementary Fig. 3)¹⁰. Furthermore, *Cry1/2* overexpression reduced effects of urocortin-3 on CRE:Luc activity in cells expressing Corticotropin-releasing factor receptor 2 β ₁₄ (Supplementary Fig. 4). Thus, these data indicate that Cry may exert more general effects on GPCR-mediated increases in Creb-Crtc2 dependent transcription.

Based on its inhibitory effects on GPCR signaling in fibroblasts, Cry may correspondingly modulate GLU effects on liver gene expression. Supporting this idea, hepatic expression of adenovirally encoded Cry1 (Ad-Cry1) reduced GLU-induced CRE:Luc activity in liver of fasted mice at ZT10–13, when endogenous *Cry1* expression is low (Fig. 3a). Moreover, Ad-Cry1 also down-regulated gluconeogenic gene expression (*G6Pase*, *Pck1*) in fasted, GLU-IP administered mice (Fig. 3b). Consequently, Ad-Cry1 expressing mice exhibited lower circulating blood glucose concentrations compared to Ad-GFP controls (Fig. 3c).

We tested whether depletion of Cry, either by targeted disruption of *Cry1* and *Cry2* genes¹⁵, or by RNAi-mediated knockdown in the liver, was sufficient to increase Creb activity. By contrast with the circadian oscillation of hepatic CRE:Luc induction in wild-type mice, GLU-stimulated CRE:Luc activity was constitutively elevated in *Cry1*^{-/-}*Cry2*^{-/-} animals (Supplementary Fig. 5). Similarly, RNAi-mediated depletion of hepatic *Cry1/2* increased GLU-stimulated CRE:Luc activity at ZT1 relative to control animals expressing unspecific RNAi (USi, Fig. 3d). As a result, hepatic mRNA amounts for gluconeogenic genes (*G6Pase*, *Pck1*) were increased in *Cry1/2*-knockdown mice compared to controls (Fig. 3e). Circulating glucose concentrations were elevated following *Cry* depletion (Supplementary Fig. 6), suggesting that circadian changes in hepatic *Cry* expression are sufficient to modulate the gluconeogenic program in part through their inhibitory effects on the Creb-Crtc2 pathway. Indeed, knockdown of *Cry1/2* also increased glucose production in primary hepatocytes following exposure to GLU, indicating that Cry effects on hepatic glucose metabolism are likely cell autonomous (Fig. 3f).

We wondered whether Cry would similarly modulate circulating glucose concentrations in the setting of insulin resistance. Supporting this idea, adenoviral expression of *Cry1* reduced fasting blood glucose concentrations and improved whole body insulin sensitivity in insulin resistant *db/db* mice, as measured by glucose tolerance testing (Fig. 3g). Pointing to an effect on hepatic gluconeogenesis, *Cry1* over-expression also lowered glucose excursions in pyruvate tolerance testing studies (Fig. 3h).

Based on the rhythmic changes in hepatic Creb phosphorylation, we suspected that Cry1 may disrupt effects of GLU on adenylyl cyclase (AC) activation during fasting (Fig. 1c).

Supporting this idea, GLU-IP administration increased hepatic cAMP concentrations at ZT10–13, when *Cry1* expression is low, and to a lesser extent at ZT22–1, when *Cry1* levels are elevated (Fig. 4a). Consistently, Ad-Cry1 expression reduced hepatic cAMP concentrations in ZT10–13 fasted mice exposed to GLU (Fig. 4b). Indeed, amounts of phosphorylated Creb were down-regulated in livers of fasted Ad-Cry1 mice compared to GFP control animals (Fig. 4c).

We evaluated effects of Cry on Creb-Crtc2 signaling in cultures of primary hepatocytes. Similar to its effects in liver, Ad-Cry1 also down-regulated CRE:Luc reporter activity, as well as gluconeogenic gene expression in cells exposed to GLU (Supplementary Fig. 7a,b). By contrast with its effect in cells exposed to GLU, Ad-Cry1 did not reduce gluconeogenic gene expression or CRE:Luc reporter activity in cells treated with forskolin (FSK), a direct activator of AC. Consistent with these results, Ad-Cry1 expression reduced cAMP accumulation in cells exposed to GLU but not to FSK (Fig. 4d). In addition, RNAi-mediated knockdown of *Cry1* and 2 increased cAMP content in primary hepatocytes exposed to GLU but not to FSK (Supplementary Fig. 7c). Pointing to a more general effect of this repressor, *Cry1/2* over-expression also reduced cAMP accumulation in response to VIP treatment in fibroblasts expressed *Vipr2* (Supplementary Fig. 8). Together, these results indicate that Cry1 acts upstream of AC to regulate hepatic Creb activity during fasting.

To evaluate further the mechanism, we performed *in vitro* reconstitution assays. Addition of isoproterenol (ISO) to plasma membranes from HEK293T cells, which express β -adrenergic receptors, increased cAMP accumulation *in vitro* (Fig. 4e,f). Co-incubation of plasma membrane fractions with cytoplasmic extracts from transfected HEK293T cells expressing *Cry1* potently inhibited cAMP increases in response to ISO relative to control (GFP) extracts. We considered that Cry1 could disrupt cAMP accumulation either directly by binding to relevant signaling components, or indirectly by stimulating the expression of an intra-cellular inhibitor. Supporting the former hypothesis, immuno-depletion of HA-tagged Cry1 with HA-specific antiserum restored cAMP accumulation in response to ISO, whereas control IgG had no effect (Fig. 4f). Taken together, these results demonstrate that Cry1 inhibits activation of the Creb-Crtc2 pathway in liver by disrupting cAMP production in response to ligand-dependent activation of Gs coupled receptors.

Based on its ability to disrupt GPCR but not FSK-mediated increases in cAMP production, Cry would be predicted to associate with either GPCRs or with the G α subunit of heterotrimeric G protein. Although it did not interact detectably with the GPCR family member *Vipr2*, Cry associated with G α in pull-down assays of membrane fractions from transfected HEK293T cells incubated with recombinant glutathione-S-transferase (GST)-Cry protein (Fig. 4g). We also observed the Cry-G α interaction in reciprocal co-immunoprecipitation assays of HEK293T cells and primary hepatocytes (Fig. 4h). Together, these data demonstrate a role for extra-nuclear Cry in regulating cAMP production.

The circadian clock has been known to modulate hepatic gluconeogenesis for over 20 years, although the underlying mechanism has remained unclear. We found that the liver clock regulates the gluconeogenic program through Cry-mediated inhibition of Creb activity during fasting. Historically regarded as a transcriptional repressor, Cry appeared to inhibit

gluconeogenic gene expression primarily by blocking GPCR mediated increases in cAMP and in the subsequent phosphorylation of Creb (Fig. 4i). These results are consistent with recent findings that hepatic Creb phosphorylation oscillates in a circadian fashion⁸. Similar to our observations in liver, cAMP concentrations have also been found to oscillate in the clock master organ, the suprachiasmatic nucleus, reaching a nadir when Cry expression is highest¹⁶. Future studies should provide further insight in to the detailed mechanism by which Cry modulates Gs effects on cAMP in liver and other tissues.

Methods

Adenoviruses and animals

We delivered adenoviruses (1×10^8 p.f.u.) Cry1, control GFP, Cry1i, Cry2i, or control USi into 8–10-week-old male C57BL/6J-Tyrc-2J/J (Jackson Laboratories) mice together with CRE:Luc (1×10^8 p.f.u.) and RSV β -gal (5×10^7 p.f.u.) by tail- vein injection. Targeted sequences for knockdown were: 5'-GGAAATTGCTCTCAAGGAAGT-3' (Cry1i) and 5'-GCTGAATTCGCGTCTGTTTGT-3' (Cry2i). All mice were housed in colony cages with 12 h light-dark cycle for 4 weeks before study.

In vivo Imaging

We fasted mice beginning at ZT10 or ZT22 and injected intraperitoneally (IP) with GLU (100 mg kg^{-1} ; Sigma) at ZT13 or ZT1, respectively, and continued fasting for 1 h. At the end of fasting (ZT14 or ZT2), we injected IP with 50 mg kg^{-1} Nembutal (Abbott Laboratories) and 100 mg kg^{-1} sterile firefly D-luciferin (Biosynth AG), and imaged them on the IVIS-100 Imaging System within 15–45 min post-injection of luciferin, and analyzed with Living Image software (Xenogen)^{7,17}.

In vivo Analysis

We sonicated mouse tissues at 4 °C, centrifuged, and reserved supernatants for β -gal activity, protein determinations, and immunoblotting analysis. Blood glucose values were determined using a LifeScan automatic glucometer.

Cell Culture

We cultured cells in DMEM with 10% FBS and transfected with plasmid DNA by using lipofectamine2000. Primary hepatocytes were isolated and cultured as described⁷.

Luminometry and luciferase assay

We grew cells harboring Vipr2 and CRE:Luc reporter to confluence in 35 mm dishes, followed by a 2 h serum-shock and released to serum-free medium. We applied synthetic VIP (2 nM in saline, American Peptide) to different dishes 15–48 h post-serum shock at 3 h intervals. We collected data in a luminometer (Actimetrics). Luciferase reporter assay using HEK293T cells was described¹⁸.

mRNA Analysis

We extracted total RNA from whole liver or primary hepatocytes using the RNeasy kit (Qiagen), and measured mRNA levels as reported⁷.

Glucose production assay

We prepared primary hepatocytes as described¹⁸. Briefly, we seeded 5×10^5 cells per well in a 6-well dish with M199 medium (Invitrogen). Adenoviruses infected (M.O.I. of 5 for each CRY1 and 10 for control US1) for 120 min prior to additional incubation for 42–45 h. We treated cells with glucagon ($10 \mu\text{g ml}^{-1}$, Sigma) for 5 h, washed, and incubated with glucose production buffer (saline supplemented with 20 mM sodium lactate and 2 mM sodium pyruvate) for additional 2 h. We collected the supernatant for measuring glucose concentration (BioVision), and normalized the readings with whole-cell protein amount.

Glucose Tolerance Tests (GTT)

We fasted mice overnight and injected IP with glucose (1 g kg^{-1} body weight, Sigma). We measured blood glucose levels before the injection, and at 15, 30, 45, 60, and 90 min after injection.

Pyruvate Tolerance Tests (PTT)

We fasted mice for 24 h and injected IP with sodium pyruvate (2 g kg^{-1} , Sigma). We measured blood glucose before the injection, and at 20, 40, 60, and 100 min after injection.

Cell fractionation

We collected and washed HEK293T cells, and resuspended them in lysis buffer (10 mM Tris-HCl, pH 7.4, 1 mM EDTA, and proteinase inhibitors), and centrifuged at $500 \times g$ for 5 min. Supernatants were ultracentrifuged at $40,000 \times g$ for 30 min. The resulting supernatant was allocated as cytosol fraction and the pellet was resuspended in extraction buffer (75 mM Tris-HCl, pH 7.4, 1 mM EDTA, 10% glycerol and 12.5 mM MgCl_2 , and proteinase inhibitors) as membrane fraction. We measured protein concentrations, adjusted to $1 \mu\text{g}\mu\text{l}^{-1}$, and stored them in -80°C freezer.

Adenyl cyclase activity assay

We mixed equal volume of membrane and cytosol fractions, incubated at 37°C for 5 min, and then mixed with one volume reaction buffer (5 mM Tris-HCl, pH 7.4, 17.5 mM MgCl_2 , 120 μM ATP, 3 μM GTP, 1.5 μM FAD, 60 mM phosphocreatine and 0.375 Unit Creatine phosphokinase) in the presence or absence of isoproterenol (100 μM) or FSK (1 μM) for reaction at 37°C for 30 min, stopped by mixing with HCl and neutralized by NaOH. We measured cAMP levels by ELISA kit (R&D). For the pre-clearance of HA-Cry1, cytosol fractions from Ad-HA-Cry1 infected cells were immuno-precipitated by HA-specific antibody or mouse IgG beads at 4°C for 1 h.

GST pull-down

We prepared GST-Cry1 and control GST-GFP proteins with Glutathione-Sepharose beads (GE Healthcare) as described¹⁹. We dissolved membrane pellets in pull-down buffer (10

mM Tris-HCl, pH 7.4, 1 mM EDTA, 150 mM NaCl, 2mM Dithiothreitol, and proteinase inhibitors), and mixed with 10 μ g GST proteins. After 2 h incubation at 4° C and washes, we suspended pellets in SDS loading buffer, and applied 1% input samples for quantification.

Immunoprecipitation

We conducted immunoprecipitation assays using whole cell lysates.

Statistical analyses

We performed all studies at least three independent occasions. We reported results as mean and s.e.m., and considered differences statistically significant when $P < 0.05$.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank L. Vera, A. Luzader, E. Rodrigo, and X. Li for adenoviral injections, J. Altarejos and M. Lindstrom for mouse blood collection and glucose measurement, and N. Goebel for preparing primary hepatocytes. We also thank M. Yamout and P. E. Wright at The Scripps Research Institute for reagents, and N. Gekakis, D. Welsh and S. Wang for reading the manuscript. This is manuscript #080809 of Genomics Institute of the Novartis Research Foundation, and was supported in part by grants from US National Institutes of Health (R01 GM074868 and R01 MH051573 to SAK; R01 DK083834 and R01 DK049777 to MM). SAK is a co-founder of ReSet Therapeutics and is a member of its Scientific Advisory Board. YL is supported by “One Hundred Talents” Program of Chinese Academy of Sciences (No. 2010OHTP08) and Shanghai Pujiang Program (No. 10PJ1411200) in China.

References

1. Nordlie RC, Foster JD, Lange AJ. Regulation of glucose production by the liver. *Annual Review of Nutrition*. 1999; 19:379–406.
2. Sreaton RA, et al. The CREB coactivator TORC2 functions as a calcium- and cAMP-sensitive coincidence detector. *Cell*. 2004; 119:61–74. [PubMed: 15454081]
3. Green CB, Takahashi JS, Bass J. The meter of metabolism. *Cell*. 2008; 134:728–742. [PubMed: 18775307]
4. Kida K, et al. The circadian change of gluconeogenesis in the liver. *The Journal of Biochemistry*. 1980; 88:1009–1013. [PubMed: 7451400]
5. Gelling RW, et al. Lower blood glucose, hyperglucagonemia, and pancreatic β cell hyperplasia in glucagon receptor knockout mice. *Proceedings of the National Academy of Sciences of the United States of America*. 2003; 100:1438–1443. [PubMed: 12552113]
6. Ramsey KM, Marcheva B, Kohsaka A, Bass J. The clockwork of metabolism. *Annual Review of Nutrition*. 2007; 27:219–240.
7. Dentin R, et al. Insulin modulates gluconeogenesis by inhibition of the coactivator TORC2. *Nature*. 2007; 449:366–369. [PubMed: 17805301]
8. Vollmers C, et al. Time of feeding and the intrinsic circadian clock drive rhythms in hepatic gene expression. *Proceedings of the National Academy of Sciences*. 2009; 106:21453–21458.
9. Panda S, et al. Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell*. 2002; 109:307–320. [PubMed: 12015981]
10. Ueda HR. A transcription factor response element for gene expression during circadian night. *Nature*. 2002; 418:534–539. [PubMed: 12152080]
11. Zhang X, et al. Genome-wide analysis of cAMP-response element binding protein occupancy, phosphorylation, and target gene activation in human tissues. *Proceedings of the National Academy of Sciences*. 2005; 102:4459–4464.

12. Aton SJ, Colwell CS, Harmar AJ, Waschek J, Herzog ED. Vasoactive intestinal polypeptide mediates circadian rhythmicity and synchrony in mammalian clock neurons. *Nat Neurosci.* 2005; 8:476–483. [PubMed: 15750589]
13. Pulivarthy SR, et al. Reciprocity between phase shifts and amplitude changes in the mammalian circadian clock. *Proceedings of the National Academy of Sciences.* 2007; 104:20356–20361.
14. Li C, Chen P, Vaughan J, Lee KF, Vale W. Urocortin 3 regulates glucose-stimulated insulin secretion and energy homeostasis. *Proceedings of the National Academy of Sciences.* 2007; 104:4206–4211.
15. van der Horst GTJ, et al. Mammalian Cry1 and Cry2 are essential for maintenance of circadian rhythms. *Nature.* 1999; 398:627–630. [PubMed: 10217146]
16. O'Neill JS, Maywood ES, Chesham JE, Takahashi JS, Hastings MH. cAMP-dependent signaling as a core component of the mammalian circadian pacemaker. *Science.* 2008; 320:949–953. [PubMed: 18487196]
17. Dentin R, Hedrick S, Xie J, Yates J III, Montminy M. Hepatic glucose sensing via the CREB coactivator CRTC2. *Science.* 2008; 319:1402–1405. [PubMed: 18323454]
18. Sato TK, et al. Feedback repression is required for mammalian circadian clock function. *Nat Genet.* 2006; 38:312–319. [PubMed: 16474406]
19. Sanada K, Harada Y, Sakai M, Todo T, Fukada Y. Serine phosphorylation of mCRY1 and mCRY2 by mitogen-activated protein kinase. *Genes to Cells.* 2004; 9:697–708. [PubMed: 15298678]




Figure 1. Hepatic Creb:Crtc2 activity is modulated by the circadian clock

(a) CRE:Luciferase reporter activity in mice fasted for 3 hours followed by intra-peritoneal (IP) GLU administration. Relative effect of fasting at ZT10–13 and ZT22–1 is indicated. Right, bar graph shows CRE:Luc activity from hepatic lysates normalized to beta galactosidase activity from co-infected Ad-RSV: β -gal adenovirus. Asterisks indicate $P < 0.01$, $n = 5$. (b) Q-PCR analysis of gluconeogenic gene expression (*G6Pase*, left; *Pck1*, right) in mice fasted at ZT10–13 or ZT22–1 and then injected IP with GLU. Asterisks indicate $P < 0.01$, $n = 5$. (c) Immunoblot of Creb and Crtc2 proteins in liver extracts from mice fasted ZT10–13 or ZT22–1 followed by IP with GLU. Relative effects ZT13 and ZT1 fasting on hepatic amounts of phospho-Crtc2 and phospho-Creb (Ser133) indicated.




Figure 2. Cry inhibits Creb activity

(a) CRE:Luciferase reporter activity in *Vipr2*-expressing mouse fibroblasts synchronized by serum shock and then exposed to VIP at indicated times. Replicate samples (a total of 24) received VIP treatment every 3 h post-synchronization from 15–48 h. For clarity, one responsive curve per time point was shown in different colour. Mouse fibroblasts harbouring *Per2*:dLuc (green dashed lines) or *Bmal1*:dLuc reporters (red dashed lines) were used to report the two distinct phases of clock gene expression (scale at right). Data are representative of three experiments. (b) Effect of Cry over-expression on CRE:Luc reporter activity. *Vipr2*/CRE:Luc cells were infected with lentiviral particles expressing either GFP, Cry1 or Cry2. Cells were treated with VIP 24 h after medium change (red arrow). (c–d) Cry represses CRE:Luc induction in transfected cells expressing *Vipr2*. CRE:Luc (20 ng) and *Vipr2* (40 ng) expression plasmids were co-transfected into HEK 293T cells along with *Per1*, Cry1 or Cry2 (40 ng) as indicated. 24 h post transfection, cells were treated with VIP

(1 nM) for 8 h to induce CRE:Luc activity. N.S., not significance; Asterisks indicate $P < 0.01$ analyzed by student *t*-test.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 3. Cry blocks induction of the gluconeogenic program by Creb and Crtc2

(a) Left, live imaging analysis of hepatic CRE:Luc reporter activity in control (Ad-GFP) and Ad-Cry1 expressing mice. CRE:Luc activity in fasted mice (ZT10–13) injected with GLU is shown. Right, quantitative analysis of CRE:Luc activity. Asterisks indicate $P < 0.01$, $n = 5$.

(b) Q-PCR analysis of gluconeogenic gene expression (*G6Pase*, *Pck1*) in control and Cry1 expressing fasted and fed mice. Asterisks indicate $P < 0.05$, $n = 5$.

(c) Effect of Cry1 over-expression on fasting blood glucose concentrations relative to control GFP expressing mice. $P < 0.05$, $n = 3$.

(d) Left, live imaging analysis of CRE:Luc reporter activity in mice expressing adenovirally encoded Cry1/2 or unspecific RNAs (USi) in liver. Right, quantitative analysis of CRE:Luc activity. Asterisk indicates $P < 0.01$, $n = 4$.

(e) Q-PCR analysis of hepatic *G6Pase* and *Pck1* mRNA amounts in control (USi) and CRYi expressing mice. $P < 0.001$ for *G6Pase* and $P < 0.01$ for *Pck1*, $n = 3$.

(f) Glucose production assay in primary hepatocytes infected with Ad-CRYi or control. Asterisks indicate $P < 0.01$, $n = 3$.

(g) Glucose tolerance testing of *db/db* mice expressing adenovirus encoded Cry1 or GFP. Asterisks indicate $P < 0.01$, $n = 5$.

(h) Pyruvate tolerance testing of WT mice expressing

adenovirus-encoded Cry1 or GFP. Fasted mice were injected with pyruvate and blood glucose concentrations were measured at indicated times. Asterisks indicate $P < 0.05$, $n = 5$.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4. Cry inhibits GPCR-dependent increases in adenylyl cyclase activity

(a) Hepatic cAMP concentrations in fed or fasted mice following IP injection of GLU as indicated. (b) Effect of Cry1 over-expression on hepatic cAMP levels in fasted mice following IP GLU administration. (c) Immunoblot showing effect of Ad-Cry1 expression on hepatic amounts of phospho-Creb (Ser133) in fed and fasted mice. (d) Effect of Adenoviral Cry1 expression on intracellular cAMP concentrations in hepatocytes exposed to FSK or GLU. (e-f) Effect of Cry1 and Cry2 on cAMP accumulation. *In vitro* reconstitution studies showing effect of cytosolic fractions from control (GFP) and Cry1 expressing cells on cAMP accumulation in reactions containing plasma membrane fractions from HEK293T cells. (g) Immunoblot showing recovery of Gsα and Vipr2 from membrane fractions of transfected HEK293T cells incubated with recombinant GST- Cry1 or GST-GFP control. Upper panels: input and pull-down signals using 293T cells overexpressed with Flag-Gsα, Vipr2-V5, Flag-Per1, and Flag-Luciferase (LUC), respectively. Lower panel: GST-tagged proteins were purified and stained with Coomassie Brilliant Blue (CBB). Asterisk points to truncated GST-Cry1 polypeptide. (h) Co-immunoprecipitation of Gsα and Cry1. Upper panels: Immunoblot showing recovery of HA-Cry1 from immunoprecipitates of Flag-Gsα prepared from HEK293T cells. Asterisks point to non-specific signals. Lower panels: Immunoblot showing recovery of endogenous Gsα from immunoprecipitates of HA-Cry1

prepared from primary mouse hepatocytes. (i) Circadian regulation of cAMP signaling in liver is Cry-dependent. Blue oval represents E-box bound transcription factors CLOCK-BMAL1, and green oval represents CRE-mediated transcription activators or co-activators such as Creb, p300/Cbp and Crtc2.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript