BMC Cancer

CORRECTION   **Open Access**

Correction to: A novel coordination complex of platinum (PT) induces cell death in colorectal cancer by altering redox balance and modulating MAPK pathway

Khayal Al-Khayal[1], Mansoor-Ali Vaali-Mohammed[1], Mohammed Elwatidy[2], Thamer Bin Traiki[1], Omar Al-Obeed[1], Mohammad Azam[3], Zahid Khan[4], Maha Abdulla[1] and Rehan Ahmad[1*]

Correction to: BMC Cancer 20, 685 (2020)
https://doi.org/10.1186/s12885-020-07165-w

Following publication of the original article [1], the authors identified a typesetting error in Fig. 2, whereby the full figure was not published. The complete Fig. 2 is published in this correction article and the original article [1] has been corrected.

Author details
[1]Colorectal Research Chair, Department of Surgery, King Saud University College of Medicine, PO Box 7805 (37), 11472 Riyadh, Saudi Arabia. [2]College of Medicine Research Center, King Saud University College of Medicine, Riyadh 11472, Saudi Arabia. [3]Department of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia. [4]Genome Research Chair, Department of Biochemistry, College of Science, King Saud University, Riyadh, Saudi Arabia.

Published online: 01 September 2020

Reference
1. Al-Khayal K, Vaali-Mohammed M, Elwatidy M, et al. A novel coordination complex of platinum (PT) induces cell death in colorectal cancer by altering redox balance and modulating MAPK pathway. BMC Cancer. 2020;20:685 https://doi.org/10.1186/s12885-020-07165-w.

---

* Correspondence: arehan@ksu.edu.sa
[1]Colorectal Research Chair, Department of Surgery, King Saud University College of Medicine, PO Box 7805 (37), 11472 Riyadh, Saudi Arabia
Full list of author information is available at the end of the article

Al-Khayal *et al. BMC Cancer*   (2020) 20:834

Page 2 of 2

Fig. 2 PT induces apoptosis and cell cycle arrest. **a** HT-29 **b** SW480 **c** SW620 cells were treated with 5 and 10 μM of PT for 24 h. Total cell death including apoptosis and necrosis was analyzed by Annexin V/ PI staining using flow cytometry. **d** HT-29 and **e** SW620 cell cycle distribution was measured by PI staining using flow cytometry and the percentage of cell population was determined in the G0/G1, S and G2/M phases. Results shown are representative of three independent experiment ($n = 3$). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ vs control