


Does the Application of International Classification of Disease Codes for the Cause of Death on Death Certificates Reduce Garbage Codes?

INQUIRY: The Journal of Health Care Organization, Provision, and Financing
Volume 59: 1–9
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/00469580221081433
journals.sagepub.com/home/inq


Soobeom Park, MD¹ and Sun Hyu Kim, MD, PhD¹ 

Abstract

Introduction: This study aimed to determine if applying International Classification of Diseases (ICD) disease codes directly as the cause of death (COD) on death certificates (DCs) instead of writing or typing the COD could reduce the use of garbage codes.

Methods: Beginning in April of 2016, a documentation process change was made, retiring the process of hand-writing or typing the COD onto DCs to directly applying ICD disease codes that were registered during the patient's course of treatment. The DCs issued at the emergency department (ED) 1 year before (Pre-code group) and after (Code group) applying ICD disease codes directly on DCs was instituted were retrospectively analyzed. The occurrence of garbage codes along with other major and minor errors was compared between the two groups. The investigation and judgment of errors were performed by four emergency physicians.

Results: The overall garbage code occurrence in the Code group (25%) was significantly lower than that in the Pre-code group (49%). Fewer garbage codes were used in the Code group with an average of .5 in the Pre-code group and .3 in the Code group. No significant difference was identified in major error occurrences except for in the garbage codes. Minor errors were more common in the Pre-code group than in the Code group.

Conclusion: The overall use of garbage codes on DCs could be reduced by changing the process by which physicians complete DCs, that is, the application of documenting ICD disease codes directly as the COD on DCs.

Keywords

death certificates, cause of death, emergency service, disease codes

¹Department of Emergency Medicine, University of Ulsan College of Medicine, Ulsan University Hospital, Dong-gu Ulsan, Korea

Corresponding Author:

Sun Hyu Kim, MD, PhD, Department of Emergency Medicine, University of Ulsan College of Medicine, Ulsan University Hospital, 877 Bangeojinsunhwando-ro, Dong-gu Ulsan 44033, Korea.

E-mail: stachyl@paran.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and

Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

What Do We Already Know About This Topic?

It is known that there were many errors in death certificates, including the use of garbage codes, which could not be used as an underlying cause of death. Reducing garbage codes is important to improving the quality of mortality statistics.

How Does Your Research Contribute to the Field?

Garbage codes can be reduced by applying disease codes as the cause of death, which can improve the quality of death certificate statistics.

What Are Your Research's Implications Toward Theory, Practice, or Policy?

To improve the quality of death statistics, the quality of death certificates should be improved. As one of the methods, if disease codes are applied to death certificates as the cause of death, the quality of death certificates can be improved by reducing the use of garbage codes.

Introduction

A death certificate (DC) is a legal record of a person's cause of death (COD). The information from a DC is collected for COD statistics, which is essential in determining the direction of healthcare policies.¹⁻⁴ Since the 1990s, researchers in the Global Burden of Diseases (GBD) study have tried to analyze the COD data derived from DCs. Despite the researchers' efforts, the validity and quality of COD data remain questionable in many countries.⁵

In 1996, Murray et al introduced the term "garbage codes" to describe diagnoses that could not be used as an underlying COD. Various studies are being conducted to classify and reduce these garbage codes to improve the quality of COD statistics.^{5,6} Although there have been studies such as educational interventions and online learning programs on how to complete a DC, and the development of algorithms to redistribute these garbage codes to improve COD data,^{3,6-10} garbage codes are still found in about 30% of all DCs.¹¹ Since garbage codes impact healthcare policy establishment, attempts should be made to reduce these errors.

Establishing a well-functioning COD registration system is challenging for a number of reasons. Physicians occasionally document CODs outside of accordance with the standardized rules of the International Classification of Diseases (ICD) proposed by the World Health Organization (WHO). These physicians utilize non-standardized terms in the COD section on a DC, often because they are handwritten or typed in by the attending physician. This informal documentation of the COD on DCs then goes through a process called coding, which converts the descriptive terms into diagnoses that suit the ICD standard. Lu et al¹² identified that the most frequent type of disagreement between the reviewers and coders involved COD nomenclature. Physicians tend to complete DCs with diagnoses familiar to them in clinical practice, without fully understanding how the coding system works. Due to these challenges, garbage codes occur frequently, making the COD statistics less reliable. To the best of our knowledge, there is a lack of studies focusing on

changing ways to complete DCs to overcome these challenges.

This study aimed to reduce garbage codes by suggesting alternative methods to complete DCs. In this study, we compared DCs that were completed by typing in the COD with those completed by selecting the COD directly from an ICD coding system and evaluated the occurrence of garbage codes.

Methods

This was a retrospective analysis study, which included DCs issued in the emergency department of Ulsan University Hospital, a university training hospital in Korea. It was reviewed by an institutional review board.

The CODs on the DCs were handwritten or typed by the attending physicians until March of 2016 in Ulsan University Hospital, where this study was conducted. Beginning in April of 2016, a documentation process change was made, retiring the process of hand-writing or typing the COD onto DCs to directly applying ICD disease codes that were registered during the patient's course of treatment (Figure 1). The disease codes that were registered during the treatment of the deceased were based on the ICD-10 published by the WHO. It was made possible that if the physician disagreed with the selected disease code or simply wanted to type in the cause, it could be deleted and typed in. However, typing in the COD was possible only after deleting the COD that was selected from the registered disease codes, so the attending physicians could not type in the COD from the beginning.

In this study, we analyzed DCs that were issued in the emergency department at Ulsan University Hospital in the years 2015 and 2017, one year before and after the 2016 documentation process change for disease codes used for the COD on DCs was applied. The DCs issued in the emergency department of the university hospital in 2015 (Pre-code group), when the COD was typed in directly, were compared with the ones issued in 2017 (Code group), when the COD was selected from the registered disease codes. This study included the DCs of the patients that died during

A

Cause of Death	A	Cause of death	Ⓐ	Time interval	
	B	Cause of A	Ⓐ		
	C	Cause of B	Ⓐ		
	D	Cause of C	Ⓐ		
	Other significant conditions not associated with to from A to D				
	Major findings of surgery			Date of surgery	YYYY/MM/DD
	Major findings of autopsy				

B

Cause of Death	A	Cause of death	Ⓐ	Time interval	
	B	Cause of A	Ⓐ		
	C	Cause of B	Ⓐ		
	D	Cause of C	Ⓐ		
	Other significant conditions not associated with to from A to D				
	Major findings of surgery			Date of surgery	YYYY/MM/DD
	Major findings of autopsy				

ⓑ

외래	2022-01-24	R57.1	Hypovolemic shock
	2022-01-24	K25.01B	Acute stomach ulcer (peptic) with hemorrhage
입원	2022-01-24	K25.01B	Acute stomach ulcer (peptic) with hemorrhage
외래	2022-01-24	C22.0A	Malignant neoplasm of hepatocellular carcinoma
입원	2022-01-24	C22.0A	Malignant neoplasm of hepatocellular carcinoma

Figure 1. Electronic record input for the COD section in DCs in the Pre-code group (A). The initial COD section in the Code group (B) was the same as in the Pre-code group. In the Pre-code group, a physician typed in the COD after clicking (a). In the Code group, upon clicking (a), a popup window containing ICD-10 codes appeared (b), and a physician could search for and select the suitable COD code.

treatment at the hospital with relatively clear CODs. Autopsy certificates issued after death and DCs of patients who died of unknown causes were excluded from the study. DCs of patients who were dead on arrival were also excluded. Among the total of 192 DCs in 2015 (Pre-code group) and 152 DCs in 2017 (Code group), 50 cases and 26 cases, respectively, were excluded from the study (Figure 2). The power for the sample sizes of 142 of in the Pre-codes group and 126 in the Code group was .99.

The garbage codes, introduced by Murray and Lopez, and further classified by Naghavi et al into four categories, were as follows.¹¹

1. Type 1: Causes that cannot or should not be considered as an underlying cause of death (UCOD).
2. Type 2: Intermediate causes of death without an underlying cause such as septicemia or pulmonary hypertension.




Figure 2. Study subjects.

Table 1. The definition of errors on death certificates.

Type of Error	Definition or Examples
Garbage codes	
Type 1	Causes that cannot or should not be considered UCODS
Type 2	Intermediate CODs without an underlying cause such as septicemia or pulmonary hypertension
Type 3	Immediate CODs, such as cardiac arrest and respiratory failure
Type 4	Insufficiently specified causes within ICD chapters
Major errors	
Improper sequence	Listing hemorrhagic shock as the cause of esophageal varix bleeding
Incompatible causal relationship	Listing DM as the cause of cerebral hemorrhage
≥2 causes of death on a single line	
Incorrect manner of death	Wrong judgment for the manner of death such as natural cause or external cause
Unacceptable cause of death	Unacceptable COD based on medical records
Minor errors	
Blank/duplication	A blank line between CODs or the same COD duplicated
No record for date of onset	No record for date of onset
Incorrect date of onset	Incorrect record of date of onset
Incorrect time interval	Incorrect or no record of time interval
Incorrect other significant conditions	Incorrect or no record of other significant conditions
Incorrect type of accident	Incorrect or no record for type of accident
Incorrect intention of external cause	Incorrect or no record for intention
Incorrect time of accident	Incorrect or no record for time of accident
Incorrect place of accident	Incorrect or no record of place of accident
Absence of accident mechanism Only the accident mechanism as COD	No record of accident mechanism Only the accident mechanism recorded as COD

UCOD: underlying cause of death; COD: cause of death; ICD: International Classification of Diseases; DM: diabetes mellitus.

- Type 3: Immediate causes of death such as cardiac arrest or respiratory failure.
- Type 4: Insufficiently specified causes of death.

Errors other than garbage codes were also divided into major and minor errors, as described in several previous studies.^{4,7,13-17} The major errors were as follows. (1) Improper sequence of time between the CODs. (2) Incompatible

causal relationship between two or more CODs. (3) Two or more CODs recorded in a single COD. (4) Incorrect manner of death (misunderstanding an external cause of disease). (5) Unacceptable COD based on the medical records. The minor errors were as follows. (1) A blank line between the COD or listing the same COD repeatedly. (2) The type of accident (ICD-10 codes V01-Y89) was recorded correctly, with no record of the mode of injury (ICD-10 code S00-T98). (3) Incorrect record of the type of accident as the UCOD. (4) No record of the date of onset. (5) Incorrect date of onset. (6) Incorrect or no record of the time interval for each COD. (7) Incorrect or no record of other significant conditions. (8) Incorrect or no record of the type of accident. (9) Incorrect intention of the accident. (10) Incorrect or no record of the date of the accident. (11) Incorrect or no record of the location of the accident (Table 1).

The number and type of garbage codes were investigated along with the number of CODs recorded on the DC, the number of major and minor errors, and the total number of errors. The judgment of errors was first conducted by three individual resident-physicians who had received education on certifying death and ICD-10 coding, and were currently practicing in the ED. Then, a senior emergency physician who had research experience on DCs performed a secondary review process. The senior emergency physician and the residents then performed a final consensus process together for CODs that needed further determinations. CODs that were hard to determine even after the final consensus process were to be excluded from the study, but no cases were excluded from this study. The independent variables in this study were Pre-code group and Code group, and the rest of the variables were dependent.

Statistical Analyses

Frequency analysis, the chi-squared test, Fisher's exact test, and the Student's t-test were used to compare the errors including garbage codes in the Pre-code group and the Code group. IBM SPSS 24.0 (IBM, Armonk, NY, USA) was used for the statistical analyses, and statistical significance was defined as $P < .05$.

Results

A total of 142 cases in the Pre-code group and 129 cases in the Code group met the inclusion criteria for the study. The average age was 63 years in the Pre-code group, and 62 in the Code group. Among the deaths, 57% were male in the Pre-code group and 68% in the Code group. Of the deaths, 76% in the Pre-code group and 73% in the Code group were disease-related, with no significant difference between the two groups. Of all the CODs, 60% of the CODs in the Pre-code group had a single COD, whereas 52% of the Code group had a single COD. There were 2 or more CODs in 40% and 48% of the Pre-Code and Code groups, respectively.

The overall garbage code occurrence in the Code group (25%) was significantly lower ($P = .000$) than that in the Pre-code group (49%). The total number of errors was 2.5 in the Code group, which was less than 3.5 in the Pre-code group ($P = .000$). There were fewer garbage codes in the Code group with an average of .500 in the Pre-code group and .278 in the Code group ($P = .001$). No significant difference was identified in major error occurrences except for the garbage codes ($P = .074$). Minor errors were more common in the

Table 2. Characteristics of death certificates according to the application of disease codes for the cause of death.

Characteristics	Pre-code (n=142)	Code (n=126)	P-value
Age, years old	62.9±21.7	61.9±18.8	.698
Sex, male (%)	81 (57.0)	86 (68.3)	.059
Manner of death (%)			.511
Disease-related	108 (76.1)	92 (73.0)	
External cause	33 (23.2)	34 (27.0)	
Number of lines used for cause of death, mean±SD	1.6±0.8	1.6±0.7	.993
Number of lines used for cause of death, n (%)			.020
One	85 (59.9)	66 (52.4)	
Two	37 (26.1)	51 (40.5)	
Three or more	20 (14.1)	9 (7.1)	
Errors of DC, n (%)			
Garbage codes	69 (48.6)	31 (24.6)	.000
Major errors	25 (17.6)	14 (9.3)	.132
Minor errors	141 (99.3)	109 (86.5)	.000
Total number of errors on death certificate	3.5±1.6	2.5±1.7	.000
Garbage codes	.500	.278	.001
Major errors	.197	.111	.074
Minor errors	2.789	2.103	.000

Table 3. Garbage codes on death certificates according to the application of disease codes for the cause of death.

Garbage Codes	Pre-code (n= 142)	Code (n=126)	P-value
Type 1	17 (12.0%)	2 (1.6%)	.001
Type 2	18 (12.7%)	18 (14.3%)	.700
Type 3	36 (25.4%)	14 (11.1%)	.003
Type 4	0 (.0%)	1 (.4%)	.288

Pre-code group than in the Code group (2.789 vs 2.103, $P = .000$) (Table 2).

Type 1 and type 3 garbage codes were found to be more frequent in the Pre-code group than in the Code group (12.0% vs 1.6%, $P = .001$), (25.4% vs 11.1%, $P = .003$). The occurrence of Type 2 and type 4 garbage codes was not significantly different between the Pre-code group and the Code group (12.7% vs 14.3%, $P = .700$ and .0% vs .4%, $P = .288$, respectively) (Table 3).

Three of the minor error types were reduced after the application of disease codes: incorrect time intervals, incorrect other significant conditions, and only the mechanism of an accident as the COD (Table 4).

Discussion

Garbage codes were found in 48.6% of the Pre-code group, which was similar to previous studies where errors related to the cause of death were found in about 48–64% of all DCs.^{4,18,19} In contrast, garbage codes in the Codes group were decreased to 24.6%.

Causes that cannot or should not be considered a UCOD, classified as a type 1 garbage code, usually occur mainly in elderly patients, where the definitive cause of death is difficult to identify, and were found on up to 10% of all DCs in previous studies.^{6,13} In this study, type 1 garbage codes decreased from 12% in the Pre-code group to 1.6% in the Code group. Senility was the most frequent cause of death accounting for type 1 garbage codes. According to Lee et al,²⁰ senility accounted for 5% of all causes of death in Korea. A diagnosis such as senility, which is not a valid cause of death, is of no use from the point of view of the cause of death statistics. There were two cases in which type 1 garbage codes were used as the cause of death in the Code group, both of which had been typed in and not selected from the ICD disease code list. It is not impossible to select senility as the cause of death from the ICD disease codes list, but it seems plausible that the process of looking up the ICD disease code list might have helped in selecting a valid disease code during the documentation process.

Intermediate CODs without an underlying cause such as septicemia or pulmonary hypertension, classified as type 2 garbage codes, ranged from 22% to 30% of all DCs in previous studies.^{3,10,21} There was no difference in the incidence of type 2 garbage codes between the Pre-code group and the Code group, as they were found in 12.7% and 14.3%, respectively.

The application of disease codes alone did not reduce the occurrence of type 2 garbage codes since intermediate CODs usually occurred due to reasons such as poor training on death certification or limited diagnostic equipment.^{6,11} In settings that do not lack diagnostic equipment, like in this study, educational interventions could be helpful in reducing the use of type 2 garbage codes as shown in the studies of Myers et al,⁷ Lakkireddy et al,⁸ and Wood et al.¹⁰

The immediate causes of death such as cardiac arrest or respiratory failure, classified as type 3 garbage codes, ranged from 5.8% to 34% on all DCs in previous studies.^{1,4,18,22} The incidence of type 3 garbage codes decreased from 25.4% in the Pre-code group to 11.1% in the Code group. Cardiac arrest was the most frequent reason for type 3 garbage codes, which occurred when the time to conduct diagnostic tests to accurately determine the patient's condition was limited. There were cases in both groups where the immediate cause of death had to be used because the UCOD could not be determined. However, as Schuppener et al³ proposed, the majority of the errors on DCs were due to the omission of known conditions or illogical sequencing of known events leading to death. Making an effort to determine the UCOD through sufficient review of the patient's medical history and test results remains crucial.

Insufficiently specified causes of death, classified as type 4 garbage codes, ranged from 4.5% to 19% in previous studies.^{1,3,18} In this study, type 4 garbage codes were not found in the Pre-code group, and there was only one case of a type 4 garbage code in the Code group. The rate of type 4 garbage codes in this study was significantly lower than in previous studies. This is because, unlike other studies, we did not consider the lack of a specific cancer site (eg, not specifying which lobe of the lung or which part of the colon) or not specifying the occluded artery in a stroke, problematic. According to Naghavi et al,⁹ these diagnoses are likely to be confined to a single disease category, having much less impact on healthcare policy implications. Listing lung cancer as the COD without specifying its specific location would still be assigned as death by lung cancer, similar to stroke death reporting.

In this study, a type 4 garbage code was found in the death of a lung cancer patient with brain metastasis. The diagnosis of secondary cancer, not lung cancer, was used as the COD. When receiving outpatient and inpatient treatment, the secondary cancer code, not lung cancer, was continuously selected, and even when the patient died, the most recently registered secondary cancer code was selected as the COD on the DC. If applying disease codes as the COD becomes common, physicians should always give careful consideration when selecting the COD to avoid making this kind of mistake.

As identified in previous studies,^{14,15} in this study, the greater the number of CODs listed, the more errors were found. Major and minor errors were found in 3.3% and 92.1% in DCs with a single COD, 20.5% and 93.2% in DCs with two CODs, and 55.2% and 100% in DCs with three or more CODs, respectively.

Table 4. Errors other than garbage codes as the underlying cause of death on death certificates according to the application of disease codes for the cause of death.

	Pre-code (n=142)	Code (n=126)	P-value
Major errors, n (%)	25 (17.6)	14 (11.1)	.132
Improper sequence	2 (1.4)	0 (0)	.500
Incompatible causal relationship	4 (2.8)	0 (0)	.125
≥2 cause of death on a single line	14 (9.9)	6 (4.8)	.113
Incorrect manner of death	3 (2.1)	3 (2.4)	1.000
Unacceptable cause of death	6 (4.2)	5 (4.0)	.916
Minor errors, n (%)	141 (99.3)	109 (86.5)	.000
Blank/duplication	11 (7.7)	11 (8.7)	.770
No record for date of onset	59 (41.5)	39 (31.0)	.072
Incorrect date of onset	26 (18.3)	22 (17.5)	.856
Incorrect time interval	133 (93.7)	98 (77.8)	.000
Incorrect other significant conditions	105 (73.9)	63 (50.0)	.000
Incorrect type of accident	6 (4.2)	3 (2.4)	.508
Incorrect intention of external cause	10 (7.0)	3 (2.4)	.092
Incorrect time of accident	4 (2.8)	5 (4.0)	.043
Incorrect place of accident	10 (7.0)	7 (5.6)	.618
Absence of accident mechanism	15 (10.6)	6 (4.8)	.078
Only accident mechanism as COD	6 (4.2)	0 (0)	.031

UCOD: underlying cause of death; COD: cause of death.

The number of GCs and the number of CODs written did not show a statistically significant relationship. Major errors were found in 17.6% and 11.1% of the Pre-code and Code groups, respectively ($P = .132$). Although the results failed to show statistical significance, the number of major errors tended to decrease in the Code group. The most common type of major error was listing more than two CODs on a single line. It would be challenging to choose the correct cause of death when two or more causes of death are recorded in a single line, which could lead to a decrease in the quality of the COD data.

As interest in COD statistics increases, research on garbage codes is being conducted. In many cases, methods to improve the quality of COD statistics by analyzing garbage codes that have already occurred are being studied. Reclassification methods, where a physician evaluates the medical records and test results to define the COD along with redistribution, which is a process of defining the UCOD of the DCs containing garbage codes based on statistical methods or analyses, have been conducted.^{6,20,23} In addition, as electronic coding systems for the causes of death have been introduced in many countries, studies that use automated coding systems such as IRIS to analyze the COD are showing promising results.^{24,25} However, since these attempts rely on the data of the DCs that have already been completed, it is necessary to increase the accuracy of DCs in advance. In addition to educational interventions on completing a DC, which were shown to be effective in several previous studies, the change in the documentation method of completing DCs attempted in this study is expected to enable physicians to certify death more accurately.

In Korea, medical doctors, dentists, and oriental medicine doctors have the authority to complete death certificates. If a death certificate is issued with a cause of death that does not require an autopsy and has no legal problems, the death certificate is registered at the administrative office based on the death certificate, and after a final review by the National Statistical Office, it is finally registered in the national death statistics. If the cause of death is an external cause or an autopsy is required, the death certificate is issued after an investigation by the police and prosecutors, and the subsequent process is the same. Despite the researchers' efforts, the validity and quality of COD data remain questionable in many countries including Korea. Chang et al and Park et al reported disappointing results regarding the validity of DCs issued in Korea.^{14,15} Although many types of errors in DCs are made by physicians every day, there is no official education or feedback program in Korea concerning how to write a death certificate.

This study analyzed DCs issued by a single training hospital's emergency department and there is a limitation on generalizing the results of this study since DC forms vary from institution to institution and the pattern of errors may be different. Furthermore, this study did not include the degree of education related to the issuer of the DC, which could have a significant impact on errors on the DC. However, even though no education related to issuing DCs was provided before and after the application of disease codes as the COD, the incidence of garbage codes decreased. It is possible that applying disease codes for the COD, aroused physicians' attention on certifying death, which led to the decreased use of garbage codes.

Conclusions

In this study, we found that the overall garbage code usage could be reduced by a relatively simple method of changing the process by which physicians complete DCs, that is, the application of documenting ICD disease codes directly as the COD on DCs. If the effectiveness is proven through continuous research related to the application of disease codes for the COD in an environment that can register the disease code according to the ICD classification, this method could be worth referencing as a model for reducing garbage codes, and the overall positive impact this could have on healthcare policies. With this result, it is important to select the correct ICD disease codes before writing the DC. If incorrect ICD disease codes are selected, the expected positive effect of reducing errors in the DC by applying ICD disease codes will be reduced. Therefore, education on selecting the correct ICD disease codes should also be carried out.

Acknowledgments

The author(s) thanks to Dr Byeong Ju Park and Dr Sangyeop Jung, resident of emergency medicine in Ulsan University Hospital for their contribution of data collection.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Sun Hyu Kim, MD, PhD  <https://orcid.org/0000-0002-4836-0940>

References

- Lu TH, Shau WY, Shih TP, Lee MC, Chou MC, Lin CK. Factors associated with errors in death certificate completion. A national study in Taiwan. *J Clin Epidemiol*. March 2001;54(3):232-238. doi:10.1016/s0895-4356(00)00299-7
- Mieno MN, Tanaka N, Arai T, et al. Accuracy of death certificates and assessment of factors for misclassification of underlying cause of death. *J Epidemiol*. 2016;26(4):191-198. doi:10.2188/jea.JE20150010
- Schuppener LM, Olson K, Brooks EG. Death certification: errors and interventions. *Clin Med Res*. March 2020;18(1):21-26. doi:10.3121/cm.2019.1496
- Filippatos G, Andriopoulos P, Panoutsopoulos G, et al. The quality of death certification practice in Greece. *Hippokratia*. Jan-Mar 2016;20(1):19-25.
- Lopez AD, Murray CC. The global burden of disease, 1990-2020. *Nat Med*. November 1998;4(11):1241-1243. doi:10.1038/3218
- Naghavi M, Makela S, Foreman K, O'Brien J, Pourmalek F, Lozano R. Algorithms for enhancing public health utility of national causes-of-death data. *Popul Health Metrics*. May 2010;8:9. doi:10.1186/1478-7954-8-9
- Myers KA, Farquhar DR. Improving the accuracy of death certification. *CMAJ (Can Med Assoc J)*. May 19 1998;158(10):1317-1323.
- Lakkireddy DR, Basarakodu KR, Vacek JL, et al. Improving death certificate completion: a trial of two training interventions. *J Gen Intern Med*. April 2007;22(4):544-548. doi:10.1007/s11606-006-0071-6
- Naghavi M, Richards N, Chowdhury H, et al. Improving the quality of cause of death data for public health policy: are all 'garbage' codes equally problematic? *BMC Med*. March 9 2020;18(1):55. doi:10.1186/s12916-020-01525-w
- Wood KA, Weinberg SH, Weinberg ML. Death certification in northern alberta: error occurrence rate and educational intervention. *Am J Forensic Med Pathol*. March 2020;41(1):11-17. doi:10.1097/paf.0000000000000527
- Mikkelsen L, Richards N, Lopez A. *Redefining 'garbage Codes' for Public Health Policy: Report on the Expert Group Meeting, 27-28 February 2017*. CRVS Technical Outcome Series Melbourne, Australia: Bloomberg Philanthropies Data For Health Initiative, Civil Registration And Vital Statistics Improvement, University of Melbourne; 2018.
- Lu TH, Lee MC, Chou MC. Accuracy of cause-of-death coding in Taiwan: types of miscoding and effects on mortality statistics. *Int J Epidemiol*. April 2000;29(2):336-343. doi:10.1093/ije/29.2.336
- Maharjan L, Shah A, Shrestha KB, Shrestha G. Errors in cause-of-death statement on death certificates in intensive care unit of Kathmandu, Nepal. *BMC Health Serv Res*. November 12 2015;15:507. doi:10.1186/s12913-015-1168-6
- Park CH, Kim SH. Errors of death certificate for poisoning related death. *Journal of The Korean Society of Clinical Toxicology*. 2020;18(1):11-17.
- Chang JH, Kim SH, Lee H, Choi B. Analysis of errors on death certificate for trauma related death. *Journal of Trauma and Injury*. 2019;32(3):127-135.
- Alipour J, Karimi A, Hayavi Haghighi MH, Hosseini Teshnizi S, and Mehdipour Y. Death certificate errors in three teaching hospitals of Zahedan, southeast of Iran. [published online ahead of print August 6, 2020]. *Death Stud*: 1-9. doi:10.1080/07481187.2020.1801893
- Alipour J, Payandeh A. Common errors in reporting cause-of-death statement on death certificates: A systematic review and meta-analysis. *J Forensic Leg Med*. August 2021;82:102220. doi:10.1016/j.jflm.2021.102220
- Katsakiori PF, Panagiotopoulou EC, Sakellaropoulos GC, Papazafropoulou A, Kardara M. Errors in death certificates in a rural area of Greece. *Rural Rem Health*. October-December 2007;7(4):822.
- Cambridge B, Cina SJ. The accuracy of death certificate completion in a suburban community. *Am J Forensic Med Pathol*. September 2010;31(3):232-235. doi:10.1097/PAF.0b013e3181e5e0e2

20. Lee YR, Kim YA, Park SY, Oh CM, Kim YE, Oh IH. Application of a modified garbage code algorithm to estimate cause-specific mortality and years of life lost in Korea. *J Kor Med Sci*. November 2016;31 (Suppl 2):S121-s128. doi:[10.3346/jkms.2016.31.S2.S121](https://doi.org/10.3346/jkms.2016.31.S2.S121)
21. Pritt BS, Hardin NJ, Richmond JA, Shapiro SL. Death certification errors at an academic institution. *Arch Pathol Lab Med*. November 2005;129(11):1476-1479. doi:[10.5858/2005-129-1476-dceaaa](https://doi.org/10.5858/2005-129-1476-dceaaa)
22. Smith Sehdev AE, Hutchins GM. Problems with proper completion and accuracy of the cause-of-death statement. *Arch Intern Med*. January 22 2001;161(2):277-284. doi:[10.1001/archinte.161.2.277](https://doi.org/10.1001/archinte.161.2.277)
23. França EB. Garbage codes assigned as cause-of-death in health statistics. *Rev Bras Epidemiol*. November 28 2019; 22(Suppl 3):e19001. supl.3. Códigos garbage declarados como causas de morte nas estatísticas de saúde. doi:[10.1590/1980-549720190001.supl.3](https://doi.org/10.1590/1980-549720190001.supl.3)
24. Harteloh P. The implementation of an automated coding system for cause-of-death statistics. *Inf Health Soc Care*. January 2020; 45(1):1-14. doi:[10.1080/17538157.2018.1496092](https://doi.org/10.1080/17538157.2018.1496092)
25. Martins RC, Buchalla CM. Automatic coding and selection of causes of death: an adaptation of Iris software for using in Brazil. *Rev Bras Epidemiol*. October-December 2015;18(4): 883-893. doi:[10.1590/1980-5497201500040016](https://doi.org/10.1590/1980-5497201500040016)