



Not all that glitters is gold

Marco Anile¹, Francesco Ferrante¹, Beatrice Zacchini¹, Angelina Pernazza², Massimiliano Bassi¹, Jacopo Vannucci³, Federico Venuta¹

¹Department of Thoracic Surgery, University of Rome Sapienza, Rome, Italy; ²Department of Radiology, Oncology and Pathology, University of Rome Sapienza, Rome, Italy; ³Department of Thoracic Surgery, University of Perugia, Perugia, Italy

Correspondence to: Marco Anile, MD, PhD. Department of Thoracic Surgery, University of Rome Sapienza, Viale del Policlinico, 155, 00161 Rome, Italy. Email: marco.anile@uniroma1.it.

Comment on: Li D, Deng C, Wang S, *et al.* Ten-Year Follow-up Results of Pure Ground-Glass Opacity-Featured Lung Adenocarcinomas After Surgery. *Ann Thorac Surg* 2023;116:230-7.

Keywords: Ground-glass opacities (GGOs); lymphadenectomy; surgery

Submitted Jan 06, 2024. Accepted for publication Mar 08, 2024. Published online Apr 12, 2024.

doi: 10.21037/jtd-24-38

View this article at: <https://dx.doi.org/10.21037/jtd-24-38>

Lung cancer (LC) is the leading cause of cancer-related death worldwide with an overall 5-year survival of 15–20%, mostly due to a delayed diagnosis encountering an elevated percentage of advanced stages (1). For this reason, in the last years, huge and onerous efforts have been made in different countries to develop computed tomography (CT) screening programs aiming at improving the detection of early-stage LC. A complementary result of this policy has been the identification of a higher number of potential cancers presenting as different forms of ground-glass opacities (GGOs) (2). They are radiologically defined as focal areas of slightly increased CT attenuation through which the normal lung parenchyma structures, airways, and vessels are visually preserved. According to several papers (3,4), the lung nodules can be simply classified using the consolidation-to-tumor ratio (CTR) obtaining three groups: pure GGOs (non-solid tumors) with CTR =0, part-solid nodules with CTR between 0 and 1, and solid nodules with CTR =1. This radiological classification has been proven to have an impact on prognosis showing a strong correlation between an increased CTR and poorer survival (5). Pure GGOs are a subset of lesions ranging from benign findings to adenocarcinoma *in situ* or minimally invasive adenocarcinoma (MIA) up to, in some circumstances, invasive adenocarcinoma. These findings imply that a correct timing and an appropriate management of these lung nodules are crucial to achieve satisfactory long-term

results. The paper by Li *et al.* (6) published in August 2023 in the *Annals of Thoracic Surgery* highlights these aspects showing excellent results about long-term follow-up in a “subgroup of 308 patients with pure GGOs”. In particular, the 10-year recurrence-free survival was 100% and the 10-year overall survival of 96.9%; furthermore, more than 60% of operative procedures were sublobar resections, with a higher prevalence of wedge resections and a quarter of patients were affected by invasive cancers. At the first glance, the strength of the numbers (308 patients with pure GGOs), a long-term follow-up (10 years), and the impressive survival rates could persuade us that this paper represents a milestone in the management of GGOs. However, not all that glitters is gold...

In fact, the paper reveals some criticisms suggesting further analysis. First, the taxonomy is crucial: “pure GGO” is well defined as a nodule with CTR =0 meaning that no solid part must be present in the nodule. Travis *et al.* (7) in 2016 showed in a table the comparison between clinical and pathological features of small lung adenocarcinoma (<3 cm) with ground glass and lepidic component. In this table, it is clearly reported that in lepidic predominant adenocarcinomas (LPAs) the solid part is constantly present as like as in case of acinar predominant adenocarcinomas (APAs) or papillary predominant adenocarcinomas (PPAs). Thus, this population (21.6%) might be excluded from the analysis because it has not a “pure GGO” lesion.

Furthermore, we should focus also on MIAs because in these lesions a solid component is radiologically often present as reported by Travis; sometimes, it is related to zones of organizing pneumonia, but in other circumstances it could be the sign of the cancer progression and in this study more the 40% of patients had a MIA.

The second point is related to the window time of surgery. As reported by authors, at first detection of a pure GGO nodule their policy is to follow it and to operate only in case of slowly and progressive growth. This could be an acceptable strategy, but it is not clear the timing of this follow-up or if some radiological guidelines are considered. Also, it is not clear which is the diameter cut-off that makes to prefer a sublobar resection respect to a lobectomy. Furthermore, no mention is dedicated to lymph node management; it is true that some recent studies (8,9) have shown that in GGO lesions a systematic lymph node dissection could have no impact on prognosis, but (I) the study population of this paper is antecedent [2007–2013] of these trials and, thus, it is not justified an absence of any form of lymph node assessment and (II) when frozen section is not performed (as in this study), the risk of an invasive adenocarcinoma must be considered and a correct intraoperative staging must be always obtained (10).

Finally, as noted by the authors the population study represents a subgroup of patients with already known favorable prognosis (high percentage of female gender and never smokers). This could make difficult to compare these results with less homogenous and selected patients worldwide and to attend same survival rates in other countries.

Although in clinical studies the number of the population and a long-term follow-up are two powerful factors, we have always to keep in mind that other parameters as methodology, accuracy of the data, correct classification, and management are crucial to leave an adequate take-home message.

Acknowledgments

We acknowledge Martina Anile (Department of Political Sciences, University of Rome Sapienza) for the linguistic revision of the manuscript.

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned

by the editorial office, *Journal of Thoracic Disease*. The article has undergone external peer review.

Peer Review File: Available at <https://jtd.amegroups.com/article/view/10.21037/jtd-24-38/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://jtd.amegroups.com/article/view/10.21037/jtd-24-38/coif>). F.V. serves as an unpaid editorial board member of *Journal of Thoracic Disease* from August 2023 to July 2025. The other authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Siegel RL, Miller KD, Wagle NS, et al. Cancer statistics, 2023. *CA Cancer J Clin* 2023;73:17-48.
2. Matsunaga T, Suzuki K, Takamochi K, et al. What is the radiological definition of part-solid tumour in lung cancer?†. *Eur J Cardiothorac Surg* 2017;51:242-7.
3. Asamura H, Hishida T, Suzuki K, et al. Radiographically determined noninvasive adenocarcinoma of the lung: survival outcomes of Japan Clinical Oncology Group 0201. *J Thorac Cardiovasc Surg* 2013;146:24-30.
4. Tsuchida H, Tanahashi M, Suzuki E, et al. Impact of ground glass opacity in a three-dimensional analysis for pathological findings and prognosis in stage IA pure solid lung cancer. *J Thorac Dis* 2023;15:3829-39.
5. Sun K, You A, Wang B, et al. Clinical T1aN0M0 lung cancer: differences in clinicopathological patterns and oncological outcomes based on the findings on high-resolution computed tomography. *Eur Radiol* 2021;31:7353-62.

6. Li D, Deng C, Wang S, et al. Ten-Year Follow-up Results of Pure Ground-Glass Opacity-Featured Lung Adenocarcinomas After Surgery. *Ann Thorac Surg* 2023;116:230-7.
7. Travis WD, Asamura H, Bankier AA, et al. The IASLC Lung Cancer Staging Project: Proposals for Coding T Categories for Subsolid Nodules and Assessment of Tumor Size in Part-Solid Tumors in the Forthcoming Eighth Edition of the TNM Classification of Lung Cancer. *J Thorac Oncol* 2016;11:1204-23.
8. Sim HJ, Choi SH, Chae EJ, et al. Surgical management of pulmonary adenocarcinoma presenting as a pure ground-glass nodule. *Eur J Cardiothorac Surg* 2014;46:632-6; discussion 636.
9. Lin YH, Chen CK, Hsieh CC, et al. Lymphadenectomy is Unnecessary for Pure Ground-Glass Opacity Pulmonary Nodules. *J Clin Med* 2020;9:672.
10. Cardillo G, Petersen RH, Ricciardi S, et al. European guidelines for the surgical management of pure ground-glass opacities and part-solid nodules: Task Force of the European Association of Cardio-Thoracic Surgery and the European Society of Thoracic Surgeons. *Eur J Cardiothorac Surg* 2023;64:ezad222. Erratum in: *Eur J Cardiothorac Surg* 2023.

Cite this article as: Anile M, Ferrante F, Zacchini B, Pernazza A, Bassi M, Vannucci J, Venuta F. Not all that glitters is gold. *J Thorac Dis* 2024;16(4):2681-2683. doi: 10.21037/jtd-24-38