

4'-Chloro-3',5'-dimethoxyacetanilide

Huiyu Li, Yanyan Zhu and Zhonghua Chen*

Department of Mathematics and Physics, Shanghai University of Electric Power, Shanghai 201300, People's Republic of China

Correspondence e-mail: chenzhonghua@shiep.edu.cn


Received 27 February 2009; accepted 7 April 2009

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.035; wR factor = 0.105; data-to-parameter ratio = 11.2.

The title compound, $\text{C}_{10}\text{H}_{12}\text{ClNO}_3$, crystallizes with four independent molecules in the asymmetric unit which are linked by intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

The natural pyranoacridone acronycine, which can be synthesized from the title compound, exhibits a broad spectrum of activity against numerous experimental tumor models, see: Nguyen *et al.* (2006). For a related structure, see: Lai *et al.* (2007).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_{12}\text{ClNO}_3$
 $M_r = 229.66$

 Monoclinic, $P2_1$
 $a = 11.1373$ (14) Å

 $b = 15.1159$ (19) Å

 $c = 14.2802$ (18) Å

 $\beta = 111.9280$ (10)°

 $V = 2230.1$ (5) Å³
 $Z = 8$

 Mo $K\alpha$ radiation

 $\mu = 0.33$ mm⁻¹
 $T = 296$ K

 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2000)

 $T_{\min} = 0.908$, $T_{\max} = 0.937$

11396 measured reflections

6377 independent reflections

 5985 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.105$
 $S = 1.07$

6377 reflections

570 parameters

38 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.25$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³

Absolute structure: Flack (1983),

2293 Friedel pairs

 Flack parameter: -0.02 (5)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H3A}\cdots\text{O11}^i$	0.86 (3)	2.03 (4)	2.886 (4)	176 (3)
$\text{N2}-\text{H2A}\cdots\text{O2}^{ii}$	0.895 (18)	1.993 (19)	2.887 (4)	178 (3)
$\text{N1}-\text{H1A}\cdots\text{O8}^{iii}$	0.85 (4)	2.06 (4)	2.906 (4)	176 (4)
$\text{N4}-\text{H4A}\cdots\text{O5}$	0.84 (4)	2.10 (4)	2.909 (4)	163 (3)

 Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + 1$; (ii) $x, y - 1, z$; (iii) $-x, y + \frac{1}{2}, -z + 1$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors gratefully acknowledge support from the Excellent Youth Teacher Fund Research Project of Shanghai (No. Z-2008-25), the Special Project of Shanghai Nanotechnology (No. 0852 nm02400) and the Special Project of the Science and Technology Commission of Shanghai Municipality (No. 071605213).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FL2238).

References

- Bruker (2000). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Lai, Q.-L., Zhao, Q.-J., Liu, Z. & Shen, J.-S. (2007). *Acta Cryst.* **E63**, o2782.
- Nguyen, T. M., Sittisombut, C., Boutefnouchet, S., Lallemand, M.-C., Michel, S., Kock, M., Tillequin, F., Mazinghien, R., Lansiaux, A., David-Cordonnier, M.-H., Pfeiffer, B., Kraus-Berthier, L., Léonce, S. & Pierré, A. (2006). *J. Med. Chem.* **49**, 3383–3394.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o1078 [doi:10.1107/S1600536809013166]

4'-Chloro-3',5'-dimethoxyacetanilide

H. Li, Y. Zhu and Z. Chen

Comment

The natural pyranoacridone acronycine was shown to exhibit a broad spectrum of activity against numerous experimental tumor models, including sarcoma, myeloma, carcinoma, and melanoma (Nguyen *et al.*, 2006). The title compound, (I, Fig. 1), can be used as an intermediate in the synthesis of acronycine derivatives. In this paper, we present the X-ray crystallographic analysis of (I), which crystallizes in the monoclinic space group P2(1) with four molecules in the asymmetric unit.

It is interesting to note that there are four independent molecules in the asymmetric unit which is different from a similar structure (Lai *et al.*, 2007) that has two molecules in the asymmetric unit. The dihedral angle between the acetamide groups and the benzene rings is 3.786 (3)°, 15.359 (4)°, 18.189 (3)° and 32.301 (6)° respectively. The torsion angles of the methoxy groups with respect to the benzene rings also vary (from 2.57 (1) to -15.43 (1)°). The molecules are linked into stacks by intermolecular N-H...O hydrogen bonding (Fig. 2 and Table 1) with packing between the stacks stabilized by van der Waals forces.

Experimental

To a solution of 3,5-dimethoxyanilide (195 mg, 1.0 mmol) in dry CH₂Cl₂ (4 ml) cooled with an ice-water bath was added *N*-chlorobutanamide (147 mg, 1.1 mmol) in portions. After 2 h, the reaction was warmed to room temperature and stirred for 30 min and then filtered. The resulting solution was evaporated to give a white solid which was further purified by column chromatography (CH₂Cl₂) to give (I) (yield 75%, m. p. 493 K). Single crystals of (I) were obtained by slow evaporation of a petroleum ether-ethyl acetate solution. After one week, single crystals suitable for X-ray diffraction were obtained respectively.

Refinement

All the H atoms were placed in geometrically idealized positions and constrained to ride their parent atoms, with N—H = 0.840–0.895 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ and C—H = 0.93 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. View of the molecule of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.




Fig. 2. The crystal packing of (I), viewed along the *a*-axis. Hydrogen bonds are shown as dashed lines.

4'-Chloro-3',5'-dimethoxyacetanilide

Crystal data

$C_{10}H_{12}ClNO_3$

$M_r = 229.66$

Monoclinic, $P2(1)$

$a = 11.1373$ (14) Å

$b = 15.1159$ (19) Å

$c = 14.2802$ (18) Å

$\beta = 111.9280$ (10)°

$V = 2230.1$ (5) Å³

$Z = 8$

$F_{000} = 960$

$D_x = 1.368$ Mg m⁻³

Melting point: 493 K

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 7547 reflections

$\theta = 2.4$ – 27.2 °

$\mu = 0.33$ mm⁻¹

$T = 296$ K

Block, colourless

$0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 296$ K

φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2000)

$T_{\min} = 0.908$, $T_{\max} = 0.937$

11396 measured reflections

6377 independent reflections

5985 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\max} = 25.0$ °

$\theta_{\min} = 2.0$ °

$h = -12 \rightarrow 13$

$k = -13 \rightarrow 17$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.035$

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0603P)^2 + 0.5909P]$

$wR(F^2) = 0.105$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\max} = 0.001$
6377 reflections	$\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$
570 parameters	$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$
38 restraints	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1+0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0059 (8)
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 2293 Friedel pairs
	Flack parameter: -0.02 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.50814 (9)	0.60141 (6)	0.49262 (8)	0.0642 (3)
O1	0.2442 (2)	0.54965 (16)	0.3810 (2)	0.0611 (6)
O2	0.1586 (3)	0.97634 (19)	0.3086 (2)	0.0662 (7)
O3	0.5250 (2)	0.78822 (18)	0.4969 (2)	0.0568 (6)
N1	0.0655 (3)	0.8428 (2)	0.3098 (2)	0.0503 (7)
C1	0.1754 (3)	0.7879 (2)	0.3532 (2)	0.0421 (7)
C2	0.2992 (3)	0.8216 (2)	0.4038 (2)	0.0424 (7)
H2	0.3133	0.8824	0.4097	0.051*
C3	0.4001 (3)	0.7634 (2)	0.4448 (2)	0.0421 (7)
C4	0.3794 (3)	0.6724 (2)	0.4367 (2)	0.0440 (7)
C5	0.2560 (3)	0.6400 (2)	0.3866 (2)	0.0455 (7)
C6	0.1529 (3)	0.6973 (2)	0.3435 (2)	0.0462 (7)
H6	0.0701	0.6755	0.3086	0.055*
C7	0.1193 (4)	0.5133 (3)	0.3338 (4)	0.0762 (12)
H7A	0.0653	0.5320	0.3691	0.114*
H7B	0.1249	0.4499	0.3350	0.114*
H7C	0.0826	0.5333	0.2651	0.114*
C8	0.0608 (3)	0.9307 (3)	0.2914 (3)	0.0535 (8)
C9	-0.0731 (4)	0.9684 (3)	0.2456 (4)	0.0721 (12)
H9A	-0.0679	1.0316	0.2405	0.108*
H9B	-0.1207	0.9541	0.2874	0.108*
H9C	-0.1163	0.9438	0.1795	0.108*

supplementary materials

C10	0.5538 (4)	0.8806 (3)	0.5012 (3)	0.0665 (11)
H10A	0.5279	0.9037	0.4339	0.100*
H10B	0.6451	0.8893	0.5361	0.100*
H10C	0.5077	0.9108	0.5366	0.100*
Cl2	0.55954 (10)	-0.15609 (7)	0.01677 (8)	0.0650 (3)
O4	0.3842 (2)	-0.20909 (15)	0.11098 (18)	0.0527 (6)
O5	0.3629 (2)	0.21934 (17)	0.1938 (2)	0.0605 (6)
O6	0.5970 (3)	0.03091 (19)	0.0383 (2)	0.0683 (7)
N2	0.3041 (3)	0.07914 (18)	0.2169 (2)	0.0441 (6)
C11	0.3702 (3)	0.0277 (2)	0.1693 (2)	0.0399 (7)
C12	0.3470 (3)	-0.0642 (2)	0.1676 (2)	0.0420 (7)
H12	0.2938	-0.0870	0.1986	0.050*
C13	0.4040 (3)	-0.1200 (2)	0.1194 (2)	0.0415 (7)
C14	0.4881 (3)	-0.0857 (2)	0.0775 (2)	0.0460 (7)
C15	0.5123 (3)	0.0043 (2)	0.0808 (2)	0.0457 (7)
C16	0.4517 (3)	0.0612 (2)	0.1258 (2)	0.0460 (7)
H16	0.4662	0.1218	0.1264	0.055*
C17	0.6186 (5)	0.1241 (3)	0.0333 (4)	0.0850 (15)
H17A	0.5369	0.1539	0.0039	0.128*
H17B	0.6689	0.1342	-0.0075	0.128*
H17C	0.6643	0.1466	0.1001	0.128*
C18	0.3005 (3)	0.1681 (2)	0.2249 (3)	0.0502 (8)
C19	0.2165 (4)	0.2013 (3)	0.2789 (3)	0.0629 (10)
H19A	0.2691	0.2130	0.3483	0.094*
H19B	0.1529	0.1573	0.2757	0.094*
H19C	0.1738	0.2547	0.2474	0.094*
C20	0.2957 (4)	-0.2457 (3)	0.1515 (3)	0.0606 (10)
H20A	0.3300	-0.2388	0.2236	0.091*
H20B	0.2834	-0.3074	0.1349	0.091*
H20C	0.2142	-0.2155	0.1233	0.091*
Cl3	0.01359 (8)	-0.00462 (6)	1.02777 (6)	0.0524 (2)
O7	0.0568 (2)	0.18003 (16)	1.00635 (18)	0.0531 (6)
O8	0.1927 (2)	0.27104 (18)	0.73842 (19)	0.0609 (7)
O9	0.1311 (3)	-0.10568 (15)	0.91356 (18)	0.0538 (6)
N3	0.3026 (3)	0.1427 (2)	0.7915 (2)	0.0451 (6)
C21	0.2327 (3)	0.1090 (2)	0.8477 (2)	0.0392 (7)
C22	0.1802 (3)	0.1657 (2)	0.8994 (2)	0.0423 (7)
H22	0.1904	0.2266	0.8972	0.051*
C23	0.1126 (3)	0.1297 (2)	0.9542 (2)	0.0412 (7)
C24	0.0979 (3)	0.0387 (2)	0.9583 (2)	0.0389 (6)
C25	0.1523 (3)	-0.0168 (2)	0.9073 (2)	0.0405 (7)
C26	0.2202 (3)	0.0183 (2)	0.8522 (2)	0.0428 (7)
H26	0.2570	-0.0189	0.8185	0.051*
C27	0.1790 (5)	-0.1637 (3)	0.8574 (3)	0.0682 (11)
H27A	0.1397	-0.1494	0.7869	0.102*
H27B	0.1584	-0.2237	0.8678	0.102*
H27C	0.2712	-0.1572	0.8795	0.102*
C28	0.2784 (3)	0.2203 (2)	0.7393 (2)	0.0454 (7)
C29	0.3613 (3)	0.2385 (3)	0.6798 (3)	0.0573 (9)

H29A	0.3143	0.2242	0.6102	0.086*
H29B	0.4383	0.2031	0.7055	0.086*
H29C	0.3846	0.3000	0.6856	0.086*
C30	0.0697 (4)	0.2730 (2)	1.0017 (3)	0.0606 (9)
H30A	0.1596	0.2887	1.0307	0.091*
H30B	0.0240	0.3014	1.0386	0.091*
H30C	0.0342	0.2919	0.9325	0.091*
Cl4	1.03856 (8)	0.29948 (8)	0.49008 (8)	0.0684 (3)
O10	0.8345 (2)	0.17496 (18)	0.3899 (2)	0.0619 (7)
O11	0.4972 (2)	0.53597 (17)	0.23430 (19)	0.0573 (6)
O12	0.9289 (2)	0.47290 (17)	0.47789 (18)	0.0566 (6)
N4	0.4909 (2)	0.38854 (19)	0.2602 (2)	0.0429 (6)
C31	0.6224 (3)	0.3698 (2)	0.3182 (2)	0.0361 (6)
C32	0.6607 (3)	0.2822 (2)	0.3235 (2)	0.0430 (7)
H32	0.6014	0.2386	0.2899	0.052*
C33	0.7887 (3)	0.2602 (2)	0.3797 (2)	0.0457 (7)
C34	0.8772 (3)	0.3252 (2)	0.4263 (2)	0.0437 (7)
C35	0.8372 (3)	0.4127 (2)	0.4242 (2)	0.0410 (7)
C36	0.7081 (3)	0.4349 (2)	0.3696 (2)	0.0403 (7)
H36	0.6803	0.4931	0.3679	0.048*
C37	0.8871 (4)	0.5614 (3)	0.4803 (3)	0.0678 (11)
H37A	0.8620	0.5870	0.4142	0.102*
H37B	0.9566	0.5953	0.5271	0.102*
H37C	0.8146	0.5618	0.5012	0.102*
C38	0.4380 (3)	0.4664 (2)	0.2209 (2)	0.0426 (7)
C39	0.2963 (3)	0.4632 (3)	0.1571 (3)	0.0570 (9)
H39A	0.2558	0.5171	0.1654	0.086*
H39B	0.2572	0.4140	0.1774	0.086*
H39C	0.2850	0.4565	0.0874	0.086*
C40	0.7423 (4)	0.1055 (3)	0.3627 (3)	0.0641 (10)
H40A	0.6810	0.1147	0.3945	0.096*
H40B	0.7856	0.0500	0.3845	0.096*
H40C	0.6977	0.1048	0.2907	0.096*
H3A	0.361 (3)	0.109 (2)	0.786 (2)	0.039 (8)*
H2A	0.261 (3)	0.047 (2)	0.247 (3)	0.055 (10)*
H1A	-0.008 (4)	0.819 (3)	0.295 (3)	0.055 (11)*
H4A	0.440 (3)	0.346 (3)	0.235 (3)	0.050 (10)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0549 (5)	0.0394 (5)	0.0880 (6)	0.0025 (4)	0.0147 (4)	0.0066 (5)
O1	0.0557 (13)	0.0326 (14)	0.0833 (16)	-0.0071 (11)	0.0123 (12)	-0.0004 (12)
O2	0.0690 (16)	0.0501 (17)	0.0882 (18)	0.0024 (13)	0.0392 (14)	0.0134 (14)
O3	0.0495 (12)	0.0405 (15)	0.0749 (15)	-0.0088 (11)	0.0167 (11)	-0.0075 (12)
N1	0.0492 (16)	0.0429 (17)	0.0614 (17)	0.0034 (14)	0.0237 (14)	0.0054 (14)
C1	0.0497 (16)	0.0368 (19)	0.0453 (16)	0.0035 (14)	0.0241 (13)	0.0041 (14)
C2	0.0546 (17)	0.0311 (17)	0.0483 (16)	-0.0007 (14)	0.0271 (14)	-0.0018 (13)

supplementary materials

C3	0.0442 (16)	0.0391 (19)	0.0463 (16)	-0.0040 (14)	0.0206 (14)	-0.0050 (14)
C4	0.0514 (17)	0.0325 (17)	0.0494 (17)	-0.0002 (14)	0.0204 (14)	-0.0004 (14)
C5	0.0557 (18)	0.0305 (17)	0.0524 (18)	-0.0046 (14)	0.0227 (15)	0.0050 (14)
C6	0.0453 (16)	0.041 (2)	0.0520 (17)	-0.0029 (15)	0.0184 (14)	-0.0011 (15)
C7	0.065 (2)	0.039 (2)	0.116 (3)	-0.0170 (19)	0.025 (2)	-0.006 (2)
C8	0.0609 (19)	0.049 (2)	0.062 (2)	0.0056 (18)	0.0365 (17)	0.0046 (18)
C9	0.071 (2)	0.058 (3)	0.095 (3)	0.019 (2)	0.041 (2)	0.018 (2)
C10	0.064 (2)	0.043 (2)	0.089 (3)	-0.0149 (18)	0.024 (2)	-0.008 (2)
Cl2	0.0837 (6)	0.0467 (5)	0.0848 (6)	0.0072 (5)	0.0548 (5)	0.0030 (5)
O4	0.0759 (15)	0.0264 (12)	0.0671 (14)	-0.0033 (11)	0.0398 (12)	-0.0016 (11)
O5	0.0622 (14)	0.0320 (14)	0.0868 (18)	-0.0095 (12)	0.0270 (13)	-0.0055 (13)
O6	0.0870 (18)	0.0430 (15)	0.0985 (19)	-0.0058 (14)	0.0619 (16)	0.0053 (14)
N2	0.0515 (14)	0.0282 (15)	0.0535 (15)	-0.0054 (12)	0.0206 (12)	-0.0033 (11)
C11	0.0403 (15)	0.0311 (17)	0.0436 (15)	-0.0012 (12)	0.0101 (12)	-0.0014 (13)
C12	0.0457 (15)	0.0331 (18)	0.0474 (16)	-0.0024 (14)	0.0175 (13)	0.0006 (14)
C13	0.0505 (16)	0.0300 (17)	0.0429 (15)	-0.0021 (13)	0.0160 (13)	0.0027 (13)
C14	0.0504 (16)	0.0415 (19)	0.0465 (16)	0.0045 (15)	0.0186 (14)	0.0042 (14)
C15	0.0467 (16)	0.0377 (18)	0.0533 (17)	-0.0019 (14)	0.0192 (14)	0.0093 (15)
C16	0.0492 (16)	0.0326 (18)	0.0502 (16)	-0.0017 (14)	0.0117 (14)	0.0044 (14)
C17	0.121 (4)	0.055 (3)	0.106 (3)	-0.027 (3)	0.074 (3)	0.002 (3)
C18	0.0464 (16)	0.0361 (19)	0.060 (2)	0.0015 (15)	0.0103 (15)	-0.0037 (16)
C19	0.074 (2)	0.040 (2)	0.078 (2)	0.0051 (18)	0.032 (2)	-0.0058 (19)
C20	0.086 (3)	0.039 (2)	0.073 (2)	-0.0149 (19)	0.048 (2)	-0.0115 (17)
Cl3	0.0530 (4)	0.0560 (6)	0.0536 (4)	-0.0101 (4)	0.0260 (4)	-0.0017 (4)
O7	0.0646 (14)	0.0405 (14)	0.0654 (14)	0.0032 (11)	0.0370 (12)	-0.0089 (11)
O8	0.0659 (15)	0.0546 (17)	0.0636 (15)	0.0218 (13)	0.0259 (12)	0.0196 (13)
O9	0.0757 (15)	0.0296 (13)	0.0619 (14)	-0.0054 (11)	0.0325 (12)	-0.0049 (10)
N3	0.0457 (14)	0.0422 (17)	0.0514 (15)	0.0054 (13)	0.0230 (12)	0.0058 (12)
C21	0.0375 (14)	0.0394 (18)	0.0391 (14)	0.0013 (13)	0.0123 (12)	0.0017 (14)
C22	0.0469 (16)	0.0300 (17)	0.0499 (17)	0.0006 (13)	0.0180 (14)	0.0005 (13)
C23	0.0390 (14)	0.0423 (19)	0.0407 (15)	0.0026 (13)	0.0132 (12)	-0.0074 (13)
C24	0.0391 (14)	0.0415 (18)	0.0357 (14)	-0.0034 (13)	0.0135 (12)	0.0009 (13)
C25	0.0463 (15)	0.0326 (17)	0.0396 (14)	-0.0005 (13)	0.0125 (12)	-0.0005 (13)
C26	0.0477 (16)	0.0401 (19)	0.0419 (15)	0.0062 (14)	0.0181 (13)	-0.0024 (13)
C27	0.102 (3)	0.035 (2)	0.078 (3)	0.005 (2)	0.046 (2)	-0.0047 (19)
C28	0.0445 (15)	0.044 (2)	0.0424 (15)	-0.0005 (15)	0.0109 (13)	0.0064 (15)
C29	0.0567 (19)	0.056 (2)	0.061 (2)	-0.0024 (18)	0.0250 (17)	0.0147 (18)
C30	0.077 (2)	0.036 (2)	0.075 (2)	0.0081 (17)	0.036 (2)	-0.0091 (18)
Cl4	0.0425 (4)	0.0693 (7)	0.0779 (6)	0.0103 (4)	0.0048 (4)	0.0099 (5)
O10	0.0555 (14)	0.0418 (15)	0.0805 (17)	0.0127 (12)	0.0163 (12)	0.0019 (13)
O11	0.0465 (12)	0.0368 (13)	0.0775 (16)	-0.0083 (11)	0.0106 (11)	0.0043 (12)
O12	0.0441 (12)	0.0495 (16)	0.0607 (13)	-0.0096 (11)	0.0018 (10)	0.0048 (11)
N4	0.0352 (12)	0.0349 (16)	0.0527 (15)	-0.0069 (12)	0.0097 (11)	-0.0019 (12)
C31	0.0365 (13)	0.0358 (18)	0.0352 (13)	-0.0004 (12)	0.0124 (11)	0.0028 (12)
C32	0.0430 (15)	0.0372 (18)	0.0457 (16)	0.0001 (13)	0.0128 (13)	-0.0035 (14)
C33	0.0473 (16)	0.043 (2)	0.0460 (16)	0.0074 (15)	0.0168 (14)	0.0003 (14)
C34	0.0389 (15)	0.048 (2)	0.0422 (15)	0.0054 (14)	0.0124 (12)	0.0085 (14)
C35	0.0404 (15)	0.0426 (19)	0.0367 (14)	-0.0075 (14)	0.0106 (12)	0.0024 (13)
C36	0.0402 (14)	0.0378 (17)	0.0393 (14)	-0.0004 (14)	0.0106 (12)	0.0056 (13)

C37	0.060 (2)	0.047 (2)	0.073 (2)	-0.0112 (18)	-0.0012 (18)	-0.0011 (19)
C38	0.0385 (15)	0.0393 (19)	0.0488 (16)	-0.0008 (14)	0.0149 (13)	-0.0010 (14)
C39	0.0402 (16)	0.047 (2)	0.075 (2)	-0.0010 (15)	0.0111 (16)	-0.0036 (18)
C40	0.074 (2)	0.045 (2)	0.070 (2)	0.008 (2)	0.0227 (19)	-0.0044 (19)

Geometric parameters (Å, °)

C11—C4	1.729 (3)	C13—C24	1.729 (3)
O1—C5	1.371 (4)	O7—C23	1.366 (4)
O1—C7	1.412 (5)	O7—C30	1.417 (4)
O2—C8	1.234 (4)	O8—C28	1.220 (4)
O3—C3	1.363 (4)	O9—C25	1.372 (4)
O3—C10	1.428 (5)	O9—C27	1.420 (4)
N1—C8	1.351 (5)	N3—C28	1.361 (4)
N1—C1	1.416 (4)	N3—C21	1.405 (4)
N1—H1A	0.85 (4)	N3—H3A	0.86 (3)
C1—C6	1.390 (5)	C21—C26	1.383 (5)
C1—C2	1.392 (4)	C21—C22	1.394 (4)
C2—C3	1.375 (4)	C22—C23	1.384 (4)
C2—H2	0.9300	C22—H22	0.9300
C3—C4	1.393 (5)	C23—C24	1.389 (5)
C4—C5	1.381 (5)	C24—C25	1.390 (4)
C5—C6	1.385 (5)	C25—C26	1.384 (4)
C6—H6	0.9300	C26—H26	0.9300
C7—H7A	0.9600	C27—H27A	0.9600
C7—H7B	0.9600	C27—H27B	0.9600
C7—H7C	0.9600	C27—H27C	0.9600
C8—C9	1.500 (5)	C28—C29	1.495 (5)
C9—H9A	0.9600	C29—H29A	0.9600
C9—H9B	0.9600	C29—H29B	0.9600
C9—H9C	0.9600	C29—H29C	0.9600
C10—H10A	0.9600	C30—H30A	0.9600
C10—H10B	0.9600	C30—H30B	0.9600
C10—H10C	0.9600	C30—H30C	0.9600
C12—C14	1.741 (3)	C14—C34	1.729 (3)
O4—C13	1.362 (4)	O10—C33	1.373 (4)
O4—C20	1.428 (4)	O10—C40	1.418 (5)
O5—C18	1.229 (4)	O11—C38	1.219 (4)
O6—C15	1.359 (4)	O12—C35	1.369 (4)
O6—C17	1.436 (5)	O12—C37	1.421 (5)
N2—C18	1.351 (4)	N4—C38	1.341 (4)
N2—C11	1.408 (4)	N4—C31	1.416 (4)
N2—H2A	0.895 (18)	N4—H4A	0.84 (4)
C11—C16	1.374 (5)	C31—C36	1.376 (4)
C11—C12	1.411 (5)	C31—C32	1.385 (5)
C12—C13	1.384 (5)	C32—C33	1.390 (4)
C12—H12	0.9300	C32—H32	0.9300
C13—C14	1.388 (4)	C33—C34	1.375 (5)
C14—C15	1.385 (5)	C34—C35	1.392 (5)

supplementary materials

C15—C16	1.390 (5)	C35—C36	1.397 (4)
C16—H16	0.9300	C36—H36	0.9300
C17—H17A	0.9600	C37—H37A	0.9600
C17—H17B	0.9600	C37—H37B	0.9600
C17—H17C	0.9600	C37—H37C	0.9600
C18—C19	1.504 (5)	C38—C39	1.500 (4)
C19—H19A	0.9600	C39—H39A	0.9600
C19—H19B	0.9600	C39—H39B	0.9600
C19—H19C	0.9600	C39—H39C	0.9600
C20—H20A	0.9600	C40—H40A	0.9600
C20—H20B	0.9600	C40—H40B	0.9600
C20—H20C	0.9600	C40—H40C	0.9600
C5—O1—C7	118.1 (3)	C23—O7—C30	116.9 (3)
C3—O3—C10	117.4 (3)	C25—O9—C27	117.2 (3)
C8—N1—C1	128.7 (3)	C28—N3—C21	125.9 (3)
C8—N1—H1A	113 (3)	C28—N3—H3A	117 (2)
C1—N1—H1A	118 (3)	C21—N3—H3A	116 (2)
C6—C1—C2	121.3 (3)	C26—C21—C22	120.9 (3)
C6—C1—N1	116.1 (3)	C26—C21—N3	118.3 (3)
C2—C1—N1	122.6 (3)	C22—C21—N3	120.7 (3)
C3—C2—C1	118.7 (3)	C23—C22—C21	118.9 (3)
C3—C2—H2	120.7	C23—C22—H22	120.6
C1—C2—H2	120.7	C21—C22—H22	120.6
O3—C3—C2	124.2 (3)	O7—C23—C22	122.9 (3)
O3—C3—C4	115.0 (3)	O7—C23—C24	116.4 (3)
C2—C3—C4	120.9 (3)	C22—C23—C24	120.7 (3)
C5—C4—C3	119.7 (3)	C23—C24—C25	119.6 (3)
C5—C4—C11	120.9 (3)	C23—C24—C13	119.8 (2)
C3—C4—C11	119.4 (2)	C25—C24—C13	120.6 (3)
O1—C5—C4	116.0 (3)	O9—C25—C26	123.8 (3)
O1—C5—C6	123.5 (3)	O9—C25—C24	115.9 (3)
C4—C5—C6	120.5 (3)	C26—C25—C24	120.2 (3)
C5—C6—C1	118.9 (3)	C21—C26—C25	119.6 (3)
C5—C6—H6	120.5	C21—C26—H26	120.2
C1—C6—H6	120.5	C25—C26—H26	120.2
O1—C7—H7A	109.5	O9—C27—H27A	109.5
O1—C7—H7B	109.5	O9—C27—H27B	109.5
H7A—C7—H7B	109.5	H27A—C27—H27B	109.5
O1—C7—H7C	109.5	O9—C27—H27C	109.5
H7A—C7—H7C	109.5	H27A—C27—H27C	109.5
H7B—C7—H7C	109.5	H27B—C27—H27C	109.5
O2—C8—N1	122.9 (3)	O8—C28—N3	122.8 (3)
O2—C8—C9	122.3 (4)	O8—C28—C29	121.6 (3)
N1—C8—C9	114.8 (3)	N3—C28—C29	115.6 (3)
C8—C9—H9A	109.5	C28—C29—H29A	109.5
C8—C9—H9B	109.5	C28—C29—H29B	109.5
H9A—C9—H9B	109.5	H29A—C29—H29B	109.5
C8—C9—H9C	109.5	C28—C29—H29C	109.5
H9A—C9—H9C	109.5	H29A—C29—H29C	109.5

H9B—C9—H9C	109.5	H29B—C29—H29C	109.5
O3—C10—H10A	109.5	O7—C30—H30A	109.5
O3—C10—H10B	109.5	O7—C30—H30B	109.5
H10A—C10—H10B	109.5	H30A—C30—H30B	109.5
O3—C10—H10C	109.5	O7—C30—H30C	109.5
H10A—C10—H10C	109.5	H30A—C30—H30C	109.5
H10B—C10—H10C	109.5	H30B—C30—H30C	109.5
C13—O4—C20	117.4 (3)	C33—O10—C40	117.6 (3)
C15—O6—C17	118.0 (3)	C35—O12—C37	117.2 (3)
C18—N2—C11	128.9 (3)	C38—N4—C31	128.1 (3)
C18—N2—H2A	118 (2)	C38—N4—H4A	112 (3)
C11—N2—H2A	113 (2)	C31—N4—H4A	119 (3)
C16—C11—N2	124.6 (3)	C36—C31—C32	121.1 (3)
C16—C11—C12	120.1 (3)	C36—C31—N4	121.9 (3)
N2—C11—C12	115.3 (3)	C32—C31—N4	117.0 (3)
C13—C12—C11	119.7 (3)	C31—C32—C33	119.3 (3)
C13—C12—H12	120.2	C31—C32—H32	120.4
C11—C12—H12	120.2	C33—C32—H32	120.4
O4—C13—C12	123.9 (3)	O10—C33—C34	116.6 (3)
O4—C13—C14	116.4 (3)	O10—C33—C32	123.1 (3)
C12—C13—C14	119.7 (3)	C34—C33—C32	120.2 (3)
C15—C14—C13	120.4 (3)	C33—C34—C35	120.2 (3)
C15—C14—C12	119.9 (2)	C33—C34—C14	120.7 (3)
C13—C14—C12	119.6 (3)	C35—C34—C14	119.1 (2)
O6—C15—C14	115.7 (3)	O12—C35—C34	117.0 (3)
O6—C15—C16	124.1 (3)	O12—C35—C36	123.4 (3)
C14—C15—C16	120.1 (3)	C34—C35—C36	119.7 (3)
C11—C16—C15	119.9 (3)	C31—C36—C35	119.3 (3)
C11—C16—H16	120.1	C31—C36—H36	120.3
C15—C16—H16	120.1	C35—C36—H36	120.3
O6—C17—H17A	109.5	O12—C37—H37A	109.5
O6—C17—H17B	109.5	O12—C37—H37B	109.5
H17A—C17—H17B	109.5	H37A—C37—H37B	109.5
O6—C17—H17C	109.5	O12—C37—H37C	109.5
H17A—C17—H17C	109.5	H37A—C37—H37C	109.5
H17B—C17—H17C	109.5	H37B—C37—H37C	109.5
O5—C18—N2	123.9 (3)	O11—C38—N4	124.3 (3)
O5—C18—C19	121.3 (3)	O11—C38—C39	120.5 (3)
N2—C18—C19	114.8 (3)	N4—C38—C39	115.1 (3)
C18—C19—H19A	109.5	C38—C39—H39A	109.5
C18—C19—H19B	109.5	C38—C39—H39B	109.5
H19A—C19—H19B	109.5	H39A—C39—H39B	109.5
C18—C19—H19C	109.5	C38—C39—H39C	109.5
H19A—C19—H19C	109.5	H39A—C39—H39C	109.5
H19B—C19—H19C	109.5	H39B—C39—H39C	109.5
O4—C20—H20A	109.5	O10—C40—H40A	109.5
O4—C20—H20B	109.5	O10—C40—H40B	109.5
H20A—C20—H20B	109.5	H40A—C40—H40B	109.5
O4—C20—H20C	109.5	O10—C40—H40C	109.5

supplementary materials

H20A—C20—H20C	109.5	H40A—C40—H40C	109.5
H20B—C20—H20C	109.5	H40B—C40—H40C	109.5

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N3—H3A···O11 ⁱ	0.86 (3)	2.03 (4)	2.886 (4)	176 (3)
N2—H2A···O2 ⁱⁱ	0.895 (18)	1.993 (19)	2.887 (4)	178 (3)
N1—H1A···O8 ⁱⁱⁱ	0.85 (4)	2.06 (4)	2.906 (4)	176 (4)
N4—H4A···O5	0.84 (4)	2.10 (4)	2.909 (4)	163 (3)

Symmetry codes: (i) $-x+1, y-1/2, -z+1$; (ii) $x, y-1, z$; (iii) $-x, y+1/2, -z+1$.

Fig. 1




Fig. 2

