



# Draft Genome Sequence of *Vibrio parahaemolyticus* Strain VP14, Isolated from a *Penaeus vannamei* Culture Farm

Ashok Kumar Jangam,<sup>a</sup> T. Bhuvaneshwari,<sup>b</sup> A. Navaneeth Krishnan,<sup>b</sup> Vinaya Kumar Katneni,<sup>a</sup> Satheesha Avunje,<sup>b</sup> Monendra Grover,<sup>c</sup> Sujeet Kumar,<sup>b</sup> S. V. Alavandi,<sup>b</sup> K. K. Vijayan<sup>b</sup>

<sup>a</sup>Nutrition Genetics and Biotechnology Division, ICAR-Central Institute of Brackishwater Aquaculture, Chennai, Tamil Nadu, India

<sup>b</sup>Aquatic Animal Health and Environment Division, ICAR-Central Institute of Brackishwater Aquaculture, Chennai, Tamil Nadu, India

<sup>c</sup>Division of Computer Applications, ICAR-Indian Agricultural Statistics Research Institute, New Delhi, India

**ABSTRACT** Here, we report the draft genome sequence of an isolate of *Vibrio parahaemolyticus*, VP14, recovered from the gut of *Penaeus vannamei* shrimp farmed in southern India. The genome of VP14 comprised 5,224,046 bp with a GC content of 45.3% and contained 5,326 genes, including 4,972 coding sequences.

*Vibrio parahaemolyticus* is a Gram-negative halophilic bacterium, abundant in brackish water and marine environments. While most strains are nonpathogenic, some cause disease in humans and aquatic animals. Acute hepatopancreatic necrosis disease (AHPND) has emerged as an important disease of concern in the shrimp farming industry, causing substantial economic losses globally (1). AHPND is caused by specific strains of *V. parahaemolyticus* harboring a plasmid that encodes PirA<sup>VP</sup> and PirB<sup>VP</sup> (2). During disease surveillance of farmed shrimp, *V. parahaemolyticus* was isolated in the Nellore District of Andhra Pradesh, India, from the gut of *P. vannamei* with reported mortality within 40 days of culture. The purified bacterial isolate was phenotypically identified as *V. parahaemolyticus* and designated VP14. Further, the species was confirmed by species-specific PCR tests based on the *vpm*, *toxR*, and *gyrB* genes (3) and 16S rRNA gene sequencing. However, this isolate did not reproduce AHPND following previously described experimental protocols (1), and it also failed to produce AHPND-specific PCR amplicons (4). The VP14 genome was found to possess a type III secretin system 1 (T3SS1) based on a positive PCR test for the *vscP* gene. However, a negative PCR test was obtained for the *vscC2*, *vopB2*, *vscS2*, and *vopC* (5) genes, which are indicative of the presence of T3SS2. Further, the VP14 isolate could not induce mortality to juvenile shrimp during infection experiments.

To generate the VP14 genome, DNA was initially isolated using a PowerLyzer UltraClean microbial DNA isolation kit (Mo Bio, USA) and sequenced on the Illumina HiSeq platform. The *de novo* assembly with 1,754,630 paired-end reads of 250 bp in the MaSuRCA assembler (6) generated a genome of 5,224,046 bp with a 45.3% GC content. The assembly contained 94 scaffolds with a coverage of 84×, an  $N_{50}$  length of 279,839 bp, and a mean length of 55,575 bp. Genome annotation was performed with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (7) and subsequently with the Rapid Annotations using Subsystems Technology (RAST) server (8). The genome contained 5,326 genes, including 4,972 coding sequences, 10 5S rRNAs, 15 16S rRNAs, 12 23S rRNAs, 161 tRNAs, 4 noncoding RNAs, and 152 pseudogenes.

The VP14 genome was found to have all 39 virulence factors listed as T3SS1 (responsible for causing cytotoxicity of host cells) in the Virulence Factors Database (VFDB, <http://www.mgc.ac.cn/VFs>), as well as the MAM7 gene (responsible for mediating attachment of bacteria to host cells). However, other virulence factors listed in the

Received 8 February 2018 Accepted 13 February 2018 Published 15 March 2018

**Citation** Jangam AK, Bhuvaneshwari T, Krishnan AN, Katneni VK, Avunje S, Grover M, Kumar S, Alavandi SV, Vijayan KK. 2018. Draft genome sequence of *Vibrio parahaemolyticus* strain VP14, isolated from a *Penaeus vannamei* culture farm. Genome Announc 6:e00149-18. <https://doi.org/10.1128/genomeA.00149-18>.

**Copyright** © 2018 Jangam et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to K. K. Vijayan, [vjciba2014@gmail.com](mailto:vjciba2014@gmail.com).

VFDB for *V. parahaemolyticus* were absent. In addition, plasmid-borne genes reported to be associated with AHPND, namely, PirA<sup>VP</sup> and PirB<sup>VP</sup>, were also found to be absent in VP14. Estimates of average nucleotide identity (ANI) computed for the VP14 genome with other completed genomes of *V. parahaemolyticus* at GenBank using PyANI (9) indicated its closeness to CHN25, a strain isolated from aquatic products in China, with an ANI value of 98.62%. Analysis of the VP14 genome using ResFinder (10) indicated the absence of any antibiotic resistance genes. The genome of this non-AHPND-causing strain isolated from a brackish water aquaculture system would be an addition to existing genomic resources of *V. parahaemolyticus* for comparative genomic studies.

**Accession number(s).** The draft genome sequence reported here has been deposited at GenBank under the accession number [PKMB00000000](https://www.ncbi.nlm.nih.gov/nuclseq/PKMB00000000).

## ACKNOWLEDGMENT

We acknowledge financial support received from the Indian Council of Agricultural Research (ICAR), New Delhi, under the Network Project on Agricultural Bioinformatics and Computational Biology (CABin).

## REFERENCES

1. Tran L, Nunan L, Redman RM, Mohney LL, Pantoja CR, Fitzsimmons K, Lightner DV. 2013. Determination of the infectious nature of the agent of acute hepatopancreatic necrosis syndrome affecting penaeid shrimp. *Dis Aquat Organ* 105:45–55. <https://doi.org/10.3354/dao02621>.
2. Lee CT, Chen IT, Yang YT, Ko TP, Huang YT, Huang JY, Huang MF, Lin SJ, Chen CY, Lin SS, Lightner DV. 2015. The opportunistic marine pathogen *Vibrio parahaemolyticus* becomes virulent by acquiring a plasmid that expresses a deadly toxin. *Proc Natl Acad Sci U S A* 112:10798–11803. <https://doi.org/10.1073/pnas.1503129112>.
3. Luan XY, Chen JX, Zhang XH, Jia JT, Sun FR, Li Y. 2007. Comparison of different primers for rapid detection of *Vibrio parahaemolyticus* using the polymerase chain reaction. *Lett Appl Microbiol* 44:242–247. <https://doi.org/10.1111/j.1472-765X.2006.02074.x>.
4. Sirikharin R, Taengchaiyaphum S, Sanguanrut P, Chi TD, Mavichak R, Proespraiwong P, Nuangsaeng B, Thitamadee S, Flegel TW, Sritunyaluck-sana K. 2015. Characterization and PCR detection of binary, Pir-like toxins from *Vibrio parahaemolyticus* isolates that cause acute hepatopancreatic necrosis disease (AHPND) in shrimp. *PLoS One* 10:e0126987. <https://doi.org/10.1371/journal.pone.0126987>.
5. Noriega NF, 3rd, Johnson CN, Griffitht KJ, Grimes DJ. 2010. Distribution of type III secretion systems in *Vibrio parahaemolyticus* from the northern Gulf of Mexico. *J Appl Microbiol* 109:953–962. <https://doi.org/10.1111/j.1365-2672.2010.04722.x>.
6. Zimin AV, Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. 2013. The MaSuRCA genome assembler. *Bioinformatics* 29:2669–2677. <https://doi.org/10.1093/bioinformatics/btt476>.
7. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufu S, Li W. 2013. Prokaryotic Genome Annotation Pipeline. NCBI, Bethesda, MD. <https://www.ncbi.nlm.nih.gov/books/NBK174280>.
8. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
9. Arahal DR. 2014. Whole-genome analyses: average nucleotide identity, p 103–122. *In* Goodfellow M, Sutcliffe I, Chun J (ed), *Methods in microbiology: new approaches to prokaryotic systematics*. Elsevier, Amsterdam, Netherlands. <https://doi.org/10.1016/bs.mim.2014.07.002>.
10. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.