

For the pursuit of oxygen and carbon dioxide channels in mitochondria

Respiratory gas exchange is a vital activity for many organisms including human being. Through respiration, carbon dioxide (CO₂) is exhaled for the exchange of oxygen (O₂) from the air. At the cellular level, O₂ is used to convert biochemical energy from nutrients into adenosine triphosphate (ATP) with the production of CO₂. In the resting state, a healthy adult consumes about 550 L of pure oxygen and produces over 2 pounds (~0.91 kg) of CO₂ every day (Jequier et al., 1987). Given that continuous production of ATP is absolute essential for both cell survival and conduction of normal cellular function, exchange of O₂ and CO₂ at both the respiratory and cellular levels is critical.

The mitochondrion is the powerhouse of cells that performs most cellular oxidations and produces majority of ATP (Cheng et al., 2015; d'Esterre et al., 2015; Jing et al., 2015; Shenoda, 2015; Alhadidi et al., 2016). Mitochondria are unusual organelles with two membranes and their own genome. It is believed that the mitochondrial outer membrane has many protein-based pores that are big enough to allow the passage of ions and molecules as large as small proteins. On the other hand, the inner membrane has much more restricted permeability as the plasma membrane of the cell (Mannella, 1992). As the site for main cellular respiration processes of citric acid cycle and oxidative phosphorylation, mitochondria matrix are not only the final destiny for O₂ but also the site for CO₂ production. O₂ and CO₂ have to be transported across the cellular membrane, the inner and outer mitochondrial membrane to reach the destiny or release out of the cell.

The classical concept of the plasma membranes is extremely permeable to gases, which has been extended to the solubility-diffusion mechanism for the gas transport across plasma membrane, including O₂ and CO₂ (Finkelstein, 1976; Harch, 2015; Katz et al., 2015; Langston and Toombs, 2015; Parra et al., 2015; Stoller, 2015). It is generally believed that both O₂ and CO₂ molecules are transport across the plasma membrane through diffusion. Nonetheless, the O₂ permeability coefficient for the plasma membrane has been found to be two times lower than that for a water layer of the same thickness as the membrane (Subczynski et al., 1992). Furthermore, plasma membrane has been found to be even less permeable for CO₂ (Boron et al., 1994; Waisbren et al., 1994; Endeward and Gros, 2005). Thus, trans-membrane CO₂ and O₂ channels might serve as an alternative mecha-

nism to increase the permeability of O₂ and CO₂ cross the plasma membrane. In the last two decades, an increasing evidence has been emerged that aquaporin (AQP) may serve as plasma membrane CO₂ channels including AQP1, AQP4, and AQP5 with varied affinities (Herrera and Garvin, 2011; Endeward et al., 2014). On the other hand, pursuing membrane O₂ channel has been even more elusive (Endeward et al., 2014).

The unique double membrane structure of mitochondria creates even a greater barrier for the cross membrane transport of CO₂ and O₂. Ironically, much less effort has been invested for the pursuit of mitochondrial membrane channels for O₂ and CO₂. Water is one of the major end products of mitochondrial oxidative phosphorylation. Therefore it might not be a surprise that the mitochondrial membrane express extensive water channels. Osmotic swelling is one of the fundamental features exhibited by mitochondria in pathological conditions (Halestrap, 1989). The mitochondrial permeability transition pore (MPTP) has been indicted to permit the passage of molecule of mass < 1.5 kDa in a nonselective manner including water, thus, has long been postulated to be the primary mediator for water movement in term of mitochondrial swelling. However the molecular identity of MPTP remains obscure. The discovery of the expression of APQ channels, AQP8 and 9, in the mitochondria inner membrane indicated that APQs may function as transporters for water and CO₂ trafficking in the mitochondria (Lee and Thevenod, 2006). Otherwise, no other potential gas channel has been identified in mitochondria for the trans-membrane transport of CO₂ and O₂. CO₂ molecule is distinct from O₂ molecule in the difference of plasma membrane permeability and solubility. The absolute solubility of CO₂ in both water and lipid are more than an order of magnitude greater than those of O₂ (Endeward et al., 2014). Therefore, it is plausible that CO₂ is transported across cellular as well as mitochondrial membrane *via* different mechanisms from O₂. Nevertheless, identification of the precise mechanism underlying the transportation of O₂ and CO₂ crossing mitochondrial membranes will provide significant insights for our understanding of mitochondria function and its role in both physiological and pathological conditions. With the limited achievement of the plasma CO₂ channels, the pursuit of mitochondrial gas channels will have a long way to go.

Shao-hua Yang*, Ran Liu

Center for Neuroscience Discovery, Institute for Healthy Aging, University of North Texas Health Science Center, Fort Worth, TX, USA

*Correspondence to: Shao-hua Yang, M.D., Ph.D.,
shaohua.yang@unthsc.edu.

orcid: 0000-0002-0405-0887 (Shao-hua Yang)

doi: 10.4103/2045-9912.196908

How to cite this article: Yang SH, Liu R (2016) For the pursuit of oxygen and carbon dioxide channels in mitochondria. *Med Gas Res* 6(4):237-238.

Open access statement: This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

REFERENCES

- Alhadidi Q, Bin Sayeed MS, Shah ZA (2016) Cofilin as a promising therapeutic target for ischemic and hemorrhagic stroke. *Transl Stroke Res* 7:33-41.
- Boron WF, Waisbren SJ, Modlin IM, Geibel JP (1994) Unique permeability barrier of the apical surface of parietal and chief cells in isolated perfused gastric glands. *J Exp Biol* 196:347-360.
- Cheng C, Jiang L, Yang Y, Wu H, Huang Z, Sun X (2015) Effect of APOE gene polymorphism on early cerebral perfusion after aneurysmal subarachnoid hemorrhage. *Transl Stroke Res* 6:446-450.
- d'Esterre CD, Aviv RI, Morrison L, Fainardi E, Lee TY (2015) Acute multi-modal neuroimaging in a porcine model of endothelin-1-induced cerebral ischemia: defining the acute infarct core. *Transl Stroke Res* 6:234-241.
- Endeward V, Gros G (2005) Low carbon dioxide permeability of the apical epithelial membrane of guinea-pig colon. *J Physiol* 567:253-265.
- Endeward V, Al-Samir S, Itel F, Gros G (2014) How does carbon dioxide permeate cell membranes? A discussion of concepts, results and methods. *Front Physiol* 4:382.
- Finkelstein A (1976) Water and nonelectrolyte permeability of lipid bilayer membranes. *J Gen Physiol* 68:127-135.
- Halestrap AP (1989) The regulation of the matrix volume of mammalian mitochondria in vivo and in vitro and its role in the control of mitochondrial metabolism. *Biochim Biophys Acta* 973:355-382.
- Harch PG (2015) Hyperbaric oxygen in chronic traumatic brain injury: oxygen, pressure, and gene therapy. *Med Gas Res* 5:9.
- Herrera M, Garvin JL (2011) Aquaporins as gas channels. *Pflugers Arch* 462:623-630.
- Jequier E, Acheson K, Schutz Y (1987) Assessment of energy expenditure and fuel utilization in man. *Annu Rev Nutr* 7:187-208.
- Jing L, Wang Y, Zhao XM, Zhao B, Han JJ, Qin SC, Sun XJ (2015) Cardioprotective effect of hydrogen-rich saline on isoproterenol-induced myocardial infarction in rats. *Heart Lung Circ* 24:602-610.
- Katz I, Murdock J, Palgen M, Pype J, Caillibotte G (2015) Pharmacokinetic analysis of the chronic administration of the inert gases Xe and Ar using a physiological based model. *Med Gas Res* 5:8.
- Langston JW, Toombs CF (2015) Defining the minimally effective dose and schedule for parenteral hydrogen sulfide: long-term benefits in a rat model of hindlimb ischemia. *Med Gas Res* 5:5.
- Lee WK, Thevenod F (2006) A role for mitochondrial aquaporins in cellular life-and-death decisions? *Am J Physiol Cell Physiol* 291:C195-202.
- Mannella CA (1992) The 'ins' and 'outs' of mitochondrial membrane channels. *Trends Biochem Sci* 17:315-320.
- Parra RS, Lopes AH, Carreira EU, Feitosa MR, Cunha FQ, Garcia SB, Cunha TM, da Rocha JJ, Feres O (2015) Hyperbaric oxygen therapy ameliorates TNBS-induced acute distal colitis in rats. *Med Gas Res* 5:6.
- Shenoda B (2015) The role of Na⁺/Ca²⁺ exchanger subtypes in neuronal ischemic injury. *Transl Stroke Res* 6:181-190.
- Stoller KP (2015) All the right moves: the need for the timely use of hyperbaric oxygen therapy for treating TBI/CTE/PTSD. *Med Gas Res* 5:7.
- Subczynski WK, Hopwood LE, Hyde JS (1992) Is the mammalian cell plasma membrane a barrier to oxygen transport? *J Gen Physiol* 100:69-87.
- Waisbren SJ, Geibel JP, Modlin IM, Boron WF (1994) Unusual permeability properties of gastric gland cells. *Nature* 368:332-335.