

# Rediscovery of Laura's glassfrog *Nymphargus laurae* (Anura: Centrolenidae) with new data on its morphology, colouration, phylogenetic position and conservation in Ecuador

María José Sánchez-Carvajal<sup>1,2</sup>, Grace C. Reyes-Ortega<sup>1,2</sup>,  
Diego F. Cisneros-Heredia<sup>3,4</sup> and H. Mauricio Ortega-Andrade<sup>1,2,4</sup>

<sup>1</sup> Ingeniería en Ecosistemas, Facultad de Ciencias de la Vida, Universidad Regional Amazónica Ikiám, Tena, Napo, Ecuador

<sup>2</sup> Grupo de Investigación en Biogeografía y Ecología Espacial, Universidad Regional Amazónica Ikiám, Tena, Napo, Ecuador

<sup>3</sup> Colegio de Ciencias Biológicas y Ambientales, Instituto de Biodiversidad Tropical iBIOTROP, Museo de Zoología, Laboratorio de Zoología Terrestre, Universidad San Francisco de Quito, Quito, Pichincha, Ecuador

<sup>4</sup> División de Herpetología, Instituto Nacional de Biodiversidad, Quito, Pichincha, Ecuador

## ABSTRACT

We report the rediscovery of Laura's Glassfrog, *Nymphargus laurae* Cisneros-Heredia & McDiarmid, 2007, based on two specimens collected at the Colonso-Chalupas Biological Reserve, province of Napo, Ecuador. The species was described and known from a single male specimen collected in 1955 at Loreto, north-eastern Andean foothills of Ecuador. Limited information was available about the colouration, systematics, ecology, and biogeography of *N. laurae*. We provide new data on the external morphology, colouration, distribution and comment on its conservation status and extinction risk. We discuss the phylogenetic relationships of *N. laurae*, which forms a clade together with *N. siren* and *N. humboldti*. The importance of research in unexplored areas must be a national priority to document the biodiversity associated, especially in protected areas.

**Subjects** Biodiversity, Biogeography, Conservation Biology, Taxonomy, Zoology

**Keywords** Phylogeny, Taxonomy, Glassfrogs, Colonso-Chalupas biological reserve, Systematics

## INTRODUCTION

*Nymphargus* Cisneros-Heredia & McDiarmid, 2007 currently includes 42 described species of glassfrogs (family Centrolenidae), and 21 of them occur in Ecuador (*Guayasamin et al., 2020; Frost, 2021*). Despite increasing efforts to better understand the diversity, natural history, ecology and distribution of glassfrogs in the tropical Andes, several species of *Nymphargus* remain known only from their type localities or few collected specimens (e.g., *Nymphargus buenaventura* (*Cisneros-Heredia & Yáñez-Muñoz, 2007; Yáñez-Muñoz et al., 2014*), *N. laurae* (*Cisneros-Heredia & McDiarmid, 2007*), *N. lindae* Guayasamin in *Guayasamin et al. (2020)*, *N. manduriacu* *Guayasamin et al. (2019)*).

Submitted 15 October 2021  
Accepted 26 November 2021  
Published 23 December 2021

Corresponding author  
H. Mauricio Ortega-Andrade,  
mauricio.ortega@ikiam.edu.ec

Academic editor  
Juan J. Morrone

Additional Information and  
Declarations can be found on  
page 14

DOI 10.7717/peerj.12644

© Copyright  
2021 Sánchez-Carvajal et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

Laura's Glassfrog *Nymphargus laurae* was described based on a male specimen collected in 1955 at Loreto, on the north-eastern foothills of the Andes of Ecuador ([Cisneros-Heredia & McDiarmid, 2007](#)). More than 60 years have passed since the collection of the holotype and single known specimen of *N. laurae*, and no additional individuals or information has become available for the species ([Guayasamin et al., 2020](#)). Since its description, numerous herpetologists have searched for *N. laurae* along the eastern slopes of the Andes of Ecuador without success. Due to its apparent rarity, restricted distribution, and extensive habitat change and loss at the type-locality, *N. laurae* was classified as Endangered at the national level ([Ortega-Andrade et al., 2021](#)) and Critically Endangered at the global level ([Cisneros-Heredia, 2008](#)).


Between 2016–2018, we collected two individuals of *Nymphargus laurae* at the Colonso-Chalupas Biological Reserve. Based in this findings, we provide new information about the morphological and chromatic variation, natural history, conservation status and extinction risk and reveal for the first time the phylogenetic position of *N. laurae* in Centrolenidae.

## MATERIALS AND METHODS

### Study area and field surveys

The Colonso-Chalupas Biological Reserve is a national protected area located on the foothills of the north-eastern Andes of Ecuador, in the province of Napo. This biological reserve protects 932.46 km<sup>2</sup>, extending between 560–4,432 m elevation and being home to a variety of ecosystems, from tropical evergreen forests to paramos ([Van der Hoek et al., 2018](#)). It is part of an ecological corridor with two neighbouring protected areas: Antisana Ecological Reserve, to the north, and Llanganates National Park, to the south ([Ramis, Álvarez-Solas & Peñuela, 2018](#); [Van der Hoek et al., 2018](#)). Montane evergreen cloud forests in Colonso-Chalupas are characterised by a great variety of trees of the families Melastomataceae, Solanaceae, Myrsinaceae, Aquifoliaceae, Araliaceae, Rubiaceae. Those trees reach up to 15–25 m in height, showing gnarled trunks and branches, and dense and compact crowns covered by epiphytes, including orchids, bromeliads, aroids, and ferns ([Ministerio del Ambiente del Ecuador, 2012, 2013](#)). This ecosystem is usually covered by mist, either constantly or during the early morning and late afternoon ([Ramis, Álvarez-Solas & Peñuela, 2018](#)). The average annual rainfall is 4,620 mm, and the average annual temperature is 28.7 °C. The rainy season extends between March and July, with 448 mm of monthly average rainfall and 23.5 °C of monthly average temperature. The dry season is between August and January, with 353 mm of average monthly precipitation and 23.9 °C of average monthly temperature ([INAMHI, 2015](#)). Nocturnal surveys for collection of amphibians and reptiles were conducted at the Colonso-Chalupas Biological Reserve from 19h00–02h00, at a stream nearby to Ikiam's Scientific Station, on 17 October 2016 (0.9348°S 77.9270°W, 1,506 m) and at the Narpa stream, on 09 June 2018 (0.9353°S 77.9268°W, 1,440 m) ([Fig. 1](#)). Environmental Ministry of Ecuador provided full approval for this research (MAE-DNB-CM-2017-0062). Specimens are deposited in the herpetological collection of the Instituto Nacional de Biodiversidad (Quito, Ecuador).





**Figure 1** (A) Map of Ecuador showing the distribution ranges of *Nymphargus laurae*: type locality (red star) and new locality (specimens INABIO15383-84; red dot), and phylogenetic sister species of *N. laurae*: *N. siren* (yellow dots) and *N. humboldti* (green dots). Interlined rectangle in A delimit the area shown in B. (B) Environmental risk surface (Ortega-Andrade et al., 2021) and protected areas in the distribution range of *N. laurae*. Numbers correspond to the following protected areas: (1) Cayambe-Coca National Park,

**Figure 1** (continued)

(2) Sumaco-Napo-Galeras National Park, (3) Antisana Ecological Reserve, (4) Colonso-Chalupas Biological Reserve, (5) Llanganates National Park. Note high risk modelled for the type locality, which is excluded from the National System of Protected Areas of Ecuador. (C) Satellite image (2019, Google Earth) of the Upper Rio Napo valley, type locality near the town of Loreto (ca. 0.666670°S, 77.316700°W, ca. 500 m elevation), slopes of the Sumaco Volcano, on the Cordillera Oriental, eastern slopes of the Andes, Provincia de Orellana, República del Ecuador.

Full-size  DOI: [10.7717/peerj.12644/fig-1](https://doi.org/10.7717/peerj.12644/fig-1)

## Morphological characteristics

Terminology, characters and measurements follow formats and definitions described by *Cisneros-Heredia & McDiarmid (2007)*, *Watters et al. (2016)* and *Guayasamin et al. (2020)*. Examined frogs were photographed in life, anaesthetised with lidocaine 2%; specimens and tissues were fixed in 96% ethanol and preserved separately in 75% ethanol. Sex and maturity were determined by directly examining gonads through dissections and noting secondary sexual characters (*i.e.*, vocal slits and nuptial pads). Colour patterns are based on photographs and annotations of living specimens taken in the field. Adjective “enamelled” is used to describe the shiny white colouration produced by the accumulation of iridophores (*Lynch & Duellman, 1973; Cisneros-Heredia & McDiarmid, 2007*).

The following measurements were taken with a digital calliper (0.02 mm accuracy, rounded to nearest 0.1 mm) under a stereomicroscope: snout-vent length (SVL); head width (HW); head length (HL), snout length (SL); interorbital distance (IOD), horizontal eye diameter (ED), internarial distance (IND), eye-nostril distance (EN), horizontal tympanum diameter (TD); upper eyelid width (UEW), forearm length (FLL), hand length (HAL), finger IV disk width (Fin4DW), thigh length (THL), tibia length (TL), foot length (FL).

Morphological data used for comparisons were obtained from direct examination of specimens deposited in the following collections: The Natural History Museum, Department of Zoology, London (BMNH); División de Herpetología, Instituto Nacional de Biodiversidad, Quito (DHMECN); Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá (ICN); University of Kansas Natural History Museum, Lawrence, KS (KU); Museum of Comparative Zoology, Harvard University, Cambridge, MA (MCZ); Museo de Zoología, Pontificia Universidad Católica del Ecuador, Quito (QCAZ); National Museum of Natural History, Smithsonian Institution, Washington, D.C. (USNM); Museo de Zoología, Universidad San Francisco de Quito, Quito (ZSFQ).

## DNA extraction, amplification, sequencing, and phylogenetic analyses

Genomic DNA was extracted from hepatic tissue preserved in 96% ethanol from specimen DHMECN 15383, using the “Isolation of Genomic DNA” protocol, Wizard Genomic DNA Purification Kit (*Promega, 2019*). Three mitochondrial gene fragments [12S ribosomal rRNA gene, 16S ribosomal rRNA gene, Cytochrome C Oxidase Subunit 1 (COI)], and one nuclear gene [Recombination Activating Protein 1 gene (RAG1)] were amplified using the Polymerase Chain Reaction (*Saiki et al., 1988*). Primers for 12S gene (12S H10-FWR 5'-CACYTTCCRGTRCRYTTACCRTGTTACGACTT-3'/12S



L4E-REV 5'-TACACATGCAAGTYTCCGC-3'), 16S gene (16Sar-L-FWR 5'-CGCCTGTT TATCAAAAACAT-3'/16Sbr-H-REV 5'-CCGGTCTGAACTCAGATCACGT-3'), COI (dgLCO-1490-FWR 5'-GGTCAACAAATCATAAAGAYATYGG-3'/dgHCO-2198-REV 5'-TAAACTTCAGGGTGACCAAARAAYCA-3') and RAG1 (RAG1-R182 5'GCCAT AACTGCTGGAGCATYAT3'/RAG1-R270 5'AGYAGATGTTGCCTGGGTCTTC3'), were used (Heinicke, Duellman & Hedges, 2007). Each PCR reaction was composed of a 25 µl reaction mix containing: 12.5 µl GoTaq Green Master Mix, 4.5 µl H<sub>2</sub>O, 1 µl on 10 µM of Forward primer and 1 µl on 10 µM Reverse primer, and 5 µl of genomic DNA. We perform amplifications on an Applied Biosystems GeneAmp PCR System 9,700 thermal cycler. The amplification program for 12S primers was set with an initial denaturation of 94 °C (5 min) followed by 38 cycles of 94 °C (30 s), 49 °C (30 s), 72 °C (1 min), with a final extension temperature of 72 °C (7 min) and 4 °C for an unlimited period; for 16S primers was set with an initial denaturation of 95 °C (5 min) followed by 30 cycles of 95 °C (30 s), 57 °C (30 s), 72 °C (45 s), with a final extension temperature of 72 °C (5 min) and 4 °C for an unlimited period; for COI primers was set with an initial denaturation of 95 °C (5 min) followed by 30 cycles of 95 °C (30 s), 48 °C (30 s), 72 °C (45 s), with a final extension temperature of 72 °C (5 min) and 4 °C for an unlimited period; and for RAG1 primers was set with an initial denaturation of 94 °C (5 min) followed by 38 cycles of 94 °C (30 s), 52 °C (30 s), 72 °C (1 min), with a final extension temperature of 72 °C (7 min) and 4 °C for an unlimited period. Amplified DNA products were visualised by electrophoresis on a 2% agarose gel and post-staining with Tris/Borate/EDTA buffer (TBE) under blue light. PCR-amplified products were purified using Illustra™ ExoProStar™ Enzymatic PCR and Sequencing Clean-Up Kit. Sequencing was performed in both DNA strain directions and undertaken by Macrogen, Seoul, South Korea (<http://www.macrogen.com>). Chromatographs resulting from sequencing were revised and edited using Geneious Prime v.2020.0.5 software (Kearse et al., 2012). New sequences were deposited in GenBank with the following accession numbers: (12S: MZ820691; 16S: MZ831508; COI: MZ828399; RAG1: MZ835991).

To infer the phylogenetic position of *Nymphargus laurae*, we included sequences for 119 species (Table S1) selected from Guayasamin et al. (2020), obtained from the NCBI GenBank database (National Center for Biotechnology Information NCBI, 2020). We included two species for Allophryniidae, a sister lineage of Centrolenidae, as outgroups (Table S1) and rooted the phylogeny with *Allophryne ruthveni*. Alignments were reviewed and edited manually to remove regions with a high proportion of missing data at the edges and hypervariable regions with Geneious Prime v.2020.0.5 software (Kearse et al., 2012). Highly variable regions into the alignments are subject to the accumulation of gaps caused by deletions, insertions and substitution mutation, commonly identified to contributed to the inaccuracy of phylogenetic inference (Dwivedi & Gadagkar, 2009). We used Mesquite v3.61 software (Maddison & Maddison, 2019) to store the sequences and to create a concatenated matrix of all genes (12S, 16S, COI and RAG1). Because our combined data set comprised two ribosomal genes (12S and 16S), one protein-coding mitochondrial gene (COI) and one nuclear gene (RAG1), we tested a matrix with 8 partitions with PartitionFinder2 v2.1.1 (Lanfear et al., 2012), in the CIPRES Science

Gateway V.3.3 (Miller, Pfeiffer & Schwartz, 2010). This process resulted in the selection of GTR + I + G as the optimal model for nucleotide substitution for all partitions.

Phylogenetic analyses were conducted using Maximum Likelihood (ML) and Bayesian Methods (BA) on the aligned matrix in the CIPRES Science Gateway V.3.3. ML analyses were performed using Garli v2.0 [Genetic Algorithm for Rapid Likelihood Inference; (Zwickl, 2006)]. We perform a total of 10 runs to reduce the probability of inferring a suboptimal solution; all the other settings were set on default. Node support was evaluated using 1,000 bootstrap pseudoreplicates. Bayesian phylogenetic analyses were performed in MrBayes v3.2.2 (Ronquist & Huelsenbeck, 2003), using five runs of the Monte Carlo Markov Chain (MCMC) algorithm for 20 million generations each, with four heated chains (0.2 heating parameter). Trees were sampled every 20,000 generations, with burning of 25% of the total trees. To evaluate the effective sampling size of the five independent, uncorrelated runs, we used the statistical number of effectively independent draws from the posterior (ESS > 200) visualised with Tracer v1.6. (Rambaut et al., 2013). Phylogenetic trees were edited using FigTree v1.4.2. (Rambaut, 2014).

### Conservation status

We assessed the conservation status of *Nymphargus laurae* based on an environmental risk surface model (0 = no threats, 1 = maximum threat value) produced for Ecuadorian amphibians by Ortega-Andrade et al. (2021), a satellite image (2019, Google Earth) from the type locality of the species, and a map of the National System of Protected Areas (<http://ide.ambiente.gob.ec/mapainteractivo/>). We classify the extinction risk of *N. laurae* based on the categories and criteria presented by IUCN (2012, 2019).

## RESULTS

Surveys at the Colonso-Chalupas Biological Reserve resulted in the collection of two specimens of *Nymphargus laurae*. A subadult female (DHMECN 15383) was found at a stream near the Ikiam's Scientific Station on 17 October 2016 by H. Mauricio Ortega-Andrade. An adult male (DHMECN 15384) was collected at Narpa stream, on 09 June 2018 by Miguel Gómez Laporta and H. Mauricio Ortega-Andrade. Both specimens were found active on leaves of riverine vegetation up to 4.5 meters, next to small creeks, at night between 22h00–23h00. *Nymphargus laurae* was found in syntopy with *N. cochranae*, *Pristimantis quaquaversus*, *P. malli*, *P. incomptus*, *P. ventrimarmoratus*, *P. aff. petersi*, and *P. aff. conspicillatus*. The female was classified as subadult by having unconvoluted oviducts and immature ovarian eggs.

Both specimens are very similar to the holotype in their anatomy and colouration, showing all diagnostic characters described for *Nymphargus laurae*: (1) vomerine teeth absent; (2) snout truncated in dorsal and lateral views; (3) tympanic annulus evident; (4) dorsal skin shagreen with slightly elevated warts corresponding to ocelli and scattered spicules, (5) ventral skin granular, without cloacal ornamentation except for a pair of large flat tubercles; (6) parietal peritoneum white, covering 2/3 of the abdomen; all other peritonea clear; (7) liver lobed, hepatic peritoneum clear; (8) humeral spine absent; (9) basal webbing between fingers I, II and III, outer fingers III  $2\frac{2}{3}$ – $2\frac{1}{3}$  IV; (10) webbing

on feet similar to holotype (see below); (11) no dermal folds or tubercles on hands, arms, feet or tarsi; (12) unpigmented nuptial pad Type-I, concealed prepollex; (13) Finger II longer than Finger I; (14) eye diameter larger than the width of disc on Finger III; (15) colouration in life, green with ocelli (yellow spots surrounded by black), and in preservative, lavender with ocelli (cream-coloured centre surrounded by dark lavender).

The new specimens show no relevant differences with the holotype, and observed differences fall within known intraspecific variation found also between specimens of other congeneric species. The male specimen (DHMECN 15384) has SVL = 22.1 mm, slightly larger than the male holotype (USNM 288453, 19.9 mm SVL), and both are smaller than the subadult female (DHMECN 15383, 22.3 mm SVL). Differences in measurements and proportions between these males are probably due to intraspecific variation (Table 1). The male has a combination of large and small spicules (visible under magnification) on the head, dorsum, and flanks, but spicules on the lower part of dorsum and eyelids are smaller. The female has smaller spicules compared to the male on the head, dorsum and flanks. A spicule is present in the centre of each ocellus, being more prominent and pointed when compared to other body spicules. The female has the tympanic annulus proportionally more covered by the supratympanic fold than in males. Hand webbing in the new specimens (basal webbing between fingers I, II and III, outer fingers III  $2\frac{2}{3}$ – $2\frac{1}{3}$  IV; Fig. 2A) is very similar to the holotype (III  $2\frac{2}{3}$ – $2\frac{1}{2}$  IV), and feet webbing shows slight variation: I  $2$ – $2\frac{3}{4}$  II  $1\frac{1}{2}$ – $2\frac{3}{4}$  III  $1\frac{1}{2}$ – $2\frac{3}{4}$  IV  $2\frac{3}{4}$ – $1\frac{1}{2}$  V in the female and I  $2\frac{3}{4}$ – $2\frac{3}{4}$  II  $1\frac{1}{2}$ – $2\frac{3}{4}$  III  $1\frac{1}{2}$ – $2\frac{3}{4}$  IV  $2\frac{1}{2}$ – $1\frac{3}{4}$  V in the male (Fig. 2B) (I  $2^-$ – $2^+$  II  $1\frac{1}{2}$ – $2^+$  III  $1^+$ – $2\frac{1}{2}$  IV  $2\frac{1}{2}$ – $1\frac{1}{2}$  V in the male holotype). The male (DHMECN 15384) has two papillae on discs of Toe I and II (Fig. 2C). The female (DHMECN 15383) lacks papillae on toes. The holotype of *N. laurae* has two papillae on each toe disc, except for Toe V.

In preservation, the new specimens show similar colourations to the holotype. However, the female shows a lavender dorsum, while the new male and the holotype have cream dorsum with lavender tones. The female has 19 ocelli on the body and eight on the head (Fig. 3A), and the male has six ocelli on the body and three on the head (Fig. 3B) (14 on the body and five on the head of the holotype). Upper eyelids are dark lavender. Fingers and toes lack melanophores. All ventral surfaces are cream. The parietal peritoneum and sclera are white, covering  $\frac{2}{3}$  of the abdomen; pericardium, digestive peritonea, hepatic peritoneum, and urogenital peritonea are clear.

The colouration in life of *Nymphargus laurae* remains known only from the brief description provided by Gustavo Orcés-Villagómez, Ecuadorian zoologists who donated the specimen to James A. Peters, USNM curator, and reported in the original description of the species: “green with yellow spots surrounded by black” (*Cisneros-Heredia & McDiarmid, 2007*). The new specimens allow for a complete description: Head green, darker than the body, lip greenish cream; dorsal surfaces of body, arms and legs green; ocelli on head and body having yellow spots surrounded by black; ocelli absent on arms and legs; upper flanks coloured as dorsum but lower flanks cream, with a sharp division between both; hands, finger, feet and toes yellowish-green, with yellow discs; nuptial pad cream (Fig. 4). Throat greenish-white, all other ventral surfaces cream white.




**Table 1** Morphometric measurements (mm) in specimens of *Nymphargus laurae*.


Character	Male (Holotype) USNM 288453	Male INABIO15384	Subadult female INABIO15383
HW	7.4	7.8	8.0
SVL	19.9	22.1	22.3
TL	11.7	13.8	13.6
IOD	3.8	3.1	2.8
HL	6.9	6.9	6.5
ED	2.9	3.4	3.6
IND	1.6	2.3	2.7
EN	1.7	2.1	3.0
FL	8.7	10.5	10.3
TD	–	0.9	0.7
THL	–	12.3	11.7
SL	–	3.1	2.8
FLL	–	5.0	4.4
UEW	–	2.1	2.9
HAL	–	7.8	7.4
Fin 4DW	–	1.5	2.0
HW/HL	1.1	1.1	1.2
HW/SVL	0.4	0.4	0.4
HL/SVL	0.4	0.3	0.3
EN/HL	0.3	0.3	0.5
ED/HL	0.4	0.5	0.5
IOD/ED	1.3	0.9	0.8
EN/ED	0.6	0.6	0.8
EN/IOD	0.5	0.7	1.1
TL/SVL	0.6	0.6	0.6
FL/SVL	0.4	0.5	0.5
HAL/SVL	–	0.4	0.3
FLL/SVL	–	0.2	0.2
ED/Fin 4DW	–	2.3	1.8

Yellow circumpupillary ring and whitish iris with thin dark reticulations and dark flecks concentrated towards the middle (Fig. 4). Nictitating membrane yellowish, without reticulations. Green bones.


We reconstructed the evolutionary tree (Fig. 5) of *Nymphargus laurae* with a dataset including 120 taxa and 2823 nucleotides (in the aligned matrix). ML and BA analyses are both congruent and recovered the phylogenetic position of *N. laurae* as sister species of *N. siren* and both forming a clade sister to *N. humboldti*. The clade *N. laurae* + *N. siren* + *N. humboldti* has low BA posterior probability (<0.9 node value in Fig. 5), while the ML bootstrap value has relative high support (0.7 node value in Fig. S1). This clade is closely related to *N. megacheirus* and *N. anomalus* (Fig. 5). Phylogenetic relationships among major groups to genus level are supported with high values, in



**Figure 2** Hand (A), foot (B) and papillae (C) of *Nymphargus laurae* (INABIO15384). Tags and background color have been digitally removed. [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02\_img.jpg\) DOI: 10.7717/peerj.12644/fig-2](https://doi.org/10.7717/peerj.12644/fig-2)



**Figure 3** Views of the body (dorsum and venter), of (A) subadult female INABIO15383 and (B) adult male INABIO15384 in specimens of *Nymphargus laurae*. Tags and background colour have been digitally removed. [Full-size !\[\]\(7c47b229ca7bdb95c18f544ee7ceb332\_img.jpg\) DOI: 10.7717/peerj.12644/fig-3](https://doi.org/10.7717/peerj.12644/fig-3)




**Figure 4** *Nymphargus laurae* (INABIO15383), (A) dorsal view, (B) side view, (C) front view and (D) ventral view. Full-size [DOI: 10.7717/peerj.12644/fig-4](https://doi.org/10.7717/peerj.12644/fig-4)

*Hyalinobatrachium*, *Centrolene*, *Cochranella*, *Espadarana*, *Rulyrana*, *Sachatamia*, *Teratohyla*, and *Vitreorana* (Fig. S1).

*Nymphargus laurae* is known from two localities in the province of Napo, on the north-eastern flanks of the Andes of Ecuador, at elevations between 700–1,500 m (Fig. 1). The type locality, Loreto, was originally covered by lowland evergreen forests, and it is located on the lower slopes of the Sumaco volcano, on the upper Napo valley. Satellite images (Fig. 1) show that less than 10% of the natural forests remains at the type locality. The new locality, Colonso-Chalupas, is still covered by evergreen montane forest (Fig. 1). The environmental risk surface (ERS) results in threat values from 0 (Colonso-Chalupas) to 0.37 (Loreto), due to habitat loss and fragmentation for cattle raising and agriculture, deforestation, roads, oil pipelines, and stochastic events related with explosions of the Sumaco Volcano.

## DISCUSSION

The records of *Nymphargus laurae* presented in this paper correspond to the first report of the species after 66 years from its original collection. The Colonso-Chalupas Biological Reserve is the second known locality of *N. laurae*, extending its geographic range in ca. 77 km SW from the type locality, at Loreto, province of Orellana, Ecuador (Cisneros-Heredia & McDiarmid, 2007). These records also extend the altitudinal range of the species from ca. 700 m (see comments on the elevation of Loreto by Urgilés, Sanchez-Nivicela & Cisneros-Heredia (2017) up to 1,500 m. Despite the altitudinal difference, both localities are in the same biogeographic region and watershed, and no



**Figure 5** Optimal maximum likelihood tree (log likelihood = -28155.635) of *Nymphargus* (the clade including *N. laurae* is highlighted by an orange rectangle) inferred from concatenated DNA sequences of fragments of the mitochondrial genes 12S, 16S, and COI, and the nuclear gen RAG1, totaling 2,823 aligned base pairs. Circles indicate significant support values for clades recovered by Bayesian (BA) and Likelihood (ML) analyses. [Full-size !\[\]\(5f471a71b78d7676bc356df190b88ab4\_img.jpg\) DOI: 10.7717/peerj.12644/fig-5](https://doi.org/10.7717/peerj.12644/fig-5)

significant biogeographic barriers separate them. *Nymphargus laurae* maybe more widespread than currently known, but possibly it is endemic to north-eastern Ecuador.

Dorsal colouration pattern showing ocelli with yellow centre surrounded by black on a green dorsum is shared by three ocellated glassfrogs: *N. cochranae*, *N. laurae* and *N. lindae*. These species share a common biogeographic pattern across the eastern Andean slopes in Ecuador, with *N. cochranae* being widespread across the eastern Andean slopes of the Andes of Ecuador and southern Colombia and sympatric with *N. laurae* (Cisneros-Heredia & McDiarmid, 2005; 2006; 2007, this paper). All three species are very similar in their morphology and colouration, and when a single known specimen was available for *N. laurae*, the differentiation between this species and *N. lindae* was weak and it was suggested that *N. cochranae* and *N. laurae* may be synonyms (Guayasamin et al., 2020). Now we can provide strong evidence for the distinctiveness of *N. laurae*, which is not closely related to *N. cochranae* nor *N. lindae*, based on morphological, chromatic, and



molecular data. Externally, *N. laurae* differs from *N. cochranae* by having much larger ocelli (ocelli in *N. cochranae* are small, and in some specimens they are so small that without close inspection, they appear to be just dark spots); ocelli with yellow centre (orange centre in *N. cochranae*), Finger II longer than Finger I (Finger I > Finger II in *N. cochranae*); distal subarticular tubercle of fourth finger bifurcate; indistinct outer metatarsal tubercle; supernumerary tubercles present; no ocelli on forearms and shanks (present in some *N. cochranae*); no vomerine teeth (present in some *N. cochranae*); and smaller body size (23.8–31.6 mm SVL in males of *N. cochranae* vs. 19.9–22.3 mm SVL in males of *N. laurae*). *Nymphargus laurae* and *N. lindae* are very similar, but *N. lindae* is diagnosable due to the presence of vomerine teeth (absent in *N. laurae*), low ulnar and tarsal folds present (absent in *N. laurae*), and slightly larger body size (19.9–22.3 mm SVL in males of *N. laurae* vs. 23.0–26.5 mm SVL in males of *N. lindae*). The condition of the papillae at the tip of toes was used as a diagnostic character in the original description of *N. lindae*, but it is not a useful taxonomic character due to its variation in *N. laurae*. Absence of papillae in the female and in some toes in the new male of *N. laurae* (Fig. 2B) suggest that papillae show intraspecific variation or is of external origin. The presence of papillae on toes discs was not used as a diagnostic character in the original description of *N. laurae*. Cisneros-Heredia & McDiarmid (2007) actually noted that intraspecific variation was observed in the presence/absence of papillae on toe discs of other congeneric species (i.e., *Nymphargus ignotus*). We have seen similar papillae in some specimens of *Chimerella*, *Vitreorana* and *Espadarana*, showing intraspecific variation (e.g., *Espadarana prosoblepon*, *Vitreorana ritae*).

Phylogenetic analyses place *Nymphargus laurae* in a clade with *N. siren* and *N. humboldti* (Fig. 5). These results are interesting due to the colouration differences among *N. laurae*, *N. humboldti* and *N. siren* and their close distribution in nearby areas at the Sumaco volcano and the Guacamayos mountain range. *Nymphargus siren* and *N. humboldti* are almost identical, the only phenotypic diagnostic character being the smaller body size of *N. siren*. However, our phylogenetic information shows that, despite their similarities, they are not sister to each other. *Nymphargus siren* is distributed on the eastern Andean slopes from southern Colombia to northern Ecuador, at elevations between 1,410–2,000 m; *N. humboldti*, is known from two localities on the eastern Andean slopes of central Ecuador, at elevations between 1,770–2,400 m (Guayasamin et al., 2020); and *N. laurae* is restricted to lowland and foothill forests along the Upper Napo River basin (Cisneros-Heredia & McDiarmid, 2007, this paper). Our results suggest the dispersal of this clade occurred in the northern Andes, along montane forest in the upper Napo River basin, Guacamayos mountain range and Sumaco volcano (Fig. 1A). Although these species have similar elevations and distributional ranges, the role of morphological, behavioural, bioacoustics and physiological features (i.e., climatic tolerances) is still intriguing, regarding their evolution and biogeographical diversification in eastern Andes of Ecuador.

Based on data provided herein, we propose the following extinction risk assessment for *Nymphargus laurae*: (1) *N. laurae* has suffered population reductions, based on the continuous decline in habitat quality at its type locality and surroundings, where no recent



record for the species have been obtained despite surveys. Habitat quality at Colonso-Chalupas is better by being part of a protected area. However, since only three specimens are known for the species, we refrain from using criterion A until more data are available to at least infer the population status of the species; (2) the species is known from just two localities with different conservation conditions, thus each one should be evaluated as a different threat-defined location; (3) an EOO cannot be estimated with two localities but the estimated AOO is 8 km<sup>2</sup>, which is within the threshold for Critically Endangered (<10 km<sup>2</sup>). However, we consider that it is possible that the geographic range of *N. laurae* is larger, closer to the threshold for Endangered (10–500 km<sup>2</sup>) under criterion B2; (4) the type locality and any potential locality outside of Colonso-Chalupas are under ongoing habitat decline due to forest loss and water pollution. This information suggests the extinction risk of *N. laurae* is relatively high and we propose that it should be classified under the IUCN category of Endangered (EN) based on criteria B2ab (iii, iv). Although *N. laurae* now is expected to have a wider distribution, urgent conservation actions are encouraged for this species and other range-restricted amphibians the eastern Andes slopes of Ecuador.

## CONCLUSIONS

We provide new information about *Nymphargus laurae*, a species previously known from a single specimen collected decades ago. Our new specimens collected at the Colonso-Chalupas Biological Reserve increase the geographic range of the species along the north-eastern slopes of the Ecuadorian Andes. New insights into the morphology, colouration, and phylogeny of *N. laurae* demonstrate its distinctiveness among other ocellated glassfrogs, with which it is not closely related because it is part of a clade with *N. siren* and *N. humboldti*. Although now known from a second locality, the geographic range of *N. laurae* is still limited and habitat loss and fragmentation are threatening the long-term survival of populations outside of protected areas, thus we suggest that the species' extinction risk should be categorised as Endangered at the global and national level and conservation actions are urgently encouraged. The importance of research in unexplored areas must be a national priority to document the biodiversity associated, especially for range-restricted species and in little-explored protected areas.

## ACKNOWLEDGEMENTS

We thank Miguel Gómez-Laporta, Michelle Guachamin, and Jimmy Velasteguí for field companion and support; Andrea Carrera for provide support for the molecular labwork; and the reviewers for their comments. We are grateful to the following people and institutions who provided access to specimens: Jeff Streicher, David Gower and Mark Wilkinson (BMNH), Mario Yáñez-Muñoz (DHMECN), John D. Lynch (ICN), Linda Trueb, William E. Duellman and John E. Simmons (KU), James Hanken and Jose Rosado (MCZ), Santiago Ron (QCAZ), George Zug, Roy McDiarmid and Ron Heyer (USNM), and Carolina Reyes-Puig and Emilia Peñaherrera (ZSFQ). New specimens of *N. laurae* were collected as part of the project “On the quest of the golden fleece in Amazonia:

The first herpetological DNA—barcoding expedition to unexplored areas on the Napo watershed, Ecuador”.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the following projects: “On the quest of the golden fleece in Amazonia: The first herpetological DNA—barcoding expedition to unexplored areas on the Napo watershed, Ecuador”, funded by the Secretaría Nacional de Ciencia y Tecnología del Ecuador (Senescyt-ENSAMBLE Grant #PIC-17-BENS-001), and The World Academy of Sciences (TWAS Grant #16-095, granted to H. Mauricio Ortega-Andrade). Research by Diego Francisco Cisneros-Heredia was supported by the Smithsonian Women’s Committee, Smithsonian Institution (2002 Research Training Program, National Museum of Natural History), World Wildlife Fund (WWF-EFN, Russel E. Train Education for Nature Program), Secretaría de Educación Superior, Ciencia, Tecnología e Innovación (SENESCYT, Programa “Becas de Excelencia”), Universidad San Francisco de Quito (Chancellor grants, COCIBA grants, Collaboration grants, projects HUBI ID 34, 36, 39, 48, 1057, 7703, 12253, 13524), and “Proyecto Descubre Napo”, an initiative of Universidad San Francisco de Quito in association with Wildlife Conservation Society and funded by the Gordon and Betty Moore Foundation as part of the project: WCS Consolidating Conservation of Critical Landscapes (mosaics) in the Andes. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Secretaría Nacional de Ciencia y Tecnología del Ecuador: PIC-17-BENS-001.

World Academy of Sciences: 16-095.

Smithsonian Institution.

World Wildlife Fund.

Secretaría de Educación Superior, Ciencia, Tecnología e Innovación.

Universidad San Francisco de Quito.

Gordon and Betty Moore Foundation.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- María José Sánchez-Carvajal conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Grace C. Reyes-Ortega conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, logistics and organization of fieldwork and expeditions, and approved the final draft.

- Diego F. Cisneros-Heredia conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- H. Mauricio Ortega-Andrade conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, logistics and organization of fieldwork and expeditions, and approved the final draft.

### Animal Ethics

The following information was supplied relating to ethical approvals (*i.e.*, approving body and any reference numbers):

Environmental Ministry of Ecuador provided full approval for this research (MAE-DNB-CM-2017-0062).

### DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The new sequences are available in GenBank: 12S [MZ820691](#); 16S [MZ831508](#); COI [MZ828399](#); RAG1 [MZ835991](#).

### Data Availability

The following information was supplied regarding data availability:

The raw data is available in the [Supplemental Files](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.12644#supplemental-information>.

## REFERENCES

- Cisneros-Heredia DF. 2008.** *Nymphargus laurae*. The IUCN Red List of threatened species 2008: e.T135996A4223953. DOI [10.2305/IUCN.UK.2008.RLTS.T135996A4223953.en](https://doi.org/10.2305/IUCN.UK.2008.RLTS.T135996A4223953.en).
- Cisneros-Heredia DF, McDiarmid RW. 2005.** Amphibia, Centrolenidae, *Centrolene peristictum*, *Centrolene prosoblepon*, *Cochranella cochranae*, *Cochranella midas*, *Cochranella resplendens*, *Cochranella spinosa*, *Hyalinobatrachium munozorum*: range extensions and new provincial records. *Check List* **1**:18–22 DOI [10.15560/1.1.18](https://doi.org/10.15560/1.1.18).
- Cisneros-Heredia DF, McDiarmid RW. 2006.** A new species of the genus *Centrolene* (Amphibia: Anura: Centrolenidae) from Ecuador with comments on the taxonomy and biogeography of Glassfrogs. *Zootaxa* **1244**:1–32 DOI [10.5281/zenodo.172896](https://doi.org/10.5281/zenodo.172896).
- Cisneros-Heredia DF, McDiarmid RW. 2007.** Revision of the characters of Centrolenidae (Amphibia: Anura: Athesphatanura), with comments on its taxonomy and the description of new taxa of glassfrogs. *Zootaxa* **1572**:1–82 DOI [10.11646/zootaxa.1572.1.1](https://doi.org/10.11646/zootaxa.1572.1.1).
- Cisneros-Heredia DF, Yáñez-Muñoz MH. 2007.** A new species of Glassfrog (Centrolenidae) from the southern Andean foothills on the west Ecuadorian region. *South American Journal of Herpetology* **2**:1–10 DOI [10.2994/1808-9798\(2007\)2\[1:ANSOGC\]2.0.CO;2](https://doi.org/10.2994/1808-9798(2007)2[1:ANSOGC]2.0.CO;2).

- Dwivedi B, Gadagkar SR. 2009.** Phylogenetic inference under varying proportions of indel-induced alignment gaps. *BMC Evolutionary Biology* **9**(1):1–18 DOI 10.1186/1471-2148-9-211.
- Frost DR. 2021.** Amphibian species of the world: an online reference. Available at <https://amphibiansoftheworld.amnh.org/index.php>.
- Guayasamin JM, Cisneros-Heredia DF, McDiarmid RW, Peña P, Hutter CR. 2020.** Glassfrogs of Ecuador: diversity, evolution, and conservation. *Diversity* **12**(6):222 DOI 10.3390/d12060222.
- Guayasamin JM, Cisneros-Heredia DF, Vieira J, Kohn S, Gavilanes G, Lynch RL, Hamilton PS, Maynard RJ. 2019.** A new glassfrog (Centrolenidae) from the chocó-andean río manduriacu reserve, Ecuador, endangered by mining. *PeerJ* **7**(4):e6400 DOI 10.7717/peerj.6400.
- Heinicke MP, Duellman WE, Hedges SB. 2007.** Major Caribbean and Central American frog faunas originated by ancient oceanic dispersal. *Proceedings of the National Academy of Sciences of the United States of America* **104**(24):10092–10097 DOI 10.1073/pnas.0611051104.
- INAMHI. 2015.** Anuario meteorológico N°52-2012. Available at <https://www.inamhi.gob.ec/wp-content/uploads/uploads/anuarios/meteorologicos/Am%202012.pdf>.
- IUCN. 2012.** *IUCN red list categories and criteria. version 3.1.* Second Edition. Gland and Cambridge: IUCN Species Survival Commission.
- IUCN. 2019.** Guidelines for using the IUCN red list categories and criteria IUCN. Species survival commission. version 14. Available at <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C. 2012.** Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**(12):1647–1649 DOI 10.1093/bioinformatics/bts199.
- Lanfear R, Calcott B, Ho SY, Guindon S. 2012.** Partitionfinder: combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* **29**:1695–1701 DOI 10.1093/molbev/mss020.
- Lynch JD, Duellman WE. 1973.** A review of the centrolenid frogs of Ecuador, with descriptions of new species. *Occasional Papers University of Kansas Museum of Natural History* **16**:1–66.
- Maddison W, Maddison DR. 2019.** Mesquite: a modular system for evolutionary analysis, ver. 3.61. Available at <http://mesquiteproject.org>.
- Miller M, Pfeiffer W, Schwartz T. 2010.** Creating the CIPRES science gateway for inference of large phylogenetic trees. In: *Gateway Computing Environments Workshop*. 1–8.
- Ministerio del Ambiente del Ecuador. 2012.** *Sistema de clasificación de ecosistemas del Ecuador continental*. Quito: Subsecretaría de Patrimonio Natural.
- Ministerio del Ambiente del Ecuador. 2013.** *Sistema de Clasificación de Ecosistemas del Ecuador Continental*. Quito: Subsecretaría de Patrimonio Natural.
- National Center for Biotechnology Information NCBI. 2020.** GenBank. Bethesda (MD): national library of medicine (US), national center for biotechnology information. Available at <https://www.ncbi.nlm.nih.gov/genbank/>.
- Ortega-Andrade HM, Rodes Blanco M, Cisneros-Heredia DF, Guerra Arévalo N, López de Vargas-Machuca KG, Sánchez-Nivicela JC, Armijos-Ojeda D, Cáceres Andrade JF, Reyes-Puig C, Quezada Riera AB, Székely P, Rojas Soto OR, Székely D, Guayasamin JM, Siavichay Pesántez FR, Amador L, Betancourt R, Ramírez-Jaramillo SM, Timbe-Borja B, Gómez Laporta M, Webster Bernal JF, Oyagata Cachimuel LA, Chávez Jácome D, Posse V, Valle-Piñuela C, Padilla Jiménez D, Reyes-Puig JP, Terán-Valdez A, Coloma LA, Pérez Lara MB, Carvajal-Endara S, Urgilés M, Yáñez Muñoz MH. 2021.** Red list assessment of

- amphibian species of Ecuador: a multidimensional approach for their conservation. *PLOS ONE* **16**:e0251027 DOI [10.1371/journal.pone.0251027](https://doi.org/10.1371/journal.pone.0251027).
- Promega. 2019.** Wizard® genomic DNA purification kit technical manual. Available at <https://worldwide.promega.com/resources/protocols/technical-manuals/0/wizard-genomic-dna-purification-kit-protocol/>.
- Rambaut A. 2014.** FigTree v1.4.2: tree figure drawing tool. Available at <http://tree.bio.ed.ac.uk/software/figtree/>.
- Rambaut A, Suchard M, Xie D, Drummond A. 2013.** *Tracer 1.6*. Edinburgh, UK: University of Edinburgh.
- Ramis L, Álvarez-Solas S, Peñuela M. 2018.** Diagnóstico preliminar de la presencia de primates que habitan el piedemonte de la Reserva Biológica Colonso-Chalupas. *Revista de Investigación Talentos* **5**(2):1–11 DOI [10.33789/talentos.5.78](https://doi.org/10.33789/talentos.5.78).
- Ronquist F, Huelsenbeck JP. 2003.** MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**(12):1572–1574 DOI [10.1093/bioinformatics/btg180](https://doi.org/10.1093/bioinformatics/btg180).
- Saiki RK, Gelfand DH, Stoffel S, Scharf SJ, Higuchi R, Horn GT, Mullis KB, Erlich HA. 1988.** Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* **239**(4839):487–491 DOI [10.1126/science.2448875](https://doi.org/10.1126/science.2448875).
- Urgilés VL, Sanchez-Nivicela JC, Cisneros-Heredia DF. 2017.** Notes on the distribution of *Gastrotheca testudinea* (Jiménez de la Espada, 1870) in Ecuador. *Check List* **13**(4):121–125 DOI [10.15560/13.4.121](https://doi.org/10.15560/13.4.121).
- Van der Hoek Y, Jensen R, Salagaje LA, Ordóñez Delgado L. 2018.** A preliminary list of the birds of the foothills and south-eastern buffer zone of Colonso Chalupas Biological. *Cotinga* **40**:12–22.
- Watters JL, Cummings ST, Flanagan RL, Siler CD. 2016.** Review of morphometric measurements used in anuran species descriptions and recommendations for a standardized approach. *Zootaxa* **4072**(4):477–495 DOI [10.11646/zootaxa.4072.4.6](https://doi.org/10.11646/zootaxa.4072.4.6).
- Yáñez-Muñoz MH, Sánchez JC, López K, Rea E, Meza-Ramos PA, Oyagata LA, Guerrero P. 2014.** Ampliaciones del rango de distribución de algunas especies de anfibios y reptiles en el suroccidente de Ecuador. *ACI Avances en Ciencias e Ingenierías* **6**(1):2–5 DOI [10.18272/aci.v6i1.151](https://doi.org/10.18272/aci.v6i1.151).
- Zwickl D. 2006.** GARLI: genetic algorithm for rapid likelihood inference. Available at <http://www.bio.utexas.edu/faculty/antisense/garli/Garli.html>.