

Correction to “Solvent-Free Nonthermal Destruction of PFAS Chemicals and PFAS in Sediment by Piezoelectric Ball Milling”

Nanyang Yang,[†] Shasha Yang,[†] Qingquan Ma, Claudia Beltran, Yunqiao Guan, Madison Morsey, Elizabeth Brown, Sujan Fernando, Thomas M. Holsen, Wen Zhang, and Yang Yang*

Environ. Sci. Technol. Lett. 2023, 10, 2, 198–203. DOI: 10.1021/acs.estlett.2c00902



Cite This: *Environ. Sci. Technol. Lett.* 2024, 11, 492–492



Read Online

ACCESS |

 Metrics & More

 Article Recommendations

In our original article, the PZ potential calculated by eq 1 should be 1070 kV (not ~3000 V).

The discussion below eq 1 should be revised as “Surprisingly, the theoretical calculation gives a high PZ potential of 1070 kV. If the midpoint of the PZ potential is arbitrarily set as the Fermi level of BN ($-3.6 \text{ V}_{\text{RHE}}$), then the 1070 kV is composed of $-535 \text{ kV}_{\text{RHE}}$ cathodic potential and $535 \text{ kV}_{\text{RHE}}$ anodic potential.”

The correction of the theoretical value of PZ potential does not affect the remaining content of the article.

AUTHOR INFORMATION

Corresponding Author

Yang Yang

Authors

Nanyang Yang

Shasha Yang;  orcid.org/0000-0002-6646-5332

Qingquan Ma;  orcid.org/0000-0001-6879-6410

Claudia Beltran

Yunqiao Guan

Madison Morsey

Elizabeth Brown

Sujan Fernando

Thomas M. Holsen

Wen Zhang;  orcid.org/0000-0001-8413-0598

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.estlett.4c00242>

Author Contributions

[†]These authors contributed equally.

Received: April 1, 2024

Accepted: April 3, 2024

Published: April 3, 2024

