

Cardiometabolic risk clusters and their reproductive correlates among Indian women

Supplementary table 1. Differences in sociodemographic and lifestyle characteristics between analytic and excluded sample

Characteristic	Weighted % or mean (95% CI)		P-value
	Analytic sample (n=644,191)	Excluded sample (n=79,924)	
<i>Sociodemographic</i>			
Age (years)	30.6 (30.6, 30.7)	28.9 (28.8, 29.0)	<.0001
Education			<.0001
None or lower than secondary	34.5 (34.3, 34.8)	31.4 (30.8, 31.9)	
Secondary	50.3 (50.1, 50.5)	49.6 (49.0, 50.1)	
Higher than secondary	15.2 (15.0, 15.4)	19.1 (18.5, 19.7)	
Wealth quintile			<.0001
Lowest	18.4 (18.2, 18.7)	19.1 (18.6, 19.5)	
Second	20.2 (20.0, 20.4)	18.3 (17.9, 18.8)	
Middle	20.8 (20.6, 21.1)	18.2 (17.7, 18.7)	
Fourth	20.9 (20.7, 21.2)	19.9 (19.3, 20.4)	
Highest	19.6 (19.3, 19.9)	24.5 (23.7, 25.4)	
Marital status			<.0001
Never married	24.4 (24.2, 24.5)	19.1 (18.6, 19.5)	
Currently married and together	71.3 (71.1, 71.4)	77.6 (77.1, 78)	
Ever married, but currently apart ^a	4.4 (4.3, 4.4)	3.4 (3.2, 3.6)	
Religion			<.0001
Hindu	82.0 (81.6, 82.4)	76.8 (76.0, 77.6)	
Muslim	13.0 (12.6, 13.4)	17.2 (16.4, 17.9)	
Christian	2.3 (2.2, 2.4)	2.5 (2.3, 2.7)	
Other ^b	2.7 (2.6, 2.8)	3.5 (3.1, 3.8)	
Place of residence			<.0001
Urban	31.9 (31.5, 32.2)	37.0 (36.2, 37.9)	
Rural	68.1 (67.8, 68.5)	63.0 (62.1, 63.8)	
<i>Health behaviors</i>			
Tobacco use	4.1 (4.0, 4.2)	3.7 (3.5, 3.8)	0.1200
Alcohol use	0.8 (0.7, 0.8)	0.7 (0.6, 0.8)	0.0047
Less than daily fruit and vegetable consumption	92.1 (92.0, 92.3)	88.4 (88.0, 88.8)	<.0001

^aEither widowed, divorced, separated, or deserted.^bOther category includes Sikh, Buddhist/Neo-Buddhist, Jain, Jewish, Parsi/Zoroastrian, and no religion.