



# Draft Genome Sequence of the 1,2-Dichloroethane-Utilizing *Micrococcus* sp. Strain NDB3Y10, Isolated from an Australian Bore Well Producing Coal Seam Gas

Joseph Adelskov,<sup>a</sup> Bharat K. C. Patel<sup>b</sup>

School of Natural Sciences, Griffith University, Brisbane, Queensland, Australia<sup>a</sup>; School of Earth, Environment and Biological Sciences, Queensland University of Technology, Brisbane, Queensland, Australia<sup>b</sup>

**ABSTRACT** *Micrococcus luteus* strain NDB3Y10, which utilizes 1,2-dichloroethane as a carbon source, was isolated from a bore well that produces coal seam gas. The draft genome size of the strain was 2.49 Mb with a G+C content of 72.97%. Genes involved in the metabolism of halogenated substrates, including halogenated hydrocarbons, were identified.

The genus *Micrococcus*, a member of the phylum *Actinobacteria*, currently consists of 17 taxonomically valid species (<http://www.bacterio.net>). Members of the genus *Micrococcus* are found in a wide range of diverse environments, including the type species *Micrococcus luteus*, a known commensal that is found in humans and can contribute to food spoilage (1). *Micrococcus* sp. strain NDB3Y10 was isolated from a water sample collected from a newly drilled (65 h) gas bore well located in the upper Surat Basin, Roma, Queensland, Australia. Strain NDB3Y10 grew optimally under aerobic conditions at 37°C and pH 8 in basal PL medium (2) supplemented with 0.2% yeast extract, glucose, and tryptone. The strain was also able to grow on PL medium supplemented with 0.01% yeast extract and 10 mM 1,2-dichloroethane (1,2-DCE), a highly toxic environmental pollutant. Partial 16S rRNA gene sequencing analysis revealed that *Micrococcus luteus* DSM 20030<sup>T</sup> was the nearest phylogenetic neighbor with a sequence identity of 96.44%. Further analysis using Biolog plates (GN2 Micro Plate, Biolog, USA) showed that the strain metabolized several different carboxylic acids, saccharides, pyruvic acid methyl ester, and succinic acid mono-methyl-ester. We report here the first genome sequence of the 1,2-DCE-utilizing *Micrococcus* sp. strain NDB3Y10, isolated from a water sample collected from a newly drilled bore well producing coal seam gas.

High-molecular-weight DNA was extracted from a late log phase culture of strain NDB3Y10 using a modification of Marmur's method (3). The DNA was sent to the Australian Genome Research Facility (AGRF) for TruSeq library prep and sequencing on the Illumina MiSeq platform. The sequencing run produced a total of 1,306,503 paired-end reads (250 bp), which were reduced to 1,296,434 reads after quality filtering. *De novo* assembly was performed using SPAdes version 3.5.0, which produced a high-quality draft genome consisting of 84 contigs ( $\geq 500$  bp in length) with an  $N_{50}$  of 64,137 bp and an average genome coverage of 247 $\times$ . The contigs had a G+C content of 72.97% and a combined length of 2,497,512 bp, which is estimated to cover 99.85% of the total genome length.

Annotation of the draft genome of strain NDB3Y10 using Prokka version 1.12b (4) revealed the presence of 2,274 protein-coding and 55 RNA gene sequences. Average nucleotide identity using BLAST (ANIb) analysis showed that strain NDB3Y10 shared the

Received 2 March 2017 Accepted 7 March 2017 Published 27 April 2017

**Citation** Adelskov J, Patel BKC. 2017. Draft genome sequence of the 1,2-dichloroethane-utilizing *Micrococcus* sp. strain NDB3Y10, isolated from an Australian bore well producing coal seam gas. *Genome Announc* 5:e00255-17. <https://doi.org/10.1128/genomeA.00255-17>.

**Copyright** © 2017 Adelskov and Patel. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Bharat K. C. Patel, [bharatgu19@gmail.com](mailto:bharatgu19@gmail.com).

highest ANIb value (>98%) with *M. luteus* strains SK58 and 773\_MLUT, which were isolated from human hosts. A comparative analysis using Roary (5) identified genes involved in the uptake of molybdenum and in the biosynthesis of molybdopterin, as well as a methane monooxygenase in strain NDB3Y10, thus excluding the other 11 genomes of *M. luteus* strains (SK85, modasa, NCTC 2665, trpE16, RIT304, RIT305, RIT324W, SUBG006, 773\_MLUT, 1058\_MLUT, and SEN31). RAST (6) annotations of the genome of strain NDB3Y10 identified haloacid dehalogenase and haloalkane dehydrogenase-like genes, a class of hydrolytic dehalogenases involved in the degradation of halogenated hydrocarbons such as 1, 2-DCE (7). This constitutes the first report of the 1,2-DCE-utilizing *Micrococcus* sp. strain NDB3Y10 in which potential hydrolytic dehalogenases have been identified.

**Accession number(s).** This whole-genome shotgun project has been deposited in GenBank under the accession number [LQAC00000000](https://www.ncbi.nlm.nih.gov/nuccore/LQAC00000000). The version described in this paper is the first version, LQAC00000000.1.

## ACKNOWLEDGMENTS

An Australian Postgraduate Award (APA) scholarship to J.A. and the IT infrastructure for genomics provided by the National eResearch Collaboration Tools and Resources (NeCTAR) Project to B.K.C.P. is gratefully acknowledged.

## REFERENCES

1. Kocur M, Pacova Z, Martinec T. 1972. Taxonomic status of *Micrococcus luteus* (Schroeter 1872) Cohn 1872, and designation of the neotype strain. *Int J Syst Bacteriol* 22:218–223. <https://doi.org/10.1099/00207713-22-4-218>.
2. Ogg CD, Patel BKC. 2009. *Caloramator australicus* sp. nov., a thermophilic, anaerobic bacterium from the Great Artesian Basin of Australia. *Int J Syst Evol Microbiol* 59:95–101. <https://doi.org/10.1099/ijs.0.000802-0>.
3. Marmur J. 1961. A procedure for the isolation of deoxyribonucleic acid from micro-organisms. *J Mol Biol* 3:208–218. [https://doi.org/10.1016/S0022-2836\(61\)80047-8](https://doi.org/10.1016/S0022-2836(61)80047-8).
4. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
5. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, Fookes M, Falush D, Keane JA, Parkhill J. 2015. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* 31:3691–3693. <https://doi.org/10.1093/bioinformatics/btv421>.
6. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
7. Olaniran AO, Pillay D, Pillay B. 2004. Haloalkane and haloacid dehalogenases from aerobic bacterial isolates indigenous to contaminated sites in Africa demonstrate diverse substrate specificities. *Chemosphere* 55: 27–33. <https://doi.org/10.1016/j.chemosphere.2003.10.067>.