



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Design and synthesis of a series of serine derivatives as small molecule inhibitors of the SARS coronavirus 3CL protease



Hiroyuki Konno^{a,*}, Masaki Wakabayashi^a, Daiki Takanuma^a, Yota Saito^a, Kenichi Akaji^{b,*}

^a Department of Biological Engineering, Graduate School of Science and Technology, Yamagata University, Yonezawa, Yamagata 992-8510, Japan

^b Department of Medicinal Chemistry, Kyoto Pharmaceutical University, Yamashina-ku, Kyoto 607-8414, Japan

ARTICLE INFO

Article history:

Received 13 January 2016

Revised 26 January 2016

Accepted 27 January 2016

Available online 29 January 2016

Keywords:

SARS CoV

SARS 3CL protease

Serine derivative

Docking simulation

Cathepsin B

Cytotoxicity

ABSTRACT

Synthesis of serine derivatives having the essential functional groups for the inhibitor of SARS 3CL protease and evaluation of their inhibitory activities using SARS 3CL R188I mutant protease are described. The lead compounds, functionalized serine derivatives, were designed based on the tetrapeptide aldehyde and Bai's cinnamoly inhibitor, and additionally performed with simulation on GOLD software. Structure activity relationship studies of the candidate compounds were given reasonable inhibitors ent-**3** and ent-**7k** against SARS 3CL R188I mutant protease. These inhibitors showed protease selectivity and no cytotoxicity.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

Severe acute respiratory syndrome (SARS) is a contagious respiratory disease to humans that is caused by the SARS coronavirus (SARS-CoV). In 2003, an atypical pneumonia due to SARS-CoV led to progressive respiratory failure in over 8,000 individuals and 800 deaths within a few months.^{1–3} Thereafter, the SARS epidemic was successfully controlled, but potential reemergence of pandemic SARS-CoV continues to be a risk, and new strains of SARS or a SARS-like virus could potentially be more infectious than the strains that led to the 2003 outbreak. Two human coronaviruses, NJ63 and HKU1, have been identified in patients around the world since 2003.^{4–6} A more recently identified human coronavirus, Middle East Respiratory Syndrome coronavirus (MERS-CoV), has infected 114 people since April 2012 with a nearly 50% mortality rate, and this number keeps rising daily.^{7,8} Two Asian bat coronaviruses, BtCoV-HKU4 and BtCoV-HKU5, have been shown to be the closest relations to MERS-CoV.⁹ The possibility of a future SARS-like pandemic remains, and no vaccines or antiviral agents have yet been developed to prevent or treat SARS-like infections.¹⁰

SARS-CoV is a novel human coronavirus featuring the largest positive-stranded RNA genome known to date (27 kb for pp1a and 31 kb for pp1b). The key enzyme in the processing of polypro-

teins translated by the viral RNA genome of SARS-CoV is a 33 kDa protease called 3C-like protease (3CL protease). SARS 3CL protease is a cysteine protease containing a Cys-His catalytic dyad. It cleaves precursor polyproteins at as many as 11 conserved sites involving a conserved Gln at the P1 position and a small amino acid (Ser, Ala, or Gly) at the P1' position. Due to its functional importance in the viral life cycle, SARS 3CL protease is considered to be an attractive target for drug design to treat SARS.

A variety of inhibitors against the SARS 3CL protease have been reported in the literature for the past decade, including substrate-based peptidomimetic,¹¹ flavonoid derivatives,¹² tannin derivatives,¹³ ML188,¹⁴ isatin derivatives,¹⁵ and decaisoquinolin,¹⁶ etc. However, no effective therapeutic drug or vaccine has been developed to date although many candidate anti-SARS CoV agents have been identified.

In the previous study, we reported that design and synthesis of the peptide aldehyde inhibitor Ac-Thr-Val-Cha-His-H (**1**) showed high inhibitory activity with an IC₅₀ value of 98 nM toward 3CL R188I mutant protease as a substrate mimetic concept.^{17,18} The inhibitor (**1**) was optimized by the screening of P1, P2 and P4 site residues of the previous reported peptide aldehyde inhibitor Ac-Ser-Ala-Val-Leu-Gln(Me)₂-H, except at the P3 site where the side-chain directed outside and made no interactions with the protease guided by the X-ray crystal structure of the lead compound that bound the R188I mutant SARS 3CL protease. The interactions of inhibitor (**1**) at the P1 and P2 sites with the protease seemed very effective. Additionally, a new synthetic method using acetal

* Corresponding authors. Tel./fax: +81 (0)238 26 3131 (H.K.).

E-mail addresses: konno@yz.yamagata-u.ac.jp (H. Konno), akaji@mb.kyoto-phu.ac.jp (K. Akaji).

to aldehyde conversion via thioacetal formation was described to afford the C-terminal peptide aldehyde.^{19,20}

Side chain structures at the P1, P2 and P4 sites and C-terminus aldehyde as the thiol capture of tetrapeptide inhibitor (**1**) were thought to have a critical role in its potent inhibitory activity. In contrast, there is generally enzymatic digestion of peptide chains and α -proton racemization and/or low specificity of aldehyde functionality. To develop non-peptidyl small molecular inhibitors of the SARS 3CL protease focusing on the P1, P2 and P4 site interactions, we aimed to design and synthesize serine derivatives having the essential functional groups and evaluated their inhibitory activity in the present study. Serine, a commercially available proteinogenic amino acid, has three variant reaction sites; alcohol, amine and carboxylic acid, which can be orthogonally connected to various functional groups (Fig. 1).

2. Results and discussion

2.1. Design

To determine whether a series of serine derivatives could adopt an energetically favorable conformation mimicking a tetrapeptide inhibitor (**1**), we performed a variety of molecular mechanics calculations with SPARTAN from Wavefunction and docking simulations of protein interactions by GOLD from CCDC. Firstly, imidazole, cyclohexyl and hydroxy groups, which were optimized functionalities with potent biological activities in the previous literature, were connected with L-serine to design the serine derivative (**2a,b**) as shown in Scheme 1. However, it did not give good coverage of the substrate-recognition pocket of 3CL protease (PDB code 3AW1), that is, the interactions with optimized motifs and the corresponding pockets were different by binding mode, contrary to expectations. The cyclohexyl group of serine derivative (**2**) occupied the S1 pocket of the 3CL protease imperatively. Thus, the imidazole and hydroxy groups of serine derivative (**2**) expected for the interactions with the S1 and S4 pockets were not effective and consequently the numbers of interaction atom pairs between the inhibitor (**2**) and SARS-CoV 3CL protease were diminished. In contrast, Bai²¹ reported that the cinnamoyl derivatives inhibited SARS-CoV 3CL protease. In this case, the Bai's inhibitor locates deep insides of the S'1, S1 and S2 pockets with appropriate cinnamoyl functionalities. This result was approximately identified with the Bai's simulation employed by the AutoDock 3.0 estimated free energy of binding for the docking of the Bai's inhibitor to the SARS 3CL protease (PDB code 1UJ1). Therefore, we examined molecular docking of the serine derivatives contained together with benzoyl and aniline moieties to give **2c**, which was an attractive ligand for the 3CL protease. Furthermore, we investigated a reasonable structure for the inhibitor and made hybrid compounds with Bai's and our functionalities on the serine derivative (**3**). As a result, we concluded that reasonable structures are *N*-cinnamoyl derivatives with benzoate for the side chain on virtual screening by GOLD. Compound **3** has the following characteristics: (a) since the sub-

strates by nature have involved small amino acids (Ser, Ala or Gly) at the P'1 position, it may be preferable to adopt the aromatic rings. (b) The proper placement of the cyclohexyl ring may be the result of development of a serine type inhibitor with functionality of the P1 position, as well as stability of interaction of other positions. (c) The S4 pocket is rather hydrophobic in nature and therefore the corresponding side chains, namely aromatic, hydrocarbon or neutral functionalities, make good contact with target regions. (d) Additionally, the region adjacent to the ester bond of **3** is catalytic thiol functionality (Cys145) of the enzyme (Scheme 1).

2.2. Chemistry

In view of the promising computational evaluation of the serine derivative (**3**) in terms of tetrapeptide mimetics, we plan to synthesize the target molecules. To evaluate inhibitory activity of the serine derivative (**3**) against SARS-CoV 3CL protease, a structure activity relationship study using the serine derivatives was attempted. As depicted in Scheme 2, the serine derivatives attached with three functionalities were prepared. Each coupling partner to introduce the functionalities is shown in Tables 1–3. Coupling of Fmoc-L-Ser(*t*Bu)-OH and amines as a P1 moiety with WSC/HOBt followed by the deprotection of Fmoc group by 20% piperidine/DMF gave the serine amide derivative (**4**) at acceptable chemical yields. The P4 moiety was introduced by coupling with carboxylic acid in the presence of coupling reagents or acylation using carboxylic anhydride or acyl chloride to afford diamide derivatives. Subsequently, treatment of TFA for the deprotection of the *t*-butyl group gave alcohol (**5**) at moderate yields. Finally, the esterification of a hydroxy group of **5** with a variety of acyl reagents was performed to give novel serine derivatives (**6**, **7** and **8**) for evaluation of the small molecular inhibitors against SARS 3CL protease. Purification of all crude compounds was performed by silica gel column chromatography. The chemical structures for synthetic compounds were mainly determined by ¹H and ¹³C NMRs, IR and mass spectra (Scheme 2).

As depicted in Table 1, five cyclic functionalities were attached to the carboxylic acid of the serine template. As mentioned above, this was to investigate the importance of cyclohexane ring defined as the P1 position. Treatment of Fmoc-L-Ser(*t*Bu)-OH and cyclohexylamine with WSC/HOBt and continuous Fmoc-deprotection afforded **4a** in 97% yield (entry 1). Under the above-mentioned conditions, Fmoc-L-Ser(*t*Bu)-OH was subjected to a coupling reaction with piperidine, morpholine, benzylamine or cyclohexylmethylamine to give **4b–e**. Despite the poor yields in entries 2 and 4, it was available for the evaluation of inhibitory activities (Table 1).

To optimize P4 functionality, a variety of carboxylic acids, especially α,β -unsaturated carboxylic acids for **4a**, were introduced. Coupling of **4a** and cinnamic acid with HATU²²/HOAt²³/DIPEA in CH₂Cl₂ (condition A) followed by treatment with TFA for deprotection of the *t*-butyl group afforded **5a** in 38% yield (entry 1). Treatment of **4a** with 3-(*p*-hydroxyphenyl)propanoic acid in the presence of HATU/HOAt/DIPEA in CH₂Cl₂ also gave **5b** in 79% yield after removal of *t*-butyl group (entry 2). Since **5c** was not isolated by the condition of HATU/HOAt, we found that DMT-MM (condition B)²⁴ as a coupling reagent was moderate to give **5c** in 37% yield over 2 steps (entry 3). In an attempt to improve the chemical yields for the coupling reaction of cinnamic acid derivative and amines, we knew that HATU or DMT-MM as the coupling reagents were useful in preliminary studies. For this reason, a series of cinnamoyl derivatives (**5d**, **5f–o**) were also prepared under identical conditions at moderate yields (entries 6–15). In addition, acylation of **4a** with benzoyl chloride (condition C) was successfully performed to yield **5e** without any problems (entry 5) (Table 2).






Figure 1. Ac-Thr-Val-Cha-His-H (**1**) and concept for serine derivative.



Scheme 1. Virtual screening of the serine derivatives for SARS-CoV 3CL protease (PDB code 3AW1) on GOLD software.



Scheme 2. Synthetic outline for the preparation of serine derivatives (**6a–o** and **7b–o**).

For the structure–activity relationship study of P1 functionality of serine derivatives, we prepared the serine derivatives (**3**), (**6a–o**) and (**7b–o**) as depicted in Table 3. Acylation of the substrates (**5a–o**) was employed the Shiina²⁵ and Yamaguchi²⁶ conditions for the coupling of carboxylic acids or acyl chloride and acetic anhydride reagents. Using commercially available acyl chlorides and Ac₂O as the acyl reagent gave the corresponding esters (**3**), (**6c**), (**6d**), (**6g**), (**6h**), (**6i**) and (**6o**) at moderate yields (entries 1, 4, 5, 8–10, 16). Esterification of **5a** with *N*-acetyl-L-histidine by the Shiina reagent NMBA/DMAP afforded the His-compound (**6a**) at an extremely low yield (entry 2). In this case, 2-nitro-6-methylbenzotriazole as a by-product was given at a moderate yield and it was difficult to improve the chemical yield even after attempting several conditions. Preparation of a pGlu-compound (**6b**), isonicotinic acid (**6e**) and nicotine (**6j**) also resulted in similar situations (entries

Table 1
Coupling of the P2 position and Fmoc-deprotection of Fmoc-Ser(tBu)-OH

Entry	Amine ^a	Product
1	Cyclohexylamine	4a (97%)
2	Piperidine	4b (26%)
3	Morpholine	4c (91%)
4	Benzylamine	4d (19%)
5	Cyclohexylmethylamine	4e (48%)

^a Reagent and conditions: (1) amine, WSC, HOBT, CH₂Cl₂, (2) 20% piperidine/CH₂Cl₂.
















3, 6 and 11). The Yamaguchi protocol was performed to give **6f** and **6k** (entries 7 and 12) and also acyl chlorides prepared by 3,4,5-trimethoxybenzoic acid, 3-phenylpropanoic acid or cinnamic acid with oxalyl chloride in the presence of DMF in situ reacted with **5a** to afford **6l**, **6m** and **6n** at low yields (entries 13–15). On the other hand, the treatment of a variety of alcohols (**5b–o**) with BzCl/Et₃N/DMAP in CH₂Cl₂ converted to benzoyl esters (**7b–o**). Chemical yields of benzoyl esters (**7b–o**) were variable by used substrates (entries 17–30) (Table 3).

2.3. Biological assay

The inhibition of SARS 3CL R188I mutant protease was determined by the previous procedure using a synthetic decapeptide with the S01 cleavage sequence as a substrate. Synthetic L-serine derivatives from several structural subclasses were investigated; **5a**, cinnamoyl-L-serine derivatives (**6a–o**) with modification at the P1 position (Table 4), benzoyl-L-serine derivatives (**7b–k**) with modification at the P4 position (Table 5).

As depicted in Table 4, inhibitory potency of the synthesized cinnamoyl-L-serine derivatives (**6a–o**) with modification at the P1 position was evaluated as IC₅₀ values. The simulated compound (**3**) as a control exhibited an IC₅₀ value of 125 μM (entry 14). Since the IC₅₀ value of a previous reported peptide aldehyde (**1**) was reported to be 98 nM, imidazole of histidine or γ-lactam of pyroglutamic acid were important functionalities. However **6a** and **6b** that had been coupled with **5a** and His or pGlu showed low inhibitory activities as expected from the docking simulations (entries 2 and 3). For optimization of the P1 functional group of **6**, different acyl units; alkyl, phenyl and pyridyl groups, were examined. Aliphatic groups-containing compounds (**6c**), (**6g**) and (**6h**) showed lower inhibitory activities than **3** (entries 4, 8 and 9). Pyridyl derivatives (**6e**), (**6j**) and (**6k**) were also similar (entries 6, 11 and 12). Aromatic rings with nitro, chloride and methoxy groups (**6f**), (**6i**) and (**6l**) gave moderate results (entries 7, 10 and 13). Phenylacetate (**6d**) also showed lower inhibitory activity than **3** (entry 5). In contrast, conversion of the phenylpropionate or cinnamoyl groups at a P1 site would be expected to have a dramatic effect on inhibitory activity. Especially, the cinnamoyl compound (**6n**) showed potent inhibitory activity at an IC₅₀ value of 85 μM (entry 16). In addition, a 2-methyl-6-nitrophenyl derivative (**6o**) showed good potential as a 3CL protease inhibitor (entry 17). The IC₅₀ value

Table 2
Coupling of P4 position and deprotection of *t*-butyl group

Entry	Subst	Conditions ^a	Product	Entry	Subst	Conditions ^a	Product
1	4a		5a (38%)	9	4a		5i (69%)
2	4a		5b (79%)	10	4a		5j (55%)
3	4a		5c (37%)	11	4a		5k (29%)
4	4a		5d (43%)	12	4b		5l (26%)
5	4a		5e (80%)	13	4c		5m (91%)
6	4a		5f (7%)	14	4d		5n (19%)
7	4a		5g (23%)	15	4e		5o (48%)
8	4a		5h (29%)				

^a Reagents and conditions: (1) conditions A, B or C, DIPEA, CH₂Cl₂, rt, (2) 50% TFA/CH₂Cl₂, rt. (A) HATU, HOAT, (B) DMT-MM, (C) Et₃N, DMAP.

of saturated compound (**6m**) was decreased in comparison with those of the fixed unsaturated derivative (**6n**) (entry 15). These results suggest that serine type derivatives containing optimized functionalities would be useful as 3CL protease inhibitors. As the correlations between IC₅₀ and ClogP, we could assume that **3**, **6m**, **6n** and **6o** with potent inhibitory activities tend relatively to have high lipophilicity with the ClogP values of 4.82–5.28 (Table 4).






























To achieve further optimization using a serine template, a structure activity relationship study of the P4 functionality (R₂) of a lead compound (**3**) was conducted. The aliphatic pivaloyl-containing compound (**7b**) substituted the cinnamoyl group of **3** showed no inhibitory activity and introduction of a phenyl-2-propionate (**7c**) and benzoyl group (**7e**) did not make a significant contribution (entries 1, 2 and 4). Next, modified cinnamoyl groups were designed because the moiety of the P4 site was replaced with the methoxy connecting cinnamoyl group to permit the moiety to fit into the S4 pocket more tightly than **3**. Although each 2- or 3-methoxy functionalities were not effective (compounds **7d**, **7f**, entries 3 and 5), the inhibitors containing 3- and/or 4-methoxy groups on a phenyl ring at the P4 site exhibited markedly increased inhibitory activity (entries 6, 7 and 8). The IC₅₀ value of **7k** was 74 μM and so the results suggested that a planar aromatic ring and its hydrophobic functionality were essential factors to produce

a reasonable 3CL protease inhibitor (entry 10). The correlations between IC₅₀ and ClogP are difficult to explain because these compounds are close in molecular formula (Table 5).

To investigate the importance of the cyclohexyl group of **3** at the P1 site, we evaluated four derivatives combined with different amines; piperidine for **7l**, morpholine for **7m**, benzylamine for **7n** and cyclohexylmethylamine for **7o**. As a result, **7l** and **7m** showed no inhibitory activities. These compounds are shorter in terms of one carbon length of the P1 site than **3** and therefore the P1 position of the inhibitor showed decreased flexibility as expected and hardly fit the S1 pocket. On the other hand, aromatic benzyl amine (compound **7n**) has a planer structure, which led to moderate inhibitory activity. Attempts to use cyclohexylmethylamine for **7o** was acceptable to show IC₅₀ = 180 μM. It is suggested that the cyclohexyl structure at a P1 site strictly interacts with the hydrophobic S1 pocket. As mentioned above, we optimized the functionalities at the P1, P1 and P4 sites of an L-serine template based on reference compound **3**, which was guided by virtual screening on GOLD. P1 and P4 sites are preferred for their similar characteristics of aromatic rings, an sp² planer structure and hydrophobicity.


Next, the inhibitory activities of the selected serine derivatives with the D-form were evaluated based on the IC₅₀ values. The IC₅₀ values and tPSA of selected inhibitors are summarized in Table 6. Preparation of D-serine derivatives was performed by synthetic

Table 3
Coupling of the P1 position for the inhibitors

Entry	Subst	Conditions ^a	Product	Entry	Subst	Conditions ^a	Product
1	5a	 A	3 (71%)	16	5a	 2A	6o (36%)
2	5a	 B	6a (3%)	17	5b	 A	7b (7%)
3	5a	 D	6b (25%)	18	5c	 A	7c (17%)
4	5a	 A	6c (78%)	19	5d	 A	7d (22%)
5	5a	 A	6d (43%)	20	5e	 A	7e (43%)
6	5a	 B	6e (47%)	21	5f	 A	7f (57%)
7	5a	 C	6f (27%)	22	5g	 A	7g (37%)
8	5a	 A	6g (70%)	23	5h	 A	7h (19%)
9	5a	Ac ₂ O	6h (64%)	24	5i	 A	7i (57%)
10	5a	 A	6i (26%)	25	5j	 A	7j (52%)
11	5a	 B	6j (47%)	26	5k	 A	7k (81%)
12	5a	 C	6k (95%)	27	5l	 A	7l (44%)
13	5a	 D	6l (4%)	28	5m	 A	7m (75%)
14	5a	 B	6m (13%)	29	5n	 A	7n (44%)
15	5a	 D	6n (26%)	30	5o	 A	7o (57%)

^a (A) carboxylic acid chloride or acetic anhydride, DMAP, Et₃N, CH₂Cl, (B) carboxylic acid, NMBA, DMAP, Et₃N, CH₂Cl₂, (C) carboxylic acid, 2,4,6-trichlorobenzoic acid chloride, DMAP, DIPEA, CH₂Cl₂, (D) carboxylic acid, (COCl)₂, DMF.

Table 4
Inhibitory activities of **5a** and cinnamoyl-L-serine derivatives (**6a–o**) with modification at the P'1 position



Entry	Compd	R ₃	IC ₅₀ ^a	ClogP ^b	Entry	Compd	R ₃	IC ₅₀ ^a	ClogP ^b
1	5a	H	1500	2.233	10	6i		250	4.312
2	6a		>3200	1.485	11	6j		180	3.382
3	6b		>1600	2.116	12	6k		175	3.482
4	6c		650	4.187	13	6l		170	4.584
5	6d		650	4.748	14	3		125	4.826
6	6e		560	3.482	15	6m		120	5.226
7	6f		550	6.085	16	6n		85	5.284
8	6g		450	4.367	17	6o		65	4.878
9	6h		400	3.129					

^a μM.


^b ClogP was calculated by ChemBio3D Ultra 12.0 (PerkinElmer).

protocols of the corresponding L-serine compounds without any problems. Fmoc-D-Ser(tBu)-OH as a starting material was selected and a 5 step sequence was performed as shown in Scheme 2. Interestingly, there were no extreme variants for IC₅₀ values between the L- and D-forms except for **6n** (entry 6). Both enantiomers of **5a**, **6a** and **7l** showed no inhibitory activities (entries 1, 2 and 3). The inhibitory activity of the L-forms of **6h**, **7o**, **6j**, **7n**, **6k** and **6o** showed a tendency to be more potent than those of the D-forms (entries 4, 5, 7–10). In contrast, **7i**, **7h**, **7j**, **7k** and **3**, which exhibited potent inhibitory activities against the 3CL protease, tended to have D-forms with more potent activities than L-forms (entries 11–15). We guessed the enantiomers of **7i**, **7h**, **7j**, **7k** and **3** fitted the active site of 3CL protease via different binding modes, that is, the P'1 site functionalities interact with the S4 pocket and consequently P4 site groups located in the S'1 pocket. These serine derivatives have high flexibility and therefore two hydrophobic spaces of the 3CL protease are recognized in a rigorous manner. The enantiomer of **3** (ent-**3**) exhibited the most potent inhibitory activity (IC₅₀ = 30 μM) in the evaluated compounds. Though there

are no correlation between the IC₅₀ value and tPSA in Table 6, these tPSA values were acceptable toward the protease inhibitor for the target of catalytic domain (Table 6).

The binding mode of **7k**, ent-**7k** and ent-**3** to SARS 3CL protease was predicted by docking simulations using GOLD from CCDC. These results suggest that reasonable inhibitors assessed by docking simulation can be designed and lead effectively to acceptable structures. Interestingly, the interactions between the enzyme and both enantiomers of serine derivatives were similar. In other words, the functionalities at the P'1, P1 and P4 sites of the D-serine template were located on the S'1, S1 and S4 pockets, respectively. This suggests the flexibility of backbones and recognition of the functionalities for serine type inhibitors. Therefore, the enantiomers of the designed compounds were also significant candidates as SARS 3CL protease inhibitors. Moreover, we may consider that D-serine plays a critical role in digestion by proteases (Fig. 2).

It is important to have enzyme selectivity for the development of protease inhibitors. We investigated inhibition of cathepsin B,²⁷

Table 5
Inhibitory activities of benzoyl-L-serine derivatives (**7b–k**) with modification at the P4 position

Entry	Compd	R ₂	IC ₅₀ ^a	ClogP ^b	Entry	Compd	R ₂	IC ₅₀ ^a	ClogP ^b
1	7b		>3200	2.955	6	7g		154	4.484
2	7c		500	4.673	7	7h		100	4.484
3	7d		240	4.745	8	7i		98	4.745
4	7e		220	3.968	9	7j		95	6.304
5	7f		155	4.745	10	7k		74	4.126

^a μM.^b ClogP was calculated by ChemBio3D Ultra 12.0 (PerkinElmer).**Table 6**
Inhibitory activities of D-serine derivatives

Entry	Compd	IC ₅₀ ^a	tPSA ^b	Entry	Compd	IC ₅₀ ^a	tPSA ^b
1	ent- 5a	>1600	78.43	9	ent- 6k	210	96.86
2	ent- 7l	>1600	75.71	10	ent- 6o	100	136.3
3	ent- 6a	1600	138.0	11	ent- 7i	85	93.73
4	ent- 6h	550	84.50	12	ent- 7h	80	103.0
5	ent- 7o	430	84.50	13	ent- 7j	68	110.8
6	ent- 6n	340	84.50	14	ent- 7k	65	112.2
7	ent- 6j	240	96.86	15	ent- 3	30	84.50
8	ent- 7n	225	84.50				

^a μM.^b tPSA was calculated by ChemBio3D Ultra 12.0 (PerkinElmer).

which is a mammalian cysteine protease involved in numerous pathological processes. The cysteine protease inhibitor E-64 was hardly inhibited against SARS 3CL protease in our previous report. E-64, miraziridine **A**²⁸ and tokaramide **A**²⁹ were reasonable inhibitors to bind the active site of cathepsin B. For this reason, the inhibitors of SARS 3CL protease are thought to show specificity against endogenous protease. Therefore, we attempted to evaluate the inhibitory activity of the serine derivatives against cathepsin B. The inhibitory activity toward cathepsin B was determined with an assay using a Z-Arg-Arg-MCA substrate developed by Hiwasa et al.^{30,31} The inhibition potency of serine derivatives was screened for cathepsin B inhibition at 1.0 mM concentration and subsequently, the IC₅₀ values were determined only for the selected compounds. The reasonable inhibitors, **3**, **7i**, **7j**, **7h** and **7k** and its enantiomers, for SARS 3CL R188I mutant protease and **6m**, **7f** and **7g**, which had good potential as cathepsin B inhibitors, are shown in Table 7. As a result, the serine derivatives that exhibited

potent inhibitory activity against SARS 3CL protease showed relatively less potent inhibitory effects against cathepsin B activity and had high protease selectivity (entries 1–10). On the other hand, three compounds were extracted by the screening for a cathepsin B inhibitor (entries 11–13). These compounds were useless for SARS 3CL protease inhibition (Table 7).

The inhibitory activities of selected serine derivatives, **6m**, **7f** and **7g**, were evaluated using the corresponding IC₅₀ values. The IC₅₀ value of synthetic **7f** was 170 μM. Comparing IC₅₀ values against SARS 3CL protease and cathepsin B, it was strongly suggested that the inhibitory activity is attributable mainly to the ester site of **7f**.³² The results are consistent with the structural data on **7f** in a complex with cathepsin B, which suggests that the P1 site of **7f** can easily adopt a conformation that is similar to the binding mode of E-64d. Although the P2 site had a rather weak effect compared with the P1 site, the inhibitory activity of **7f** was about 10 times that of **3**. To estimate the inhibitory mechanism, an inhibitory kinetics experiment with **7f** was performed by making a Lineweaver–Burk plot. The rate of cleavage for different amounts of substrate by cathepsin B in the absence or presence of **7f** (100, 500 μM) was monitored using Fluorescence spectrometer and consequently competitive inhibition toward cathepsin B was assessed.

2.4. Cytotoxicity against HeLa cells

The cytotoxic ability of serine derivatives was also important because any protease inhibitor needs to have an orthogonal relationship between anti-virus activity and cytotoxicity. A cytotoxicity assay was performed against HeLa cells using synthesized serine derivatives for cytotoxicity alongside rotenone, which




Figure 2. Docking simulation of selected inhibitors bound to SARS 3CL protease (PDB code 3AW1) using GOLD from CCDC. Molecular graphic image shown using PyMOL from Schrödinger; oxygen (red) and nitrogen (blue) of inhibitors; Cys145 (red) of SARS 3CL protease. (A) Surface mode with **7k**, (B) with ent-**7k**, (C) with ent-**3**, (D) model of interaction with ent-**3**.

Table 7
Inhibitory activities against cathepsin B

Entry	Compd	Inhibition ^a (%)	Entry	Compd	Inhibition ^a (%)
1	3	34	8	ent- 7h	22
2	ent- 3	NI ^b	9	7k	21
3	7i	36	10	ent- 7k	70
4	ent- 7i	44	11	6m	82
5	7j	35	12	7f	67
6	ent- 7j	19	13	7g	88
7	7h	NI ^b			

^a Inhibitory activity was measured by 1 mM.

^b NI: no inhibition.

served as the control. The cytotoxicities of these compounds were determined by measuring live-cell succinate-tetrazolium reductase activity (MTT assay). Although the serine derivatives, both enantiomers of **3**, **7i**, **7j**, **7h** and **7k** at 200 μM concentrations were evaluated against Hela cells (5000 cells), the cell death ratio caused by these compounds was less than 20% and therefore showed practically no cytotoxicity.

3. Conclusion

Inhibitory activities of ent-**3** against SARS 3CL protease and cathepsin B showed an IC_{50} value of 30 μM and no inhibition at 1 mM concentration, respectively. Moreover, ent-**7k** has inhibitory effects with $\text{IC}_{50} = 65 \mu\text{M}$ for SARS 3CL protease and $\text{IC}_{50} = 500 \mu\text{M}$ for cathepsin B. These results indicate high specificity between the two proteases. In addition, the ent-**3** and ent-**7k** inhibitors showed no cytotoxicity and therefore have good potential as drugs. Ent-**3** and ent-**7k** were chosen for the next stages and designated as **SK23** and **SK69**, respectively (Fig. 3).




Figure 3. Optimized inhibitors ent-**3** (SK23) and ent-**7k** (SK69).

In conclusion, we found effective serine derivatives toward the SARS 3CL protease inhibitor by the combination of docking simulation and a structure activity relationship study. Consequently, two attractive inhibitors, ent-**3** (SK23) and ent-**7k** (SK69), which were derived from the tetrapeptide aldehyde (**1**) and Bai's inhibitor were produced. These inhibitors showed protease selectivity and no cytotoxicity and therefore a useful methodology using a serine template will facilitate drug design. Especially, we testified that ent-**3** (SK23) derived from the virtual screening was most reasonable inhibitor. To improve the inhibitory activity against the SARS 3CL protease, molecular design and evaluation are now underway.

4. Experimental

4.1. General information

All solvents were reagent grade. CH_2Cl_2 was distilled from CaH_2 . All commercial reagents were of the highest purity available. Amino acid derivatives were purchased from Watanabe Chemical Industries. Other chemicals were purchased from Aldrich, Wako chemical and Nacal tesque. Optical rotations were determined

with a JASCO DIP-371 polarimeter at the sodium D line. IR spectra of sample were obtained as films with a HORIBA FT-720 spectrometer. ^1H (400 and 500 MHz) and ^{13}C NMR (100 and 125 MHz) were determined on a JNM-ECX400 and JNM-ECX500. Chemical shifts are reported in ppm with reference to tetramethylsilane [^1H NMR: TMS (0.00)], or solvent signals [^1H NMR: CDCl_3 (7.26), $\text{MeOH-}d_4$ (3.30); ^{13}C NMR: CDCl_3 (77.16), $\text{MeOH-}d_4$ (50.05)]. Mass spectra were recorded on a JEOL AccuTOF JMS-T100LC (ESI). Analytical TLC was performed on Merck Silica gel 60F₂₅₄. Crude products were purified by column chromatography on Silica Gel 60 N [Kanto, particle size, (spherical, neutral) 63–210 μm or 100–200 μm]. HPLC system (monitored at 220 nm) equipped with Cosmosil 5C18 AR-II column (4.6 \times 150 mm) using MeCN in 0.1% aqueous TFA. Absorbance for enzyme inhibition assay was measured with the HITACHI F-7000. Absorbance for cytotoxicity assay was measured with the CORONA MTP-310Lab.

4.2. Coupling of P4 position and tBu-deprotection of amine

4.2.1. (S)-N-Cinnamoylserinecyclohexylamide (5a)

To a solution of amine (**4a**) (4.45 g, 11.3 mmol) in CH_2Cl_2 were added cinnamic acid (3.35 g, 22.6 mmol), HATU (8.59 g, 22.6 mmol), HOAt (3.08 g, 22.6 mmol) and DIPEA (7.80 ml, 45.2 mmol) and the mixture was stirred at room temperature for 8 h. After CH_2Cl_2 and H_2O were added to the solution, the organic layer was washed with brine and dried over MgSO_4 . Organic layer was evaporated in vacuo and the residue was purified by silica gel column chromatography (hexane/ AcOEt = 4:1) to give L-serine derivative (6.16 g, 16.5 mmol). To a solution of L-serine derivative was added TFA (3.0 ml) at room temperature. After being stirred for 2 h, the TFA was carried out azeotropic removal with MeOH and CH_2Cl_2 in vacuo. The residue was purified by silica gel column chromatography (CHCl_3) to give alcohol (**5a**) (1.36 g, 4.30 mmol, 38%) as a white powder, mp 185–188 °C. [α] $^{25}_D$ -78° (c 0.13, CHCl_3). IR (film) ν_{max} cm^{-1} : 3276, 2930, 1645, 1618, 1542, 1215, 1052, 975, 752, 666. ^1H NMR (500 MHz, CDCl_3) δ : 1.15–1.36 (6H, m), 1.56–1.72 (3H, m), 1.82–1.91 (2H, m), 3.69–3.77 (2H, m), 4.17 (1H, dd, J = 12.0, 3.0 Hz), 4.57 (1H, t, J = 3.5 Hz), 6.51 (1H, d, J = 15.0 Hz), 7.36–7.37 (3H, m), 7.49–7.51 (2H, m), 7.65 (1H, d, J = 16.0 Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.7, 25.6, 32.8, 32.9, 48.5, 53.8, 63.1, 90.6, 119.7, 128.1, 129.1, 130.3, 134.6, 142.5, 166.9, 170.4. HR-MS (ESI) m/z . Calcd for $\text{C}_{18}\text{H}_{24}\text{N}_2\text{O}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 339.1658. Found 339.1700. ent-**5a**: mp 183–185 °C. [α] $^{27}_D$ $+79^\circ$ (c 0.27, CHCl_3), HRMS (ESI) Found 339.1668.

4.2.2. (S)-N-(p-Hydroxyphenyl)propionylserinecyclohexylamide (5b)

(0.261 g, 0.780 mmol, 79%) as an oil. [α] $^{28}_D$ -2.7° (c 0.75, MeOH). IR (film) ν_{max} cm^{-1} : 3290, 2935, 2858, 1671, 1516, 1452, 1352, 1202, 1132, 1026, 892, 833, 758. ^1H NMR (500 MHz, CD_3OD) δ : 1.20–1.85 (10H, m), 2.61 (2H, t, J = 8.0 Hz), 2.79 (2H, t, J = 8.0 Hz), 3.61–3.88 (5H, m), 4.04–4.06 (1H, m), 4.30–4.33 (1H, m), 4.42–4.46 (1H, m), 6.65 (2H, d, J = 8.5 Hz), 6.97 (2H, d, J = 8.0 Hz). ^{13}C NMR (100 MHz, CD_3OD) δ : 24.8, 25.2, 30.5, 32.3, 37.7, 48.3, 55.4, 61.8, 114.9, 129.0, 131.5, 155.5, 169.8, 174.1. HR-MS (ESI) m/z : calcd for $\text{C}_{18}\text{H}_{26}\text{N}_2\text{O}_4\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 357.1790. Found 357.1800.

4.2.3. (S)-N-(E)-4-phenylbut-2-enoicserinecyclohexylamide (5c)

(0.278 g, 0.841 mmol, 37%) as a white powder, mp 150–152 °C. [α] $^{26}_D$ -3.0° (c 0.10, CHCl_3). IR (film) ν_{max} cm^{-1} : 3282, 3059, 2933, 2854, 1749, 1655, 1583, 1545, 1475, 1456. ^1H NMR (400 MHz, CD_3OD) δ : 1.11–1.88 (11H, m), 4.58–4.67 (1H, m), 3.93 (2H, s), 4.36–4.42 (1H, m), 4.51–4.56 (1H, m), 6.33 (1H, d, J = 16.0 Hz), 6.53 (1H, d, J = 16.0 Hz), 7.17–7.21 (1H, m), 7.26–7.29 (2H, m), 7.37 (2H, d, J = 7.2 Hz). ^{13}C NMR (100 MHz, CD_3OD) δ : 24.8, 25.3, 26.3,

32.3, 37.8, 39.5, 54.3, 55.5, 57.4, 61.8, 122.5, 126.0, 127.2, 128.2, 133.5, 167.6, 170.0, 172.7. HR-MS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_3$ Na [$\text{M}+\text{Na}$] $^+$ 353.1841. Found 353.1805.

4.2.4. (S)-N-(E)-4'-Methoxy-3-phenylprop-2-enoicserinecyclohexylamide (5d)

(0.228 g, 0.659 mmol, 43%) as a yellow powder, mp 226–228 °C. [α] $^{23}_D$ $+1.0^\circ$ (c 0.10, CHCl_3). IR (film) ν_{max} cm^{-1} : 3282, 2933, 2856, 2777, 1724, 1657, 1603, 1552, 1485, 1412. ^1H NMR (400 MHz, CDCl_3) δ : 1.10–1.92 (11H, m), 4.02 (3H, s), 4.08–4.22 (2H, m), 4.51–4.54 (1H, m), 6.37 (1H, d, J = 15.6 Hz), 6.89 (2H, d, J = 9.2 Hz), 7.45 (2H, d, J = 8.7 Hz), 7.60 (1H, d, J = 15.6 Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.3, 32.3, 48.7, 54.5, 55.6, 57.5, 61.8, 62.0, 114.0, 117.6, 129.2, 129.2, 140.8, 150.2, 161.4, 167.8, 169.8, 169.9, 170.2. HR-MS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_4\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 369.1790. Found 369.1783.

4.2.5. (S)-N-Benzoylserinecyclohexylamide (5e)

(0.247 g, 0.851 mmol, 80%) as a white powder, mp 189–191 °C. [α] $^{28}_D$ $+26^\circ$ (c 0.10, CHCl_3). IR (film) ν_{max} cm^{-1} : 3290, 3072, 2933, 2854, 2355, 2326, 1635, 1548, 1489, 1448. ^1H NMR (400 MHz, CDCl_3) δ : 1.10–1.95 (11H, m), 3.48–3.79 (2H, m), 4.23–4.28 (1H, m), 4.52–4.56 (1H, m), 7.43–7.56 (3H, m), 7.81 (2H, d, J = 11.6 Hz); ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.3, 32.3, 48.3, 56.0, 62.0, 127.2, 128.3, 131.6, 133.8, 168.7, 170.1. HR-MS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{22}\text{N}_2\text{O}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 313.1528. Found 313.1514.

4.2.6. (S)-N-(E)-2'-Methoxy-3-phenylprop-2-enoicserinecyclohexylamide (5f)

(34.0 mg, 98.2 μmol , 7.0%) as a white powder, mp 189–192 °C. [α] $^{25}_D$ $+6.0^\circ$ (c 0.10, MeOH). IR (film) ν_{max} cm^{-1} : 3282, 2933, 2846, 1649, 1539, 1489, 1452, 1248, 1201, 1180; ^1H NMR (400 MHz, CDCl_3) δ : 1.07–1.95 (11H, m), 3.62–3.78 (2H, m), 3.89 (3H, s), 4.19 (1H, dd, J = 12.0, 2.8 Hz), 4.48–4.54 (1H, m), 6.65 (1H, d, J = 16.0 Hz), 6.86–6.99 (2H, m), 7.30–7.37 (1H, m), 7.47 (1H, d, J = 8.0 Hz), 7.88 (1H, d, J = 16.0 Hz); ^{13}C NMR (100 MHz, CDCl_3) δ : 22.9, 24.8, 25.3, 29.3, 32.3, 54.7, 55.6, 62.0, 111.1, 120.4, 123.5, 128.1, 131.0, 136.4, 158.4, 167.8, 170.1. HR-MS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_4\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 369.1790. Found 369.1799.

4.2.7. (S)-N-(E)-3',4'-Dimethoxy-3-phenylprop-2-enoicserinecyclohexylamide (5g)

(0.100 g, 0.266 mmol, 23%) as a yellow powder, mp 220 °C. [α] $^{26}_D$ $+28^\circ$ (c 0.10, MeOH). IR (film) ν_{max} cm^{-1} : 3276, 3099, 3001, 2931, 2846, 1643, 1610, 1543, 1514, 1452. ^1H NMR (400 MHz, CDCl_3) δ : 1.09–1.94 (11H, m), 3.66–3.81 (1H, m), 3.89 (6H, s), 6.43 (1H, d, J = 16.0 Hz), 6.80–6.89 (1H, m), 7.00–7.04 (1H, m), 7.08 (1H, d, J = 6.4 Hz), 7.58 (1H, d, J = 16.0 Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.7, 25.5, 32.6, 38.8, 48.4, 54.0, 56.1, 63.2, 109.6, 111.1, 117.3, 122.5, 127.4, 142.4, 149.3, 151.0, 167.3; HR-MS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_5\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 399.1896. Found 399.1907.

4.2.8. (S)-N-(E)-3',5'-Dimethoxy-3-phenylprop-2-enoicserinecyclohexylamide (5h)

(0.236 g, 0.627 mmol, 29%) as a white powder, mp 175–177 °C. [α] $^{26}_D$ $+22^\circ$ (c 0.10, CHCl_3). IR (film) ν_{max} cm^{-1} : 3269, 2924, 2846, 2355, 2326, 1647, 1606, 1558, 1541, 1522. ^1H NMR (400 MHz, CDCl_3) δ : 1.10–1.92 (11H, m), 3.62–3.76 (1H, m), 3.83 (6H, s), 4.12–4.21 (1H, m), 4.48–4.56 (1H, m), 6.41 (1H, d, J = 16.0 Hz), 6.47 (1H, s), 6.65 (2H, s), 7.56 (1H, d, J = 15.3 Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.7, 25.5, 29.8, 32.8, 35.1, 48.3, 53.4, 55.5, 57.1, 63.0, 100.0, 102.0, 104.2, 105.9, 120.1, 126.3, 136.4, 142.3, 143.0, 161.1. HR-MS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_5\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 399.1896. Found 399.1872.

4.2.9. (S)-N-(E)-3'-Methoxy-3-phenylprop-2-enoicserinecyclohexylamide (5i)

(0.290 g, 0.838 mmol, 69% in 2 steps) as a white powder, mp 180–181 °C. $[\alpha]_D^{26} +25^\circ$ (c 0.10, MeOH). IR (film) ν_{\max} cm^{-1} : 3269, 3086, 2931, 2854, 1647, 1620, 1577, 1547, 1491, 1448; ^1H NMR (400 MHz, CDCl_3) δ : 1.11–2.11 (11H, m), 3.63–3.68 (1H, m), 3.83 (3H, s), 4.19–4.25 (1H, m), 4.46–4.50 (1H, m), 6.45 (1H, d, $J = 16.0$ Hz), 6.76–6.83 (1H, m), 6.92 (1H, dd, $J = 8.8, 2.8$ Hz), 7.03 (1H, s), 7.11 (1H, d, $J = 8.0$ Hz), 7.62 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, $\text{MeOH}-d_4$) δ : 24.8, 25.3, 32.3, 54.4, 55.7, 62.0, 112.6, 115.3, 120.1, 120.5, 129.6, 136.3, 140.9, 160.2, 161.4, 167.1, 170.1. HR-MS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 369.1790. Found 369.1798.

4.2.10. (S)-N-(E)-4'-hydroxy-3-phenylprop-2-enoicserinecyclohexylamide (5j)

(0.170 g, 0.511 mmol, 55%) as a white powder, mp 220–222 °C. IR (film) ν_{\max} cm^{-1} : 3338, 3269, 2918, 2854, 2495, 2432, 1655, 1641, 1606, 1570. ^1H NMR (400 MHz, $\text{MeOH}-d_4$) δ : 1.10–1.90 (11H, m), 3.60–3.68 (1H, m), 3.76 (1H, d, $J = 5.2$ Hz), 4.49 (1H, t, $J = 6.0$ Hz), 6.52 (1H, d, $J = 15.6$ Hz), 6.77 (2H, d, $J = 8.7$ Hz), 7.41 (2H, d, $J = 8.7$ Hz), 7.46 (1H, d, $J = 15.6$ Hz). ^{13}C NMR (100 MHz, $\text{MeOH}-d_4$) δ : 24.8, 25.3, 32.3, 48.6, 55.6, 62.0, 115.4, 116.6, 126.2, 129.4, 141.2, 159.4, 167.8, 170.1. HR-MS (ESI) m/z : calcd for $\text{C}_{18}\text{H}_{24}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 355.1634. Found 355.1639.

4.2.11. (S)-N-(E)-3',4',5'-Trimethoxy-3-phenylprop-2-enoicserinecyclohexylamide (5k)

(0.104 g, 0.256 mmol, 29%) as a white powder, mp 156–162 °C. $[\alpha]_D^{26} +31^\circ$ (c 0.10, MeOH). IR (film) ν_{\max} cm^{-1} : 3298, 2933, 2846, 1649, 1616, 1583, 1542, 1506, 1452, 1419. ^1H NMR (400 MHz, CDCl_3) δ : 1.09–1.96 (11H, m), 3.60–3.78 (4H, m), 3.84 (9H, s), 4.10 (1H, dd, $J = 8.4, 4.0$ Hz), 4.57–4.64 (1H, m), 6.44 (1H, d, $J = 16.0$ Hz), 6.70 (2H, s), 7.04 (1H, d, $J = 8.0$ Hz), 7.12 (1H, d, $J = 5.6$ Hz), 7.52 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.7, 25.5, 32.8, 48.6, 54.4, 56.2, 61.0, 63.4, 105.2, 119.3, 130.0, 140.0, 142.1, 153.5, 166.8, 169.9. HR-MS (ESI) m/z : calcd for $\text{C}_{21}\text{H}_{30}\text{N}_2\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 429.2002. Found 429.2009.

4.2.12. (S)-N-Cinnamoylserinepiperidineamide (5l)

(95.4 mg, 26%) as an oil, $[\alpha]_D^{25} +16^\circ$ (c 0.40, MeOH). IR (film) ν_{\max} cm^{-1} : 3286, 3060, 3027, 2939, 2860, 2245, 1660, 1608, 1549, 1450. ^1H NMR (400 MHz, CDCl_3) δ : 1.53–1.69 (6H, m), 3.50–3.56 (2H, m), 3.58–3.66 (1H, m), 3.82 (2H, s), 5.06–5.11 (1H, m), 6.50 (1H, d, $J = 16.0$ Hz), 7.32–7.38 (3H, m), 7.46–7.51 (2H, m), 7.62 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.4, 25.6, 26.5, 43.6, 47.0, 50.9, 52.0, 119.9, 128.0, 128.9, 130.0, 142.1, 166.5, 168.2. HR-MS (ESI) m/z : calcd for $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 325.1528. Found 325.1540. ent-5l: $[\alpha]_D^{25} -16^\circ$ (c 0.20, MeOH). HR-MS (ESI) m/z : found 325.1514.

4.2.13. (S)-N-Cinnamoylserinemorpholineamide (5m)

(196 mg, 0.644 mmol, 91%) as an oil, $[\alpha]_D^{25} +8.5^\circ$ (c 0.55, CHCl_3). IR (film) ν_{\max} cm^{-1} : 3303, 3059, 3016, 2966, 2931, 2862, 2368, 2305, 1736, 1620. ^1H NMR (400 MHz, CDCl_3) δ : 3.57–3.74 (8H, m), 3.79 (1H, dd, $J = 11.2, 4.0$ Hz), 3.88 (1H, dd, $J = 11.6, 4.8$ Hz), 4.38 (1H, s), 5.06–5.15 (1H, m), 6.52 (1H, d, $J = 16.0$ Hz), 7.30–7.40 (3H, m), 7.44–7.52 (2H, m), 7.62 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 42.8, 46.4, 51.1, 64.1, 66.7, 119.9, 128.7, 130.1, 134.5, 142.3, 166.4, 169.3. HR-MS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{20}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 327.1321. Found 327.1286. ent-5m: $[\alpha]_D^{27} -7.1^\circ$ (c 0.20, CHCl_3). HR-MS (ESI) m/z : found 327.1302.

4.2.14. (S)-N-Cinnamoylserinebenzylamide (5n)

(111 mg, 0.342 mmol, 19%) as an oil. $[\alpha]_D^{26} +46^\circ$ (c 0.10, MeOH). IR (film) ν_{\max} cm^{-1} : 3435, 3400, 3369, 3278, 1645, 1618, 1215,

1051. ^1H NMR (400 MHz, CDCl_3) δ : 3.64–3.72 (1H, m), 4.25–4.31 (1H, m), 4.41–4.49 (2H, m), 4.54–4.60 (1H, m), 4.46 (1H, d, $J = 16.0$ Hz), 6.78 (1H, s), 7.26–7.35 (4H, m), 7.38 (3H, t, $J = 3.2$ Hz), 7.50 (2H, d, $J = 5.6$ Hz), 7.65 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, $\text{MeOH}-d_4$) δ : 42.8, 55.9, 61.9, 120.4, 126.8, 127.6, 128.6, 129.6, 134.9, 138.4, 141.1, 167.4, 171.2. HR-MS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 347.1372. Found 347.1353. ent-5n: mp 171–176 °C. HR-MS (ESI) m/z : found 347.1361.

4.2.15. (S)-N-Cinnamoylserinecyclohexanemethylamide (5o)

(399 mg, 1.21 mmol, 48%) as a white powder, mp 183–184 °C. $[\alpha]_D^{26} +21^\circ$ (c 0.10, MeOH), IR (film) ν_{\max} cm^{-1} : 3273, 3107, 3047, 2922, 2850, 2368, 2326, 1954, 1884, 1651. ^1H NMR (400 MHz, CDCl_3) δ : 0.83–0.97 (2H, m), 1.05–1.29 (3H, m), 1.40–1.53 (1H, m), 1.59–1.74 (6H, m), 3.05–3.14 (2H, m), 3.75–3.85 (2H, m), 3.75–3.84 (1H, m), 4.18 (1H, dd, $J = 12.0, 4.0$ Hz), 4.64–4.71 (1H, m), 6.57 (1H, d, $J = 16.0$ Hz), 7.32–7.40 (3H, m), 7.45–7.53 (2H, m), 7.65 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 25.8, 26.4, 30.8, 37.8, 38.8, 46.0, 54.2, 63.0, 119.3, 128.1, 129.0, 130.3, 134.3, 142.9, 166.1, 167.4, 171.2. HR-MS (ESI) m/z : calcd for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 353.1841. Found 353.1818. ent-5o: mp 181–183 °C. $[\alpha]_D^{27} -24^\circ$ (c 0.10, MeOH), HR-MS (ESI) m/z : found 353.1818.

4.3. Coupling of P1' position for the inhibitors

4.3.1. (S)-N-Cinnamoyl serine(benzoyl)cyclohexylamide (3)

To a solution of 5a (19.6 mg, 0.0619 mmol) in CH_2Cl_2 were added Benzoyl chloride (14.3 μl , 0.124 mmol), DMAP (15.1 mg, 0.124 mmol) and DIPEA (53.4 μl , 0.310 mmol) at 0 °C, and the mixture was stirred at room temperature for 3 h. After CH_2Cl_2 and H_2O were added to the solution, the organic layer was washed with brine and dried over MgSO_4 . The solvent was evaporated in vacuo and the residue was purified by silica gel column chromatography (hexane/AcOEt = 4:1) to give 3 (18.6 mg, 0.0442 mmol, 71%) as a white powder, mp 200–203 °C. $[\alpha]_D^{24} -1.5^\circ$ (c 0.76, CHCl_3). IR (film) ν_{\max} cm^{-1} : 3282, 2930, 2849, 2364, 1727, 1646, 1617, 1541, 1450, 1270, 1112, 1025, 975, 710. ^1H NMR (400 MHz, CDCl_3) δ : 1.09–1.39 (6H, m), 1.66 (2H, m), 1.88 (2H, m), 3.77 (1H, m), 4.60 (1H, dd, $J = 11.2, 5.6$ Hz), 4.71 (1H, dd, $J = 11.2, 6.4$ Hz), 5.00 (1H, q, $J = 6.0$ Hz), 6.50 (1H, d, $J = 7.2$ Hz), 6.52 (1H, d, $J = 15.6$ Hz), 6.97 (1H, d, $J = 7.6$ Hz), 7.36–7.61 (8H, m), 7.64 (1H, d, $J = 15.6$ Hz), 8.02 (2H, d, $J = 7.6$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.8, 25.5, 32.9, 48.8, 52.8, 64.2, 120.1, 128.1, 128.6, 129.0, 129.9, 130.1, 130.3, 133.4, 133.6, 134.7, 136.8, 141.8, 142.2, 166.3, 166.5, 168.2, 170.4. HR-MS (ESI) m/z : calcd for $\text{C}_{25}\text{H}_{28}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 443.1947. Found 443.1913. ent-3: mp 205–206 °C. $[\alpha]_D^{27} +1.4^\circ$ (c 0.14, CHCl_3). HR-MS (ESI) m/z : found 443.1906.

4.3.2. (S)-N-Cinnamoylserine(3-(R)-pyroglutamoyl)cyclohexylamide (6b)

(10.9 mg, 0.0255 mmol, 25%). $[\alpha]_D^{26} +0.87^\circ$ (c 0.23, CHCl_3). IR (film) ν_{\max} cm^{-1} : 3418, 2933, 2854, 2360, 2341, 1749, 1647, 1558, 1541, 1456, 1192. ^1H NMR (400 MHz, CDCl_3) δ : 1.15–1.88 (10H, m), 2.29–2.44 (4H, m), 3.74 (1H, br), 4.28–4.39 (2H, m), 4.53–4.57 (1H, m), 4.96 (1H, br), 6.51 (1H, d, $J = 15.6$ Hz), 6.79 (1H, d, $J = 7.2$ Hz), 7.12 (1H, d, $J = 7.2$ Hz), 7.31–7.50 (6H, m), 7.66 (1H, d, $J = 15.6$ Hz), 7.72 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.3, 24.4, 24.9, 25.5, 29.5, 32.9, 48.9, 52.1, 52.3, 65.3, 119.8, 128.0, 129.0, 130.2, 134.5, 142.5, 166.3, 167.5, 171.8, 178.4. HR-MS (ESI) m/z : calcd for $\text{C}_{23}\text{H}_{29}\text{N}_3\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 450.2005. Found 450.2041.

4.3.3. (S)-N-Cinnamoylserine(butanoyl)cyclohexylamide (6c)

(23.6 mg, 0.0611 mmol, 78%) as a white powder, mp 176–182 °C. $[\alpha]_D^{25} -0.6^\circ$ (c 0.17, CHCl_3). IR (film) ν_{\max} cm^{-1} : 3282,

3069, 2933, 2855, 1741, 1649, 1620, 1546, 1449, 1344, 1172, 1092, 977. ^1H NMR (500 MHz, CDCl_3) δ : 1.13–1.37 (1H, m), 1.60–1.72 (4H, m), 1.91 (2H, m), 3.73–3.79 (1H, m), 4.29 (1H, dd, $J = 11.5$, 5.0 Hz), 4.49 (1H, dd, $J = 11.5$, 6.0 Hz), 4.79 (1H, q, $J = 6.3$ Hz), 6.13 (1H, br), 6.45 (1H, d, $J = 15.5$ Hz), 6.59 (1H, br), 7.37–7.39 (3H, m), 7.51–7.52 (2H, m), 7.64 (1H, d, $J = 15.5$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.9, 25.6, 27.3, 33.1, 39.0, 48.8, 53.0, 64.1, 119.9, 128.1, 129.0, 130.2, 134.6, 142.3, 166.1, 167.8, 178.8. HR-MS (ESI) m/z : calcd for $\text{C}_{22}\text{H}_{30}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 409.2103. Found 409.2080.

4.3.4. (S)-N-Cinnamoylserine(phenylacetyl)cyclohexylamide (6d)

(18.0 mg, 0.0414 mmol, 67%) as a white powder, mp 172–177 °C. $[\alpha]_{\text{D}}^{27}$ 0° (c 0.10, MeOH). IR (film) ν_{max} cm^{-1} : 3649, 3583, 3282, 3064, 3030, 2931, 2854, 2362, 2312, 1749. ^1H NMR (400 MHz, CDCl_3) δ : 0.95–1.20 (5H, m), 1.21–1.40 (4H, m), 1.52–1.73 (4H, m), 1.76–1.89 (3H, m), 3.67–3.77 (1H, m), 4.27–4.35 (1H, m), 4.46–4.54 (1H, m), 4.73–4.82 (1H, m), 6.09–6.19 (1H, m), 6.34 (1H, d, $J = 16.0$ Hz), 6.49 (1H, d, $J = 8.0$ Hz), 7.26–7.33 (5H, m), 7.34–7.40 (3H, m), 7.46–7.52 (1H, m), 7.60 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 32.9, 41.3, 48.7, 52.5, 64.3, 119.9, 127.3, 128.0, 128.8, 129.0, 129.4, 130.1, 133.7, 134.6, 142.1, 166.0, 167.6, 171.5. HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 457.2103. Found 457.2079.

4.3.5. (S)-N-Cinnamoylserine(isonicotinoyl)cyclohexylamide (6e)

(13.2 mg, 0.0313 mmol, 47%) as a white powder, mp 210–212 °C. $[\alpha]_{\text{D}}^{24}$ -6.1° (c 0.08, CHCl_3). IR (film) ν_{max} cm^{-1} : 3282, 2929, 2854, 2360, 2341, 1741, 1645, 1616, 1560, 1539, 1211, 1144, 1066, 974. ^1H NMR (400 MHz, CDCl_3) δ : 1.09–1.39 (4H, m), 1.59–1.71 (4H, m), 1.89 (2H, m), 3.74–3.81 (1H, m), 4.64–4.72 (2H, m), 5.05 (1H, q, $J = 6.4$ Hz), 6.50 (1H, d, $J = 15.6$ Hz), 6.66 (1H, d, $J = 7.6$ Hz), 6.86 (1H, d, $J = 7.6$ Hz), 7.36–7.37 (3H, m), 7.48–7.50 (2H, m), 7.63 (1H, d, $J = 15.6$ Hz), 7.82 (2H, d, $J = 6.0$ Hz), 8.76 (2H, d, $J = 5.6$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.8, 25.5, 33.0, 48.9, 52.6, 65.4, 119.8, 123.0, 128.1, 129.1, 130.3, 134.5, 136.9, 142.5, 150.8, 165.1, 166.2, 167.5. HR-MS (ESI) m/z : calcd for $\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 444.1899. Found 444.1864.

4.3.6. (S)-N-Cinnamoylserine(2,4,6-trichlorobenzoyl)cyclohexylamide (6f)

(9.00 mg, 0.0172 mmol, 27%) as a white powder, mp 208–211 °C. $[\alpha]_{\text{D}}^{26}$ -19° (c 0.67, CHCl_3). IR (film) ν_{max} cm^{-1} : 3275, 2931, 2854, 2360, 2330, 1747, 1651, 1620, 1558, 1520, 1458, 1273, 1119, 1057, 972, 852, 752. ^1H NMR (500 MHz, CDCl_3) δ : 1.13–1.38 (4H, m), 1.58–1.69 (4H, m), 1.86–1.90 (2H, m), 3.70–3.79 (1H, m), 4.61 (1H, dd, $J = 11.5$, 5.0 Hz), 4.87 (1H, dd, $J = 11.5$, 6.5 Hz), 4.93 (1H, q, $J = 6.3$ Hz), 6.33 (1H, d, $J = 8.5$ Hz), 6.45 (1H, d, $J = 16.0$ Hz), 6.62 (1H, d, $J = 8.0$ Hz), 7.34–7.38 (5H, m), 7.49–7.51 (2H, m), 7.65 (1H, d, $J = 15.5$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.8, 25.6, 32.9, 48.8, 52.6, 65.4, 119.7, 128.1, 128.3, 129.0, 130.3, 132.8, 134.5, 134.7, 142.6, 164.1, 166.3, 167.5. HR-MS (ESI) m/z : calcd for $\text{C}_{25}\text{H}_{25}\text{Cl}_3\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 545.0778. Found 545.0815.

4.3.7. (S)-N-Cinnamoylserine(pivaloyl)cyclohexylamide (6g)

(20.6 mg, 0.0514 mmol, 70%) as a white powder, mp 183–186 °C. $[\alpha]_{\text{D}}^{28}$ $+0.7^\circ$ (c 0.28, CHCl_3). IR (film) ν_{max} cm^{-1} : 3282, 2932, 2856, 2364, 1737, 1646, 1620, 1547, 1450, 1215, 1155, 979; ^1H NMR (400 MHz, CDCl_3) δ : 0.94 (3H, t, $J = 7.4$ Hz), 1.13–1.41 (5H, m), 1.60–1.69 (7H, m), 1.89 (2H, m), 2.32 (2H, t, $J = 7.2$ Hz), 3.78 (1H, m), 4.30 (1H, dd, $J = 11.4$, 5.5 Hz), 4.79 (1H, q, $J = 6.3$ Hz), 6.22 (1H, br), 6.46 (1H, d, $J = 16.0$ Hz), 6.64 (1H, br), 7.37–7.38 (3H, m), 7.51–7.52 (2H, m), 7.64 (1H, d, $J = 15.6$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 13.8, 18.5, 24.8, 25.6, 33.0, 36.1,

48.7, 52.9, 63.9, 119.9, 128.1, 129.0, 130.2, 134.6, 142.4, 167.8, 173.9. HR-MS (ESI) m/z : calcd for $\text{C}_{23}\text{H}_{32}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 423.2260. Found 423.2227.

4.3.8. (S)-N-Cinnamoylserine(acetyl)cyclohexylamide (6h)

(18.4 mg, 0.0513 mmol, 64%) as a white powder, mp 191–194 °C. $[\alpha]_{\text{D}}^{26}$ -3.0° (c 0.81, CHCl_3). IR (film) ν_{max} cm^{-1} : 3298, 3098, 3093, 2929, 2854, 2360, 2332, 1743, 1647, 1616, 1558, 1540, 1448, 1375, 1246, 1219, 1080, 1049, 974. ^1H NMR (500 MHz, CDCl_3) δ : 1.11–1.25 (3H, m), 1.29–1.40 (2H, m), 1.59–1.61 (1H, m), 1.68–1.73 (2H, m), 1.84–1.94 (2H, m), 2.07 (3H, s), 3.77 (1H, m), 4.34 (1H, dd, $J = 11.5$, 5.5 Hz), 4.48 (1H, dd, $J = 11.5$, 6.0 Hz), 4.90 (1H, q, $J = 6.0$ Hz), 6.53 (1H, d, $J = 16.5$ Hz), 6.66 (1H, d, $J = 8.0$ Hz), 6.95 (1H, d, $J = 8.0$ Hz), 7.34–7.36 (3H, m), 7.47–7.50 (2H, m), 7.47–7.50 (2H, m), 7.64 (1H, d, $J = 15.5$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 21.0, 24.9, 25.6, 33.0, 48.7, 52.6, 64.2, 120.1, 129.0, 130.1, 134.7, 142.1, 166.2, 167.9, 171.0. HR-MS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 381.1790. Found 381.1793. ent-6h: mp 185–187 °C. $[\alpha]_{\text{D}}^{27}$ $+2.4^\circ$ (c 0.17, CHCl_3). HRMS (ESI). Found 381.1749.

4.3.9. (S)-N-Cinnamoylserine(benzoyl)cyclohexylamide (6i)

(8.2 mg, 0.0161 mmol, 26%). $[\alpha]_{\text{D}}^{28}$ -11° (c 0.23, CHCl_3). IR (film) ν_{max} cm^{-1} : 3583, 3282, 2924, 2854, 2360, 2332, 1728, 1645, 1616, 1541, 1458, 1348, 1176. ^1H NMR (500 MHz, CDCl_3) δ : 1.25 (6H, m), 1.71–1.74 (2H, m), 1.94 (2H, m), 3.80 (1H, m), 4.72 (1H, dd, $J = 11.5$, 7.0 Hz), 4.76 (1H, dd, $J = 11.5$, 5.5 Hz), 5.01 (1H, q, $J = 6.7$ Hz), 6.16 (1H, d, $J = 9.0$ Hz), 6.47 (1H, d, $J = 15.0$ Hz), 6.59 (1H, d, $J = 8.0$ Hz), 7.37–7.39 (3H, m), 7.50–7.52 (2H, m), 7.65 (1H, d, $J = 16.5$ Hz), 9.13 (2H, d, $J = 1.5$ Hz), 9.23 (1H, m). HR-MS (ESI) m/z : calcd for $\text{C}_{25}\text{H}_{26}\text{N}_4\text{O}_8\text{Na}$ $[\text{M}+\text{Na}]^+$ 533.1648. Found 533.1600.

4.3.10. (S)-N-Cinnamoylserine(nicotinoyl)cyclohexylamide (6j)

(13.2 mg, 0.0313 mmol, 47%) as a white powder, mp 201–202 °C. $[\alpha]_{\text{D}}^{26}$ -4.0° (c 0.45, CHCl_3). IR (film) ν_{max} cm^{-1} : 3400, 2931, 2854, 2360, 2332, 1733, 1647, 1618, 1541, 1292, 1211, 1144, 974. ^1H NMR (400 MHz, CDCl_3) δ : 1.13–1.35 (6H, m), 1.68–1.71 (2H, m), 1.87–1.90 (2H, m), 3.73–3.82 (1H, m), 4.64 (1H, dd, $J = 12.0$, 5.6 Hz), 4.73 (1H, dd, $J = 11.6$, 6.0 Hz), 4.98 (1H, q, $J = 6.4$ Hz), 6.30 (1H, d, $J = 8.4$ Hz), 6.48 (1H, d, $J = 16.0$ Hz), 6.71 (1H, d, $J = 7.6$ Hz), 7.35–7.38 (4H, m), 7.48–7.51 (2H, m), 7.64 (1H, d, $J = 15.6$ Hz), 8.28 (1H, dt, $J = 8.0$, 2.0 Hz), 8.78 (1H, dd, $J = 4.8$, 1.6 Hz), 9.20 (1H, d, $J = 1.2$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.9, 25.5, 33.0, 48.8, 52.7, 65.0, 119.9, 123.5, 125.7, 128.1, 129.0, 130.2, 134.6, 137.4, 142.5, 151.1, 153.9, 165.2, 166.2, 167.7. HR-MS (ESI) m/z : calcd for $\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 444.1899. Found 444.1861. ent-6j: mp 197–201 °C. $[\alpha]_{\text{D}}^{27}$ $+3.0^\circ$ (c 0.38, CHCl_3). HRMS (ESI). Found 444.1866.

4.3.11. (S)-N-Cinnamoylserine(picolinoyl)cyclohexylamide (6k)

(25.8 mg, 0.0612 mmol, 95%) as a white powder, mp 202–204 °C. $[\alpha]_{\text{D}}^{28}$ -6.0° (c 0.19, CHCl_3). IR (film) ν_{max} cm^{-1} : 3260, 2933, 2854, 2360, 2341, 1734, 1653, 1622, 1558, 1541, 1290, 1217, 1130, 1090, 978. ^1H NMR (500 MHz, CDCl_3) δ : 1.10–1.36 (6H, m), 1.56–1.71 (2H, m), 1.83–1.90 (2H, m), 3.74–3.81 (1H, m), 4.51 (1H, q, $J = 5.8$ Hz), 4.86 (1H, q, $J = 5.7$ Hz), 4.97 (1H, q, $J = 6.5$ Hz), 6.50 (1H, d, $J = 15.5$ Hz), 6.66 (1H, d, $J = 8.0$ Hz), 7.02 (1H, d, $J = 7.5$ Hz), 7.36–7.38 (3H, m), 7.49–7.52 (3H, m), 7.64 (1H, d, $J = 16.5$ Hz), 7.86 (1H, dd, $J = 7.5$, 2.0 Hz), 8.14 (1H, d, $J = 7.5$ Hz), 8.72 (1H, dd, $J = 4.0$, 2.0 Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.8, 25.6, 32.8, 32.9, 48.8, 52.4, 65.0, 120.1, 125.6, 127.5, 128.1, 129.0, 130.1, 134.7, 137.4, 142.1, 147.6, 149.8, 164.9, 166.2, 167.8. HR-MS (ESI) m/z : calcd for $\text{C}_{24}\text{H}_{27}\text{N}_3\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 444.1899. Found 444.1870. ent-6k: mp 205–208 °C. HRMS (ESI) Found 444.1866.

4.3.12. (S)-N-Cinnamoylserine((E)-3',4',5'-trimethoxy-3-phenylprop-2-enyl)cyclohexylamide (6l)

Oxalylchloride (27.3 μl , 0.316 mmol) and DMF (cat.) were added to (E)-3',4',5'-trimethoxy-3-phenylprop-2-enoic acid (37.6 mg, 0.158 mmol) at room temperature in CH_2Cl_2 under nitrogen atmosphere. The mixture was stirred at room temperature for 10 min, the solvent was evaporated in vacuo and the residue was added to the mixture of **5a** (50.0 mg, 0.158 mmol), TEA (132 ml, 0.738 mmol) in CH_2Cl_2 and stirred at room temperature for 2 h. After CH_2Cl_2 and H_2O were added to the solution, the organic layer was dried over MgSO_4 . The solvent was evaporated in vacuo and the residue was purified by silica-gel column chromatography (CHCl_3) to give **6l** (3.00 mg, 5.59 μmol , 3.9%) as a white powder, mp 227–229 °C. $[\alpha]_D^{27} -9.8^\circ$ (c 0.25, CHCl_3). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3283, 3073, 2931, 2841, 2370, 2337, 1720, 1640, 1621, 1582. ^1H NMR (400 MHz, CDCl_3) δ : 1.10–1.95 (11H, m), 3.87 (9H, s), 4.47–4.51 (1H, m), 4.60–4.67 (1H, m), 4.81–4.89 (1H, m), 6.21 (1H, d, $J = 9.6$ Hz), 6.34 (1H, d, $J = 16.0$ Hz), 6.47 (1H, d, $J = 16.0$ Hz), 6.68 (1H, d, $J = 1.6$ Hz), 6.73 (2H, s), 7.37 (3H, s), 7.50 (2H, s), 7.60–7.67 (2H, m). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 33.0, 48.6, 53.0, 56.2, 61.1, 64.2, 105.4, 116.4, 119.9, 128.0, 129.0, 129.7, 130.1, 134.5, 142.2, 146.0, 150.9, 153.5, 166.2, 167.0, 167.8. HR-MS (ESI) m/z : calcd for $\text{C}_{30}\text{H}_{37}\text{N}_2\text{O}_7$ $[\text{M}+\text{H}]^+$ 537.2601 Found 537.2612.

4.3.13. (S)-N-Cinnamoylserine(phenylpropanoyl)cyclohexylamide (6m)

(4.2 mg, 9.36 μmol , 13%) as an oil. $[\alpha]_D^{27} -19^\circ$ (c 0.16, CHCl_3). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3286, 2931, 2854, 2360, 1736, 1651, 1624, 1543, 1454, 1362, 1215, 1173, 1080, 976. ^1H NMR (500 MHz, CDCl_3) δ : 1.10–1.39 (6H, m), 1.68–1.71 (2H, m), 1.87–1.89 (2H, m), 2.67 (2H, t, $J = 7.8$ Hz), 2.95 (2H, t, $J = 7.8$ Hz), 3.75–3.76 (1H, m), 4.29 (1H, dd, $J = 12.0, 5.5$ Hz), 4.50 (1H, dd, $J = 11.5, 6.5$ Hz), 4.73 (1H, q, $J = 6.3$ Hz), 6.06 (1H, d, $J = 8.0$ Hz), 6.39 (1H, d, $J = 15.5$ Hz), 6.44 (1H, d, $J = 7.0$ Hz), 7.18–7.21 (2H, m), 7.27–7.30 (3H, m), 7.38–7.40 (3H, m), 7.50–7.52 (2H, m), 7.64 (1H, d, $J = 15.5$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.8, 25.6, 30.9, 33.0, 35.7, 48.7, 52.8, 64.1, 119.8, 126.5, 128.1, 128.4, 128.7, 129.0, 134.6, 140.4, 142.4, 166.1, 167.7. HR-MS (ESI) m/z : calcd for $\text{C}_{27}\text{H}_{32}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 471.2260. Found 471.2264.

4.3.14. (S)-N-Cinnamoylserine(cinnamoyl)cyclohexylamide (6n)

(4.00 mg, 0.00896 mmol, 26%) as a white powder, mp 192–198 °C. $[\alpha]_D^{27} -6.9^\circ$ (c 0.18, CHCl_3). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3279, 2929, 2856, 1716, 1643, 1617, 1541, 1447, 1311, 1172, 975, 769. ^1H NMR (500 MHz, CDCl_3) δ : 1.25–1.90 (10H, m), 3.80 (1H, m), 4.47 (1H, dd, $J = 11.5, 5.0$ Hz), 4.63 (1H, dd, $J = 12.0, 6.5$ Hz), 4.84 (1H, q, $J = 6.2$ Hz), 6.21 (1H, br), 6.44 (1H, d, $J = 16.0$ Hz), 6.47 (1H, d, $J = 15.0$ Hz), 6.69 (1H, br), 7.37–7.40 (6H, m), 7.51–7.52 (4H, m), 7.66 (1H, d, $J = 16.0$ Hz), 7.72 (1H, d, $J = 16.0$ Hz), 7.83 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (125 MHz, CDCl_3) δ : 24.8, 25.6, 29.9, 33.0, 48.7, 53.1, 64.2, 117.2, 119.9, 128.1, 128.4, 129.0, 129.1, 130.2, 130.8, 134.3, 134.7, 142.4, 146.3, 166.2, 167.8. HR-MS (ESI) m/z : calcd for $\text{C}_{27}\text{H}_{30}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 469.2104. Found 469.2078. ent-**6n**: mp 207–211 °C. $[\alpha]_D^{27} +9.0^\circ$ (c 0.14, CHCl_3). HRMS (ESI) Found 469.2064.

4.3.15. (S)-N-Cinnamoylserine(2-methyl-6-nitrobenzoyl)cyclohexylamide (6o)

(17.8 mg, 0.0371 mmol, 36%) as a white powder, mp 205–208 °C. $[\alpha]_D^{28} -11^\circ$ (c 0.23, CHCl_3). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3287, 2930, 2849, 2359, 1742, 1645, 1620, 1538, 1450, 1344, 1267, 1215, 1112, 1075, 975. ^1H NMR (400 MHz, CDCl_3) δ : 1.13–1.89 (10H, m), 2.42 (3H, s), 3.74 (1H, m), 4.73 (1H, dd, $J = 11.6$ Hz), 4.82 (1H, dd, $J = 11.2, 4.8$ Hz), 4.89–4.93 (1H, m), 6.32 (1H, m),

6.52 (1H, d, $J = 15.2$ Hz), 6.60 (1H, m), 7.37–7.39 (3H, m), 7.47–7.57 (4H, m), 7.68 (1H, d, $J = 16.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 19.4, 24.8, 25.5, 32.9, 48.7, 52.7, 65.4, 119.7, 121.9, 128.2, 129.0, 130.0, 130.3, 136.5, 142.6, 166.4, 167.6. HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{29}\text{N}_3\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 502.1954 Found 502.1971. ent-**6o**: mp 205–208 °C. $[\alpha]_D^{27} +12^\circ$ (c 0.15, CHCl_3). HRMS (ESI) Found 502.1926.

4.3.16. (S)-N-(E)-4-Phenylbut-2-enoicserine(benzoyl)cyclohexylamide (7c)

(11.1 mg, 25.5 μmol , 17%) as a white powder, mp 184–186 °C. $[\alpha]_D^{28} +10^\circ$ (c 0.10, CHCl_3). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3286, 3064, 3004, 2933, 2854, 2359, 1724, 1647, 1579, 1550. ^1H NMR (400 MHz, CDCl_3) δ : 1.02–1.89 (11H, m), 4.72–4.83 (1H, m), 3.92 (2H, s), 4.58–4.73 (2H, m), 4.88–4.96 (1H, m), 6.06 (1H, d, $J = 7.6$ Hz), 6.36 (1H, d, $J = 7.6$ Hz), 7.27–7.37 (1H, m), 7.38–7.43 (2H, m), 7.53–7.59 (1H, m), 7.93–8.01 (2H, m). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.4, 33.0, 48.7, 54.4, 55.0, 64.7, 128.6, 129.3, 129.8, 133.6, 166.4, 167.5, 167.9, 172.7. HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 457.2103. Found 457.2103.

4.3.17. (S)-N-(E)-4'-Methoxy-3-phenylprop-2-enoicserine(benzoyl)cyclohexylamide (7d)

(5.8 mg, 12.9 μmol , 22%) as a white powder, mp 185–187 °C. $[\alpha]_D^{23} +13^\circ$ (c 0.10, CHCl_3). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3282, 2925, 2852, 2360, 2333, 1724, 1647, 1577, 1545, 1481. ^1H NMR (400 MHz, CDCl_3) δ : 1.02–1.90 (11H, m), 3.93 (3H, s), 4.60–4.73 (2H, m), 4.90–4.96 (1H, m), 6.03 (1H, d, $J = 8.2$ Hz), 6.28–6.35 (2H, m), 6.88 (1H, d, $J = 8.7$ Hz), 7.42–7.45 (3H, m), 7.56–7.61 (2H, m), 7.96–8.02 (2H, m). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.4, 33.0, 48.7, 54.4, 55.0, 64.7, 128.6, 129.3, 133.6, 166.4, 167.4, 167.9, 172.4, 172.6. HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_5\text{Na}$ $[\text{M}+\text{Na}]^+$ 473.2052. Found 473.2099.

4.3.18. (S)-N-Benzoylserine(benzoyl)cyclohexylamide (7e)

(11.5 mg, 0.0292 mmol, 43%) as a white powder, mp 202–203 °C. $[\alpha]_D^{25} +1.0^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3292, 3072, 2924, 2852, 2359, 1734, 1649, 1635, 1558, 1541. ^1H NMR (400 MHz, CDCl_3) δ : 1.03–1.91 (11H, m), 3.70–4.81 (1H, m), 4.61–4.68 (1H, m), 4.73–4.81 (1H, m), 5.02 (1H, s), 6.50 (1H, s), 7.33–7.59 (6H, m), 7.81 (2H, d, $J = 7.3$ Hz), 8.01 (2H, d, $J = 6.9$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 32.9, 48.7, 53.2, 64.6, 127.3, 128.6, 128.8, 129.2, 129.9, 132.2, 133.5, 166.8, 167.6, 167.7. HR-MS (ESI) m/z : calcd for $\text{C}_{23}\text{H}_{25}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 417.1790. Found 417.1794.

4.3.19. (S)-N-(E)-2'-Methoxy-3-phenylprop-2-enoicserine(benzoyl)cyclohexylamide (7f)

(16.6 mg, 36.9 μmol , 57%) (*E/Z* = 5:2) as a white powder, mp 201–204 °C. $[\alpha]_D^{25} +11^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{cm}^{-1}$: 3282, 3072, 2931, 2862, 1726, 1643, 1610, 1542, 1491, 1462. ^1H NMR (400 MHz, CDCl_3) δ : 1.04–1.94 (11H, m), 3.78 (1H, s), 3.87 (1H, s), 4.30 (0.33H, dd, $J = 11.2, 5.2$ Hz), 4.49 (0.33H, dd, $J = 12.0, 5.6$ Hz), 4.59 (0.67H, dd, $J = 11.2, 5.2$ Hz), 4.70 (0.67H, dd, $J = 12.0, 6.4$ Hz), 4.75–4.81 (0.17H, d, $J = 16.0$ Hz), 4.90–4.98 (0.83H, m), 6.03 (0.29H, d, $J = 12.4$ Hz), 6.60 (0.71H, d, $J = 16.0$ Hz), 6.76 (0.23H, d, $J = 8.8$ Hz), 6.81 (0.23H, t, $J = 7.2$ Hz), 6.81 (0.23H, t, $J = 7.2$ Hz), 6.88–6.98 (0.42H, m), 7.02 (0.15H, d, $J = 12.4$ Hz), 7.15–7.20 (0.21H, m), 7.32 (0.77H, t, $J = 8.8$ Hz), 7.39–7.48 (2.58 H, m), 7.56 (0.79H, t, $J = 7.2$ Hz), 7.86–7.92 (1.79H, m), 8.02 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 32.9, 48.7, 52.8, 55.9, 56.0, 64.6, 109.7, 110.6, 111.1, 117.8, 121.0, 122.3, 123.2, 127.5, 128.5, 129.6, 129.7, 129.8, 133.4, 139.1, 142.0, 149.2, 150.9, 166.4, 166.6, 167.9. HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_5\text{Na}$ $[\text{M}+\text{Na}]^+$ 473.2052. Found 473.2051.

4.3.20. (S)-N-(E)-3',4'-Dimethoxy-3-phenylprop-2-enoicserine (benzoyl)cyclohexylamide (7g)

(E:Z = 3:1) (9.3 mg, 19.4 μmol , 37%) as a white powder, mp 199–202 °C. $[\alpha]_D^{25} +18^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3282, 3080, 3022, 2931, 2846, 1728, 1643, 1614, 1516, 1452. ^1H NMR (400 MHz, CDCl_3) δ : 1.09–1.94 (11H, m), 3.66–3.81 (1H, m), 3.89 (6H, s), 4.46 (0.25H, dd, $J = 12.0, 5.2$ Hz), 4.56–4.63 (1H, m), 4.71 (0.75H, dd, $J = 11.2, 6.0$ Hz), 4.81–4.87 (0.25H, m), 4.93 (0.75H, dd, $J = 12.8, 6.0$ Hz), 5.87 (0.25H, d, $J = 8.0$ Hz), 6.07 (0.25H, d, $J = 9.6$ Hz), 6.26 (0.75H, d, $J = 8.0$ Hz), 6.34 (0.75H, d, $J = 16.0$ Hz), 6.53–6.59 (0.25H, m), 6.65 (0.75H, d, $J = 8.8$ Hz), 6.70–6.77 (0.5H, m), 6.84 (0.75H, d, $J = 8.0$ Hz), 7.02 (0.75H, s), 7.07 (1H, d, $J = 8.0$ Hz), 7.30 (0.25H, s), 7.43 (2H, t, $J = 8.0$ Hz), 7.53–7.61 (1.5H, m), 7.93 (0.5H, d, $J = 8.0$ Hz), 8.02 (1.5H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 32.9, 48.7, 52.8, 55.9, 56.0, 64.6, 109.7, 111.1, 121.1, 122.3, 123.2, 127.6, 128.5, 129.6, 129.7, 129.8, 133.4, 142.0, 149.2, 150.9, 166.4, 166.6, 167.9. HR-MS (ESI) m/z : calcd for $\text{C}_{27}\text{H}_{32}\text{N}_2\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 503.2158. Found 503.2150.

4.3.21. (S)-N-(E)-3',5'-Dimethoxy-3-phenylprop-2-enoicserine (benzoyl)cyclohexylamide (7h)

(12.9 mg, 0.0194 mmol, 19%) as a white powder, mp 175–177 °C. $[\alpha]_D^{23} +3.0^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3396, 3276, 2924, 2846, 2360, 2333, 1726, 1645, 1614, 1558; ^1H NMR (400 MHz, CDCl_3) δ : 1.08–1.91 (11H, m), 3.96 (6H, s), 4.59 (1H, dd, $J = 11.3, 6.4$ Hz), 4.69 (1H, dd, $J = 11.3, 6.0$ Hz), 4.95–4.99 (1H, m), 6.43–6.47 (2H, m), 6.62 (2H, s), 6.80 (1H, d, $J = 7.7$ Hz), 7.42 (2H, t, $J = 8.0$ Hz), 7.52–7.57 (2H, m), 8.01 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 33.0, 48.7, 52.8, 55.5, 64.6, 102.3, 105.9, 120.5, 128.6, 129.5, 129.9, 133.5, 136.4, 142.2, 161.0, 166.0, 166.6, 167.7. HR-MS (ESI) m/z : calcd for $\text{C}_{27}\text{H}_{32}\text{N}_2\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 503.2158. Found 503.2170. ent-**7h**: mp 174–175 °C. $[\alpha]_D^{28} -2.4^\circ$ (c 0.25, MeOH). HR-MS (ESI) m/z : calcd for $\text{C}_{27}\text{H}_{33}\text{N}_2\text{O}_6$ $[\text{M}+\text{H}]^+$ 481.2338. Found 481.2361.

4.3.22. (S)-N-(E)-3'-Methoxy-3-phenylprop-2-enoicserine (benzoyl)cyclohexylamide (7i)

(16.6 mg, 0.0369 mmol, 57%) as a white powder, mp 192–194 °C. $[\alpha]_D^{27} +1.8^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3438, 3282, 2933, 2854, 2360, 2326, 1722, 1647, 1558, 1541. ^1H NMR (400 MHz, CDCl_3) δ : 1.06–1.96 (1H, m), 3.82 (3H, s), 4.56–4.62 (1H, m), 4.70 (1H, dd, $J = 12.0, 6.0$ Hz), 4.88–4.94 (1H, m), 6.17 (1H, s), 6.45 (1H, d, $J = 16.0$ Hz), 6.68 (1H, d, $J = 7.2$ Hz), 6.89–6.93 (1H, m), 7.00–7.02 (1H, m), 7.09 (1H, d, $J = 8.0$ Hz), 7.43 (2H, t, $J = 8.0$ Hz), 7.54–7.64 (2H, m), 8.02 (2H, d, $J = 7.2$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 15.4, 24.8, 25.5, 32.9, 48.7, 52.8, 55.3, 64.6, 65.9, 113.0, 115.9, 120.4, 120.6, 128.5, 129.6, 129.8, 129.9, 133.4, 136.0, 142.0, 160.0, 166.1, 166.5, 167.9; HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_5\text{Na}$ $[\text{M}+\text{Na}]^+$ 473.2052. Found 473.2046. ent-**7i**: mp 194–196 °C. $[\alpha]_D^{28} -2.4^\circ$ (c 0.25, MeOH). HR-MS (ESI) m/z : found 473.2060.

4.3.23. (S)-N-(E)-4'-Benzoyloxy-3-phenylprop-2-enoicserine (benzoyl)cyclohexylamine (7j)

(17.0 mg, 0.0314 mmol, 52%) as a white powder, mp 220–222 °C. $[\alpha]_D^{28} -1.0^\circ$ (c 0.40, CHCl_3). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3284, 2929, 2854, 2355, 2326, 1734, 1645, 1616, 1556, 1541. ^1H NMR (400 MHz, CDCl_3) δ : 1.08–1.93 (11H, m), 3.72–3.73 (1H, m), 3.93 (1H, s), 4.57–4.62 (1H, m), 4.69–4.73 (1H, m), 4.90–4.96 (1H, m), 6.21 (1H, d, $J = 8.8$ Hz), 6.45 (1H, d, $J = 15.6$ Hz), 6.73 (1H, d, $J = 7.3$ Hz), 7.43 (2H, t, $J = 8.0$ Hz), 7.49–5.59 (6H, m), 7.63–7.67 (2H, m), 8.02 (2H, d, $J = 8.0$ Hz), 8.19 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 32.9, 48.7, 52.8, 64.8, 120.1, 122.4, 128.6, 128.7, 129.2, 129.9, 130.3, 132.5, 133.5, 133.9, 141.2, 152.2, 165.0, 166.0, 166.6, 167.8, 176.2. HR-MS (ESI) m/z :

calcd for $\text{C}_{32}\text{H}_{32}\text{N}_2\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 563.2158. Found 563.2171. ent-**7j**: mp 238–240 °C. $[\alpha]_D^{28} +3.0^\circ$ (c 0.13, CHCl_3). HR-MS (ESI) m/z : calcd for $\text{C}_{32}\text{H}_{33}\text{N}_2\text{O}_6$ $[\text{M}+\text{H}]^+$ 541.2338. Found 541.2307.

4.3.24. (S)-N-(E)-3',4',5'-Trimethoxy-3-phenylprop-2-enoicserine (benzoyl)cyclohexylamide (7k)

(22.0 mg, 0.0431 mmol, 81%) as a white powder, mp 222–223 °C. $[\alpha]_D^{26} +13^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3583, 3290, 2931, 2854, 2355, 1722, 1651, 1583, 1506, 1271. ^1H NMR (400 MHz, CDCl_3) δ : 1.05–1.92 (11H, m), 3.84 (9H, s), 4.60 (1H, dd, $J = 11.2, 5.2$ Hz), 4.69 (1H, dd, $J = 11.2, 6.0$ Hz), 4.99–5.01 (1H, m), 6.41 (1H, d, $J = 16.0$ Hz), 6.65 (1H, d, $J = 8.0$ Hz), 6.69 (2H, s), 6.92 (1H, d, $J = 8.0$ Hz), 7.40 (2H, t, $J = 8.0$ Hz), 7.48–7.57 (2H, m), 8.01 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 24.8, 25.5, 32.9, 48.7, 52.7, 56.2, 61.0, 64.7, 105.1, 119.4, 128.5, 129.8, 130.1, 133.4, 142.1, 153.5, 166.0, 166.5, 167.8. HR-MS (ESI) m/z : calcd for $\text{C}_{28}\text{H}_{34}\text{N}_2\text{O}_7\text{Na}$ $[\text{M}+\text{Na}]^+$ 533.2264. Found 533.2269. ent-**7k**: mp 211–213 °C. $[\alpha]_D^{27} -12^\circ$ (c 0.15, MeOH). HR-MS (ESI) m/z : found 533.2274.

4.3.25. (S)-N-Cinnamoylserine(benzoyl)piperidineamide (7l)

(19.5 mg, 49.9 μmol , 75%) as an oil. $[\alpha]_D^{26} -2.5^\circ$ (c 0.40, MeOH). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3288, 3060, 3025, 3006, 2941, 2857, 2360, 2331, 1722, 1616. ^1H NMR (400 MHz, CDCl_3) δ : 1.52–1.74 (6H, m), 3.47–3.71 (4H, m), 4.46 (1H, dd, $J = 12.0, 5.6$ Hz), 4.58 (1H, dd, $J = 12.0, 4.0$ Hz), 5.46–5.52 (1H, m), 6.48 (1H, d, $J = 16.0$ Hz), 6.92 (1H, d, $J = 8.0$ Hz), 7.32–7.57 (7H, m), 7.60 (1H, d, $J = 16.0$ Hz), 8.00 (2H, d, $J = 6.8$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 14.3, 24.5, 25.6, 26.5, 43.7, 46.9, 48.7, 60.5, 64.7, 120.4, 128.0, 128.5, 128.9, 129.7, 129.9, 133.3, 141.7, 165.4, 166.8. HR-MS (ESI) m/z : calcd for $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 429.1793. Found 429.1782. ent-**7l**: $[\alpha]_D^{26} +1.6^\circ$ (c 0.15, MeOH). HR-MS (ESI) m/z : found 429.1770.

4.3.26. (S)-N-Cinnamoylserine(benzoyl)morpholineamide (7m)

(13.0 mg, 0.0318 mmol, 31%) as an oil. $[\alpha]_D^{25} +43^\circ$ (c 0.11, CHCl_3). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3649, 3583, 2362, 1718, 1558, 1541, 1508, 1473, 1419, 1313. ^1H NMR (400 MHz, CDCl_3) δ : 3.65–3.78 (8H, m), 4.49 (1H, dd, $J = 12.0, 6.4$ Hz), 4.58 (1H, dd, $J = 11.6, 4.4$ Hz), 5.44–5.52 (1H, m), 6.48 (1H, d, $J = 16.0$ Hz), 6.81 (1H, d, $J = 8.0$ Hz), 7.32–7.58 (5H, m), 7.61 (1H, d, $J = 16.0$ Hz), 8.00 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 42.8, 46.3, 48.5, 64.5, 66.7, 120.1, 128.0, 128.6, 128.9, 129.9, 133.4, 134.7, 142.0, 165.5, 167.5. HR-MS (ESI) m/z : calcd for $\text{C}_{23}\text{H}_{24}\text{N}_2\text{O}_5\text{Na}$ $[\text{M}+\text{Na}]^+$ 431.1583. Found 431.1557. ent-**7m**: $[\alpha]_D^{28} -31^\circ$ (c 0.20, CHCl_3). HR-MS (ESI) m/z : found 431.1569.

4.3.27. (S)-N-Cinnamoylserine(benzoyl)benzylamide (7n)

(11.4 mg, 0.0292 mmol, 44%) as a white powder, mp 207–208 °C. $[\alpha]_D^{27} +26^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3585, 3286, 2360, 1728, 1647, 1624, 1540, 1450, 1394, 1338. ^1H NMR (400 MHz, CDCl_3) δ : 4.47 (2H, d, $J = 6.0$ Hz), 4.64 (1H, dd, $J = 11.6, 6.0$ Hz), 6.45 (1H, d, $J = 16.0$ Hz), 6.74 (2H, d, $J = 6.8$ Hz), 7.22 (4H, s), 7.34–7.44 (5H, m), 7.46–7.50 (2H, m), 7.62 (1H, d, $J = 16.0$ Hz), 7.97 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 43.8, 53.0, 64.5, 119.7, 127.7, 128.1, 128.6, 128.8, 128.9, 129.9, 130.1, 133.5, 142.5, 166.3, 168.8. HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{24}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 451.1634. Found 451.1603. ent-**7n**: mp 207–209 °C. $[\alpha]_D^{23} -25^\circ$ (c 0.11, MeOH). HR-MS (ESI) m/z : found 451.1594.

4.3.28. (S)-N-Cinnamoylserine(benzoyl)cyclohexanemethylamide (7o)

(15.0 mg, 0.0345 mmol, 57%) as a white powder, mp 208 °C. $[\alpha]_D^{27} +27^\circ$ (c 0.10, MeOH). IR (film) $\nu_{\text{max}} \text{ cm}^{-1}$: 3583, 3290, 2918, 2852, 2360, 2341, 1730, 1647, 1622, 1545. ^1H NMR (400 MHz, CDCl_3) δ : 0.81–0.94 (2H, m), 1.03–1.18 (3H, m), 1.36–1.48 (1H, m),

1.55–1.69 (6H, m), 3.05–3.21 (2H, m), 4.62 (1H, dd, $J = 12.0, 4.8$ Hz), 4.73 (1H, dd, $J = 11.2, 6.0$ Hz), 4.97–5.04 (1H, m), 6.48 (1H, d, $J = 16.0$ Hz), 7.33–7.38 (3H, m), 7.42 (2H, t, $J = 8.0$ Hz), 7.46–7.52 (2H, m), 7.55 (1H, t, $J = 8.0$ Hz), 7.63 (1H, d, $J = 16.0$ Hz), 8.01 (2H, d, $J = 8.0$ Hz). ^{13}C NMR (100 MHz, CDCl_3) δ : 25.8, 26.3, 30.8, 37.9, 46.0, 53.0, 64.6, 119.8, 128.1, 128.6, 129.0, 130.1, 133.5, 134.5, 142.4, 166.2, 166.8, 168.8. HR-MS (ESI) m/z : calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 457.2103. Found 457.2153. ent-**7o**: mp 202–203 °C. $[\alpha]_D^{27}$ –27° (c 0.15, MeOH). HR-MS (ESI) m/z : found 457.2052.

4.4. Measurement of inhibitory activity against SARS 3CL R188I mutant protease

Enzyme assays were carried out using recombinant SARS 3CL R188I mutant protease at an enzyme concentration of 120 nM. The reaction mixture (40 mM AcONa buffer, pH 4.0, containing 10% glycerol, 10 mM DTT and 4 M NaCl) was analyzed on a Cosmosil 5C18-AR-II column (4.6 × 150 mm), employing a linear gradient of MeCN (10–40%, 30 min) in aq 0.1% TFA. Each IC_{50} value was obtained from a sigmoidal dose–response curve obtained from the decrease of the substrate in the reaction mixture. Each experiment was repeated three times.

4.5. Measurement of inhibitory activity against cathepsin B

Bovine spleen cathepsin B was purchased from Sigma Chemical company (St. Louis, MO). 10 mU of the cathepsin B was preincubated at 40 °C for 10 min in 90 μl of 50 mM MES (pH 6.0), 2 mM DTT, and 0.1% Brij-35. The solution was then mixed with 20 mM Z-Arg-Arg-MCA and each inhibitor (1 nM to 1 mM) in DMSO (10 μl), and the mixture was incubated at 40 °C for 10 min. The reaction was stopped by adding 100 μl of 100 mM sodium monochloroacetate, 30 mM sodium acetate, and 70 mM acetic acid (pH 4.3). Each IC_{50} value was obtained from a sigmoidal dose–response curve obtained from the decrease of the substrate in the reaction mixture. Each experiment was repeated three times.

4.6. Evaluation of the cytotoxicity by MTT assay

HeLa cells were maintained in a suspension culture of DMEM supplemented with 5% FBS (Fetal Bovine Serum) containing 1% of a penicillin–streptomycin mixture. A 100 μl aliquot of HeLa cells (5000 cells/ml) was added to a 96 well plate and incubated for 24 h at 37 °C in a humidified incubator containing 5% CO_2 in air. After 24 h, a 10 μl aliquot of compound **serine derivatives** (concentrations varying in the range of 5, 4, 3, 2, 1, 0.1 μM) and rotenone as a control was added to each of the 96 wells and incubated for 24 h. A 10 μl WST-8 solution (mixture of WST-8 and 1-Methoxy PMS) was added to each well and the incubation continued for 2 h. The visible absorbance at 450 nm and 630 nm as the reference wavelength of each well was quantified using a microplate reader.

Acknowledgement

This work was supported in part by Adaptable and Seamless Technology transfer Program (A-STEP) through target driven R&D, Japan Science and Technology Agency.

Supplementary data

Supplementary data (additional experimental procedures, analytical data for library compounds) associated with this article

can be found, in the online version, at <http://dx.doi.org/10.1016/j.bmc.2016.01.052>.

References and notes

- Lee, N.; Hui, D.; Wu, A.; Chan, P.; Cameron, P.; Joynt, G. M.; Ahuja, A.; Yung, M. Y.; Leung, C. B.; To, K. F. *N. Engl. J. Med.* **2003**, *348*, 1986.
- Drosten, C.; Gunther, S.; Preiser, W.; Ven der Werf, S.; Brodt, H. R.; Becker, S.; Rabenau, H.; Panning, M.; Kolensnikova, L.; Fouchier, R. A. M.; Berger, A.; Burguiere, A. M.; Cinatl, J.; Eickmann, M.; Escriou, N.; Grywna, K.; Kramme, S.; Manuguerra, J.; Muller, S.; Rickerts, V.; Sturmer, M.; Vieth, S.; Klenk, H. D.; Osterhaus, A. D. E. E.; Schemitz, H.; Doerr, H. W. *N. Engl. J. Med.* **2003**, *348*, 1967.
- Ksiazek, T. G.; Erdman, D.; Goldsmith, C. S.; Zaki, S. R.; Peret, T.; Emery, S.; Tong, S.; Urbani, C.; Comer, J. A.; Lim, W.; Rollin, R. E.; Dowell, S. F.; Ling, A. E.; Humphrey, C. D.; Shieh, G. J.; Guarner, J. Y.; Paddock, C. D.; Rota, P.; Fields, B.; DeRisi, L. J.; Yang, J. Y.; Cox, N.; Hughes, J. M.; LeDuc, J. W.; Bellinim, W. J.; Anderson, J. W. *N. Engl. J. Med.* **2003**, *348*, 1953.
- Pyrk, K.; Berkhout, B.; Van der Hoek, L. *J. Virol.* **2007**, *81*, 3051.
- MacBride, R.; Fielding, B. C. *Future Microbiol.* **2011**, *6*, 153.
- Cui, L.-J.; Zhang, C.; Zhang, T.; Lu, R.-J.; Xie, Z.-D.; Zhang, L.-L.; Liu, C.-Y.; Zhou, W.-M.; Ma, X.-J.; Tan, W.-J. *Adv. Virol.* **2011**, No. 129134.
- Zaki, A. M.; van Boheemen, S.; Bestebroer, T. M.; Osterhaus, A. D.; Fouchier, R. A. *N. Engl. J. Med.* **2014**, *2012*, 367.
- Birmingham, A.; Brown, C. S.; Aarons, E.; Tong, C.; Langrish, C.; Hoschler, K.; Brown, K.; Galiano, M.; Myers, R.; Pebody, R. G.; Green, H. K.; Boddington, N. L.; Gopal, R.; Newsholme, W.; Drosten, C.; Fouchier, R. A.; Zambon, M. *Euro. Surveill.* **2012**, *17*, 20290.
- Van Boheemen, S.; Graaf, M.; Lauber, C.; Bestebroer, T. M.; Raj, V. S.; Zaki, A. M.; Osterhaus, A. D. M.; Haagmans, B. L.; Gorbalenya, A. E.; Snijder, E. J.; Fouchier, R. A. M. *MBio* **2012**, *3*, e00473–12.
- Thiel, V.; Ivanov, K. A.; Putics, A.; Hertzog, T.; Schelle, B.; Bayer, S.; Weißbrich, B.; Snijder, E. J.; Rabenau, H.; Doerr, H. W.; Gorbalenya, A. E.; Ziebuhr, J. J. *J. Gen. Virol.* **2003**, *84*, 2305.
- Thanigaimaalaai, P.; Konno, S.; Yamamoto, T.; Koiwai, Y.; Taguchi, A.; Takayama, K.; Yakushiji, F.; Akaji, K.; Kiso, Y.; Kawasaki, Y.; Chen, S.-E.; Naser-Tavakolian, A.; Scheön, A.; Freire, E.; Hayashi, Y. *Eur. J. Med. Chem.* **2013**, *65*, 436.
- Nguyen, T. T. H.; Woo, H.-J.; Kang, H.-K.; Nguyen, V. D.; Kim, Y.-M.; Kim, D.-W.; Ahn, S.-A.; Xia, Y.; Kim, D. *Biotechnol. Lett.* **2012**, *34*, 831.
- Park, J.-Y.; Kim, J. H.; Kwon, J. M.; Kwon, H.-J.; Jeong, H. J.; Kim, Y. M.; Kim, D.; Lee, W. S.; Ryu, Y. B. *Bioorg. Med. Chem.* **2013**, *21*, 3730.
- Jacobs, J.; Grum-Tokars, V.; Zhou, Y.; Turlington, M.; Saldanha, S. A.; Chase, P.; Eggler, A.; Dawson, E. C.; Baez-Santos, Y. M.; Tomar, S.; Mielech, A. M.; Baker, S. C.; Lindsley, C. W.; Hodder, P.; Mesecar, A.; Stauffer, S. R. *J. Med. Chem.* **2013**, *56*, 534.
- Liu, W.; Zhu, H.-M.; Niu, G.-J.; Shi, E.-Z.; Chen, J.; Sun, B.; Chen, W.-Q.; Zhou, H.-G.; Yang, C. *Bioorg. Med. Chem.* **2014**, *22*, 292.
- Shimamoto, Y.; Hattori, Y.; Kobayashi, K.; Teruya, K.; Sanjoh, A.; Nakagawa, A.; Yamashita, E.; Akaji, K. *Bioorg. Med. Chem.* **2015**, *23*, 876.
- Akaji, K.; Konno, H.; Onozuka, M.; Makino, A.; Nosaka, K. *Bioorg. Med. Chem.* **2008**, *16*, 9400.
- Akaji, K.; Konno, H.; Mitsui, H.; Teruya, K.; Shimamoto, Y.; Hattori, Y.; Ozaki, T.; Kusunoki, M.; Sanjoh, A. *J. Med. Chem.* **2011**, *54*, 7962.
- Konno, H.; Sema, Y.; Ishii, M.; Hattori, Y.; Nosaka, K.; Akaji, K. *Tetrahedron Lett.* **2013**, *54*, 4848.
- Konno, H.; Sema, Y.; Tokairin, Y. *Tetrahedron* **2015**, *71*, 3433.
- Bai, D.; Yang, Q.; Chem, L.; He, X.; Gao, Z.; Shen, X. *Chem. Pharm. Bull.* **2008**, *56*, 1400.
- Carpino, L. A. *J. Am. Chem. Soc.* **1993**, *115*, 4397.
- Carpino, L. A.; Imazumi, H.; El-Faham, A.; Ferrer, F. J.; Zhang, C.; Lee, Y.; Foxman, B. M.; Henklein, P.; Hanay, C.; Mugg, C.; Wenschuh, H.; Klose, J.; Beyerman, M.; Bienert, M. *Angew. Chem., Int. Ed.* **2002**, *41*, 442.
- Kunishima, M.; Wamachi, C.; Hioki, K.; Terao, K.; Tani, S. *Tetrahedron* **2001**, *57*, 1551.
- Shiina, I. *Bull. Chem. Soc. Jpn.* **2014**, *87*, 196.
- Inanaga, J.; Hirata, K.; Saeki, H.; Katsuki, Y.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 1989.
- Jacobs, J.; Grum-Tokars, V.; Zhou, Y.; Turlington, M.; Saldanha, S. A.; Chase, P.; Eggler, A.; Dawson, E. S.; Baez-Santos, Y. M.; Tomar, S.; Mielech, A. M.; Baker, S. C.; Lindsley, C. W.; Hodder, P.; Mesecar, A.; Stauffer, S. R. *J. Med. Chem.* **2013**, *56*, 534.
- Konno, H.; Nosaka, K.; Akaji, K. *Tetrahedron* **2011**, *67*, 9067.
- Konno, K.; Kubo, K.; Makabe, H.; Toshiro, E.; Nosaka, K.; Akaji, K. *Tetrahedron* **2007**, *63*, 9502.
- Hiwasa, T.; Fujita-Yoshigaki, J.; Shirouzu, M.; Koide, H.; Sawada, T.; Sakiyama, S.; Yokoyama, S. *Cancer Lett.* **1993**, *69*, 161.
- Barrett, A. J.; Kirschke, H. *Method Enzymol.* **1981**, *380*, 535.
- Lim, I. T.; Meroueh, S. O.; Lee, M.; Heeg, M. J.; Mobashery, S. *J. Am. Chem. Soc.* **2004**, *126*, 10271.