



IgGs from patients with amyotrophic lateral sclerosis and diabetes target $\text{Ca}_v\alpha_2\delta_1$ subunits impairing islet cell function and survival

Yue Shi^{a,1,2}, Kyoung Sun Park^{b,1}, Seung Hyun Kim^c, Jia Yu^a, Kaixuan Zhao^a, Lina Yu^a, Ki Wook Oh^c, Kayoung Lee^b, Jaeyoon Kim^b, Kanchan Chaggar^d, Yuxin Li^e, Annette C. Dolphin^d, William A. Catterall^{f,2}, Sung Ho Ryu^b, Shao-Nian Yang^{a,e,2,3}, and Per-Olof Berggren^{a,b,g,h,2,3}

^aThe Rolf Luft Research Center for Diabetes and Endocrinology, Karolinska Institutet, Karolinska University Hospital L1, SE-171 76 Stockholm, Sweden; ^bDivision of Integrative Biosciences and Biotechnology, Pohang University of Science and Technology, Pohang 37673, Korea; ^cDepartment of Neurology, Hanyang University Hospital, Seoul 04763, Korea; ^dDepartment of Neuroscience, Physiology and Pharmacology, University College London, London WC1E 6BT, United Kingdom; ^eNational Engineering Laboratory for Druggable Gene and Protein Screening, Northeast Normal University, Changchun 130024, China; ^fDepartment of Pharmacology, School of Medicine, University of Washington, Seattle, WA 98195-7280; ^gLee Kong Chian School of Medicine, Nanyang Technological University, Singapore 637553; and ^hDiabetes Research Institute, Miller School of Medicine, University of Miami, Miami, FL 33136

Contributed by William A. Catterall, November 6, 2019 (sent for review July 12, 2019; reviewed by Veit Flockerzi, Jeffrey M. Friedman, and Solomon H. Snyder)

Patients with amyotrophic lateral sclerosis (ALS) often show hallmarks of type 2 diabetes mellitus (T2DM). However, the causal link between ALS and T2DM has remained a mystery. We now demonstrate that 60% of ALS patients with T2DM (ALS-T2DM) have sera that exaggerated K^+ -induced increases in cytosolic free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) in mouse islet cells. The effect was attributed to the presence of pathogenic immunoglobulin Gs (IgGs) in ALS-T2DM sera. The pathogenic IgGs immunocaptured the voltage-dependent Ca^{2+} (Ca_v) channel subunit $\text{Ca}_v\alpha_2\delta_1$ in the plasma membrane enhancing Ca_v1 channel-mediated Ca^{2+} influx and $[\text{Ca}^{2+}]_i$, resulting in impaired mitochondrial function. Consequently, impairments in $[\text{Ca}^{2+}]_i$ dynamics, insulin secretion, and cell viability occurred. These data reveal that patients with ALS-T2DM carry cytotoxic ALS-T2DM-IgG autoantibodies that serve as a causal link between ALS and T2DM by immunoattacking $\text{Ca}_v\alpha_2\delta_1$ subunits. Our findings may lay the foundation for a pharmacological treatment strategy for patients suffering from a combination of these diseases.

amyotrophic lateral sclerosis | calcium channel | cytosolic free Ca^{2+} concentration | diabetes | immunoglobulin

Patients with amyotrophic lateral sclerosis (ALS) show progressive dysfunction and degeneration of motor neurons in the brainstem and spinal cord (1). There are no effective therapeutics available for ALS (1). Clinically, ALS patients manifest advanced muscular weakness and paralysis and die within 2 to 5 y from the onset of the disease (1). Although the exact pathogenic mechanisms of ALS are not clarified, dysregulation of voltage-dependent Ca^{2+} (Ca_v) channels, cytosolic free Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$), and synaptic plasticity induced by altered humoral immunity has been proposed to participate in the development of the disease (2, 3). Exposure to purified immunoglobulin G (IgG) from the serum of ALS patients produced ultrastructural abnormalities with Ca^{2+} accumulation in and increased transmitter release from rodent motor neurons (4). It has long been recognized that intracellular accumulation of Ca^{2+} is cytotoxic and causes mitochondrial dysfunction, free radical damage, and Ca^{2+} -dependent cell death (5–7).

A series of neurodegenerative diseases, like ALS, is known to be associated with hallmarks of type 2 diabetes mellitus (T2DM), such as impaired glucose homeostasis, but their causal links are not known (8–10). Blood glucose homeostasis is under strict control of hormone release from pancreatic islet cells. T2DM and its characteristic symptom hyperglycemia occur in people with inadequate islet cell mass and function. Islet cells and neurons share a series of physiological and pathological mechanisms, such as Ca^{2+} -dependent exocytosis and Ca^{2+} -triggered cell death, for their function/dysfunction and survival/death (11–14). Therefore,

the present study hypothesized that IgG from ALS patients with T2DM (ALS-T2DM) may recognize similar targets in islet cells, as revealed in motor neurons, and thereby impair islet cell function and viability by disturbing Ca^{2+} signaling. Indeed, the present study reveals that a subgroup of ALS-T2DM patients has sera that enhance K^+ -induced $[\text{Ca}^{2+}]_i$ responses in islet cells via cytotoxic IgGs. Moreover, it demonstrates that ALS-T2DM-IgGs immunocapture $\text{Ca}_v\alpha_2\delta_1$ subunits and thereby enhance Ca_v1 channel-mediated Ca^{2+} influx, resulting in altered $[\text{Ca}^{2+}]_i$ dynamics and, consequently, impaired mitochondrial function, insulin secretion, and cell viability.

Results

A Subgroup of ALS-T2DM Patients Has Sera That Exaggerate K^+ -Induced $[\text{Ca}^{2+}]_i$ Responses in Mouse Islet Cells. ALS patient sera accommodate altered humoral immunity that results in pathological exaggeration of voltage-dependent Ca^{2+} entry and $[\text{Ca}^{2+}]_i$

Significance

We provide evidence of a mechanistic link between ALS and T2DM. Our data show that a subgroup of ALS-T2DM patients have sera that enhance Ca_v1 channel-mediated Ca^{2+} influx and exaggerate $[\text{Ca}^{2+}]_i$. These effects occur because the sera accommodate cytotoxic IgG autoantibodies that immunocapture $\text{Ca}_v\alpha_2\delta_1$ subunits. As a consequence, impairments in $[\text{Ca}^{2+}]_i$ dynamics, mitochondrial function, insulin secretion, and cell viability appear. We could clarify not only the identity of this serum factor but also the molecular mechanisms underlying its effects on the islet cells. Our findings may lay the foundation for a treatment strategy for this complex and severe group of diabetic patients.

Author contributions: Y.S., K.S.P., W.A.C., S.-N.Y., and P.-O.B. designed research; Y.S., K.S.P., J.Y., K.Z., L.Y., K.L., J.K., and Y.L. performed research; S.H.K., K.W.O., K.C., A.C.D., W.A.C., and S.H.R. contributed new reagents/analytic tools; Y.S., K.S.P., J.Y., K.Z., L.Y., K.L., J.K., and Y.L. analyzed data; and Y.S., K.S.P., W.A.C., S.-N.Y., and P.-O.B. wrote the paper.

Reviewers: V.F., Saarland University; J.M.F., The Rockefeller University; and S.H.S., Johns Hopkins University School of Medicine.

The authors declare no competing interest.

This open access article is distributed under [Creative Commons Attribution License 4.0 \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).

¹Y.S. and K.S.P. contributed equally to this work.

²To whom correspondence may be addressed. Email: yue.shi@ki.se, wcatt@uw.edu, shao-nian.yang@ki.se, or per-olof.berggren@ki.se.

³S.-N.Y. and P.-O.B. contributed equally to this work.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1911956116/-DCSupplemental>.

First published December 11, 2019.

in motor neurons, thereby damaging these cells in a Ca^{2+} -dependent manner (2–4). This prompted us to explore if ALS-T2DM serum drives similar pathological events in mouse islet cells. To implement such an exploration, we collected 4 types of sera from healthy human subjects (HSs) and patients with ALS, T2DM, and ALS-T2DM (SI Appendix, Table S1). During the course of the present study, 2 separate batches of sera were collected. The first batch of sera was obtained from 12 patients with ALS-T2DM as well as 12 HSs, 8 patients with ALS, and 8 patients with T2DM as controls (SI Appendix, Table S1). With each individual serum we treated dissociated islet cells and conducted high-throughput measurements of K^+ depolarization-induced $[Ca^{2+}]_i$ responses. General analysis of pooled data revealed that average $[Ca^{2+}]_i$ response to stimulation with 25 mM KCl in the ALS-T2DM serum-treated group was significantly greater than that in groups subjected to treatment with HS, ALS, or T2DM sera (Fig. 1A). Furthermore, the 3 latter groups did not significantly differ in this parameter (Fig. 1A). Interestingly, thorough analysis of individual data showed that 7 out of 12 ALS-T2DM sera produced significant increases in $[Ca^{2+}]_i$ in response to 25 mM KCl, whereas the rest did not, in comparison to HS, ALS, and T2DM sera (Fig. 1B and C). To corroborate the results obtained with the first batch of sera, we repeated $[Ca^{2+}]_i$ measurements with the second batch of sera donated by 5 patients with ALS-T2DM and 6 T2DM patients (SI Appendix, Table S1 and Fig. 1D and E). Consistent with the first batch of sera, the ALS-T2DM group displayed a significant increase in mean $[Ca^{2+}]_i$ response to KCl depolarization compared to the T2DM group, and 3 out of 5 ALS-T2DM sera gave rise to

significant elevations in K^+ -induced $[Ca^{2+}]_i$ responses in comparison to T2DM sera (Fig. 1D and E). Moreover, the specific Ca_v1 channel blocker nifedipine almost completely ablated K^+ -induced $[Ca^{2+}]_i$ responses in mouse islet cells (SI Appendix, Fig. S1). In addition, there was no significant difference in basal $[Ca^{2+}]_i$ between T2DM and ALS-T2DM groups (Fura-2 F340/F380 ratio for the T2DM group of 0.537 ± 0.028 vs. Fura-2 F340/F380 ratio for the ALS-T2DM group of 0.541 ± 0.017 , $P > 0.05$). Taken together, 60% of ALS-T2DM patients have sera that authentically exaggerate K^+ -induced $[Ca^{2+}]_i$ responses in islet cells. These ALS-T2DM sera were defined as positive ALS-T2DM sera and randomly chosen for subsequent experiments.

Pathogenic IgGs Present in Positive ALS-T2DM Sera Enhance K^+ -Induced $[Ca^{2+}]_i$ Responses in Mouse Islet Cells. It has been demonstrated that pathogenic IgGs reside in sera of ALS patients and account for a great deal of Ca^{2+} -dependent destruction of motor neurons and skeletal muscle cells (4, 15–18). This raised the question of whether IgGs in positive ALS-T2DM sera (ALS-T2DM-IgGs) also serve as molecular pathogenic factors to impair islet cell function and survival by perturbing $[Ca^{2+}]_i$ homeostasis. To tackle this question, we purified IgGs from positive ALS-T2DM sera and T2DM sera. Thereafter, we measured the effects of these purified IgGs on K^+ -induced $[Ca^{2+}]_i$ responses in islet cells. Incubation with these individual ALS-T2DM-IgGs induced significantly stronger $[Ca^{2+}]_i$ responses to 25 mM KCl in comparison to exposure to IgGs from T2DM sera (T2DM-IgGs) in mouse islet cells (Fig. 2A and B). Furthermore, the effect of ALS-T2DM-IgGs was lost when boiled (Fig. 2C). Cells exposed to either boiled ALS-T2DM-IgGs or T2DM-IgGs responded similarly to KCl stimulation with regard to increases in $[Ca^{2+}]_i$ (Fig. 2C). These data demonstrate that ALS-T2DM-IgG in ALS-T2DM sera enhances K^+ -induced $[Ca^{2+}]_i$ responses.

Positive ALS-T2DM Sera Up-regulate Ca_v1 Channels through Direct Interaction with $Ca_v\alpha_2\delta_1$ Subunits in Mouse Islet Cells. Autoantibodies against $Ca_v1.1$, $Ca_v2.1$, and $Ca_v2.2$ subunits have been demonstrated to be present in ALS patients (15, 16). Importantly, these autoantibodies enhance Ca^{2+} conductivity of these Ca^{2+} -conducting pores, resulting in excessively high $[Ca^{2+}]_i$ and, consequently, Ca^{2+} -dependent cytotoxicity in skeletal muscle cells and neurons (4, 15–19). Of particular importance is that selective $Ca_v1.1$, $Ca_v2.1$, and $Ca_v2.2$ channel blockers substantially improve defects in neuromuscular activity and viability induced by IgGs from ALS patients (2, 18, 20–23). Our finding that both ALS-T2DM serum and ALS-T2DM-IgG promote K^+ -evoked $[Ca^{2+}]_i$ responses suggests that $Ca_v1.2$ channels might serve as downstream targets of ALS-T2DM serum and ALS-T2DM-IgG. It is well known that depolarization-evoked $[Ca^{2+}]_i$ responses in mouse islet cells primarily result from Ca^{2+} influx through $Ca_v1.2$ channels (24).

To clarify if ALS-T2DM serum affects β cell Ca_v channels, we examined the effect of positive ALS-T2DM serum on β cell Ca_v channel currents. Indeed, whole-cell patch-clamp analysis revealed that treatment with individual positive ALS-T2DM sera obtained in the first and second batches significantly elevated whole-cell Ca_v channel currents in mouse β cells, as manifested by representative whole-cell Ca^{2+} current traces and average Ca_v channel current density, in comparison to T2DM serum exposure (Fig. 3A–C). These data verify that ALS-T2DM serum up-regulates Ca_v1 channels, causing pathologically exaggerated $[Ca^{2+}]_i$ responses.

The up-regulation of Ca_v1 channels by ALS-T2DM serum raises the possibility that IgGs in ALS-T2DM serum may target Ca_v channel subunits in β cells. We chose the most important pore-forming subunit $Ca_v1.2$ as a starting point. Immunoprecipitation assays followed by immunoblot analysis showed that antibodies against $Ca_v1.2$ subunits efficiently pulled down $Ca_v1.2$




Fig. 1. Effects of ALS-T2DM sera on KCl-induced $[Ca^{2+}]_i$ responses in mouse islet cells. (A) Δ Fura-2 F340/F380 ratios showing average net increases in $[Ca^{2+}]_i$ evoked by K^+ depolarization in groups subjected to 10 h of treatment with ALS-T2DM ($n = 12$), HS ($n = 12$), ALS ($n = 8$), or T2DM sera ($n = 8$) obtained from the first batch of patients. (B) Δ Fura-2 F340/F380 ratios illustrating net increases in $[Ca^{2+}]_i$ induced by K^+ stimulation in cells treated for 10 h with 12 individual ALS-T2DM sera as well as 12 HS, 8 ALS, and 8 T2DM sera collected from the first batch of patients. (C) Example recordings of Fura-2 F340/F380 ratios showing $[Ca^{2+}]_i$ responses to 25 mM KCl in cells following 10 h of incubation with ALS-T2DM serum (Right) or T2DM serum (Left). (D) Δ Fura-2 F340/F380 ratios illustrating mean net increases in $[Ca^{2+}]_i$ induced by K^+ stimulation in groups treated for 10 h with ALS-T2DM ($n = 5$) and T2DM sera ($n = 6$) donated by the second batch of patients. (E) Δ Fura-2 F340/F380 ratios showing net increases in $[Ca^{2+}]_i$ evoked by K^+ depolarization in cells treated for 10 h with 5 individual ALS-T2DM and 6 T2DM sera obtained from the second batch of patients. Ten out of 17 positive ALS-T2DM sera significantly enhance KCl-induced $[Ca^{2+}]_i$ responses in comparison to HS, ALS, and T2DM sera. * $P < 0.05$ and ** $P < 0.01$ vs. T2DM, ALS, or HS group.




Fig. 2. Effects of IgGs purified from positive ALS-T2DM sera on KCl-induced $[Ca^{2+}]_i$ responses in mouse islet cells. (A) Representative $[Ca^{2+}]_i$ responses to 25 mM KCl in islet cells following exposure to T2DM-IgG (Left) and ALS-T2DM-IgG (Right). (B) Δ Fura-2 F340/F380 ratios showing net increases in $[Ca^{2+}]_i$ evoked by 25 mM KCl in islet cells treated with IgGs purified from 8 T2DM sera or 6 positive ALS-T2DM sera. * $P < 0.05$ and ** $P < 0.01$ vs. the T2DM-IgG group. (C) Δ Fura-2 F340/F380 ratios showing net increases in $[Ca^{2+}]_i$ in islet cells treated with T2DM-IgG, ALS-T2DM-IgG, or boiled ALS-T2DM-IgG. Experiments were done with 8 T2DM-IgGs, 6 ALS-T2DM-IgGs, or 6 boiled ALS-T2DM-IgGs. ** $P < 0.01$ vs. the T2DM-IgG group and # $P < 0.05$ vs. the boiled ALS-T2DM-IgG group.

subunits from the membrane fraction of insulin-secreting RINm5f cells (SI Appendix, Fig. S2A). However, neither ALS-T2DM-IgGs nor T2DM-IgGs could recognize the immunoprecipitated $Ca_v1.2$ subunits under denaturing or renaturing conditions (SI Appendix, Fig. S2B and C). Furthermore, both ALS-T2DM-IgGs and T2DM-IgGs could not specifically fish out $Ca_v1.2$ subunits and additional proteins (SI Appendix, Fig. S2D and E). These results show that ALS-T2DM-IgGs could not strongly bind to immunoprecipitated $Ca_v1.2$ subunits under such experimental conditions.

The $Ca_v\alpha_2\delta_1$ subunit, an important constituent of Ca_v channel complexes, including the β cell $Ca_v1.2$ channel complex, is critical for the surface expression of functional Ca_v channels (12, 25–27). Moreover, the entire $Ca_v\alpha_2\delta_1$ is exposed extracellularly (28). Among all β cell $Ca_v1.2$ channel components, they have the highest likelihood of serving as targets for factors, such as ALS-T2DM-IgGs, of positive ALS-T2DM sera. In addition, polyclonal anti- $Ca_v\alpha_2\delta_1$ antibodies selectively recognize extracellular $Ca_v\alpha_2\delta_1$ subunits associated with the plasma membrane of living cells bathed in a physiological solution (28, 29). This prompted us to clarify if ALS-T2DM-IgGs interacts with $Ca_v\alpha_2\delta_1$ subunits under physiological conditions by using anti- $Ca_v\alpha_2\delta_1$ antibodies. We carried out 4-(2-[6-(Diocetyl-amino)-2-naphthalenyl]ethenyl)-1-(3-sulfopropyl)pyridinium inner salt (di-8-ANEPPS) labeling of the plasma membrane and immunofluorescence staining of $Ca_v\alpha_2\delta_1$ subunits in intact living mouse islet cells and

tsA-201 cells stably expressing $Ca_v\alpha_2\delta_1$ subunits coincubated with antibodies against $Ca_v\alpha_2\delta_1$ subunits and ALS-T2DM-IgGs. $Ca_v\alpha_2\delta_1$ -specific immunofluorescence was clearly localized in the di-8-ANEPPS-labeled plasma membrane (Fig. 3D–R and SI Appendix, Fig. S3). Interestingly, ALS-T2DM-IgGs effectively competed with the anti- $Ca_v\alpha_2\delta_1$ antibodies for the extracellular $Ca_v\alpha_2\delta_1$ subunits, resulting in a significant reduction in the immunofluorescence intensity of the anti- $Ca_v\alpha_2\delta_1$ antibodies in comparison to HS-IgGs or T2DM-IgGs (Fig. 3S and T). This verifies that ALS-T2DM-IgGs are capable of directly interacting with $Ca_v\alpha_2\delta_1$ subunits in living cells in the absence of interferences from detergents, high ionic strengths, and substantial rinsing, which are unavoidable in immunoprecipitation and immunoblot analyses of association between ALS-T2DM-IgGs and $Ca_v1.2$ subunits.

Positive ALS-T2DM Sera Interfere with Mitochondrial Function in Mouse Islet Cells. Translation of an excessive elevation of $[Ca^{2+}]_i$ into mitochondrial Ca^{2+} overload results in mitochondrial membrane depolarization, concomitant mitochondrial dysfunction, and eventual apoptosis, thus playing an important role in driving Ca^{2+} -dependent cell death (30). This made us wonder whether such a mitochondrial mechanism is able to convert the ALS-T2DM serum-induced exaggeration of $[Ca^{2+}]_i$ to mitochondrial dysfunction in mouse islet cells. Therefore, we measured mitochondrial membrane potential in mouse islet cells using rhodamine 123. As shown in Fig. 4A–C, ALS-T2DM serum-treated cells displayed a significant decrease not only in basal fluorescence intensity of rhodamine 123 (Fig. 4A and B) but also in the glucose-induced quenching of rhodamine 123 (Fig. 4A and C), compared to those cells incubated with T2DM serum. The effects are attributed to reduced amounts of rhodamine 123 loaded into mitochondria due to less negative mitochondrial membrane potential, i.e., mitochondrial dysfunction, induced by ALS-T2DM serum treatment (31). Our results demonstrate that mouse islet cells insulted by exaggerated $[Ca^{2+}]_i$ resulting from exposure to positive ALS-T2DM sera undergo mitochondrial dysfunction and suggest that ALS-T2DM serum-induced mitochondrial dysfunction is most likely to drive islet cell death.

Positive ALS-T2DM Sera Impair $[Ca^{2+}]_i$ Dynamics and Insulin Secretion in Mouse Islets. Normal glucose homeostasis critically relies on adequately functioning β cells (32). The function of β cells is under the control of exquisitely fine-tuned $[Ca^{2+}]_i$ dynamics that serves as fingerprints for β cell well-being (12, 13, 24, 33, 34). This made us wonder if ALS-T2DM serum drives disorganized $[Ca^{2+}]_i$ dynamics and impaired insulin secretion in islets, thereby accounting for aberrant glucose homeostasis often observed in ALS patients.

We characterized $[Ca^{2+}]_i$ dynamics in β cells situated within intact islets during glucose stimulation. As shown in photomicrographs of Fura-2-loaded mouse islets, ALS-T2DM serum treatment made islets become irregular and disintegrated (Fig. 5A, Lower). In striking contrast, incubation with T2DM serum did not alter the morphology of islets that were intact with spherical shapes and smooth boundaries (Fig. 5A, Upper). Indeed, ALS-T2DM serum-treated islets showed chaotic $[Ca^{2+}]_i$ dynamics manifested as a relatively steady increase in $[Ca^{2+}]_i$ with tiny amplitude oscillations in response to 11.1 mM glucose (Fig. 5B, Lower and SI Appendix, Fig. S4, Lower). However, islets exposed to T2DM serum displayed a normal $[Ca^{2+}]_i$ profile, characterized by fast oscillations superimposed on slow oscillations, following stimulation with 11.1 mM glucose (Fig. 5B, Upper and SI Appendix, Fig. S4, Upper). These results reveal that ALS-T2DM serum does indeed potently derange $[Ca^{2+}]_i$ handling in β cells.

The primary function of β cells is glucose-stimulated insulin secretion that crucially depends on Ca_v channel-mediated Ca^{2+} influx and complex $[Ca^{2+}]_i$ dynamics (12, 13, 24, 33–35).




Fig. 3. Effects of positive ALS-T2DM sera on whole-cell Ca_V currents in mouse islet cells and influences of ALS-T2DM-IgGs on $\text{Ca}_V\alpha_2\delta_1$ immunofluorescence in the plasma membrane of living mouse islet cells and tsA-201 cells stably expressing $\text{Ca}_V\alpha_2\delta_1$ subunits. (A) Representative whole-cell Ca^{2+} currents in islet β cells incubated with T2DM serum (Upper) or ALS-T2DM serum (Lower) for 10 h. (B and C) Whole-cell current–voltage relationships of Ca_V channels in cells treated with the first batch (B) of one T2DM or one ALS-T2DM serum and with the second batch (C) of 3 T2DM or 3 ALS-T2DM sera. * $P < 0.05$ for the ALS-T2DM₁ group vs. the T2DM₃ group in B; * $P < 0.05$ for the ALS-T2DM₁₄, ALS-T2DM₁₆, or ALS-T2DM₁₇ group vs. the T2DM_{9,10,12} group and # $P < 0.05$ for the ALS-T2DM₁₄ or ALS-T2DM₁₆ group vs. the T2DM_{9,10,12} group in (C). $\text{CD}_{\text{Ca}^{2+}}$, Ca^{2+} current density. (D–R) Representative di-8-ANEPPS fluorescence (D, E, J, K, and L), $\text{Ca}_V\alpha_2\delta_1$ immunofluorescence (F, G, M, N, and O), and their overlay images (H, I, P, Q, and R) of living mouse islet cells (D–I) and tsA-201 cells stably expressing $\text{Ca}_V\alpha_2\delta_1$ subunits (J–R) incubated with rabbit polyclonal anti- $\text{Ca}_V\alpha_2\delta_1$ antibodies in the presence of HS-IgGs (J, M, and P), T2DM-IgGs (D, F, H, K, N, and Q), and ALS-T2DM-IgGs (E, G, I, L, O, and R). (Scale bar, 10 μm .) (S and T) Mean $\text{Ca}_V\alpha_2\delta_1$ immunofluorescence/di-8-ANEPPS fluorescence ratio in living mouse islet cells (S) and tsA-201 cells stably expressing $\text{Ca}_V\alpha_2\delta_1$ subunits (T) incubated with $\text{Ca}_V\alpha_2\delta_1$ antibodies plus HS-IgGs ($\text{Ca}_V\alpha_2\delta_1/\text{HS-IgG}$), T2DM-IgGs ($\text{Ca}_V\alpha_2\delta_1/\text{T2DM-IgG}$), and ALS-T2DM-IgGs ($\text{Ca}_V\alpha_2\delta_1/\text{ALS-T2DM-IgG}$). Experiments were done with 3 T2DM and 3 ALS-T2DM sera. * $P < 0.05$ vs. the $\text{Ca}_V\alpha_2\delta_1/\text{HS-IgG}$ group or $\text{Ca}_V\alpha_2\delta_1/\text{T2DM-IgG}$ group.

ALS-T2DM serum-induced defects in $[\text{Ca}^{2+}]_i$ dynamics should cause impaired glucose-stimulated insulin secretion. T2DM serum- and ALS-T2DM serum-treated islets released a similar amount of insulin following incubation with 11.1 mM glucose (Fig. 5 C and D). However, insulin secreted from T2DM serum-treated islets at 11.1 mM glucose was significantly greater than that at 3.3 mM glucose, whereas insulin released from ALS-T2DM serum-treated islets at 11.1 mM glucose did not significantly differ from that at 3.3 mM glucose due to increased basal insulin release (Fig. 5 C and D). In addition, the insulin content of ALS-T2DM serum-treated islets was significantly lower than that of islets exposed to T2DM serum (Fig. 5E). These data suggest that exposure to ALS-T2DM serum interferes with the ability of the β cell to maintain adequate insulin release.

Positive ALS-T2DM Sera Reduce Mouse Islet Cell Viability in an IgG- and Ca_V1 Channel-Dependent Manner. The exaggerated Ca_V1 channel-mediated Ca^{2+} influx, increased $[\text{Ca}^{2+}]_i$, and disturbed $[\text{Ca}^{2+}]_i$ dynamics in islet cells exposed to ALS-T2DM serum might explain the destructive action of this serum on islet integrity and islet insulin content. Therefore, we examined the possible effects of ALS-T2DM serum on islet cell survival. WST-1 assay showed that ALS-T2DM serum exposure significantly decreased islet cell viability, as reflected by significantly reduced WST-1 absorbance, in comparison to treatment with T2DM serum (Fig. 6A). Furthermore, cell death imaging with SYTOX Orange nucleic acid stain revealed that SYTOX Orange–positive profiles, representing dead nuclei, were significantly greater in dissociated islet cells incubated with ALS-T2DM serum compared to T2DM serum-treated ones (Fig. 6 B and C). Furthermore, the selective Ca_V1

channel blocker nifedipine fully ablated ALS-T2DM serum-induced reduction of islet cell viability (Fig. 6D). In addition, treatment with ALS-T2DM-IgGs significantly reduced mouse islet cell viability in comparison to incubation with T2DM-IgGs. The effects of ALS-T2DM-IgG on islet cell viability were effectively ablated by boiling (Fig. 6E). The results demonstrate that positive ALS-T2DM sera interfere with islet cell survival in an IgG- and Ca_V1 channel-dependent manner. Taken together, our results suggest that ALS-T2DM serum treatment destroys islet cells by excessively increasing Ca_V1 channel-mediated Ca^{2+} influx and $[\text{Ca}^{2+}]_i$, and then pathologically translating the exaggerated $[\text{Ca}^{2+}]_i$ to eventual mitochondrial dysfunction.

Discussion

We have identified a subgroup of ALS-T2DM patients who have positive sera that exaggerate $[\text{Ca}^{2+}]_i$ in pancreatic islet cells upon depolarization. This suggests that these positive sera are likely to interfere with Ca_V channels via serum molecular constituent(s). Indeed, we demonstrate that pathogenic IgG is accommodated in the sera of this subgroup of ALS-T2DM patients. This not only establishes a mechanistic link between ALS and T2DM but also suggests a potential role of altered humoral immunity in the development of ALS-associated T2DM.

Importantly, we reveal that ALS-T2DM serum significantly increases whole-cell Ca^{2+} currents predominantly passing through $\text{Ca}_V1.2$ channels in mouse β cells (24). This mechanistically explains how ALS-T2DM serum and ALS-T2DM-IgG promote K^+ -evoked $[\text{Ca}^{2+}]_i$ responses and pinpoints that $\text{Ca}_V1.2$ channels most likely serve as downstream targets of ALS-T2DM serum and ALS-T2DM-IgG. This finding is intriguing since ALS-T2DM serum




Fig. 4. Effects of positive ALS-T2DM sera on mouse islet cell mitochondrial membrane potential. (A) Sample rhodamine 123 fluorescence traces registered in a T2DM serum-treated cell (black trace) and in a cell exposed to ALS-T2DM serum (red trace) following glucose (G) and FCCP stimulation. FCCP: carbonyl cyanide *p*-trifluoromethoxyphenylhydrazone, AU: arbitrary units. (B) Basal relative rhodamine 123 fluorescence intensities (bRFI) in cells incubated with T2DM, ALS-T2DM₁₄, and ALS-T2DM₁₇ sera for 24 h as well as T2DM and ALS-T2DM₁₆ sera for 12 h. (C) Net changes in glucose-induced quenching of rhodamine 123 (Δ FI) in cells treated with T2DM, ALS-T2DM₁₄, and ALS-T2DM₁₇ sera for 24 h as well as T2DM and ALS-T2DM₁₆ sera for 12 h. Experiments were done with 2 T2DM and 3 ALS-T2DM sera. * $P < 0.05$ and ** $P < 0.01$ vs. the T2DM group.

is verified to functionally interfere with mouse β cell $Ca_v1.2$ channels, which almost exclusively mediate the nifedipine-sensitive Ca^{2+} currents (11–13, 36, 37).

Interestingly, we found that ALS-T2DM-IgGs are strong enough to compete with an IgG rabbit polyclonal antibody specific to the extracellular epitope of $Ca_v\alpha_2\delta_1$ subunits in living islet cells and in tsA-201 cells stably expressing $Ca_v\alpha_2\delta_1$ subunits. This is in accordance with the fact that the $Ca_v\alpha_2\delta_1$ subunits are entirely exposed to the extracellular space and thereby the most accessible to serum components among all β cell $Ca_v1.2$ channel subunits. In addition, the $Ca_v\alpha_2\delta_1$ subunit serves as an indispensable building element of the β cell $Ca_v1.2$ channel complex to up-regulate the conductivity and surface expression of functional Ca_v channels (12, 25–27). Based on our results, we propose that ALS-T2DM-IgGs serve as autoantibodies that immunocapture $Ca_v\alpha_2\delta_1$ subunits in the plasma membrane, thereby enhancing Ca_v1 channel-mediated Ca^{2+} influx and $[Ca^{2+}]_i$ in islet cells. This process likely occurs through allosteric activation and/or gradual accumulation of Ca_v1 channels in the β cell plasma membrane since antibodies in some cases activate and accumulate rather than inhibit and neutralize their binding partners (38, 39). This autoimmune mechanism is particularly interesting since ALS autoantibodies have been shown to target only $Ca_v1.1$, $Ca_v2.1$, and $Ca_v2.2$ channels prior to the present work (2, 18, 20–23). Now the immunocapture of β cell $Ca_v\alpha_2\delta_1$ subunits by ALS-T2DM-IgGs and consequent up-regulation of β cell $Ca_v1.2$ channels come into the picture. Importantly, the present work reveals that humoral autoimmunity arises as the pathogenic machinery leading to this subset of diabetes. It is intriguing to explore if such a humoral mechanism not only operates in the pathogenesis of T2DM but also in that of type 1 diabetes in addition to a T cell-mediated autoimmune destruction of β cells (40). Moreover, our findings offer a causal link between ALS and T2DM and shed light on

potential therapeutic targets for prevention and treatment of a subgroup of ALS-T2DM patients.

The mitochondrion is a vulnerable target downstream of excessive accumulation of $[Ca^{2+}]_i$ to mediate Ca^{2+} -dependent impairments in cell function and viability (30, 41). Consequently, islet cells exposed to positive ALS-T2DM sera not only display exaggerated $[Ca^{2+}]_i$ but also mitochondrial dysfunction. These findings give a strong rationale for the ALS-T2DM serum-induced islet cell dysfunction and death. We found that positive ALS-T2DM sera derange $[Ca^{2+}]_i$ dynamics, impair insulin secretion, and drive islet cell death in a Ca_v1 channel- and IgG-dependent manner. The occurrence of these pathological phenotypes is well accounted for by impaired mitochondrial function. Our data thus suggest that IgG autoantibodies in ALS-T2DM sera immunocapture $Ca_v\alpha_2\delta_1$ subunits in the plasma membrane, resulting in a destructive exaggeration of $[Ca^{2+}]_i$ followed by its pathological translation into mitochondrial dysfunction, subsequent impairment of insulin secretion, eventual islet cell death, and diabetes. Of note, in vivo ALS-T2DM-IgGs may target peripheral insulin-sensitive tissues of patients, leading to insulin resistance (8).

The exact reason why only a fraction of ALS patients develop diabetes is unclear but is most likely due to a heterogeneous β cell sensitivity to the evoked $[Ca^{2+}]_i$ challenges. In addition, both ALS and T2DM represent an etiologically heterogeneous group of disorders where patients are likely to experience particular environmental challenges on top of their specific genetic




Fig. 5. Effects of positive ALS-T2DM sera on mouse islet $[Ca^{2+}]_i$ dynamics and insulin secretion. (A) Sample photomicrographs of Fura-2-loaded mouse islets following 10 h of treatment with T2DM (Upper) and ALS-T2DM serum (Lower). (Scale bar, 50 μ m.) (B) Representative $[Ca^{2+}]_i$ traces out of a total of 72 and 40 islets correspondingly subjected to 10 h of incubation with T2DM (Upper) or ALS-T2DM serum (Lower), followed by perfusion with 3.3 or 11.1 mM glucose (G), respectively. (C and D) Insulin secretion from islets treated with T2DM or ALS-T2DM serum for 10 h followed by exposure to 3.3 or 11.1 mM G for 30 min. * $P < 0.05$ vs. the T2DM/3.3 mM G group (C). Net insulin secretion induced by 11 mM G in T2DM and ALS-T2DM groups. *** $P < 0.01$ vs. the T2DM group (D). Experiments were done with 6 T2DM and 6 ALS-T2DM sera in triplicate. (E) Insulin content in islets exposed to T2DM or ALS-T2DM serum for 10 h. * $P < 0.05$ vs. the T2DM group. Experiments were done with 4 T2DM and 3 ALS-T2DM sera in triplicate.




Fig. 6. Effects of positive ALS-T2DM sera on mouse islet cell viability. (A) Mean WST-1 absorbance showing the viability of dissociated islet cells exposed to T2DM or ALS-T2DM serum. Experiments were done with 5 T2DM and 7 ALS-T2DM sera. $**P < 0.01$ vs. the T2DM group. (B) Example SYTOX Orange fluorescence (Left), transmitted light (Middle), and their overlay images (Right) of dissociated islet cells incubated with T2DM (Upper) and ALS-T2DM serum (Lower) followed by exposure to SYTOX Orange nucleic acid stain. (Scale bar, 50 μ m.) (C) Mean percentage of dead cells labeled with SYTOX Orange nucleic acid stain in T2DM and ALS-T2DM groups. Experiments were done with 3 T2DM and 3 ALS-T2DM sera in triplicate. $*P < 0.05$ vs. the T2DM group. (D) Mean WST-1 absorbance showing the viability of dissociated islet cells exposed to 3 T2DM sera or 3 ALS-T2DM sera in the absence and presence of the Ca_v1 channel blocker nifedipine. $**P < 0.01$ vs. the T2DM group and $^{\#}P < 0.05$ vs. the ALS-T2DM/nifedipine group. (E) Mean WST-1 absorbance in islet cells following 24 h exposure to T2DM-IgG, ALS-T2DM-IgG, or boiled ALS-T2DM-IgG. Experiments were done with 3 T2DM-IgGs, 3 ALS-T2DM-IgGs, or 3 boiled ALS-T2DM-IgGs. $**P < 0.01$ vs. the T2DM-IgG group and $^{\#\#}P < 0.01$ vs. the boiled ALS-T2DM-IgG group.

predisposition. In this context, a specific combination of genetic and environmental factors may trigger the production of autoantibodies to the autoantigen $Ca_v\alpha_2\delta_1$ subunit. This leads to a pathological Ca_v channel conductivity with resulting damage to motor neurons and islet cells.

In conclusion, the present work demonstrates that a subgroup of ALS-T2DM patients have sera that enhance Ca_v1 channel-mediated Ca^{2+} influx, resulting in exaggerated $[Ca^{2+}]_i$. These effects are attributed to the fact that the sera accommodate cytotoxic IgG autoantibodies that immunocapture $Ca_v\alpha_2\delta_1$ subunits. As a consequence, impairments in $[Ca^{2+}]_i$ dynamics, mitochondrial function, insulin secretion, and cell viability occur. This suggests that cytotoxic ALS-T2DM-IgG autoantibodies serve as a causal link between ALS and T2DM by interacting with and modulating the $Ca_v\alpha_2\delta_1$ - Ca_v1 channel complex, which may lay the foundation for a pharmacological treatment strategy for patients suffering from a combination of these severe diseases.

Methods

Animals. Male C57BL/6J mice aged from 8 to 10 wk were purchased from The Jackson Laboratory (Bar Harbor, ME). All animal experiments were conducted according to the guidelines of the Ethics Committee at Pohang University of Science and Technology (2011-0001) and the Animal Experiment Ethics Committee at Karolinska Institutet (N183/13).

Enrollment of Patients. Seventeen patients with ALS-T2DM (11 males and 6 females, age: 55.0 ± 2.1), 12 HSs, 9 patients with ALS, and 14 patients with T2DM (7 males and 7 females, age: 60.4 ± 2.5) were randomly selected (SI Appendix, Table S1). One ALS patient carrying a SOD1 mutation was excluded from the study (SI Appendix, Table S1). Ethical approval to use serum from patients was obtained from Hanyang University Hospital in Seoul, Korea (HYUH 2006-04-001-004). Blood samples were collected after obtaining written informed consent. Actual patients had no known family history of neuromuscular disease or wasting. All subjects diagnosed with T2DM were well controlled with hypoglycemic agents or insulin.

Additional experimental procedures are presented in the SI Appendix.

Data Availability. All of the data, associated protocols, code, and materials for this study are available within the paper and its SI Appendix.

ACKNOWLEDGMENTS. This research was supported by the Basic Science Research Program (Grant NRF-2016K1A1A2912722) and Brain Research Program (Grant NRF-2018M3C7A1056512) of the National Research Foundation of Korea (NRF), funded by the Ministry of Science and Information and Communications Technology; the Swedish Diabetes Association; Funds of Karolinska Institutet; the Swedish Research Council; the Novo Nordisk Foundation; the Erling-Persson Family Foundation; the Strategic Research Program in Diabetes at Karolinska Institutet; the European Research Council (Grant ERC-2013-AdG 338936-BetalImage); the Knut and Alice Wallenberg Foundation; Skandia Insurance Company, Ltd.; the Diabetes and Wellness Foundation; the Bert von Kantzow Foundation; the Lee Kong Chian School of Medicine; Nanyang Technological University (Start-Up Grant for P.-O.B.); and the Stichting af Jochnick Foundation. The work of A.C.D. was supported by a Wellcome Trust Investigator Award (098360/Z/12/Z).

1. B. R. Brooks, Natural history of ALS: Symptoms, strength, pulmonary function, and disability. *Neurology* **47** (suppl. 2), 571–582 (1996).
2. R. G. Smith *et al.*, Cytotoxicity of immunoglobulins from amyotrophic lateral sclerosis patients on a hybrid motoneuron cell line. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 3393–3397 (1994).
3. L. T. Tran, B. J. Gentil, K. E. Sullivan, H. D. Durham, The voltage-gated calcium channel blocker lomerizine is neuroprotective in motor neurons expressing mutant SOD1, but not TDP-43. *J. Neurochem.* **130**, 455–466 (2014).
4. A. H. Pullen, M. Demestre, R. S. Howard, R. W. Orrell, Passive transfer of purified IgG from patients with amyotrophic lateral sclerosis to mice results in degeneration of motor neurons accompanied by Ca^{2+} enhancement. *Acta Neuropathol.* **107**, 35–46 (2004).
5. B. Billups, I. D. Forsythe, Presynaptic mitochondrial calcium sequestration influences transmission at mammalian central synapses. *J. Neurosci.* **22**, 5840–5847 (2002).
6. P. Shi, J. Gal, D. M. Kwinter, X. Liu, H. Zhu, Mitochondrial dysfunction in amyotrophic lateral sclerosis. *Biochim. Biophys. Acta* **1802**, 45–51 (2010).
7. S. C. Barber, P. J. Shaw, Oxidative stress in ALS: Key role in motor neuron injury and therapeutic target. *Free Radic. Biol. Med.* **48**, 629–641 (2010).
8. P. F. Pradat *et al.*, Impaired glucose tolerance in patients with amyotrophic lateral sclerosis. *Amyotroph. Lateral Scler.* **11**, 166–171 (2010).
9. K. Araki *et al.*, TDP-43 regulates early-phase insulin secretion via $Ca_v1.2$ -mediated exocytosis in islets. *J. Clin. Invest.* **130**, 3578–3593 (2019).
10. A. Joardar, E. Manzo, D. C. Zarnescu, Metabolic dysregulation in amyotrophic lateral sclerosis: Challenges and opportunities. *Curr. Genet. Med. Rep.* **5**, 108–114 (2017).
11. S. N. Yang, P. O. Berggren, β -cell Ca_v channel regulation in physiology and pathophysiology. *Am. J. Physiol. Endocrinol. Metab.* **288**, E16–E28 (2005).
12. S. N. Yang, P. O. Berggren, The role of voltage-gated calcium channels in pancreatic β -cell physiology and pathophysiology. *Endocr. Rev.* **27**, 621–676 (2006).
13. S. N. Yang *et al.*, Ionic mechanisms in pancreatic β cell signaling. *Cell. Mol. Life Sci.* **71**, 4149–4177 (2014).
14. E. Nanou, W. A. Catterall, Calcium channels, synaptic plasticity, and neuropsychiatric disease. *Neuron* **98**, 466–481 (2018).
15. R. G. Smith *et al.*, Serum antibodies to L-type calcium channels in patients with amyotrophic lateral sclerosis. *N. Engl. J. Med.* **327**, 1721–1728 (1992).
16. F. Kimura *et al.*, Amyotrophic lateral sclerosis patient antibodies label Ca^{2+} channel α_1 subunit. *Ann. Neurol.* **35**, 164–171 (1994).
17. S. A. Frattantoni, G. Weisz, A. M. Pardal, R. C. Reisin, O. D. Uchitel, Amyotrophic lateral sclerosis IgG-treated neuromuscular junctions develop sensitivity to L-type calcium channel blocker. *Muscle Nerve* **23**, 543–550 (2000).
18. L. E. Gonzalez *et al.*, Amyotrophic lateral sclerosis-immunoglobulins selectively interact with neuromuscular junctions expressing P/Q-type calcium channels. *J. Neurochem.* **119**, 826–838 (2011).
19. M. Milošević *et al.*, Immunoglobulins G from patients with sporadic amyotrophic lateral sclerosis affects cytosolic Ca^{2+} homeostasis in cultured rat astrocytes. *Cell Calcium* **54**, 17–25 (2013).

20. R. Llinás *et al.*, IgG from amyotrophic lateral sclerosis patients increases current through P-type calcium channels in mammalian cerebellar Purkinje cells and in isolated channel protein in lipid bilayer. *Proc. Natl. Acad. Sci. U.S.A.* **90**, 11743–11747 (1993).
21. D. R. Mosier *et al.*, Amyotrophic lateral sclerosis immunoglobulins increase Ca²⁺ currents in a motoneuron cell line. *Ann. Neurol.* **37**, 102–109 (1995).
22. A. Losavio, S. Muchnik, Spontaneous acetylcholine release in mammalian neuromuscular junctions. *Am. J. Physiol.* **273**, C1835–C1841 (1997).
23. S. Muchnik, A. Losavio, S. De Lorenzo, Effect of amyotrophic lateral sclerosis serum on calcium channels related to spontaneous acetylcholine release. *Clin. Neurophysiol.* **113**, 1066–1071 (2002).
24. P. Rorsman, M. Braun, Q. Zhang, Regulation of calcium in pancreatic α - and β -cells in health and disease. *Cell Calcium* **51**, 300–308 (2012).
25. V. Mastrolia *et al.*, Loss of $\alpha 2\delta$ -1 calcium channel subunit function increases the susceptibility for diabetes. *Diabetes* **66**, 897–907 (2017).
26. W. A. Catterall, Structure and regulation of voltage-gated Ca²⁺ channels. *Annu. Rev. Cell Dev. Biol.* **16**, 521–555 (2000).
27. W. A. Catterall, Voltage-gated calcium channels. *Cold Spring Harb. Perspect. Biol.* **3**, a003947 (2011).
28. A. Davies *et al.*, The $\alpha 2\delta$ subunits of voltage-gated calcium channels form GPI-anchored proteins, a posttranslational modification essential for function. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 1654–1659 (2010).
29. A. Favereaux *et al.*, Bidirectional integrative regulation of Cav1.2 calcium channel by microRNA miR-103: Role in pain. *EMBO J.* **30**, 3830–3841 (2011).
30. S. W. Tait, D. R. Green, Mitochondria and cell death: Outer membrane permeabilization and beyond. *Nat. Rev. Mol. Cell Biol.* **11**, 621–632 (2010).
31. L. B. Chen, Mitochondrial membrane potential in living cells. *Annu. Rev. Cell Biol.* **4**, 155–181 (1988).
32. D. Mathis, L. Vence, C. Benoist, β -Cell death during progression to diabetes. *Nature* **414**, 792–798 (2001).
33. P. O. Berggren *et al.*, Removal of Ca²⁺ channel β_3 subunit enhances Ca²⁺ oscillation frequency and insulin exocytosis. *Cell* **119**, 273–284 (2004).
34. L. E. Fridlyand, N. Tamarina, L. H. Philipson, Bursting and calcium oscillations in pancreatic beta-cells: Specific pacemakers for specific mechanisms. *Am. J. Physiol. Endocrinol. Metab.* **299**, E517–E532 (2010).
35. S. N. Yang, P. O. Berggren, Ca_v2.3 channel and PKC λ : New players in insulin secretion. *J. Clin. Invest.* **115**, 16–20 (2005).
36. V. Schulla *et al.*, Impaired insulin secretion and glucose tolerance in β cell-selective Ca_v1.2 Ca²⁺ channel null mice. *EMBO J.* **22**, 3844–3854 (2003).
37. M. J. Sinnegger-Brauns *et al.*, Isoform-specific regulation of mood behavior and pancreatic β cell and cardiovascular function by L-type Ca²⁺ channels. *J. Clin. Invest.* **113**, 1430–1439 (2004).
38. H. Wulff, P. Christophersen, P. Colussi, K. G. Chandy, V. Yarov-Yarovoy, Antibodies and venom peptides: New modalities for ion channels. *Nat. Rev. Drug Discov.* **18**, 339–357 (2019).
39. C. T. Chan *et al.*, Antibodies in the pathogenesis of hypertension. *BioMed Res. Int.* **2014**, 504045 (2014).
40. S. N. Yang, P. O. Berggren, The eye as a novel imaging site in diabetes research. *Pharmacol. Ther.* **197**, 103–121 (2019).
41. R. Rizzuto, D. De Stefani, A. Raffaello, C. Mammucari, Mitochondria as sensors and regulators of calcium signalling. *Nat. Rev. Mol. Cell Biol.* **13**, 566–578 (2012).