

Rethinking Endothelial Dysfunction as a Crucial Target in Fighting Heart Failure

Courtney Premer, PhD; Anthony J. Kanelidis, MD; Joshua M. Hare, MD; and Ivonne Hernandez Schulman, MD

Abstract

Endothelial dysfunction is characterized by nitric oxide dysregulation and an altered redox state. Oxidative stress and inflammatory markers prevail, thus promoting atherogenesis and hypertension, important risk factors for the development and progression of heart failure. There has been a reemerging interest in the role that endothelial dysfunction plays in the failing circulation. Accordingly, patients with heart failure are being clinically assessed for endothelial dysfunction via various methods, including flow-mediated vasodilation, peripheral arterial tonometry, quantification of circulating endothelial progenitor cells, and early and late endothelial progenitor cell outgrowth measurements. Although the mechanisms underlying endothelial dysfunction are intimately related to cardiovascular disease and heart failure, it remains unclear whether targeting endothelial dysfunction is a feasible strategy for ameliorating heart failure progression. This review focuses on the pathophysiology of endothelial dysfunction, the mechanisms linking endothelial dysfunction and heart failure, and the various diagnostic methods currently used to measure endothelial function, ultimately highlighting the therapeutic implications of targeting endothelial dysfunction for the treatment of heart failure.

© 2019 THE AUTHORS. Published by Elsevier Inc on behalf of Mayo Foundation for Medical Education and Research. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>) ■ Mayo Clin Proc Inn Qual Out 2019;3(1):1-13

Heart failure (HF) incidence is increasing worldwide as risk factors such as coronary artery disease (CAD), hypertension, diabetes, obesity, and lack of physical activity remain a pervasive problem.¹ Endothelial dysfunction plays an important role in the development of HF.^{2,3} It is characterized by nitric oxide (NO) dysregulation, inflammation, and oxidative stress, which impair the capacity of the vascular endothelium to perform its many functions, such as regulation of vascular tone via smooth muscle relaxation, antifibrinolysis, the clotting cascade, and inflammatory processes. All of these mechanisms play a key role in cardiovascular diseases (CVDs) and notably HF pathophysiology (Figure 1). Endothelial dysfunction was a large area of focus during the 1990s and early 2000s and has resurfaced as an emerging area of interest as a potential therapeutic target in patients with HF. In recent clinical trials, results suggest that improving endothelial function in patients with HF concordantly improves heart biomechanics.^{4,5}

As clinical trials progress, our understanding of the pathophysiology of endothelial function

in patients with HF, as well as our understanding of different diagnostic methodologies, needs to be further elucidated and validated. In this review, we discuss the current understanding of the pathophysiology and molecular mechanisms of endothelial dysfunction in patients with HF. Additionally, we review the different methods currently available to assess endothelial function, highlighting the advantages and disadvantages of each. Lastly, we review relevant clinical trials and the implications of targeting endothelial dysfunction in this patient population.

THE ROLE OF THE ENDOTHELIUM

The endothelium is comprised of a monolayer of endothelial cells that acts as a semiselective barrier between the bloodstream and the wall of the blood vessels and modulates blood flow, blood clotting, inflammation, and plasma permeability. The vascular endothelium regulates vasomotor tone, clotting factors, and inflammation.³ More specifically, the healthy vascular endothelium regulates vascular tone by balancing the production of vasodilators and vasoconstrictors, which affect smooth muscle dilation. Nitric

From the Interdisciplinary Stem Cell Institute (C.P., J.M.H., I.H.S.) and Katz Family Division of Nephrology and Hypertension (I.H.S.), University of Miami Miller School of Medicine, Miami, FL; and Department of Medicine, University of Chicago, Chicago, IL (A.J.K.).

ARTICLE HIGHLIGHTS


- The pathophysiology of endothelial dysfunction plays a significant role in the development and progression of heart failure and cardiovascular disease.
- There are numerous methods for measuring endothelial function, most notably being intracoronary arterial infusions combined with quantitative coronary angiography and Doppler imaging, flow-mediated vasodilation, peripheral arterial tonometry, endothelial progenitor cell-colony forming unit assay, and circulating endothelial progenitor cells, each with distinct advantages and disadvantages.
- Pharmacologic and nonpharmacologic therapies are under clinical investigation for targeting endothelial dysfunction as a means to improve heart failure and cardiovascular disease outcomes.

oxide is the predominant mediator of vasodilatation, but other endogenous factors such as bradykinin, acetylcholine (ACh), catecholamines, endothelium-derived hyperpolarizing factor

(EDHF), and prostacyclin also play a role. In fact, emerging research suggests that diminished EDHF-mediated vasodilation is a major contributor to endothelial dysfunction, since EDHF is crucial for small resistance artery relaxation, especially in the setting of high cholesterol or diabetes.⁶⁻⁸ Furthermore, the endothelium produces antiproliferative and anti-inflammatory cytokines as well as regulates the coagulation cascade through a balanced production of anti-coagulant and procoagulant factors. The function of the endothelium is impaired in patients with cardiovascular risk factors or CVD, and this is associated with increased oxidative stress and impaired NO balance, which contribute to the progression of the disease.

Molecular Mechanisms: The Role of NO Signaling

Because NO is fundamental to maintaining vascular homeostasis, its dysregulation is strongly implicated in endothelial dysfunction. Nitric oxide is synthesized when endothelial NO synthase (eNOS) converts L-arginine to L-citrulline. Once produced,



NO activates soluble guanylyl cyclase and thereby generates cyclic guanosine monophosphate (cGMP) to produce multiple downstream effects. The effects of NO are also mediated via the S-nitrosylation, a posttranslational modification, of effector proteins.⁹ Under normal conditions, NO is released from the endothelium and stimulates the relaxation of smooth muscle cells. This release of NO is most notably triggered by shear stress induced by blood flowing over the surface of the vessel; however, NO release can also be stimulated by agonists and vasoactive mediators (eg, Ach, histamine, thrombin, serotonin, adenosine diphosphate, bradykinin, norepinephrine, substance P, and isoproterenol, among others).¹⁰ Moreover, NO has anti-inflammatory (inhibiting leukocyte adhesion and infiltration), antithrombotic (increasing cGMP in platelets), and antioxidant (scavenging free radicals) effects.^{10,11}

With regard to the heart, physiologic NO signaling participates in myocardial contractility and relaxation as well as mitochondrial respiration.¹² These effects are due to interactions between NO and Ca²⁺-handling proteins, including the L-type Ca²⁺ channel, ryanodine receptor/Ca²⁺-release channel, and sarcoendoplasmic reticulum Ca²⁺ ATPase, the contractile myofilaments, and respiratory complexes. Importantly, NO signaling is regulated by its site of production, which is determined by the spatial localization of the NO synthase (NOS) enzymes.¹³ Nitric oxide modulates calcium signaling through S-nitrosylation of calcium channels, thereby regulating excitation-contraction coupling.^{9,14-16} In the setting of myocardial infarction or HF, the spatial localization of the NOS enzymes is disrupted, leading to impaired myocardial NO signaling. For instance, myocardial NO has been found to inhibit the positive inotropic responses to β -adrenergic receptor stimulation¹⁷ and further impair the failing heart,¹⁸ thus highlighting the key physiologic and pathophysiologic role NO signaling has in cardiac function.

Molecular Mechanisms: Reactive Oxygen Species and Inflammation

Increased reactive oxygen species (ROS) and inflammation, combined with diminished NO production and bioavailability, are the principal

mechanisms underlying the pathophysiology of endothelial dysfunction.¹⁹ The ROS are highly reactive chemical species that are formed both physiologically as a natural by-product of oxygen metabolism, as well as pathologically during environmental and oxidative stress. The ROS can be generated during normal cellular metabolism, as intermediates during redox reactions, or enzymatically by uncoupled eNOS, xanthine oxidoreductase, mitochondrial enzymes, and nicotinamide adenine dinucleotide phosphate oxidase.²⁰ The uncoupled state of eNOS refers to the dissociation of the ferrous-dioxygen complex, which is when the ROS superoxide anion is generated from the oxygenase domain.²¹ Superoxide anion is the primary ROS in the endothelium, which when generated, signals the contraction of vascular smooth muscle and detrimentally scavenges NO within the vascular wall.²² When superoxide anion reacts with NO by a diffusion-limited reaction, it forms a much more powerful oxidant called peroxynitrite.²³ Most notably, peroxynitrite oxidizes proteins involved in contractility and ion channels, which inhibit mitochondrial energy production. Another important ROS is hydrogen peroxide (H₂O₂), an uncharged, stable, nonpolar ROS that has a longer half-life compared to superoxide and the ability to readily diffuse between cells.²⁴ Physiologically, H₂O₂ is an important intracellular messenger that regulates endothelial cell growth and proliferation, vasorelaxation, vascular remodeling, and inflammatory responses.²⁵ However, excessive H₂O₂ is implicated in the alteration of vascular function, induction of vascular smooth muscle cell inflammation and calcification, and the progression of CVD.²⁶

At low concentrations, ROS are beneficial and physiologic, contributing to host defense mechanisms against infectious agents, regulating cell growth, cell adhesion, cell senescence, and cellular apoptosis.²⁰ However, overproduction of ROS has deleterious effects, causing oxidative stress, DNA and cellular damage, vascular cell damage—notably to endothelial cells and vascular smooth muscle cells, ultimately inhibiting proper lipid and protein function—and contributing to systemic inflammation.²⁷ Elevated levels of ROS are damaging to endothelial function, largely due to disruption of the balance of NO signaling. Oxidants stimulate inflammatory cytokines to further

induce endothelial dysfunction. Inflammatory activation of endothelial cells causes increased expression of selectins, adhesion molecules, and monocyte activation.²⁸ Ultimately, higher levels of ROS cause systemic inflammation, which can in turn lead to endothelial dysfunction and worsening of CVD.²⁹ If not addressed, endothelial dysfunction has severe implications for the heart and vascular system.

ENDOTHELIAL DYSFUNCTION AND HF


Endothelial dysfunction is implicated in HF, and the mechanistic relationship between them remains an important subject of investigation. Heart failure is a complex syndrome that results from structural and/or functional impairment of ventricular filling (diastolic dysfunction) or ejection of blood (systolic dysfunction), known as HF with preserved ejection fraction (HFpEF) and HF with reduced ejection fraction (HFrEF), respectively. Heart failure with preserved ejection fraction, characterized by a left ventricular ejection fraction (LVEF) greater than 50%, is an emerging epidemic and thought to represent up to 50% of patients with HF.³⁰ Importantly, the recommended HF medications (eg, β -blockers, angiotensin-converting enzyme inhibitors, angiotensin II receptor blockers, angiotensin-receptor-neprilysin inhibitors, and aldosterone antagonists) target HFrEF, or LVEF of 40% or less, and have yielded disappointing results in patients with HFpEF. These recommended medications, or guideline-directed medical therapy, improve morbidity and mortality in patients with HFrEF, whereas guidelines recommend diuretics and aldosterone antagonists in selected patients to ameliorate symptoms and control blood pressure in HFpEF pending future clinical trials.^{31,32}

Heart failure with reduced ejection fraction develops as a result of cardiac injury, which can be ischemic from a myocardial infarction or non-ischemic (genetic or acquired), leading to systolic dysfunction and eccentric hypertrophy. The standard of care HF medications are antagonists of the renin-angiotensin-aldosterone and adrenergic systems, which help attenuate direct myocardial injury, cell necrosis, and apoptosis. Blunting the neuroendocrine system allows for reverse remodeling of the left ventricle, leading to improvements in LVEF and clinical outcomes. Because HFrEF is a progressive disease, without

attenuation of the neuroendocrine system and reverse remodeling, systolic function will continue to decline, leading to advanced HF. Heart failure with reduced ejection fraction can promote endothelial dysfunction via neurohormonal activation, altered shear stress, increased oxidative stress, and a decrease in the production of NO.^{2,3} Ultimately, HFrEF fosters an altered redox state in which oxidative stress and inflammatory markers prevail. With this imbalance of NO and oxidative stress (nitroso-redox imbalance),^{33,34} there is a subsequent decrease in coronary endothelium-dependent vasodilator capacity, which impairs myocardial perfusion, reduces coronary flow, and worsens ventricular function.³ These processes culminate in reduced NO bioavailability and worsening endothelial dysfunction, which in turn propagates progression of chronic HFrEF.³

In HFpEF, there is no direct cardiac injury, but instead a multitude of comorbidities that may underlie the etiology (eg, hypertension, obesity, diabetes, chronic kidney disease, chronic obstructive pulmonary disease, anemia, and iron deficiency).³⁵ It has been well documented that patients with HFpEF have impaired endothelial function.³⁶⁻³⁸ In this inflammatory state, there is transformation of endothelial cells into fibroblasts, also known as epithelial-mesenchymal transition.³⁹ The inflammatory state seen with these comorbidities can also cause decreased NO bioavailability leading to decreased cGMP, altered phosphorylation of titin, and microvascular ischemia.³⁵ As previously discussed, NO modulates the activity of calcium by increasing contractile calcium responsiveness and thereby regulates excitation-contraction coupling.¹⁴ With endothelial dysfunction and decreased NO bioavailability, in combination with increased ROS such as peroxynitrite, there are lower levels of intracellular cGMP and protein kinase G activity—kinases that regulate smooth muscle relaxation, platelet function, cell division, and nucleic acid synthesis.²³ The reduction in cGMP and protein kinase G activity results in increased diastolic cytosolic calcium and thus delayed myocardial relaxation.

Nitric oxide imbalance also affects endothelial progenitor cells (EPCs), impairing endothelial repair and regeneration as well as altering matrix metalloproteinases, which affects cell migration, cardiac hypertrophy, and



atherosclerotic plaque stability. Additionally, alterations in matrix metalloproteinases can cause excess production and accumulation of extracellular matrix structural proteins, leading to fibrosis that results in enhanced myocardial stiffness.⁴⁰ Also, endothelial dysfunction allows for transmigration of monocytes into the myocardium, leading to interstitial fibrosis causing diastolic dysfunction.⁴¹ All of these mechanisms can lead to concentric left ventricular hypertrophy, a main contributor to HFpEF. Because current standard of care HF treatments are less effective for HFpEF, evidenced by the results from phase 3 clinical trials—I-PRESERVE (Irbesartan in Patients with Heart Failure and Preserved Ejection Fraction),⁴² PEP-CHF (Perindopril in Elderly People with Chronic Heart Failure),⁴³ TOPCAT (Treatment of Preserved Cardiac Function Heart Failure With an Aldosterone Antagonist),⁴⁴ and CHARM-Preserved (Candesartan in Heart Failure: Assessment of Reduction in Mortality and Morbidity)⁴⁵—novel approaches need to be explored that target endothelial dysfunction in order to ameliorate HFpEF and prevent adverse CVD outcomes.

METHODS FOR MEASURING ENDOTHELIAL FUNCTION

There are numerous methods for measuring endothelial function, most notably intracoronary arterial infusions combined with quantitative coronary angiography (QCA) and Doppler imaging, flow-mediated vasodilation (FMD), peripheral arterial tonometry (PAT), EPC—colony-forming unit (EPC-CFU) assay, and circulating EPCs (Figure 2). Intracoronary ACh infusion was the first method discovered for measuring endothelial function; however, due to its invasive nature, FMD has largely replaced ACh as the most widely used method. PAT is a newly emerging noninvasive method that is currently under investigation as a supplement to FMD. Lastly, EPC measurements are generally considered a less direct method of measuring dysfunction, but studies have shown their important prognostic role in CVD.⁴⁶

Intracoronary Artery Infusions Using ACh and Other Vasoactive Substances

In 1986, QCA combined with ACh infusions was the first method utilized to detect endothelial

function and is considered the criterion standard for early detection of dysfunction.⁴⁷ Intracoronary arterial infusion is the most direct, yet most invasive and expensive, method of measuring endothelial function. During QCA, a contrast agent is injected into the coronary artery and vessel diameter is measured from multiple planes of view. After obtaining baseline measurements, either ACh or other vasoactive substances (ie, substance P, bradykinin, serotonin) are infused into the artery—typically as multiple doses infused for 1 to 2 minutes—and coronary blood flow is continuously measured using Doppler techniques. Instead of QCA, intravascular ultrasound (IVUS) can also be employed along with ACh or other vasoactive infusions to measure coronary blood flow velocity using a Doppler-guided wire to obtain vessel measurements. Regardless of QCA or IVUS, endothelial function is then calculated by determining volumetric coronary blood flow and evaluating it on a dose-response curve.⁴⁸ In a healthy endothelium, ACh causes the coronary arteries to dilate. However, in a patient with endothelial dysfunction, ACh instead causes a paradoxical vasoconstriction of the arteries.⁴⁷

Quantitative coronary angiography and IVUS are both widely validated techniques that have documented utility in studying endothelial function in patients with CVD and HF.⁴⁹⁻⁵¹ Schächinger et al⁴⁹ found that coronary endothelial vasodilation dysfunction predicted long-term (median 7.7-year follow-up in 147 patients) atherosclerotic disease progression and risk of cardiovascular events. Reriani et al⁵² recently reported that coronary endothelial function can be an invaluable tool to reclassify risk stratification of patients with early CAD, highlighting its superior discrimination over standard methods such as Framingham Risk Score. However, due to the invasiveness of the method combined with the high risk associated with coronary catheterization, intracoronary studies are not feasible for screening large populations. Consequently, alternatives to this invasive method emerged in the 1990s, and numerous studies since then have affirmed similar results between using coronary artery response and brachial FMD.⁵³

Flow-Mediated Vasodilation

In 1992, FMD was first introduced as a noninvasive method of measuring endothelial

dysfunction and has gained popularity as the most robust and widely validated technique.⁵⁴ To measure FMD, the patient is positioned supine and a blood pressure cuff is placed above the antecubital fossa. After a baseline image of the brachial radial artery is obtained, the cuff is inflated to approximately 50 mm Hg above systolic pressure in order to occlude arterial inflow for a short period. Subsequently, the cuff is deflated, which creates a potent shear stress enabling the induced reactive hyperemia to dilate the brachial artery. This shear stress is mediated by NO released from endothelial cells along with other vasoactive mediators. The change in diameter of the artery is continuously recorded for up to 2 minutes after the cuff deflation via ultrasound, and the FMD is reported as a percent change in poststimulus diameter compared to the baseline diameter.

Flow-mediated vasodilation has gained popularity in clinical trials because of its ease of use, efficiency, and noninvasive nature. However, various factors including food intake, coffee intake, ambient temperature, drugs, vitamins, exercise, tobacco use, and menstrual cycle can affect a patient's FMD, potentially confounding study results. Therefore, it is essential that patients fast before the assessment and abstain from taking medicines, vitamins, exercising, and using tobacco the day of the test. Furthermore, it is important that the study room conditions are held constant throughout the trial, meaning a consistent, comfortable room temperature, same lighting, and minimal environmental stress. Despite these potential limitations, Ghiadoni et al⁵⁵ performed a nationwide, multicenter study and found that standardized and rigorous methodology among centers resulted in accurate and homogeneous results, highlighting the reproducibility of FMD assessments.

Abundant cardiovascular studies using FMD have generated insight into the role endothelial dysfunction plays in CVD and HF. Flow-mediated vasodilation has been reported to be significantly impaired in patients with atherosclerosis, hypertension, dilated cardiomyopathy, ischemic cardiomyopathy, CAD, peripheral artery disease, and congestive HF.^{56,57} Furthermore, studies utilizing FMD have found that it serves as an important prognostic tool for CVD diagnosis.^{57,58} In a cohort of 3026 individuals, Yeboah et al⁵⁸ found that FMD was significantly

diminished and inversely associated with incident cardiovascular events, enabling FMD to be used as a means to classify patients into low-, intermediate-, and high-risk categories.

Peripheral Arterial Tone

With the increasing recognition of the role endothelial dysfunction plays in HF and the powerful predictive role proper assessment plays in discerning cardiovascular events, there is a necessity for novel noninvasive techniques. This need led the way for the emergence of reactive hyperemia-PAT technology. In the early 2000s, the idea of measuring peripheral microvascular endothelial dysfunction surfaced, and EndoPAT—a device developed by Itamar Medical Ltd—was introduced as a more rapid, cheaper, and easier method for assessing endothelial function, relying on peripheral arterial circulation in the finger. Similar to brachial FMD, hyperemia is induced by occluding blood flow through the brachial artery utilizing a cuff for 5 minutes and measuring the arterial tone changes in peripheral arterial beds generally in the index finger. On release of the cuff, reactive hyperemia is automatically calculated by the device as a postocclusion to preocclusion signal ratio, therefore making it non-operator dependent. In addition, a baseline measurement is obtained in the other arm, serving as an internal control.

One of the major drawbacks of PAT currently is that the physiology remains unclear. PAT measures microvessel dilatation, as opposed to FMD that measures macrovascular dilatation, and is thus only partly dependent on NO. Notably, inhibiting NO and subsequently measuring endothelial function via PAT attenuates the response by less than 50%, whereas using FMD nearly completely inhibits the response.⁵⁹ Because PAT is a newer technology and the goal of both technologies is to measure endothelial function, many studies have tried to prove PAT reliability by validating the correlation between FMD and PAT.⁶⁰ However, studies have yielded mixed results in this comparison. For example, both Hamburg et al⁶¹ and Lee et al⁶² found that there was no correlation between PAT and FMD, questioning the reliability of PAT. Conversely, Kuvin et al⁶³ found a significant correlation between reactive hyperemia measured by FMD and PAT. Irrespective of the variable results comparing

FMD and PAT, PAT has been reported to be a suitable diagnostic tool for patients with HF.^{64,65}

EPC and Hematopoietic and Endothelial Lineage Assessment Methodologies

Endothelial progenitor cells are vital to proper endothelial function, playing a pivotal role in maintaining vascular homeostasis and mediating vascular repair in damaged endothelium by stimulating the release of NO from the endothelium as well as by incorporating into old and damaged endothelium. Endothelial progenitor cells can be measured via flow cytometry as well as by cell culture techniques. When measuring EPCs via cell culture methodologies, there are 2 different populations of EPCs, early and late outgrowth. Recently, there has been a large debate in the literature regarding proper nomenclature, lineage, and function. Early outgrowth EPCs, identified by the classic EPC-CFU method of Hill et al,⁶⁶ are now being considered myeloid angiogenic cells and circulating angiogenic cells, which have a strong paracrine ability that stimulates angiogenesis and enhances endothelial tube network formation.⁶⁷ Conversely, late outgrowth EPCs are now identified as endothelial colony-forming cells (ECFCs), outgrowth endothelial cells, and blood outgrowth endothelial cells, which have an intrinsic tube-forming capacity and actively participate and incorporate into new blood vessel formation and vascular repair.⁶⁷ Both early and late outgrowth EPCs have been found to be prognostic of endothelial dysfunction and furthermore correlate with CVD and HF.^{68,69} Accordingly, assays for both early and late EPCs can be used to diagnose endothelial dysfunction.

To measure early outgrowth EPC-CFUs, a blood sample is obtained from the patient and the mononuclear cell layer is isolated on a Ficoll-Paque (GE Healthcare Life Sciences) density gradient. After multiple washes, the cells are plated on fibronectin-coated dishes. Two days later, these cells are replated onto smaller dishes, and on day 5, EPC-CFUs are counted. This technique is straightforward, minimally invasive, and cost effective. However, it is heavily operator dependent—and therefore colony reads can vary across investigator—and furthermore, the link between EPC-CFUs and

endothelial function is less clear than with FMD. However, low EPC-CFUs have been found to be hallmarks of endothelial dysfunction, and numerous studies have reported that EPC-CFUs are inversely associated with cardiovascular events and disease progression.^{46,56} Hill et al⁴⁶ conducted a landmark study in which they documented that EPC-CFUs were a better predictor of endothelial function than conventional risk factors, and furthermore, EPC-CFUs are a potential biomarker for vascular function and increasing cardiovascular risk. Along this line, our laboratory specifically documented that patients with dilated cardiomyopathy and endothelial dysfunction had severely impaired EPC-CFUs, and allogeneic mesenchymal stem cell treatment restored these numbers.⁵⁶

Late outgrowth EPCs, or ECFCs, are also obtained from the blood by isolating the mononuclear cell layer on a Ficoll-Paque density gradient. However, these cells are plated on collagen using endothelial growth medium and can be cultured for weeks as well as passaged for expansion. Colonies are assessed anywhere from 5 to 26 days after being plated. Like EPC-CFUs, ECFCs have also been found to be prognostic of CVD⁷⁰ and mortality in HF.⁷¹ Additionally, this technique is also minimally invasive and cost effective, and cells can be expanded and utilized for further studies.

Alternatively, circulating EPCs can be enumerated using flow cytometry, specifically staining for CD133, CD34, and KDR. However, this method has produced mixed results, potentially because high circulating EPC number does not ascertain that these cells are functionally contributing to endothelial function. Endothelial progenitor cells are a heterogeneous population, and accordingly, measuring EPC-CFUs and ECFCs appears to assess endothelial dysfunction more accurately.^{66,72}

CLINICAL TRIALS AND ENDOTHELIAL FUNCTION

Case-control studies have clearly documented a link between endothelial dysfunction and the progression of CVD and HF, highlighting how endothelial dysfunction can be a vital prognostic tool for evaluating CVD stages.^{49,54} These correlations have contributed to the hypothesis that targeting endothelial dysfunction could lead to the amelioration of CVD.

In recent clinical trials, results have suggested that improving endothelial function in patients with HF may concordantly improve cardiac biomechanics.^{5,73} Recently, we found that allogeneic mesenchymal stem cell treatment significantly improved endothelial function in patients with HF due to nonischemic dilated cardiomyopathy.^{5,56} Mesenchymal stem cell treatment also resulted in increased ejection fraction, 6-minute walk distance test results, and Minnesota Living with Heart Failure questionnaire scores, along with tumor necrosis factor α reduction.^{5,55} Our study utilized FMD and EPC-CFU assays to assess endothelial function.

Exercise has been reported to increase blood flow and shear stress, therefore increasing endothelial NOS activity and NO production and reducing inflammation, which are direct components of endothelial function,⁷⁴ making it an appealing candidate for improving endothelial function. A growing number of clinical trials are investigating the effect of exercise on endothelial function in patients with HF and HFpEF.⁷⁵ In a 458 patient meta-analysis, Pearson and Smart⁷⁶ revealed that aerobic exercise significantly improves FMD in patients with HF, suggesting that improving endothelial function via cardiac rehabilitation may be an important therapy for patients with HF. Indeed, Tanaka et al⁷⁷ recently reported that cardiac rehabilitation improved prognosis in patients with HF, although with a greater effect on patients with HF_rEF than HF_pEF, and this improvement was attributed to the impact on improving endothelial function. Ozasa et al⁷⁸ utilized EndoPAT to demonstrate that machine-assisted cycling and conventional exercise training improved endothelial function and 6-minute walk distance in 27 patients with HF. Erbs et al⁷⁹ found that bicycle training for 12 weeks in patients with advanced chronic HF improved FMD and maximum oxygen consumption and notably increased ejection fraction by approximately 10%. Sandri et al⁸⁰ assessed EPC numbers and function and found that 4 weeks of exercise was efficacious in improving endothelial function in 60 patients with chronic HF. Conversely, Angadi et al⁸¹ investigated the effect of high-intensity interval training vs moderate-intensity aerobic continuous training in patients with HFpEF and

found that although peak oxygen consumption improved, FMD did not. Kitzman et al⁸² evaluated the effect of endurance training on FMD and carotid artery stiffness in 63 patients with HFpEF and found results similar to those of Dall et al,⁸³ with improvement in peak oxygen consumption but no change in either endothelial function via FMD or arterial stiffness. Nevertheless, current guidelines strongly recommend appropriate amounts of exercise for primary and secondary prevention of CVD and HF.^{84,85}

Two statin clinical trials yielded encouraging results, reporting that statins improve endothelial function and hemodynamics in patients with HF, CAD, and atherosclerosis.⁸⁶⁻⁸⁸ Antoniadis et al⁸⁸ found that statins improve endothelial function, assessed by FMD, in patients with CAD. Additionally, they reported a potential mechanism for this improvement, namely, an increase in NO bioavailability caused by a reduction in uncoupled endothelial NOS-derived superoxide and an increase in vascular BH4 bioavailability via the up-regulation of guanosine triphosphate cyclohydrolase 1. The same group later demonstrated that 4 weeks of atorvastatin treatment significantly improved endothelial function (measured by FMD and endothelial-independent vasodilation) and EPC mobilization and reduced tumor necrosis factor α in patients with ischemic HF.⁸⁶ Erbs et al⁸⁷ found that high-dose rosuvastatin in 42 patients with HF resulted in a dramatic improvement in FMD, LVEF, vascular endothelial growth factor, circulating EPCs, and capillary density and that these effects were accompanied by decreased oxidized low-density lipoprotein.

Blood pressure medications have also been explored, yielding mixed results. Specifically, nebivolol, a β_1 -selective receptor blocker with NO-potentiating effects, demonstrated efficacy in improving endothelial function.^{89,90} Conversely, metoprolol, a β_1 -selective receptor blocker, and spironolactone, an aldosterone antagonist, had minimal to no effects on endothelial function.^{91,92}

There are abundant ongoing clinical trials assessing endothelial function in patients with CVD. Albumin, nutraceutical therapies, new oral anticoagulants, sildenafil, beetroot juice, tetrahydrobiopterin (BH4), sitagliptin, bioresorbable stents, remote ischemic preconditioning, and yoga are interventions currently under investigation for improving endothelial

function. More specifically, the Endothelial Function After Cardiac Surgery study ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02882074) Identifier: NCT02882074) is evaluating whether administration of albumin during cardiac surgery improves postoperative endothelial function compared to hydroxyethyl starch solutions using EndoPAT. In the NUTRENDO (Effects of Nutraceutical Therapies on Endothelial Function, Platelet Aggregation, and Coronary Flow Reserve) trial ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02969070) Identifier: NCT02969070), Trimarco et al are investigating the effects of nutraceutical therapies on endothelial function, platelet aggregation, and coronary flow reserve in patients with hypercholesterolemia, specifically measuring reactive hyperemia index. Kim et al are testing the effect of new oral anticoagulants (dabigatran, rivaroxaban, warfarin) for improving endothelial dysfunction—assessed via EndoPAT—and therefore preventing the progression of atherosclerosis in patients with atrial fibrillation ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02544932) Identifier: NCT02544932). Liang et al are recruiting patients with rheumatoid arthritis to see if sildenafil improves endothelial function, measured by FMD and biomarkers, and is thereby an effective strategy for CVD prevention in patients with rheumatoid arthritis—a patient population associated with a 2-fold risk of CVD ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02908490) Identifier: NCT02908490). The NITRATE-OCT (Investigating the Effects of Dietary Nitrate on Vascular Function, Platelet Reactivity and Restenosis in Stable Angina; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02529189) Identifier: NCT02529189) trial is investigating whether beetroot juice reduces angina symptoms as well as long-term risk of myocardial infarction in patients with CVD, utilizing FMD as an outcome measurement. Alongside this trial, the NITRATE-TOD (Investigation of Dietary Nitrate Effects in Hypertension-induced Target Organ Damage; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03088514) Identifier: NCT03088514) study is assessing beetroot juice in patients with hypertension. In the COPD-LT (Long Term Nitric Oxide Bioavailability on Vascular Health in Chronic Obstructive Pulmonary Disease; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02774226) Identifier: NCT02774226) trial, patients with chronic obstructive pulmonary disease are being treated with BH4 to see if endothelial dysfunction is improved (measured as an increase in FMD and change in biomarkers of oxidative stress)

and therefore future atherosclerotic CVD and events. The SAVORO (Sitagliptin Effects on Arterial Vasculature and Inflammation in Obesity; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02576288) Identifier: NCT02576288) trial is administering sitagliptin in patients with atherosclerosis and inflammation to see if FMD and carotid stiffness improve, hypothesizing that treatment will halt atherosclerosis progression to serious CVD events. Testing the efficacy of bioresorbable and metallic stents in patients treated for stable angina, Gomez-Lara et al ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02738658) Identifier: NCT02738658) are using endothelial dysfunction measured by Ach infusions and QCA as a secondary outcome measurement. The PIXIE (Prevention of Myocardial Injury in Non-cardiac Surgery; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02344797) Identifier: NCT02344797) trial is studying whether remote ischemic preconditioning prevents myocardial injury and infarction, specifically assessing endothelial dysfunction via EndoPAT as an outcome measurement. Hunter et al are exploring the effect of Bikram yoga on arterial stiffness and endothelial dysfunction as measured by FMD ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02488148) Identifier: NCT02488148), hypothesizing that yoga will improve vascular endothelial dysfunction and therefore reduce risk factors for CAD.

In addition to interventional testing, numerous clinical trials are testing the prognostic value of measuring endothelial function in patients with CVD. Bairey Merz et al ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT00573027) Identifier: NCT00573027) are studying microvascular disease in HF and cardiac syndrome X, assessing the predictive value of the noninvasive PAT, and furthermore elucidating risk factors and treatment using PAT as a primary outcome and coronary flow measurements with Ach as a secondary outcome. Thomas et al ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02636062) Identifier: NCT02636062) are aiming to define the rate of index atherosclerotic plaque developments and identify new markers of CAD progression in patients with CAD and endothelial dysfunction utilizing EndoPAT combined with blood biomarker assessments. O'Neal et al hypothesized that the pathophysiology of delirium after cardiac surgery is due to the stress caused by systemic inflammation causing endothelial dysfunction, making change in endothelial function their primary end point ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02688179) Identifier:

NCT02688179). The Mainz Registry of Flow-Mediated Constriction ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01787370) Identifier: NCT01787370) is investigating the prognostic value of FMD in patients undergoing coronary angiography. The POETRYabd (Perioperative Endothelial Dysfunction in Patients Undergoing Major Acute Abdominal Surgery; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03010969) Identifier: NCT03010969) trial is examining the correlation between perioperative endothelial dysfunction—measured by EndoPAT—and major adverse cardiovascular events as well as noncardiovascular complications. Tibirica et al are evaluating the predictive efficacy of measuring systemic microvascular endothelial dysfunction via microvascular reactivity in patients with infective endocarditis ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT02940340) Identifier: NCT02940340).

In summary, a growing number of clinical trials support the notion that drugs targeting endothelial dysfunction in patients with HF or CVD may be a feasible therapeutic option. However, it is notable that many of these interventions have only been tested in one clinical trial with low enrollment numbers. Currently, there are ongoing clinical trials assessing not only various interventions for treating endothelial dysfunction in patients with CVD but also assessing the prognostic value of endothelial dysfunction for CVD progression.

CONCLUSION

Endothelial dysfunction has reemerged as a disease process or biomarker warranting further clinical investigation. As reviewed herein, the pathophysiology of endothelial function plays a significant role in the development and progression of HF and CVD. Methods for studying endothelial function clinically have improved, been validated, and correlate with CVD outcomes, but more effective therapies for targeting endothelial dysfunction in order to ameliorate HF and CVD outcomes are needed. As endothelial function is recognized as a crucial component underlying HF and CVD, we propose that targeting endothelial dysfunction will provide, at least in part, the breakthrough therapy needed.

ACKNOWLEDGMENTS

The funding organizations had no role in the design and execution of the study, in the

collection, analyses, and interpretation of the data, or in the preparation, review, or approval of the submitted manuscript.

Drs Premer and Kanelidis contributed equally to this work.

Abbreviations and Acronyms: Ach = acetylcholine; CAD = coronary artery disease; cGMP = cyclic guanosine monophosphate; CVD = cardiovascular disease; ECFC = endothelial colony-forming cell; EDHF = endothelium-derived hyperpolarizing factor; eNOS = endothelial nitric oxide synthase; EPC = endothelial progenitor cell; EPC-CFU = EPC—colony-forming unit; FMD = flow-mediated vasodilation; HF = heart failure; HFpEF = HF with preserved ejection fraction; HFREF = HF with reduced ejection fraction; H₂O₂ = hydrogen peroxide; IVUS = intravascular ultrasound; LVEF = left ventricular ejection fraction; NO = nitric oxide; NOS = NO synthase; PAT = peripheral arterial tonometry; QCA = quantitative coronary angiography; ROS = reactive oxygen species

Grant Support: This work was supported in part by grants R01 HL R0110737 (C.P., J.M.H.), IR01HL134558-01 (J.M.H., I.H.S.), 5UM1HL113460 (J.M.H., I.H.S.), R01HL137355 (J.M.H., I.H.S.), and R01HL107110 (J.M.H.) from the National Institutes of Health. Dr Premer was also funded by American Heart Association Predoctoral Fellowship 15PRE25730025. Dr Hare is also funded by the Starr, Marcus, and Soffer Family Foundations.

Potential Competing Interests: Drs Premer and Hare report having a materials and methods patent for endothelial dysfunction. Dr Hare reports having a patent for cardiac cell-based therapy; holds equity in and maintains a professional relationship with Vestion, Inc. as a consultant and member of the Board of Directors and Scientific Advisory Board; is the chief scientific officer, a compensated consultant, and advisory board member for Longeveron LLC, holds equity in Longeveron, and is the coinventor of intellectual property licensed to Longeveron. The other authors report no competing interests.

Correspondence: Address to Ivonne H. Schulman, MD, Interdisciplinary Stem Cell Institute, Katz Family Division of Nephrology and Hypertension, University of Miami Miller School of Medicine, Biomedical Research Building, 1501 NW 10th Ave, Room 811, Miami, FL 33101 (ischulman@miami.edu).

REFERENCES

1. Benjamin EJ, Virani SS, Callaway CVW, et al: American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2018 update: a report from the American Heart Association [published correction appears in *Circulation*. 2018;137(12):e493]. *Circulation*. 2018;137(12):e67-e492.
2. Tobar SA, Umpierre D, Andrades M, Clausell N. Endothelial alterations in heart failure—mechanisms and molecular basis. In: da Luz PL, Libby P, Chagas ACP, Laurindo FRM, eds. *Endothelium and Cardiovascular Diseases*. Cambridge, MA: Academic Press; 2018:565-573.

3. Marti CN, Gheorghide M, Kalogeropoulos AP, Georgiopoulou VV, Quyyumi AA, Butler J. Endothelial dysfunction, arterial stiffness, and heart failure. *J Am Coll Cardiol*. 2012;60(16):1455-1469.
4. Jedlickova L, Merkovska L, Jackova L, et al. Effect of ivabradine on endothelial function in patients with stable angina pectoris: assessment with the Endo-PAT 2000 device. *Adv Ther*. 2015;32(10):962-970.
5. Hare JM, DiFede DL, Rieger AC, et al. Randomized comparison of allogeneic versus autologous mesenchymal stem cells for nonischemic dilated cardiomyopathy: POSEIDON-DCM Trial. *J Am Coll Cardiol*. 2017;69(5):526-537.
6. Jia G, Durante W, Sowers JR. Endothelium-derived hyperpolarizing factors: a potential therapeutic target for vascular dysfunction in obesity and insulin resistance. *Diabetes*. 2016;65(8):2118-2120.
7. Ozkor MA, Hayek SS, Rahman AM, et al. Contribution of endothelium-derived hyperpolarizing factor to exercise-induced vasodilation in health and hypercholesterolemia. *Vasc Med*. 2015;20(1):14-22.
8. Dunn SM, Hilgers RHP, Das KC. Decreased EDHF-mediated relaxation is a major mechanism in endothelial dysfunction in resistance arteries in aged mice on prolonged high-fat sucrose diet. *Physiol Rep*. 2017;5(23):e13502.
9. Schulman IH, Hare JM. Regulation of cardiovascular cellular processes by S-nitrosylation. *Biochim Biophys Acta*. 2012;1820(6):752-762.
10. Tousoulis D, Kampoli AM, Tentolouris C, Papageorgiou N, Stefanadis C. The role of nitric oxide on endothelial function. *Curr Vasc Pharmacol*. 2012;10(1):4-18.
11. Yamagishi S, Nakamura K, Matsui T, Ueda S, Noda Y, Imaizumi T. Inhibitors of advanced glycation end products (AGEs): potential utility for the treatment of cardiovascular disease. *Cardiovasc Drug Rev*. 2008;26(1):50-58.
12. Dulce RA, Kulandavelu S, Hernandez Schulman I, Fritsch J, Hare JM. Nitric oxide regulation of cardiovascular physiology and pathophysiology. In: Ignarro LJ, Freeman BA, eds. *Nitric Oxide: Biology and Pathobiology*. 3rd ed. Cambridge, MA: Academic Press; 2017:313-338.
13. Barouch LA, Harrison RW, Skaf MW, et al. Nitric oxide regulates the heart by spatial confinement of nitric oxide synthase isoforms. *Nature*. 2002;416(6878):337-339.
14. Hare JM. Nitric oxide and excitation-contraction coupling. *J Mol Cell Cardiol*. 2003;35(7):719-729.
15. Gonzalez DR, Treuer AV, Castellanos J, Dulce RA, Hare JM. Impaired S-nitrosylation of the ryanodine receptor caused by xanthine oxidase activity contributes to calcium leak in heart failure. *J Biol Chem*. 2010;285(37):28938-28945.
16. Beigi F, Gonzalez DR, Minhas KM, et al. Dynamic denitrosylation via S-nitrosoglutathione reductase regulates cardiovascular function. *Proc Natl Acad Sci U S A*. 2012;109(11):4314-4319.
17. Keaney JF Jr, Hare JM, Balligand JL, Loscalzo J, Smith TW, Colucci WS. Inhibition of nitric oxide synthase augments myocardial contractile responses to beta-adrenergic stimulation. *Am J Physiol*. 1996;271(6, pt 2):H2646-H2652.
18. Hare JM, Loh E, Creager MA, Colucci WS. Nitric oxide inhibits the positive inotropic response to beta-adrenergic stimulation in humans with left ventricular dysfunction. *Circulation*. 1995;92(8):2198-2203.
19. Cai H, Harrison DG. Endothelial dysfunction in cardiovascular diseases: the role of oxidant stress. *Circ Res*. 2000;87(10):840-844.
20. Montezano AC, Touyz RM. Reactive oxygen species and endothelial function—role of nitric oxide synthase uncoupling and Nox family nicotinamide adenine dinucleotide phosphate oxidases. *Basic Clin Pharmacol Toxicol*. 2012;110(1):87-94.
21. Yokoyama M, Hirata K. Endothelial nitric oxide synthase uncoupling: is it a physiological mechanism of endothelium-dependent relaxation in cerebral artery [editorial]? *Cardiovasc Res*. 2007;73(1):8-9.

22. Bharadwaj LA, Prasad K. Mechanism of superoxide anion-induced modulation of vascular tone. *Int J Angiol.* 2002;11(1):23-29.
23. Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev.* 2007;87(1):315-424.
24. Fridovich I. Superoxide anion radical (O_2^-), superoxide dismutases, and related matters. *J Biol Chem.* 1997;272(30):18515-18517.
25. Bretón-Romero R, Lamas S. Hydrogen peroxide signaling in vascular endothelial cells. *Redox Biol.* 2014;2:529-534.
26. Byon CH, Heath JM, Chen Y. Redox signaling in cardiovascular pathophysiology: a focus on hydrogen peroxide and vascular smooth muscle cells. *Redox Biol.* 2016;9:244-253.
27. Chen Q, Wang Q, Zhu J, Xiao Q, Zhang L. Reactive oxygen species: key regulators in vascular health and diseases. *Br J Pharmacol.* 2018;175(8):1279-1292.
28. Dröge W. Free radicals in the physiological control of cell function. *Physiol Rev.* 2002;82(1):47-95.
29. Cervantes Gracia K, Llanas-Cornejo D, Husi H. CVD and oxidative stress. *J Clin Med.* 2017;6(2):E22.
30. Oktay AA, Rich JD, Shah SJ. The emerging epidemic of heart failure with preserved ejection fraction. *Curr Heart Fail Rep.* 2013;10(4):401-410.
31. Lewis EF, Claggett B, Shah AM, et al. Racial differences in characteristics and outcomes of patients with heart failure and preserved ejection fraction in the Treatment of Preserved Cardiac Function Heart Failure trial. *Circ Heart Fail.* 2018;11(3):e004457.
32. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA focused update of the 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Am Coll Cardiol.* 2017;70(6):776-803.
33. Hare JM. Nitroso-redox balance in the cardiovascular system [editorial]. *N Engl J Med.* 2004;351(20):2112-2114.
34. Karantalis V, Schulman IH, Hare JM. Nitroso-redox imbalance affects cardiac structure and function [editorial]. *J Am Coll Cardiol.* 2013;61(9):933-935.
35. Lam CS, Lund LH. Microvascular endothelial dysfunction in heart failure with preserved ejection fraction [editorial]. *Heart.* 2016;102(4):257-259.
36. Maréchaux S, Samson R, van Belle E, et al. Vascular and microvascular endothelial function in heart failure with preserved ejection fraction. *J Card Fail.* 2016;22(1):3-11.
37. Borlaug BA, Olson TP, Lam CS, et al. Global cardiovascular reserve dysfunction in heart failure with preserved ejection fraction. *J Am Coll Cardiol.* 2010;56(11):845-854.
38. Akiyama E, Sugiyama S, Matsuzawa Y, et al. Incremental prognostic significance of peripheral endothelial dysfunction in patients with heart failure with normal left ventricular ejection fraction. *J Am Coll Cardiol.* 2012;60(19):1778-1786.
39. Zeisberg EM, Tamavski O, Zeisberg M, et al. Endothelial-to-mesenchymal transition contributes to cardiac fibrosis. *Nat Med.* 2007;13(8):952-961.
40. Fan D, Takawale A, Lee J, Kassiri Z. Cardiac fibroblasts, fibrosis and extracellular matrix remodeling in heart disease. *Fibrogenesis Tissue Repair.* 2012;5(1):15.
41. Glezeva N, Baugh JA. Role of inflammation in the pathogenesis of heart failure with preserved ejection fraction and its potential as a therapeutic target. *Heart Fail Rev.* 2014;19(5):681-694.
42. Massie BM, Carson PE, McMurray JJ, et al. I-PRESERVE Investigators. Irbesartan in patients with heart failure and preserved ejection fraction. *N Engl J Med.* 2008;359(23):2456-2467.
43. Cleland JG, Tendera M, Adamus J, Freemantle N, Polonski L, Taylor J; PEP-CHF Investigators. The Perindopril in Elderly People with Chronic Heart Failure (PEP-CHF) study. *Eur Heart J.* 2006;27(19):2338-2345.
44. Pitt B, Pfeffer MA, Assmann SF, et al; TOPCAT Investigators. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med.* 2014;370(15):1383-1392.
45. Yusuf S, Pfeffer MA, Swedberg K, et al; CHARM Investigators and Committees. Effects of candesartan in patients with chronic heart failure and preserved left-ventricular ejection fraction: the CHARM-Preserved Trial. *Lancet.* 2003;362(9386):777-781.
46. Hill JM, Finkel T, Quyyumi AA. Endothelial progenitor cells and endothelial dysfunction. *Vox Sang.* 2004;87(suppl 2):31-37.
47. Ludmer PL, Selwyn AP, Shook TL, et al. Paradoxical vasoconstriction induced by acetylcholine in atherosclerotic coronary arteries. *N Engl J Med.* 1986;315(17):1046-1051.
48. Doucette JW, Corl PD, Payne HM, et al. Validation of a Doppler guide wire for intravascular measurement of coronary artery flow velocity. *Circulation.* 1992;85(5):1899-1911.
49. Schächinger V, Britten MB, Zeiher AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation.* 2000;101(16):1899-1906.
50. Hollenberg SM, Tamburro P, Johnson MR, et al. Simultaneous intracoronary ultrasound and Doppler flow studies distinguish flow-mediated from receptor-mediated endothelial responses. *Catheter Cardiovasc Interv.* 1999;46(3):282-288.
51. Carrick D, Haig C, Rauhalmi S, et al. Prognostic significance of infarct core pathology revealed by quantitative non-contrast in comparison with contrast cardiac magnetic resonance imaging in reperfused ST-elevation myocardial infarction survivors. *Eur Heart J.* 2016;37(13):1044-1059.
52. Reniani M, Sara JD, Flammer AJ, et al. Coronary endothelial function testing provides superior discrimination compared with standard clinical risk scoring in prediction of cardiovascular events. *Coron Artery Dis.* 2016;27(3):213-220.
53. Anderson TJ, Uehata A, Gerhard MD, et al. Close relation of endothelial function in the human coronary and peripheral circulations. *J Am Coll Cardiol.* 1995;26(5):1235-1241.
54. Celermajer DS, Sorensen KE, Gooch VM, et al. Non-invasive detection of endothelial dysfunction in children and adults at risk of atherosclerosis. *Lancet.* 1992;340(8828):1111-1115.
55. Ghiadoni L, Fata F, Salvetti M, et al. Assessment of flow-mediated dilation reproducibility: a nationwide multicenter study. *J Hypertens.* 2012;30(7):1399-1405.
56. Premer C, Blum A, Bellio MA, et al. Allogeneic mesenchymal stem cells restore endothelial function in heart failure by stimulating endothelial progenitor cells. *EBioMedicine.* 2015;2(5):467-475.
57. Gokce N, Keaney JF Jr, Hunter LM, et al. Predictive value of noninvasively determined endothelial dysfunction for long-term cardiovascular events in patients with peripheral vascular disease. *J Am Coll Cardiol.* 2003;41(10):1769-1775.
58. Yeboah J, Folsom AR, Burke GL, et al. Predictive value of brachial flow-mediated dilation for incident cardiovascular events in a population-based study: the Multi-Ethnic Study of Atherosclerosis. *Circulation.* 2009;120(6):502-509.
59. Bruno RM, Gori T, Ghiadoni L. Endothelial function testing and cardiovascular disease: focus on peripheral arterial tonometry. *Vasc Health Risk Manag.* 2014;10:577-584.
60. Allan RB, Vun SV, Spark JI. A comparison of measures of endothelial function in patients with peripheral arterial disease and age and gender matched controls. *Int J Vasc Med.* 2016;2016:2969740.
61. Hamburg NM, Palmisano J, Larson MG, et al. Relation of brachial and digital measures of vascular function in the community: the Framingham Heart Study. *Hypertension.* 2011;57(3):390-396.
62. Lee CR, Bass A, Ellis K, et al. Relation between digital peripheral arterial tonometry and brachial artery ultrasound measures of vascular function in patients with coronary artery disease and in healthy volunteers. *Am J Cardiol.* 2012;109(5):651-657.
63. Kuvin JT, Patel AR, Sliney KA, et al. Assessment of peripheral vascular endothelial function with finger arterial pulse wave amplitude. *Am Heart J.* 2003;146(1):168-174.
64. Weisrock F, Fritschka M, Beckmann S, et al. Reliability of peripheral arterial tonometry in patients with heart failure, diabetic

- nephropathy and arterial hypertension. *Vasc Med*. 2017;22(4):292-300.
65. Fujisue K, Sugiyama S, Matsuzawa Y, et al. Prognostic significance of peripheral microvascular endothelial dysfunction in heart failure with reduced left ventricular ejection fraction. *Circ J*. 2015;79(12):2623-2631.
 66. Hill JM, Zalos G, Halcox JP, et al. Circulating endothelial progenitor cells, vascular function, and cardiovascular risk. *N Engl J Med*. 2003;348(7):593-600.
 67. Medina RJ, Barber CL, Sabatier F, et al. Endothelial progenitors: a consensus statement on nomenclature. *Stem Cells Transl Med*. 2017;6(5):1316-1320.
 68. Lee PS, Poh KK. Endothelial progenitor cells in cardiovascular diseases. *World J Stem Cells*. 2014;6(3):355-366.
 69. Samman Tahhan A, Hammadah M, Sandesara PB, et al. Progenitor cells and clinical outcomes in patients with heart failure. *Circ Heart Fail*. 2017;10(8):e004106.
 70. Padfield GJ, Newby DE, Mills NL. Understanding the role of endothelial progenitor cells in percutaneous coronary intervention. *J Am Coll Cardiol*. 2010;55(15):1553-1565.
 71. Michowitz Y, Goldstein E, Wexler D, Sheps D, Keren G, George J. Circulating endothelial progenitor cells and clinical outcome in patients with congestive heart failure. *Heart*. 2007;93(9):1046-1050.
 72. George J, Shmilovich H, Deutsch V, Miller H, Keren G, Roth A. Comparative analysis of methods for assessment of circulating endothelial progenitor cells. *Tissue Eng*. 2006;12(2):331-335.
 73. Legallois D, Belin A, Nesterov SV, et al. Cardiac rehabilitation improves coronary endothelial function in patients with heart failure due to dilated cardiomyopathy: a positron emission tomography study. *Eur J Prev Cardiol*. 2014;23(2):129-136.
 74. Nosarev AV, Smagliy LV, Anfinogenova Y, Popov SV, Kapilevich LV. Exercise and NO production: relevance and implications in the cardiopulmonary system. *Front Cell Dev Biol*. 2015;2:73.
 75. Gevaert AB, Lemmens K, Vrints CJ, Van Craenenbroeck EM. Targeting endothelial function to treat heart failure with preserved ejection fraction: the promise of exercise training. *Oxid Med Cell Longev*. 2017;2017:4865756.
 76. Pearson MJ, Smart NA. Aerobic training intensity for improved endothelial function in heart failure patients: a systematic review and meta-analysis. *Cardiol Res Pract*. 2017;2017:2450202.
 77. Tanaka S, Sanuki Y, Ozumi K, Harada T, Tasaki H. Heart failure with preserved vs reduced ejection fraction following cardiac rehabilitation: impact of endothelial function. *Heart Vessels*. 2018;33(8):836-892.
 78. Ozasa N, Morimoto T, Bao B, Shioi T, Kimura T. Effects of machine-assisted cycling on exercise capacity and endothelial function in elderly patients with heart failure. *Circ J*. 2012;76(8):1889-1894.
 79. Erbs S, Höllriegel R, Linke A, et al. Exercise training in patients with advanced chronic heart failure (NYHA IIIb) promotes restoration of peripheral vasomotor function, induction of endogenous regeneration, and improvement of left ventricular function. *Circ Heart Fail*. 2010;3(4):486-494.
 80. Sandri M, Viehmann M, Adams V, et al. Chronic heart failure and aging—effects of exercise training on endothelial function and mechanisms of endothelial regeneration: results from the Leipzig Exercise Intervention in Chronic Heart Failure and Aging (LEICA) study. *Eur J Prev Cardiol*. 2016;23(4):349-358.
 81. Angadi SS, Mookadam F, Lee CD, Tucker WJ, Haykowsky MJ, Gaesser GA. High-intensity interval training vs. moderate-intensity continuous exercise training in heart failure with preserved ejection fraction: a pilot study. *J Appl Physiol (1985)*. 2015;119(6):753-758.
 82. Kitzman DW, Brubaker PH, Herrington DM, et al. Effect of endurance exercise training on endothelial function and arterial stiffness in older patients with heart failure and preserved ejection fraction: a randomized, controlled, single-blind trial. *J Am Coll Cardiol*. 2013;62(7):584-592.
 83. Dall CH, Gustafsson F, Christensen SB, et al. Effect of moderate- versus high-intensity exercise on vascular function, biomarkers and quality of life in heart transplant recipients: a randomized, crossover trial. *J Heart Lung Transplant*. 2015;34(8):1033-1041.
 84. Piercy KL, Troiano RP, Ballard RM, et al. The Physical Activity Guidelines for Americans. *JAMA*. 2018;320(19):2020-2028.
 85. Ponikowski P, Voors AA, Anker SD, et al; ESC Scientific Document Group. 2016 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure of the European Society of Cardiology (ESC); developed with the special contribution of the Heart Failure Association (HFA) of the ESC [published correction appears in *Eur Heart J*. 2018;39(10):860]. *Eur Heart J*. 2016;37(27):2129-2200.
 86. Oikonomou E, Siasos G, Zaromitidou M, et al. Atorvastatin treatment improves endothelial function through endothelial progenitor cells mobilization in ischemic heart failure patients. *Atherosclerosis*. 2015;238(2):159-164.
 87. Erbs S, Beck EB, Linke A, et al. High-dose rosuvastatin in chronic heart failure promotes vasculogenesis, corrects endothelial function, and improves cardiac remodeling—results from a randomized, double-blind, and placebo-controlled study. *Int J Cardiol*. 2011;146(1):56-63.
 88. Antoniadou C, Bakogiannis C, Leeson P, et al. Rapid, direct effects of statin treatment on arterial redox state and nitric oxide bioavailability in human atherosclerosis via tetrahydrobiopterin-mediated endothelial nitric oxide synthase coupling. *Circulation*. 2011;124(3):335-345.
 89. Neuman RB, Hayek SS, Poole JC, et al. Nitric oxide contributes to vasomotor tone in hypertensive African Americans treated with nebivolol and metoprolol. *J Clin Hypertens (Greenwich)*. 2016;18(3):223-231.
 90. Khan BV, Rahman ST, Haque T, et al. Vascular effects of nebivolol added to hydrochlorothiazide in African Americans with hypertension and echocardiographic evidence of diastolic dysfunction: the NASAA study. *J Cardiovasc Pharmacol Ther*. 2012;17(3):291-297.
 91. Falskov B, Hermann TS, Raunsø J, et al. Endothelial function is unaffected by changing between carvedilol and metoprolol in patients with heart failure—a randomized study. *Cardiovasc Diabetol*. 2011;10:91.
 92. Mahle WT, Wang A, Quyyumi AA, McConnell ME, Book WM. Impact of spironolactone on endothelial function in patients with single ventricle heart. *Congenit Heart Dis*. 2009;4(1):12-16.