

# Exenatide Affects Circulating Cardiovascular Risk Biomarkers Independently of Changes in Body Composition

MATHIJS C. BUNCK, MD<sup>1</sup>  
 MICHAELA DIAMANT, MD, PHD<sup>1</sup>  
 BJORN ELIASSON, MD, PHD<sup>2</sup>  
 ANJA CORNÉR, MD<sup>3,4</sup>  
 RIMMA M. SHAGINIAN, MD<sup>5</sup>

ROBERT J. HEINE, MD, PHD, FRCP<sup>1,6</sup>  
 MARJA-RIITTA TASKINEN, MD, PHD<sup>3</sup>  
 HANNELE YKI-JÄRVINEN, MD, PHD, FRCP<sup>3</sup>  
 ULF SMITH, MD, PHD, FRCP<sup>2</sup>

**OBJECTIVE** — To study the effect of exenatide on body composition and circulating cardiovascular risk biomarkers.

**RESEARCH DESIGN AND METHODS** — Metformin-treated patients with type 2 diabetes ( $N = 69$ ) were randomized to exenatide or insulin glargine and treated for 1 year. Body composition was evaluated by dual-energy X-ray absorptiometry. Additionally, body weight, waist circumference, and cardiovascular biomarkers were measured.

**RESULTS** — Treatment with exenatide for 1 year significantly reduced body weight, waist circumference, and total body and trunkal fat mass by 6, 5, 11, and 13%, respectively. In addition, exenatide increased total adiponectin by 12% and reduced high-sensitivity C-reactive protein by 61%. Insulin glargine significantly reduced endothelin-1 by 7%. These changes were statistically independent of the change in total body fat mass and body weight.

**CONCLUSIONS** — Exenatide treatment for 1 year reduced body fat mass and improved the profile of circulating biomarkers of cardiovascular risk. No significant changes were seen with insulin glargine except a trend for reduced endothelin-1 levels.

*Diabetes Care* 33:1734–1737, 2010

Abdominal obesity is associated with both type 2 diabetes and metabolic complications (1), including elevations in several circulating biomarkers of cardiovascular risk (2). Most pharmacological glucose-lowering treatments increase body weight (3). Therefore, treatments that not only reduce A1C, but also improve other associated changes such as abdominal obesity are urgently needed (4).

We previously reported in *Diabetes Care* that exenatide improves glycemic

control to the same extent as insulin glargine, although exenatide decreased and insulin glargine raised body weight (5). Herein we present additional data on associated changes in body composition and circulating levels of biomarkers of cardiovascular risk after 1 year of treatment.

## RESEARCH DESIGN AND METHODS

Details on study design were reported previously (5). Patients were randomized to exenatide ( $n = 36$ ) or

insulin glargine ( $n = 33$ ) added to their ongoing metformin therapy (baseline characteristics and patient disposition are shown in supplemental Fig. 1 in the online appendix available at <http://care.diabetesjournals.org/cgi/content/full/dc09-2361/DC1>). The study protocol was approved by each site's ethics review committee and was in accordance with the principles described in the Declaration of Helsinki. All participating patients gave their written informed consent prior to screening.

## Dual-energy X-ray absorptiometry scan

Lean body and fat mass was assessed using dual-energy X-ray absorptiometry (DEXA) scans (Delphi A; Hologic, Waltham, MA) at baseline and after treatment. Trunk (abdominal) and limb (hip/leg) regions of interest were determined from a total body scan. Waist circumference was measured at the midline of the interval between the iliac crest and the lowest rib using the mean of two measurements prior to the DEXA scan.

## Biochemical analyses

Cardiovascular risk biomarkers were collected at baseline and after 1 year of treatment. Serum was separated by centrifugation and stored at  $-80^{\circ}\text{C}$  until analysis. All serum samples were analyzed in the Lundberg Laboratory for Diabetes Research using a single batch. Total adiponectin, high molecular weight (HMW) adiponectin, resistin, leptin, high-sensitivity C-reactive protein (hs-CRP), interleukin (IL)-6, monocyte chemoattractant protein (MCP)-1, and endothelin-1 were determined by commercial ELISAs (R&D Systems, Abingdon, U.K.).

## Statistical analysis

Non-normally distributed data were log-transformed prior to statistical analysis, after which they approximated the normal distribution. All outcome measures are compared between the two treatment groups using an ANCOVA model including factors for treatment, investigative

From the <sup>1</sup>Department of Internal Medicine, Section of Endocrinology, Diabetes Center, VU University Medical Center, Amsterdam, the Netherlands; the <sup>2</sup>Lundberg Laboratory for Diabetes Research, Sahlgrenska University Hospital, Goteborg, Sweden; the <sup>3</sup>Department of Medicine, Helsinki University Central Hospital, Helsinki, Finland; the <sup>4</sup>Minerva Medical Research Institute, Helsinki, Finland; the <sup>5</sup>Eli Lilly and Company, Houten, the Netherlands; <sup>6</sup>Eli Lilly and Company, Indianapolis, Indiana.

Corresponding author: Mathijs C. Bunck, [mcbunck@vumc.nl](mailto:mcbunck@vumc.nl).

Received 25 December 2009 and accepted 20 April 2010. Published ahead of print at <http://care.diabetesjournals.org> on 27 April 2010. DOI: 10.2337/dc09-2361. Clinical trial reg. no. NCT00097500, [clinicaltrials.gov](http://clinicaltrials.gov).

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Body composition, circulating cardiovascular risk biomarkers and percentage change from baseline

	n	Baseline	Endpoint	Percentage change from baseline		
				LS mean	Between-treatment group difference	P
Total fat mass (kg)						
Insulin glargine	28	29.9 ± 1.6	28.5 ± 1.9	−1% (−7% to +5%)		
Exenatide	29	27.8 ± 1.4	25.4 ± 1.6	−11% (−18% to −5%)	−10% (−16% to −4%)	0.003
Total lean mass (kg)						
Insulin glargine	28	60.1 ± 1.7	60.6 ± 1.8	0% (−1% to +2%)		
Exenatide	29	57.8 ± 2.1	58.1 ± 2.4	0% (−2% to +1%)	−1% (−3% to +1%)	0.480
Trunk fat mass (kg)						
Insulin glargine	28	17.8 ± 0.9	16.6 ± 1.1	−1% (−8% to +5%)		
Exenatide	29	16.3 ± 0.8	14.8 ± 1.0	−13% (−18% to −7%)	−11% (−18% to −4%)	0.002
Body weight (kg)						
Insulin glargine	29	94.1 ± 2.5	93.8 ± 2.7	−1% (−3% to +1%)		
Exenatide	30	90.3 ± 2.4	86.4 ± 2.6	−6% (−8% to −3%)	−5% (−7% to −2%)	0.001
Waist circumference (cm)						
Insulin glargine	29	106.9 ± 1.9	107.4 ± 2.0	+1% (−1% to +3%)		
Exenatide	30	106.1 ± 1.9	100.6 ± 2.1	−5% (−7% to −3%)	−6% (−8% to −4%)	<0.001
Leptin (μg/l)						
Insulin glargine	29	7.79 ± 1.29	8.41 ± 1.53	+7% (−11% to +29%)		
Exenatide	30	8.50 ± 1.32	7.45 ± 1.17	−14% (−27% to +2%)	−19% (−34% to −1%)	0.045
Total adiponectin (ng/ml)						
Insulin glargine	29	4,648 ± 461	4,508 ± 436	−5% (−13% to +5%)		
Exenatide	30	4,848 ± 432	5,314 ± 466	+12% (+3% to +21%)	+17% (+6% to +30%)	0.004
HMW adiponectin (ng/ml)						
Insulin glargine	29	1,277 ± 221	1,321 ± 236	−0% (−24% to +31%)		
Exenatide	30	1,571 ± 255	1,850 ± 273	+19% (−6% to +51%)	+19% (−12% to +61%)	0.253
hs-CRP (mg/l)						
Insulin glargine	29	1.42 ± 0.27	1.38 ± 0.35	−20% (−50% to +27%)		
Exenatide	30	1.81 ± 0.25	1.30 ± 0.22	−61% (−74% to −42%)	−52% (−71% to −19%)	0.008
IL-6 (pg/ml)						
Insulin glargine	29	1.96 ± 0.21	2.17 ± 0.20	−4% (−26% to +25%)		
Exenatide	30	2.11 ± 0.22	2.10 ± 0.25	−10% (−28% to +14%)	−6% (−30% to +26%)	0.670
MCP-1 (pg/ml)						
Insulin glargine	29	1.22 ± 0.07	1.24 ± 0.07	−1% (−12% to +11%)		
Exenatide	30	1.18 ± 0.09	1.21 ± 0.11	−4% (−13% to +7%)	−2% (−14% to +12%)	0.728
Resistin (ng/ml)						
Insulin glargine	29	330 ± 15	329 ± 20	−3% (−13% to +7%)		
Exenatide	30	316 ± 14	311 ± 16	−0% (−9% to +9%)	+3% (−8% to +16%)	0.577
Endothelin-1 (ng/ml)						
Insulin glargine	29	2.57 ± 0.18	2.46 ± 0.19	−7% (−11% to −2%)		
Exenatide	30	2.53 ± 0.19	2.53 ± 0.19	−1% (−5% to +3%)	+6% (−1% to +12%)	0.045

Data are means ± SEM (body composition measures) or geometric means ± SEM (cardiovascular biomarkers) and body weight change—adjusted least-squares mean percentage change (95% CI) from baseline. LS, least-squares.

site, and baseline A1C stratum ( $\leq 8.5\%$  or  $> 8.5\%$ ), and baseline values of corresponding outcome measure as a covariate (5). Statistical analysis was performed using SPSS 16.0 for Mac OS X (SPSS, Chicago, IL). All inferential statistical tests were conducted at a significance level of 0.05 (two-sided).

**RESULTS**— Treatment for 1 year with exenatide resulted in a statistically significant reduction in total body fat mass (Table 1), mainly in the abdominal region, as

illustrated by the decrease in trunk fat mass and waist circumference, in contrast to insulin glargine. Neither treatment significantly affected lean body mass.

In univariate analysis, the reduction in body weight in the exenatide arm was significantly correlated with the changes in leptin ( $r = 0.580$ ,  $P = 0.001$ ) and hs-CRP ( $r = -0.590$ ,  $P = 0.001$ ). No statistically significant univariate correlation was found between changes in body weight and other biomarkers. Interestingly, changes in all circulating biomark-

ers did not correlate with the changes in total body fat mass (total adiponectin: Pearson  $\chi^2$  test,  $r = -0.224$ ,  $P = 0.106$ ; HMW adiponectin:  $r = 0.057$ ,  $P = 0.694$ ; leptin:  $r = 0.229$ ,  $P = 0.106$ ; hs-CRP:  $r = -0.023$ ,  $P = 0.872$ ).

After multivariate analysis and statistical adjustment for body weight change, exenatide increased total adiponectin and decreased hs-CRP concentrations, whereas insulin glargine did not (Table 1). Insulin glargine reduced endothelin-1 concentrations, whereas exenatide did

not. No statistically significant effect of either treatment on HMW adiponectin, IL-6, MCP-1, and resistin was observed.

The crude between-treatment group differences remained statistically significant after additional multivariate adjustment for total body fat mass change: total adiponectin +16% (95% CI: +5% to +28%),  $P = 0.004$ ; leptin -20% (-34% to -2%),  $P = 0.028$ ; hs-CRP -48% (-69% to -13%),  $P = 0.015$ ; and body weight change (Table 1): total adiponectin +17% (95% CI +6% to +30%),  $P = 0.004$ ; leptin -19% (-34% to 0%),  $P = 0.045$ ; hs-CRP -52% (-71% to -19%),  $P = 0.008$ .

**CONCLUSIONS**— This study showed that exenatide reduced body fat mass and improved the profile of circulating cardiovascular biomarkers. The changes in the different biomarkers could not be fully attributed to the observed changes in body fat mass and body weight. Direct effects of glucagon-like peptide 1 (GLP-1) receptor agonists on adipocyte function have been described in both animal experimental studies and in vitro studies in normal human adipocytes (rev. in 6); however, as a significant univariate correlation between change in body weight (not with fat mass) and cardiovascular biomarkers was present, our relatively small population may influence the statistical power of our study.

Animal studies have also demonstrated beneficial effects of exenatide on visceral fat mass (7) and circulating adiponectin (8), leptin (9), and CRP (10) concentrations. However, to the best of our knowledge, controlled clinical studies on the long-term effects of GLP-1 receptor agonists on body composition and biomarkers of cardiovascular risk have not previously been reported.

A recent 3-month study comparing exenatide to insulin glargine in 56 patients with type 2 diabetes has a design comparable to our 1-year study. Similar to our findings, this study showed that exenatide treatment was associated with reduced hs-CRP, without affecting the IL-6 levels (11).

Subanalysis of the Liraglutide Effect and Action in Diabetes (LEAD)-3 study data reported that liraglutide treatment for 52 weeks compared with treatment with glimiperide reduced DEXA-measured total fat tissue mass (12). Lean tissue mass was also reduced after 1 year of treatment, but as glimiperide also reduced lean tissue mass, this reduction was

not statistically significantly different between the groups. Twenty-six-week data from the LEAD-2 study was used to show that the observed reduction in fat mass was mainly a result of a reduction in visceral fat (12). Unfortunately, this study did not report the effects of body composition on circulating biomarkers. Serum leptin, hs-CRP, and IL-6 concentrations did not change in a 14-week placebo-controlled study with liraglutide 1.9 mg (13).

Of particular interest in our study was the finding that the changes in biomarkers of cardiovascular risk appeared to be independent of the changes in body fat mass. Recently, Chung et al. reported exendin-4 directly increased adiponectin mRNA levels and secretion in 3T3-L1 adipocytes (14). In that study, exendin-4 also decreased mRNA levels of IL-6 and MCP-1 (14). Additionally, we (15) and others (10) have previously reported beneficial effects of exenatide on hepatic steatosis, which also may contribute to a reduction in CRP.

In conclusion, we found that exenatide treatment for 1 year led to a reduced total fat mass, including visceral fat, while lean body mass was not significantly altered. Additionally, the circulating levels of adiponectin, leptin, and hs-CRP showed an improved profile that appeared to be independent of the changes in fat mass. In contrast, no significant changes in body composition or circulating biomarkers were seen with insulin glargine.

**Acknowledgments**— This study was sponsored by Amylin Pharmaceuticals, Inc., and Eli Lilly and Company.

M.C.B. is a speaker for Eli Lilly and Company. M.D. is a consultant and speaker for Eli Lilly and Company. Through M.D., the VU University Medical Center in Amsterdam has received research grants from Amylin Pharmaceuticals and Eli Lilly and Company. B.E. has served on the advisory board and is a speaker for Eli Lilly Sweden. R.M.S. is an employee of Eli Lilly and Company. R.J.H. is an employee and stockholder of Eli Lilly and Company. M.R.T. is a speaker for Eli Lilly and Company. H.Y.J. serves as a consultant for Amylin Pharmaceuticals. Through M.R.T. and H.Y.J. the Helsinki University Central Hospital has received research grants from Amylin Pharmaceuticals and Eli Lilly and Company. U.S. served on the advisory board and is a speaker for Amylin Pharmaceuticals, Inc. Through U.S. the Sahlgrenska University Hospital has received research grants from Amylin Pharmaceuticals and Eli Lilly and Company.

No other potential conflicts of interest relevant to this article were reported.

The study was collectively initiated and designed by the investigators from the three study sites. The investigators had full access to the trial data and had control over the statistical analysis and interpretation of the study results. M.C.B. collected and researched data, wrote the manuscript, and contributed to the discussion. M.D. researched data, contributed to the discussion, and reviewed/edited the manuscript. B.E. collected data, contributed to the discussion, and reviewed/edited the manuscript. A.C. collected data and reviewed/edited the manuscript. R.M.S. contributed to discussion and reviewed/edited the manuscript. R.J.H. researched data, contributed to the discussion, and reviewed/edited the manuscript. M.R.T. contributed to the discussion and reviewed/edited the manuscript. H.Y.J. contributed to the discussion and reviewed/edited the manuscript. U.S. researched data, contributed to the discussion, and reviewed/edited the manuscript.

Parts of this study were presented in abstract form at the 69th Scientific Sessions of the American Diabetes Association, New Orleans, Louisiana, 5–9 June 2009.

The authors thank the subjects for participating in the study.

## References

1. Jensen MD. Role of body fat distribution and the metabolic complications of obesity. *J Clin Endocrinol Metab* 2008;93:557–63
2. Diamant M, Lamb HJ, van de Ree MA, Enderit EL, Groeneveld Y, Bots ML, Kostense PJ, Radder JK. The association between abdominal visceral fat and carotid stiffness is mediated by circulating inflammatory markers in uncomplicated type 2 diabetes. *J Clin Endocrinol Metab* 2005;90:1495–1501
3. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet* 1998;352:837–853
4. DeFronzo RA. Banting Lecture. From the triumvirate to the ominous octet: a new paradigm for the treatment of type 2 diabetes mellitus. *Diabetes* 2009;58:773–795
5. Bunck MC, Diamant M, Cornér A, Eliasson B, Malloy JL, Shaginin RM, Deng W, Kendall DM, Taskinen MR, Smith U, Yki-Järvinen H, Heine RJ. One-year treatment with exenatide improves beta-cell function, compared with insulin glargine, in metformin-treated type 2 diabetic patients: a randomized, controlled trial. *Diabetes Care* 2009;32:762–768
6. Kim W, Egan JM. The role of incretins in glucose homeostasis and diabetes treat-

- ment. *Pharmacol Rev* 2008;60:470–512
7. Szayna M, Doyle ME, Betkey JA, Holloway HW, Spencer RG, Greig NH, Egan JM. Exendin-4 decelerates food intake, weight gain, and fat deposition in Zucker rats. *Endocrinology* 2000;141:1936–1941
  8. Li L, Yang G, Li Q, Tan X, Liu H, Tang Y, Boden G. Exenatide prevents fat-induced insulin resistance and raises adiponectin expression and plasma levels. *Diabetes Obes Metab* 2008;10:921–930
  9. Pérez-Tilve D, González-Matías L, Alvarez-Crespo M, Leiras R, Tovar S, Diéguez C, Mallo F. Exendin-4 potently decreases ghrelin levels in fasting rats. *Diabetes* 2007;56:143–151
  10. Ding X, Saxena NK, Lin S, Gupta NA, Gupta N, Anania FA. Exendin-4, a glucagon-like protein-1 (GLP-1) receptor agonist, reverses hepatic steatosis in ob/ob mice. *Hepatology* 2006;43:173–181
  11. Horton ES, Cohen A, Gibson H, Lamparello B, Herzlinger S, McFarland L. Effects of exenatide vs insulin glargine on cardiovascular risk factors in subjects with type 2 diabetes. *Diabetologia* 2009;52:S298–S299
  12. Jendle J, Nauck MA, Matthews DR, Frid A, Hermansen K, Düring M, Zdravkovic M, Strauss BJ, Garber AJ, LEAD-2 and LEAD-3 Study Groups. Weight loss with liraglutide, a once-daily human glucagon-like peptide-1 analogue for type 2 diabetes treatment as monotherapy or added to metformin, is primarily as a result of a reduction in fat tissue. *Diabetes Obes Metab* 2009;11:1163–1172
  13. Courrèges JP, Vilsbøll T, Zdravkovic M, Le-Thi T, Krarup T, Schmitz O, Verhoeven R, Bugáňová I, Madsbad S. Beneficial effects of once-daily liraglutide, a human glucagon-like peptide-1 analogue, on cardiovascular risk biomarkers in patients with Type 2 diabetes. *Diabet Med* 2008;25:1129–1131
  14. Kim Chung le T, Hosaka T, Yoshida M, Harada N, Sakaue H, Sakai T, Nakaya Y. Exendin-4, a GLP-1 receptor agonist, directly induces adiponectin expression through protein kinase A pathway and prevents inflammatory adipokine expression. *Biochem Biophys Res Commun* 2009;390:613–618
  15. Tushuizen ME, Bunck MC, Pouwels PJ, van Waesberghe JH, Diamant M, Heine RJ. Incretin mimetics as a novel therapeutic option for hepatic steatosis. *Liver Int* 2006;26:1015–1017