

Research paper

Juvenile hormone affects the development and strength of circadian rhythms in young bumble bee (*Bombus terrestris*) workers

Atul Pandey^a, Uzi Motro^{a,b}, Guy Bloch^{a,b,*}

^a Department of Ecology, Evolution, and Behavior, The Hebrew University of Jerusalem, Jerusalem, Israel

^b The Federmann Center for the Study of Rationality, The Hebrew University of Jerusalem, Jerusalem, Israel

ARTICLE INFO

Keywords:

Hormone
Juvenile hormone
Bumble bee
Bombus terrestris
Circadian rhythms
Locomotor activity
Sleep
Gonadotropin

ABSTRACT

The circadian and endocrine systems influence many physiological processes in animals, but little is known on the ways they interact in insects. We tested the hypothesis that juvenile hormone (JH) influences circadian rhythms in the social bumble bee *Bombus terrestris*. JH is the major gonadotropin in this species coordinating processes such as vitellogenesis, oogenesis, wax production, and behaviors associated with reproduction. It is unknown however, whether it also influences circadian processes. We topically treated newly-emerged bees with the allatoxin Precocene-I (P-I) to reduce circulating JH titers and applied the natural JH (JH-III) for replacement therapy. We repeated this experiment in three trials, each with bees from different source colonies. Measurements of ovarian activity suggest that our JH manipulations were effective; bees treated with P-I had inactive ovaries, and this effect was fully recovered by subsequent JH treatment. We found that JH augments the strength of circadian rhythms and the pace of rhythm development in individually isolated newly emerged worker bees. JH manipulation did not affect the free-running circadian period, overall level of locomotor activity, sleep amount, or sleep structure. Given that acute manipulation at an early age produced relatively long-lasting effects, we propose that JH effects on circadian rhythms are mostly organizational, accelerating the development or integration of the circadian system.

1. Introduction

The circadian and endocrine systems are pivotal for the integration of external and internal information and for coordinating processes in multiple tissues (Bedrosian et al., 2016; Neumann et al., 2019; Tsang et al., 2013). In vertebrates, particularly in mammals, there is good evidence that these two regulatory systems interact. The circadian system influences endocrine tissues and processes, resulting in circadian rhythms in the circulating levels of many vertebrate hormones (Bedrosian et al., 2016; Hastings et al., 2007; Kriegsfeld et al., 2002; Neumann et al., 2019; Tsang et al., 2013). Hormones also influence circadian functions. For example, gonadal hormones were shown to lengthen the free-running period, decrease period precision, or reduce the duration of daily activity bouts (e.g., Albers, 1981; Daan et al., 1975; Iwahana et al., 2008; Jechura et al., 2000; Karatsoreos et al., 2011, 2007). Many of these effects are at least partially mediated by hormone receptors in the suprachiasmatic nucleus (SCN), the central brain clock of mammals (He et al., 2007; Iwahana et al., 2008; Karatsoreos et al., 2011, 2007; Sellix et al., 2004). Steroid hormones, including progesterins,

corticosteroids, estrogens, and androgens, were also shown to influence circadian rhythms in locomotor activity and transcription levels of core circadian clock genes in fishes (Zhao et al., 2018).

The interplay between the circadian and endocrine systems is relatively little explored in adult insects (Bloch et al., 2013). Only a few studies recorded hormone titers throughout the day under constant conditions. Nevertheless, these measurements, together with indirect evidence for circadian modulation of hormone biosynthesis rate, and the expression of genes encoding proteins involved in hormone biosynthesis, hormone binding, or hormone degradation, suggest that the circadian system influences the circulating levels of many insect hormones. There is also little evidence for hormonal regulation of circadian rhythms in insects (reviewed in Bloch et al., 2013). This includes the best-studied insect hormone, juvenile hormone (JH), which functions as a gonadotropin in many insects. There is some evidence that JH and ovarian activity influences circadian rhythms in the cockroach *Blattella germanica*, although the data is quite perplexing. Females of this species show strong circadian rhythms during the vitellogenic phase of the reproductive cycle when JH titers are expected to be high, but not in

* Corresponding author. Department of Ecology, Evolution, and Behavior, The Hebrew University of Jerusalem, Jerusalem, Israel.

E-mail address: guy.bloch@mail.huji.ac.il (G. Bloch).

<https://doi.org/10.1016/j.nbscr.2020.100056>

Received 18 May 2020; Received in revised form 13 August 2020; Accepted 17 August 2020

Available online 24 August 2020

2451-9944/© 2020 The Authors.

Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

sexually receptive females during the first gonadotropic cycle (Lee and Wu, 1994). Active ovaries mask the expression of circadian rhythms, and allatectomy abolished the strong circadian rhythms that are shown by ovariectomized females. However, replacement therapy with a JH analog did not restore circadian rhythmicity (Lin and Lee, 1998).

Mutations in the JH receptors Met, and the JH biosynthesis pathway enzyme JH acid O-methyltransferase (JHAMT) attenuated the strength of circadian rhythms in *D. melanogaster*, suggesting that JH affects circadian rhythms in this species too (Wu et al., 2018). Studies with the Milkweed bug *Oncopeltus fasciatus*, in which JH levels were decreased by treatment with Precocene-II, and increased by JH supplementation produced conflicting evidence concerning the effects of JH on the circadian rhythm of feeding and mating behavior (Walker, 1977; Woodard and Rankin, 1980). JH does not seem to be involved in the regulation of circadian rhythms in the cockroach *Blaberus discoidalis* because circadian rhythms in locomotor activity were similar in the allatectomized and control individuals (Shepard and Keeley, 1972). The influence of JH on circadian rhythms was also studied in the Western honey bee *Apis mellifera* in which JH does not function as a major gonadotropin (Bloch et al., 2002a). JH manipulation by allatectomy and replacement therapy with the JH analog methoprene, which successfully affected the age of first foraging (Sullivan et al., 2000), failed to affect circadian rhythms in locomotor activity (Bloch et al., 2002a). Similar JH manipulations did not have a consistent influence on the circadian brain expression of the canonical clock gene *Period*. However, there was a trend towards aberrant cycling in allatectomized bees (Bloch and Meshi, 2007).

Taken together, the available studies revealed significant variability in the effects of JH manipulation on circadian rhythms in insects and suggest that some of this variability may relate to whether or not JH functions as a gonadotropin. To test this hypothesis, we studied the influence of JH on circadian rhythms in the social bumble bee *Bombus terrestris*. Bumble bees are taxonomically related to honey bees, but live in smaller annual colonies, showing a simpler form of social organization (Michener, 1974). By contrast to honey bees, in bumble bees, JH is the major gonadotropin (Shpigler et al., 2014), and does not affect task performance (i.e., brood care vs. foraging activity; Shpigler et al., 2016). However, in both species, the division of labor is similarly correlated with the expression of circadian activity rhythms; foragers have strong circadian rhythms, whereas nurse bees are typically active around the clock with attenuated rhythms (reviewed in Bloch, 2010; Eban-Rothschild and Bloch, 2012a). We manipulated JH levels in young bumble bee workers using topical treatments with the allatoxin precocene 1 (P-I) to reduce hemolymph JH titers and the natural JH of bumble bees (JH-III; Bloch et al., 2000, 1996) for replacement therapy. Our results suggest that JH affects the strength and development of circadian rhythm in locomotor activity in *Bombus terrestris*.

2. Materials and methods

2.1. Bees

Colonies of *Bombus terrestris* containing a queen, 5–10 workers, and brood at various developmental stages (typically 2–4 days post first worker emergence) were purchased from Polyam Pollination Services, Kibbutz Yad-Mordechai, Israel (Trial 1 & 2) or BioBee Biological Systems Ltd. Kibbutz Sde Eliyahu, Israel (Trial 3). We housed each colony in a wooden nesting box inside an environmental chamber (29 ± 1 °C; $55\% \pm 10\%$ RH) in constant darkness at the Bee Research Facility at the Edmond J. Safra Campus of the Hebrew University of Jerusalem, Givat Ram, Jerusalem. The nest boxes ($21 \times 21 \times 12$ cm) were made of wood and were fitted with a top and front wall made of transparent Plexiglas panels. The colonies were fed *ad libitum* with commercial sugar syrup and pollen cakes made of honey bee collected pollen that was purchased from Polyam Pollination Services. We performed all treatments and feeding under dim red light (Energizer headlights covered with red

filters) that bumble bees do not see well (Peitsch et al., 1992) and minimized the production of substrate-borne vibrations.

2.2. General experimental outline

Our experimental design is summarized in Fig. 1. At day-1 of each trial, we collected 120–130 newly-emerged worker bees (<24 h of age) from a pool of 12–15 ‘donor colonies’, chilled them on ice, measured their head width, treated them as detailed below (~30 bees per treatment under dim red light), and placed them in groups of four of the same treatment in small wooden cages ($12 \times 5 \times 8$ cm). The bees were collected in two batches such that a bee was chilled on ice for no longer than 90 min, and the entire procedure took 2–3 h from collection in the colony until the end of the first treatment (precocene-I, castor oil or control handling, see Fig. 1 and below). On Day-3, approximately 24–26 h after the second treatment, we transferred the bees to an environmental chamber, isolated each one of them in an individual cage, and monitored their locomotor activity for 5–9 consecutive days (five days in Trial 1, eight days in Trial 2, and nine days in Trial 3, see Fig. 1) under controlled environmental conditions (29 ± 1 °C; $55\% \pm 5\%$ RH). At the end of the monitoring session, we stored all the live bees in a -20 °C until assessing their ovarian state. We repeated this experiment three times: in April 2017 (Trial 1), May 2017 (Trial 2), and April–May 2019 (Trial 3).

2.3. Manipulating circulating JH titers

We subjected the callow bees to one of four treatment groups. Control bees (“Control”), were handled and chilled on ice (20–25 min) similarly to bees from the other experimental groups, on both Day-1 and Day-2, but not treated with any drug or vehicle. Sham-treated bees (“Sham”), were chilled on ice and topically treated with castor oil on Day-1, and with dimethylformamide (DMF, J.T Backers, cat # 7032; 3.5 μ l/bee, irrespective of body sizes) on Day-2. The amount of castor oil (Sigma-Aldrich, cat # 259,853; 4.0–5.2 μ l/bee) was adjusted according to body size, as detailed in Table 1. To reduce JH levels, we treated the bees on Day-1 with the allatoxin Precocene-I (“P-I”; Sigma-Aldrich, cat # 195,855) suspended in castor oil vehicle solution. We thoroughly mixed the P-I, and castor oil by repeated pipetting followed by vortexing the mixture at high speed for 2–3 min. We adjusted the P-I treatment to body size (200–260 μ g/4.0–5.2 μ l castor oil/bee; we confirmed that our measurements of head width are strongly correlated with other measures of body size, data not shown; Table 1), as previously reported (Pandey et al., 2020). Replacement therapy (“P-I + JH”) treated bees were topically treated with P-I on Day-1, as described above. On the following day, we topically treated the bees with 50 μ g JH-III (Sigma-Aldrich, cat # J2000) dissolved in 3.5 μ l DMF vehicle solution. We handled and treated the control and sham control bees similarly to their respective treatment groups. Following the P-I, and JH-III treatments, we kept the bees on ice-chilled glass plates immobilized for ~10 min in order to improve drug absorption and minimize possible wiping-off. Additional details on our JH manipulation procedures, and the validation of our assays, can be found in Shpigler et al. (2016) and Pandey et al. (2020).

2.4. Monitoring locomotor activity

We placed each bee individually in a transparent monitoring cage (made of a modified 90 mm Petri dish). The cages with the focal bees were transferred to the monitoring environmental chamber inside a lightproof box to avoid any light exposure. The monitoring chamber was illuminated with dim red light that the bees cannot see (Edison Federal EFEF 1AE1 Far [Cherry] Red LED; mean wavelength = 740 nm, maximum and minimum wavelengths were 750 and 730, respectively). The location of each focal bee was recorded with one of four CCD cameras (Panasonic WV-BP334, 0.08 lux CCD video cameras) and an




Fig. 1. Schematic description of the general experimental outline. The numbers represent the days of the experiment measured as time passed after collecting the callow bees. On Day-1 between 14.00 and 15.00 h, we collected newly emerged (0–24hr of age) worker bees from 12 to 15 queenright “donor” colonies and subjected them to the first treatment at around 16:00. About 24 h later, on the afternoon of Day-2, we subjected the bees to the second treatment. In the afternoon (16:00–18:00) of Day-3, we transferred each bee into an individual monitoring cage and then monitored locomotor activity for 5–9 successive days under constant laboratory conditions. At the end of the monitoring session, we collected the bees and recorded their ovarian state. N = number, ①-first treatment, ②-second treatment, ◀-start, and ▶-end of monitoring session trial; T1 = trial-1, T2 = trial-2, and T3-trial-3. All treatments, observations, and manipulations were conducted under dim red light.

Table 1

Body size adjusted amounts of Precocene-I used to reduce JH titers in callow worker bees.

Size	Head Width (mm)	Precocene-I (µg/bee)	Castor oil (µl/bee)
Small	2.4–2.6	200	4.0
Small-medium	2.7–2.9	220	4.4
Large-medium	3.0–3.5	240	4.8
Large	3.5–4.2	260	5.2

image acquisition board (IMAQ 1409, National Instruments, Austin, TX) at a frequency of 1 Hz over the entire monitoring session as previously described (Shemesh et al., 2007; Yerushalmi et al., 2006). The distance traveled by the bee in pixels is calculated by comparing the location on successive images on the camera field of view. Each camera recorded activity in 30 arenas (i.e., monitoring cages) on a single tray. Four cages, one on each tray, were left empty as controls, recording background noise.

2.5. Analyses of circadian rhythms and sleep

We used the ClockLab circadian analysis software package (version 6; Actimetrics, Wilmette, IL) for all circadian rhythms and sleep analyses. We used 10-min bins to generate actograms and for the periodogram analyses. We performed χ^2 periodogram analyses to the activity data collected on days 4–7 (Trial 1), days 4–9 (Trial 2), and days 4–10 (Trial 3). Each day was defined as starting at 06:00 of the morning following transfer to the monitoring chamber. We used the ‘Power’, as an index for the strength of circadian rhythms. The Power was calculated as the height (in arbitrary units) of the periodogram peak relative to a point with a similar period length on a significance threshold line equal to $p = 0.01$ (for more details, see Klarsfeld et al., 2003; Yerushalmi et al., 2006). Bees with a periodogram peak below the $p = 0.01$ threshold line were assigned a zero power value. The free-running period (FRP, τ) was determined as the period length of the prominent peak of the χ^2 periodogram. The age of first showing circadian rhythmicity was determined as the first 3-day sliding window in which the periodogram analysis produced a statistically significant rhythm with a period between 20 and 28 h (Bloch et al., 2002a; Yerushalmi et al., 2006). The overall level of locomotor activity was determined using ClockLab analysis software and represented as total counts. A sleep state was defined as a bout of 5 min or longer with no movement. This sleep proxy is based on detailed video analyses of the sleep-like behavior of individually isolated *B. terrestris* workers (Nagari et al., 2019). This index is similar to the sleep indices used for honey bees (Eban-Rothschild and

Bloch, 2008) and *Drosophila melanogaster* fruit flies (Shaw et al., 2000). The analyses of sleep were done using the ClockLab software, following preliminary analyses showing that the measurements are comparable to those obtained with the BeeSleep and Sleepograms algorithms that we used in previous studies (Eban-Rothschild and Bloch, 2008; Nagari et al., 2019).

2.6. Assessing ovarian state

We thawed the stored bee samples and immersed them in bee saline (Huang et al., 1991) on a wax-filled dissecting plate under a stereomicroscope (Nikon SMZ645). We cut three incisions through the lateral and proximal-ventral abdominal cuticle using fine scissors and immersed the internal organs in honey bee saline. We used fine forceps to gently remove the ovaries into a drop of saline on a microscope slide and use the ocular ruler to measure the length of the four largest terminal oocytes from both ovarioles.

2.7. Statistical analyses

Assessing the influence of JH manipulation on the Free Running Period (FRP, τ), power, locomotor activity, sleep amount and structure, and oocyte size, is commonly done using a two-way ANOVA with Treatment and Trial as factors. However, assessments of our data indicate that our variables do not meet the presumption necessary for running this test. Using Kolmogorov-Smirnov tests we found that not all variables are normally distributed. Moreover, not all variables exhibit equality of variances, and not each Treatment \times Trial combination contains the same number of observations. Thus, we had to resort to different tools. Based on the Kolmogorov-Smirnov results, we determined whether the distribution of the abovementioned variables is the same across all three trials using either a one-way ANOVA for the normally distributed variables or the Kruskal-Wallis test for the non-normally distributed variables. Given that these analyses revealed differences between the trials, we analyzed each trial separately and then combined the p -values of each trial to obtain an overall p -value. The combined p -values were calculated either by the Winer method of adding t 's (for parametric analyses) or by the Mosteller-Bush method of adding weighted z 's (for non-parametric analyses; see Rosenthal, 1978). We used Cohen's d as an approximation for the overall effect size in the pooled data.

We first compared the Control and Sham treatments, using either the parametric independent samples t -test or the non-parametric Mann-Whitney U test for pair comparison within each trial, and then combined the results of each trial to obtain an overall p -value. Given that for all

variables tested, the Control and Sham groups did not differ statistically, we used in subsequent analyses the Sham treatment group as the single control for the P-I and P-I + JH treatments. Our analyses further show that our variables do not meet the assumptions needed for combining trials in the non-parametric Friedman's two-way ANOVA (with Trial and Treatment as factors). Thus, taking into account these limitations, and the low power of Kruskal-Wallis tests for the analyses of each trial (due to the relatively low sample size and expected similarity between the Sham and the replacement therapy), we chose an alternative approach that is more appropriate and sensitive. We used pairwise comparisons between the Sham, P-I, and P-I + JH treatments, using either the parametric independent samples *t*-test or the non-parametric Mann-Whitney *U* test for each trial, as appropriate. We then combined the *p*-values of the three trials using either of the procedures mentioned above. Given that in each analysis we have three comparisons, the Bonferroni corrected significance level was set at $p = 0.017$. Only differences that are significant after Bonferroni corrections are depicted as statistically different in our figures.

For analyzing the rate of rhythm development, we considered only Trials 2 and 3; during days 4–8 (Trial 1, had only two 3-day sliding windows). In these two trials, each bee has five dependent observations, which are the Power value of the sliding windows on days 4–8. Thus, each bee can be represented by the slope of the linear regression of Power over time. Note that this step of the analysis does not require any assumptions of normality (since we do not test the significance of each slope). Next, we calculated the group means of the slopes of the individual bees for each Treatment. We further tested for normality of these slope samples. If these slope samples are normally distributed, we can obtain the combined *p*-value over both trials by using the Winer method of adding *t*'s. Given that *B. terrestris* workers develop circadian rhythms in locomotor activity with age (Yerushalmi et al., 2006), we expected the slopes to be positive, allowing us to present one-tailed *p*-values.

We used IBM SPSS Version 25 and GraphPad Prism, version 5.01 for the statistical analyses and graphics, respectively.




Fig. 2. The influence of JH manipulation on ovarian state. Sham - Sham treatment with only the vehicles used to deliver the drugs; P-I - Precocene-I treatment to reduce JH levels; P-I + JH - replacement therapy with JH-III, the natural JH of bumble bees. The box plots show the median (line) and mean (+), the box frame spans over the first to the third quartile. The whiskers depict the 5th/95th percentile, outliers are depicted with black dots. The numbers within brackets shows sample size. Treatments connected with a horizontal bar and asterisks are statistically different in independent samples *t*-test followed by Bonferroni correction. For the combined analysis ("All trials" bottom right panel), we used the Winer method of adding *t*'s after applying the Bonferroni correction. * $p < 0.017$, ** $p < 0.001$.

3. Results

3.1. The influence of JH on ovarian state

Given that previous studies show that JH is the major gonadotropin regulating oogenesis in *B. terrestris* (Amsalem et al., 2014; Pandey et al., 2020; Shpigler et al., 2014; Shpigler et al. 2016), we used ovarian activity as a proxy for JH levels (Bloch et al., 2000, 1996). We found that bees treated with P-I have less developed ovaries compared to sham treated bees, an effect that was reverted by replacement therapy with JH-III. The distribution of oocyte length did not differ from normal (Kolmogorov-Smirnov test, $p = 0.134$), but equality of variances among the trials was rejected (Levin's test, $p < 0.001$). For the sake of consistency with the other analyses (see below), we treated here each trial separately, and then combined the results of each trial to obtain an overall *p*-value. We found that the Control and the Sham treatments did not differ statistically (Supplementary Fig. S1, independent samples *t*-test: Trial 1, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 18$, $t_{35} = -0.837$, $p = 0.408$; Trial 2, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 20$, $t_{37} = 1.435$, $p = 0.160$; Trial 3, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 22$, $t_{39} = -0.228$, $p = 0.821$). The combined significance level using the Winer method of adding *t*'s, was $z = 0.208$, with an overall $p = 0.835$, and an overall effect size = 0.007. Based on these analyses, we used only the Sham treatment as the control group, and carried out pairwise comparisons between Sham, P-I and P-I + JH treatments, using independent samples *t*-tests. The P-I treated bees had less developed ovaries compared to both the Sham and Replacement Therapy treatments in all three trials (Fig. 2; independent samples *t*-test for comparing the means; Sham vs. P-I: Trial 1, $n_{\text{Sham}} = 18$, $n_{\text{P-I}} = 16$, $t_{23} = 4.865$, $p < 0.001$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I}} = 22$, $t_{28} = 4.550$, $p < 0.001$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I}} = 7$, $t_{27} = 2.527$, $p = 0.018$; overall $z = 6.622$, $p < 0.001$, effect size = 0.850; P-I vs. P-I + JH: Trial 1, $n_{\text{P-I}} = 16$, $n_{\text{P-I+JH}} = 24$, $t_{37} = -7.632$, $p < 0.001$; Trial 2, $n_{\text{P-I}} = 22$, $n_{\text{P-I+JH}} = 20$, $t_{32} = -8.373$, $p < 0.001$; Trial 3, $n_{\text{P-I}} = 7$, $n_{\text{P-I+JH}} = 15$, $t_{20} = -4.397$, $p < 0.001$; overall $z = -11.345$, $p < 0.001$, effect size = 1.704 (*p*-values in bold are significant after Bonferroni correction, i.e., $p < 0.017$). These analyses of ovarian state suggest that our treatments effectively manipulated JH levels. There were no significant differences between the Sham vs. P-I + JH treatments indicating that our replacement therapy successfully

reverted the effect of JH reduction by the P-I treatment (Trial 1, $n_{\text{Sham}} = 18$, $n_{\text{P-I+JH}} = 24$, $t_{40} = -0.674$, $p = 0.504$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I+JH}} = 20$, $t_{38} = -2.042$, $p = 0.048$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I+JH}} = 15$, $t_{33} = -1.321$, $p = 0.198$; overall $z = -2.267$, $p = 0.023$, effect size = 0.266).

3.2. The influence of JH on the overall level of locomotor activity

To test whether JH affect the overall level of locomotor activity, we calculated for each bee the mean locomotor activity per hour during days 1–4. The distribution of these individual means did not differ from normal distribution (Kolmogorov-Smirnov test for normality: $p = 0.239$, $p = 0.334$ and $p = 0.198$ for Trials 1, 2 and 3, respectively) allowing us to use parametric statistics. We used the Sham as our control group following a preliminary analysis showing that the Control and the Sham treatments did not differ statistically (independent samples t -test: Trial 1, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 19$, $t_{30} = -0.513$, $p = 0.611$; Trial 2, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 20$, $t_{37} = 1.681$, $p = 0.101$; Trial 3, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 22$, $t_{31} = 0.012$, $p = 0.991$; with an overall $p = 0.509$, using the Winer method of adding t 's, the combined $z = 0.660$, and an overall effect size = 0.057). Table 2 summarizes the means of the individual mean activity per hour for each of the three trials. Pairwise independent samples t -tests between Sham, P-I and P-I + JH treatments, suggest that JH manipulation did not affect the overall level of locomotor activity. Sham vs. P-I: Trial 1, $n_{\text{Sham}} = 19$, $n_{\text{P-I}} = 16$, $t_{33} = 0.658$, $p = 0.515$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I}} = 22$, $t_{40} = -0.411$, $p = 0.683$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I}} = 7$, $t_{27} = -0.314$, $p = 0.756$; overall $z = -0.037$, $p = 0.970$, effect size = 0.067. Sham vs. P-I + JH: Trial 1, $n_{\text{Sham}} = 19$, $n_{\text{P-I+JH}} = 24$, $t_{41} = -1.293$, $p = 0.203$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I+JH}} = 20$, $t_{38} = 0.200$, $p = 0.842$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I+JH}} = 15$, $t_{35} = 1.398$, $p = 0.171$; overall $z = 0.171$, $p = 0.864$, effect size = 0.021. P-I vs. P-I + JH: Trial 1, $n_{\text{P-I}} = 16$, $n_{\text{P-I+JH}} = 24$, $t_{38} = -1.820$, $p = 0.077$; Trial 2, $n_{\text{P-I}} = 22$, $n_{\text{P-I+JH}} = 20$, $t_{40} = 0.562$, $p = 0.577$; Trial 3, $n_{\text{P-I}} = 7$, $n_{\text{P-I+JH}} = 15$, $t_{20} = 1.118$, $p = 0.277$; overall $z = -0.078$, $p = 0.938$, effect size = 0.080.

3.3. The influence of JH on the strength of circadian rhythms

Using the Power as an index for the strength of circadian rhythms in locomotor activity (Klarsfeld et al., 2003; Yerushalmi et al., 2006) we found that JH manipulation has a significant effect (Fig. 3). Given that the Power was not normally distributed (Kolmogorov-Smirnov test, $p < 0.001$), and the distribution of Power differed among the three trials (Kruskal-Wallis test, $p < 0.001$), we used non-parametric statistics, and treated each trial separately. We used the Sham as our control group following a preliminary test showing no difference between the Control and the Sham treatments in all three trials (Supplementary Fig. S2, Mann-Whitney U test: Trial 1, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 19$, $z = -0.394$, $p = 0.707$; Trial 2, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 20$, $z = -0.141$, $p = 0.895$; Trial 3, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 22$, $z = -0.013$, $p = 0.995$; Mosteller-Bush method of adding weighted z 's, $z = -0.131$, all trials combined, significance level $p = 0.896$, and an effect size = 0.011). In all three trials there was a consistent trend of attenuated circadian rhythms for the P-I treated bees compared to the other groups, but the differences were statistically significant after Bonferroni corrections only in the first trial for the comparison of the P-I and Sham treatments. The differences were statistically significant after correction in a pooled analyses that combined the p -values of all three trials (Fig. 3; Sham vs. P-I: Trial 1, $n_{\text{Sham}} = 19$, $n_{\text{P-I}} = 16$, $z = 2.563$, $p = 0.009$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I}} = 22$, $z = 2.242$,

Table 2

Manipulating JH titers did not affect the overall level of locomotor activity. The values are mean \pm se movement/hour, with sample size in parentheses.

Treatment	Trial 1	Trial 2	Trial 3	All trials
Sham	704 \pm 76 (19)	512 \pm 48 (20)	704 \pm 77 (22)	640 \pm 64 (61)
P-I	633 \pm 77 (16)	544 \pm 61 (22)	761 \pm 213 (7)	646 \pm 63 (45)
P-I + JH	858 \pm 87 (24)	497 \pm 57 (20)	529 \pm 102 (15)	628 \pm 115 (59)

$p = 0.025$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I}} = 7$, $z = 0.561$, $p = 0.600$; All trials combined, $z = 3.234$, $p = 0.001$, effect size = 0.554). Likewise, there was a consistent trend of lower Power for the P-I compared to P-I + JH treated bees, but the differences were statistically significant after Bonferroni corrections only in the pooled analysis (Trial 1, $n_{\text{P-I}} = 16$, $n_{\text{P-I+JH}} = 24$, $z = -1.453$, $p = 0.149$; Trial 2, $n_{\text{P-I}} = 22$, $n_{\text{P-I+JH}} = 20$, $z = -2.040$, $p = 0.042$; Trial 3, $n_{\text{P-I}} = 7$, $n_{\text{P-I+JH}} = 15$, $z = -0.390$, $p = 0.729$; all trials combined, $z = -2.456$, $p = 0.014$, effect size = 0.459). The Sham and replacement therapy groups did not differ in any of the trials, or in the pooled analysis, indicating that treatment with the natural JH fully recovered the P-I effects (Trial 1, $n_{\text{Sham}} = 19$, $n_{\text{P-I+JH}} = 24$, $z = 1.095$, $p = 0.279$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I+JH}} = 20$, $z = -0.839$, $p = 0.409$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I+JH}} = 15$, $z = 0.449$, $p = 0.663$ (Trial 3); All trials combined, $z = 0.434$, $p = 0.664$, effect size = 0.006).

3.4. The influence of JH on the free-running period (Tau)

Our analyses suggest that JH does not affect the free-running period of circadian rhythms in locomotor activity (Fig. 4). Given that the distribution of τ was not normal (Kolmogorov-Smirnov test, $p = 0.001$), we used non-parametric statistics. The distribution differs for the three trials (Kruskal-Wallis test, $p < 0.001$), and therefore we treated each trial separately, and then combined the obtained p -values to an overall p -value. We used the Sham as a single control group following a preliminary analysis showing that the FRP is not different between the Control and the Sham treatments in all three trials (Supplementary Fig. S3, Mann-Whitney U test for each trial: Trial 1, $n_{\text{Control}} = 19$, $n_{\text{Sham}} = 16$, $z = -0.695$, $p = 0.497$; Trial 2, $n_{\text{Control}} = 16$, $n_{\text{Sham}} = 17$, $z = 0.525$, $p = 0.611$; Trial 3, $n_{\text{Control}} = 18$, $n_{\text{Sham}} = 21$, $z = 1.089$, $p = 0.283$) as well as in a Mosteller-Bush method for combining all trials combined (weighted z 's, we got $z = 0.573$, $p = 0.567$, effect size = 0.110). The τ did not differ between the Sham, P-I, and Replacement Therapy treatment groups in any of the three trials or in the pooled analysis (Fig. 4; Mann-Whitney U test for pairwise comparisons: Trial 1, Sham vs. P-I: $n_{\text{Sham}} = 16$, $n_{\text{P-I}} = 8$, $z = 0.858$, $p = 0.408$; Trial 2, $n_{\text{Sham}} = 17$, $n_{\text{P-I}} = 19$, $z = 0.666$, $p = 0.515$; Trial 3, $n_{\text{Sham}} = 21$, $n_{\text{P-I}} = 5$, $z = -1.374$, $p = 0.180$; All trials combined, $z = 0.175$, $p = 0.861$, effect size = 0.006. Sham vs. P-I + JH: Trial 1, $n_{\text{Sham}} = 16$, $n_{\text{P-I+JH}} = 17$, $z = 1.459$, $p = 0.149$; Trial 2, $n_{\text{Sham}} = 17$, $n_{\text{P-I+JH}} = 18$, $z = -0.976$, $p = 0.338$; Trial 3, $n_{\text{Sham}} = 21$, $n_{\text{P-I+JH}} = 12$, $z = -2.196$, $p = 0.027$; All trials combined, $z = -1.002$, $p = 0.316$, effect size = 0.135. P-I vs. P-I + JH: Trial 1, $n_{\text{P-I}} = 8$, $n_{\text{P-I+JH}} = 17$, $z = -0.321$, $p = 0.765$; Trial 2, $n_{\text{P-I}} = 19$, $n_{\text{P-I+JH}} = 18$, $z = -1.933$, $p = 0.054$; Trial 3, $n_{\text{P-I}} = 5$, $n_{\text{P-I+JH}} = 12$, $z = -0.159$, $p = 0.898$; All trials combined, $z = -1.721$, $p = 0.085$, effect size = 0.115).

3.5. The influence of JH on the ontogeny of circadian rhythms

The circadian rhythms of many workers showed an overall increase in strength with age. As in previous studies, many individuals did not show clear circadian rhythms in the first few days after emerging from the pupa, but later developed robust circadian rhythms (Fig. 5). For the analyses below, we considered only the second and third trials during days 4–8; Trial 1 was not included because we monitored bees for only seven days, which is not sufficient for developmental analyses. In order to compare the development of circadian rhythms for bees subjected to the different treatments, we measured for each bee, the Power for consecutive 3-day sliding windows, and used these values to calculate the slope of the linear regression of Power as a function of age (days 4–9). The average slope was positive for all treatment groups in both trials, consistent with stronger circadian rhythms at later ages, but the slope was statistically significant only in the pooled analysis (using the Winer method of adding t 's) of the replacement therapy treatment group (Table 3).

The distribution of the slope values did not differ from normal distribution (Kolmogorov-Smirnov test for normality: $p = 0.241$ and $p = 0.591$ for Trials 2 and 3, respectively) allowing us to use t -tests for




Fig. 3. The effect of manipulating JH levels on the strength of circadian rhythms in locomotor activity. The Y-axis shows the Power, which we used as an index for the strength of circadian rhythms (arbitrary units). Treatments connected with a horizontal bar and asterisks are statistically different in a Mann-Whitney U test and after applying the Bonferroni correction for each trial, and in Mosteller-Bush method, in the pooled analyses ('All trials', bottom right panel). See Fig. 2 for additional details.




Fig. 4. The effect of manipulating JH levels on the free running period (τ). The Y-axis shows the circadian period length under constant conditions. Using Mann-Whitney U test we found that the τ did not differ between the Sham, P-I, and Replacement Therapy (P-I + JH) treatment groups in any of the three trials or in the pooled analysis. See Fig. 2 for additional details.

testing the influence of JH manipulation on the slope. These analyses suggested a faster rhythms development for replacement therapy treated bees (P-I + JH) compared to the Sham control (Table 4). There was a similar trend of faster development for the P-I + JH compared to the P-I treatment, but the p -value ($p = 0.051$) did not cross our statistical

significance threshold. These analyses are consistent with the hypothesis that JH influences the development of circadian rhythms in young bumble bee workers, although the evidence is relatively weak because the P-I treatment did not slow rhythm development relative to the sham treatment.




Fig. 5. Representative double-plot actograms for worker bees subjected to different treatments. Each panel shows locomotor activity for an individual bee monitored in constant laboratory environment. The y-axis depicts age of the bee; the height of the bars within each day corresponds to the level of locomotor activity in a 10-min bin. The x-axis shows the time of day in hours (double plotted). The control, sham, and replacement therapy (PI + JH) treated bees are from Trial 3, the precocene (P-I) treated bee is from Trial 2. Note the strengthening of circadian rhythms over time for the Control, Sham, and P-I treated bees.

Table 3

The slopes of regression lines for the strength of circadian rhythms (Power) as a function of age. The values are means \pm se, sample sizes in parentheses. The fourth column presents the combined *p*-value obtained by applying the Winer method. The rightmost column presents the combined effect size for each treatment. An asterisk denotes statistically significant difference in a one-tailed test.

Treatment	Trial 2	Trial 3	Combined	Effect Size
Sham	1.491 \pm 3.415 (20)	3.013 \pm 5.162 (22)	<i>p</i> = 0.247	0.113
P-I	4.129 \pm 2.474 (22)	5.789 \pm 10.351 (7)	<i>p</i> = 0.084	0.280
P-I + JH	8.679 \pm 2.524 (20)	17.311 \pm 5.118 (15)	<i>p</i> < 0.001 *	0.780

3.6. The influence of JH on sleep in individually isolated bees

The amount of sleep for bees in the Sham, P-I and replacement therapy treatment groups was similar (Table 5). For each bee, we calculated the mean proportion of time asleep (inactivity bouts of 5 min or more) during days 1–4. The distribution of these individual means did not differ from normal distribution (Kolmogorov-Smirnov test for normality: *p* = 0.812, *p* = 0.450 and *p* = 0.256 for Trials 1, 2 and 3,

Table 4

The influence of JH manipulation on the rate of development of circadian rhythms in locomotor activity. The table shows the *t* statistic of the comparisons between the mean slopes for each pair of treatments. A negative sign indicates that the mean of the first treatment is smaller than that of the second treatment in comparison. The fourth and rightmost columns summarized the combined *p*-value, and effect size, respectively. An asterisk denotes statistically significance difference in a one-tailed test.

Comparison	Trial 2	Trial 3	Combined	Effect Size
Sham vs. P-I	<i>t</i> ₄₀ = -0.634	<i>t</i> ₂₇ = -0.256	<i>p</i> = 0.271	0.087
Sham vs. P-I + JH	<i>t</i> ₃₈ = -1.692	<i>t</i> ₃₅ = -1.893	<i>p</i> = 0.007 *	0.393
P-I vs. P-I + JH	<i>t</i> ₄₀ = -1.285	<i>t</i> ₂₀ = -1.126	<i>p</i> = 0.051	0.346

Table 5

The influence of JH on the amount of sleep of individually isolated workers. The values are mean \pm se proportion of sleep during the day, with the sample size in parentheses. None of the comparison is statistically significant after Bonferroni correction (see text for details).

Treatment	Trial 1	Trial 2	Trial 3
Sham	0.366 \pm 0.024 (19)	0.401 \pm 0.031 (20)	0.291 \pm 0.026 (22)
P-I	0.429 \pm 0.034 (16)	0.350 \pm 0.037 (22)	0.322 \pm 0.056 (7)
P-I + JH	0.381 \pm 0.028 (24)	0.453 \pm 0.019 (20)	0.300 \pm 0.034 (15)

respectively). The Control and the Sham treatments did not differ statistically in any of the trials (independent samples *t*-test: Trial 1, $n_{\text{Control}} = 19$, mean \pm SE 0.38 ± 0.01 , $n_{\text{Sham}} = 19$, mean \pm SE = 0.37 ± 0.02 , $t_{36} = 0.329$, $p = 0.744$; Trial 2, $n_{\text{Control}} = 19$, mean \pm SE = 0.41 ± 0.009 , $n_{\text{Sham}} = 20$, mean \pm SE = 0.4 ± 0.03 , $t_{37} = -0.628$, $p = 0.534$; Trial 3, $n_{\text{Control}} = 19$, mean \pm SE = 0.31 ± 0.007 , $n_{\text{Sham}} = 22$, mean \pm SE = 0.29 ± 0.026 , $t_{39} = -0.505$, $p = 0.616$), as well as in a pooled analysis using the Winer method of adding *t*'s (combined $z = -0.452$, overall $p = 0.652$, overall effect size = 0.053). Thus, we used only the Sham treatment as the control group, and carried out pairwise comparisons between Sham, P-I and P-I + JH treatments, using independent samples *t*-tests. The statistics for this analysis are: Sham vs. P-I: Trial 1, $n_{\text{Sham}} = 19$, $n_{\text{P-I}} = 16$, $t_{33} = -1.530$, $p = 0.136$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I}} = 22$, $t_{40} = 1.041$, $p = 0.304$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I}} = 7$, $t_{27} = -0.565$, $p = 0.577$; overall $z = -0.589$, $p = 0.556$, effect size = 0.113. Sham vs. P-I + JH: Trial 1, $n_{\text{Sham}} = 19$, $n_{\text{P-I+JH}} = 24$, $t_{41} = -0.382$, $p = 0.704$; Trial 2, $n_{\text{Sham}} = 20$, $n_{\text{P-I+JH}} = 20$, $t_{31} = -1.443$, $p = 0.159$; Trial 3, $n_{\text{Sham}} = 22$, $n_{\text{P-I+JH}} = 15$, $t_{35} = -0.213$, $p = 0.832$; overall $z = -1.112$, $p = 0.266$, effect size = 0.186. P-I vs. P-I + JH: Trial 1, $n_{\text{P-I}} = 16$, $n_{\text{P-I+JH}} = 24$, $t_{38} = 1.095$, $p = 0.280$; Trial 2, $n_{\text{P-I}} = 22$, $n_{\text{P-I+JH}} = 20$, $t_{31} = -2.474$, $p = 0.019$; Trial 3, $n_{\text{P-I}} = 7$, $n_{\text{P-I+JH}} = 15$, $t_{20} = 0.354$, $p = 0.727$; overall $z = -0.570$, $p = 0.569$, effect size = 0.053. The treatment also did not have a statistically significant effect on the number of sleep bouts per day (Kruskal-Wallis *H* test, $H = 1.596$, $df = 3$, $p = 0.660$; Supplementary Fig. S4).

4. Discussion

JH is the major gonadotropin in many insects, including bumble bees in which it regulates physiological processes such as vitellogenesis, oogenesis, and wax production, and behaviors such as dominance and aggression. Little is known however, on the influences of JH on circadian rhythms and sleep. We used a pharmacological approach to test the hypothesis that JH influences circadian rhythms or sleep in young individually isolated bumble bee workers. Our measurements of ovarian activity suggest that our manipulations were effective; bees treated with P-I to reduce JH titers had inactive ovaries, and this effect was fully reverted by replacement therapy with JH-III, the natural JH of this species (Bloch et al., 2000, 1996). We found that bees with reduced JH levels have weaker circadian rhythms compared to control bees, an effect that was reverted by replacement therapy. The slope of circadian rhythm strengthening with age was steeper for bees of the replacement therapy treatment relative to the sham control, and a similar trend was seen relative to the P-I treatment. These developmental analyses are consistent with the hypothesis that developmental effects of JH may contribute to its influence on the strength of circadian rhythms, but the evidence is weak because there were no differences between the P-I and Sham treatments. To our knowledge, our results provide the strongest support for the hypothesis that JH augments circadian rhythms in an insect.

There is good evidence that gonadotropic hormones influence circadian rhythms in vertebrates (reviewed in Bedrosian et al., 2016; Gamble et al., 2014), which makes functional sense given that gonadotropins coordinate processes in many tissues involved in reproduction. The evidence for modulation of circadian processes by gonadotropic hormones in insects is more ambiguous (see Introduction; Bloch et al., 2013). Here we show that JH, the major gonadotropic hormone in the bumble bee *B. terrestris* (Pandey et al., 2020; Shpigler et al., 2014, 2016), influences the development and strength of circadian rhythms. Notably, these findings contrast with a study with honey bees in which allatectomy and replacement therapy manipulations did not affect circadian rhythms in locomotor activity in individually isolated workers (Bloch et al., 2002a). This incongruity is consistent with other evidence that JH has different functions in these two species of social bees (Bloch et al., 2002b; Hartfelder, 2000; Robinson and Vargo, 1997).

Our circadian analyses suggest that the most significant influence of JH manipulation was on the strength of circadian rhythms (Fig. 3), with

no effect on the endogenous circadian period (Fig. 4). Hormones may affect the strength of circadian rhythms in locomotor activity by several not mutually exclusive mechanisms. First, JH may act downstream of the circadian clock to regulate locomotor activity such that oscillations in circulating JH levels control daily changes in the level of locomotor activity. There is indeed evidence for circadian modulation of JH in insects, including honey bees (reviewed in Bloch et al., 2013). Moreover, in some insects, including honey bees, JH augments metabolic rates (Denlinger et al., 1984; Garcera et al., 1991; Sláma and Ilona, 1979; Sullivan et al., 2003). In some species such as the cockroach *B. germanica* there is even direct evidence that JH augments the overall level of activity (Lin and Lee, 1998). However, we did not find a reduction in overall locomotor activity in P-I treated bees or a significant increase in the replacement therapy (Table 2). JH similarly did not affect the level of locomotor activity in honey bees (Bloch et al., 2002a). The main inconsistency of this model with our findings is that it implies that insects with no JH will not show circadian rhythms. By contrast, we found that bees treated with P-I, that is assumed to disable the CA, nevertheless, show robust rhythms (although attenuated relative to control bees). Moreover, a single acute (not cyclic) treatment with JH fully recovered the rhythm attenuating effect of P-I. A second possible explanation for our findings is that oscillations in JH titers augment, but are not imperative for, circadian rhythms in locomotor activity. According to this model, central clocks in the brain concurrently influence both locomotor activity controlling centers and JH signaling (e.g., regulating JH biosynthesis in the CA; Bloch et al., 2013). Circadian rhythms in JH signaling in turn act on locomotor activity centers to augment exciting circadian rhythms in locomotor activity. As explained above for the first hypothesis, it is not easy to reconcile this proposed mechanism with our finding that a single acute treatment with JH at early age successfully reverted the rhythm attenuating effect of the P-I treatment. A third hypothesis states that JH has activation effects regulating functions in the central circadian network or in downstream output pathways controlling locomotion. This idea is consistent with studies with rodents showing that sex steroids affect various circadian parameters of locomotor activity, including the phase, FRP, amplitude (i.e., strength), and splitting of locomotor activity rhythms (Daan et al., 1975; Morin, 1980; Morin et al., 1977; reviewed in Hatcher et al., 2020). Many of these effects could be achieved by acute pharmacological treatments acting on hormone receptors indicating that the hormonal influence is activation rather than developmental (e.g., Karatsoreos et al., 2011; Model et al., 2015). In addition, treatments with steroid hormones were shown to regulate the expression of genes that are important for the generation or expression of circadian rhythms within specific cells, or by enhancing coupling within the timekeeping system (e.g., Karatsoreos et al., 2011; Nakamura et al., 2008, 2005; reviewed in Hatcher et al., 2020). Activation effects may also be mediated by the hormones acting on peripheral clocks or effector tissues (He et al., 2007; Sellix et al., 2004). Our findings that a single JH treatment at an early age produced long-lasting effects on the strength of circadian rhythms is not consistent with this hypothesis. The last hypothesis is that JH regulates the organization (development) of the circadian system. In mammals, it is well-established that sex steroid hormones have various organizational effects regulating the development of the circadian neural network and the expression of estrogen and androgen receptors which underlie sexual dimorphism in circadian activity (Hagenauer et al., 2011a, 2011b; Hummer et al., 2012; Melo et al., 2010; Royston et al., 2016; Sellix et al., 2013; for a recent review see Hatcher et al., 2020). Our findings that a single acute manipulation of JH levels shortly after eclosion resulted in relatively long-lasting effects on the strength of circadian rhythms best fit with the fourth hypothesis. The steeper age-related increase in the strength of circadian rhythms in bees subjected to replacement therapy provides additional support for this hypothesis (Tables 3 and 4, see also Fig. 5). However, this evidence is relatively weak because the P-I treated bees did not show slower rhythm development relative to the Sham treatment. A possible explanation for the difference between the slopes

of the P-I group compared to the replacement therapy but not to the sham treatment is that at the early age in which we treated with P-I, JH titers were low in the Sham group making it similar to the P-I treated bees. The amount of JH applied to the replacement therapy was very high compared to the natural levels at this age, and therefore producing a rapid response in the replacement therapy but not in the other groups. At later age, JH titers naturally increased in the Sham, but not in the P-I treated bees (because their CA is not functional) making the sham bees more similar to the replacement therapy at later ages, as seen in Fig. 3. It should be noted that we cannot exclude the possibility that activational effects (third hypothesis above) also account for at least some of the effects of JH on circadian rhythms.

Our findings that JH augments circadian rhythms in workers are puzzling because reproduction in *B. terrestris*, which is regulated by JH, is typically associated with activity around-the-clock and attenuated circadian rhythms in locomotor activity. Bumble bee gynes show robust circadian rhythms but later switch to activity around-the-clock when they establish colonies and reproduce (Eban-Rothschild et al., 2011). Similarly, nest workers, that typically have better-developed ovaries and higher JH titers (Shpigler et al., 2016; van Honk et al., 1981) are typically active around-the-clock in the presence of brood (Nagari et al., 2019; Yerushalmi et al., 2006). A plausible explanation for this apparent discrepancy is that we studied individually isolated bees, and rhythm manifestation in nurse bees is context dependent (Eban-Rothschild and Bloch, 2012b; Yerushalmi et al., 2006). Our finding that JH augments circadian rhythms in individually isolated workers does seem to reflect organizational effects on the development or integration of the circadian systems. The faster development of the circadian system in bees with high JH titers are not manifested in the colony because nurse workers and egg-laying queens are typically active around-the-clock in the presence of brood (Fig. 1.2 in Eban-Rothschild and Bloch, 2012b). Perhaps, a functional circadian clock system improves the coordination of internal processes related to reproduction. This idea is speculative at this stage. However, it is consistent with studies with mammals in which circadian clocks in the gonads and endocrine organs are imperative for fertility, and are sensitive to gonadotrophic hormones (reviewed in Sellix and Menaker, 2011). This may also be the case in male flies: *Drosophila melanogaster* flies with mutations in each of the clock genes *Period*, *Timeless*, *Cycle*, and *Clock*, show reduced fertility. The low-fertility phenotype was reversed in flies with rescued *Period* or *Timeless* function confirming the importance of these canonical clock genes. Further crosses between mutant and wild-type flies indicate that clock mutations compromised the fertility of both the male and female flies (Beaver et al., 2002). In males, peripheral circadian oscillators in the testis vas deferens complex are necessary for fertility (Beaver et al., 2002). However, in females, in which clock gene levels do not cycle in the ovary, the effects of clock gene mutations on fertility is due to non-circadian functions of the *Period* and *Timeless* in the ovary (Beaver et al., 2003). There is also evidence that JH interacts with clock genes in other insects. For example, in the linden bug, *Pyrrhocoris apterus*, JH regulates gene expression in the gut through interactions of its receptor Met with the circadian proteins Clock and Cycle (Bajgar et al., 2013).

JH manipulation did not affect sleep amount or structure. These findings contrast with studies in *Drosophila*, in which JH effects on sleep differ between male and female flies (Wu et al., 2018). Similarly, sex differences in sleep are reduced or absent in menopausal women and elderly men with reduced gonadal functions suggesting that gonadal hormones influence sleep in human. In rodents, gonadectomy in both sexes reduced or eliminated the majority of sex differences in sleep architecture and homeostasis further supporting the notion that gonadal hormones influence sleep (for a recent review see Hajali et al., 2019).

The interplay between gonadotrophic hormones, circadian clocks, and sleep is an important and little explored theme not only in bees, but also more generally in insects and vertebrates. Our study and the literature we review hint for species specific differences that should be addressed by hormone manipulation studies in diverse animal species.

Nevertheless, we believe that the available evidence is sufficient to suggest that hormonal state should be taken into account in studies on circadian rhythms and sleep.

CRediT authorship contribution statement

Atul Pandey: Conceptualization, Methodology, Data curation, Writing - original draft, Visualization, Formal analysis, Investigation.
Uzi Motro: Formal analysis, Visualization, Writing - review & editing.
Guy Bloch: Supervision, Conceptualization, Writing - review & editing, Project administration, Resources, Funding acquisition.

Declaration of competing interest

There is no conflict of interest for any of the authors.

Acknowledgments

This study was supported by grants from the US-Israel Binational Agricultural Research and Development Fund (BARD, number IS-4418-11, and IS-5077-18, to G.B.), the Israeli Science Foundation (ISF, grant No. 1274/15 to G.B.) and the Planning and Budgeting Committee (PBC) Fellowship Program for Outstanding Chinese and Indian Post-Doctoral Fellows (to A.P.).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.nbscr.2020.100056>.

Reference

- Albers, H.E., 1981. Gonadal hormones organize and modulate the circadian system of the rat. *Am. J. Physiol.* <https://doi.org/10.1152/ajpregu.1981.241.1.r62>.
- Amsalem, E., Teal, P., Grozinger, C.M., Hefetz, A., 2014. Precocene-I inhibits juvenile hormone biosynthesis, ovarian activation, aggression and alters sterility signal production in bumble bee (*Bombus terrestris*) workers. *J. Exp. Biol.* 217, 3178–3185. <https://doi.org/10.1242/jeb.107250>.
- Bajgar, A., Jindra, M., Dolezel, D., 2013. Autonomous regulation of the insect gut by circadian genes acting downstream of juvenile hormone signaling. *Proc. Natl. Acad. Sci. U. S. A.* 110, 4416–4421. <https://doi.org/10.1073/pnas.1217060110>.
- Beaver, L.M., Gvakharia, B.O., Vollintine, T.S., Hege, D.M., Stanewsky, R., Giebultowicz, J.M., 2002. Loss of circadian clock function decreases reproductive fitness in males of *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U. S. A.* 99, 2134–2139. <https://doi.org/10.1073/pnas.032426699>.
- Beaver, L.M., Rush, B.L., Gvakharia, B.O., Giebultowicz, J.M., 2003. Noncircadian regulation and function of clock genes period and timeless in oogenesis of *Drosophila melanogaster*. *J. Biol. Rhythm.* 18, 463–472. <https://doi.org/10.1177/0748730403259108>.
- Bedrosian, T.A., Fonken, L.K., Nelson, R.J., 2016. Endocrine effects of circadian disruption. *Annu. Rev. Physiol.* 78, 109–131. <https://doi.org/10.1146/annurev-physiol-021115-105102>.
- Bloch, G., 2010. The social clock of the honeybee. *J. Biol. Rhythm.* 25, 307–317. <https://doi.org/10.1177/0748730410380149>.
- Bloch, G., Borst, D.W., Huang, Z.-Y., Robinson, G.E., Cnaani, J., Hefetz, A., 2000. Juvenile hormone titers, juvenile hormone biosynthesis, ovarian development and social environment in *Bombus terrestris*. *J. Insect Physiol.* 46, 47–57. [https://doi.org/10.1016/S0022-1910\(99\)00101-8](https://doi.org/10.1016/S0022-1910(99)00101-8).
- Bloch, G., Borst, D.W., Huang, Z.-Y., Robinson, G.E., Hefetz, A., 1996. Effects of social conditions on juvenile hormone mediated reproductive development in *Bombus terrestris* workers. *Physiol. Entomol.* 21, 257–267. <https://doi.org/10.1111/j.1365-3032.1996.tb00863.x>.
- Bloch, G., Hazan, E., Rafaeli, A., 2013. Circadian rhythms and endocrine functions in adult insects. *J. Insect Physiol.* 59, 56–69. <https://doi.org/10.1016/j.jinsphys.2012.10.012>.
- Bloch, G., Meshi, A., 2007. Influences of octopamine and juvenile hormone on locomotor behavior and period gene expression in the honeybee, *Apis mellifera*. *J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol.* 193, 181–199. <https://doi.org/10.1007/s00359-006-0179-5>.
- Bloch, G., Sullivan, J.P., Robinson, G.E., 2002a. Juvenile hormone and circadian locomotor activity in the honey bee *Apis mellifera*. *J. Insect Physiol.* 48, 1123–1131. [https://doi.org/10.1016/S0022-1910\(02\)00205-6](https://doi.org/10.1016/S0022-1910(02)00205-6).
- Bloch, G., Wheeler, D.E., Robinson, G.E., 2002b. Endocrine influences on the organization of insect societies. In: Pfaff, Donald W., Arnold, Arthur P., et al. (Eds.), *Hormones, Brain and Behavior*. Academic Press, pp. 195–235.

- Daan, S., Damassa, D., Pittendrigh, C.S., Smith, E.R., 1975. An effect of castration and testosterone replacement on a circadian pacemaker in mice (*Mus musculus*). Proc. Natl. Acad. Sci. U. S. A. 72, 3744–3747. <https://doi.org/10.1073/pnas.72.9.3744>.
- Denlinger, D., Shukla, M., Faustini, D., 1984. Juvenile hormone involvement in pupal diapause of the flesh fly *Sarcophaga Crassipalpis*: Regulation of infradian cycles of O₂ consumption. J. Exp. Biol. 109, 191–199.
- Eban-Rothschild, A., Belluci, S., Bloch, G., 2011. Maternity-related plasticity in circadian rhythms of bumble-bee queens. Proc. R. Soc. B Biol. Sci. 278, 3510–3516. <https://doi.org/10.1098/rspb.2011.0579>.
- Eban-Rothschild, A., Bloch, G., 2012. Circadian rhythms and sleep in honey bees. In: Galizia, G., Eisenhardt, D., Giurfa, M. (Eds.), Honeybee Neurobiology and Behavior. Springer Nature, pp. 31–45. https://doi.org/10.1007/978-94-007-2099-2_3.
- Eban-Rothschild, A.D., Bloch, G., 2008. Differences in the sleep architecture of forager and young honeybees (*Apis mellifera*). J. Exp. Biol. 211, 2408–2416. <https://doi.org/10.1242/jeb.016915>.
- Eban-Rothschild, A., Bloch, G., 2012. Social influences on circadian rhythms and sleep in insects. In: Sokolowski, Marla, Goodwin, Stephen (Eds.), Advances in Genetics, 77. Elsevier, pp. 1–32. <https://doi.org/10.1016/B978-0-12-387687-4.00001-5>.
- Gamble, K.L., Berry, R., Frank, S.J., Young, M.E., 2014. Circadian clock control of endocrine factors. Nat. Rev. Endocrinol. 10, 466–475. <https://doi.org/10.1038/nrendo.2014.78>.
- Garcera, M.D., Ibanez, P., Martinez, R., Cunat, P., 1991. Variations induced by juvenile hormone and precocene II in the metabolic rate of three species of insects. Rev. Esp. Fisiol. 47, 31–36.
- Hagenauer, M.H., King, A.F., Possidente, B., McGinnis, M.Y., Lumia, A.R., Peckham, E. M., Lee, T.M., 2011a. Changes in circadian rhythms during puberty in *Rattus norvegicus*: developmental time course and gonadal dependency. Horm. Behav. 60, 46–57. <https://doi.org/10.1016/j.yhbeh.2011.03.001>.
- Hagenauer, M.H., Ku, J.H.Y., Lee, T.M., 2011b. Chronotype changes during puberty depend on gonadal hormones in the slow-developing rodent, *Octodon degus*. Horm. Behav. 60, 37–45. <https://doi.org/10.1016/j.yhbeh.2011.02.004>.
- Hajali, V., Andersen, M.L., Negah, S.S., Sheibani, V., 2019. Sex differences in sleep and sleep loss-induced cognitive deficits: the influence of gonadal hormones. Horm. Behav. 108, 50–61. <https://doi.org/10.1016/j.yhbeh.2018.12.013>.
- Hartfelder, K., 2000. Insect juvenile hormone: From “status quo” to high society. Braz. J. Med. Biol. Res. 33, 157–177. <https://doi.org/10.1590/S0100-879X2000000200003>.
- Hastings, M., O'Neill, J.S., Maywood, E.S., 2007. Circadian clocks: Regulators of endocrine and metabolic rhythms. J. Endocrinol. 195, 187–198. <https://doi.org/10.1677/JOE-07-0378>.
- Hatcher, K.M., Royston, S.E., Mahoney, M.M., 2020. Modulation of circadian rhythms through estrogen receptor signaling. Eur. J. Neurosci. 51, 217–228. <https://doi.org/10.1111/ejn.14184>.
- He, P.J., Hirata, M., Yamauchi, N., Hashimoto, S., Hattori, M.A., 2007. Gonadotropic regulation of circadian clockwork in rat granulosa cells. Mol. Cell. Biochem. 302, 111–118. <https://doi.org/10.1007/s11010-007-9432-7>.
- Huang, Z.Y., Robinson, G.E., Tobe, S.S., Yagi, K.J., Strambi, C., Strambi, A., Stay, B., 1991. Hormonal regulation of behavioural development in the honey bee is based on changes in the rate of juvenile hormone biosynthesis. J. Insect Physiol. 37, 733–741. [https://doi.org/10.1016/0022-1910\(91\)90107-B](https://doi.org/10.1016/0022-1910(91)90107-B).
- Hummer, D.L., Peckham, E.M., Lee, T.M., 2012. Estradiol acts during a post-pubertal sensitive period to shorten free-running circadian period in male *Octodon degus*. Eur. J. Neurosci. 36, 3051–3058. <https://doi.org/10.1111/j.1460-9568.2012.08228.x>.
- Iwahana, E., Karatsoreos, I., Shibata, S., Silver, R., 2008. Gonadectomy reveals sex differences in circadian rhythms and suprachiasmatic nucleus androgen receptors in mice. Horm. Behav. 27, 432–443. <https://doi.org/10.1016/j.yhbeh.2007.11.014>.
- Jechura, T.J., Walsh, J.M., Lee, T.M., 2000. Testicular hormones modulate circadian rhythms of the diurnal rodent, *Octodon degus*. Horm. Behav. 38, 243–249. <https://doi.org/10.1006/hbeh.2000.1624>.
- Karatsoreos, I.N., Butler, M.P., Lesauter, J., Silver, R., 2011. Androgens modulate structure and function of the suprachiasmatic nucleus brain clock. Endocrinology 152, 1970. <https://doi.org/10.1210/en.2010-1398>.
- Karatsoreos, I.N., Wang, A., Sasanian, J., Silver, R., 2007. A role for androgens in regulating circadian behavior and the suprachiasmatic nucleus. Endocrinology 148, 5487–5495. <https://doi.org/10.1210/en.2007-0775>.
- Klarsfeld, A., Leloup, J.C., Rouyer, F., 2003. Circadian rhythms of locomotor activity in *Drosophila*. Behav. Process. 64, 161–175. [https://doi.org/10.1016/S0376-6357\(03\)00133-5](https://doi.org/10.1016/S0376-6357(03)00133-5).
- Kriegsfeld, et al., 2002. Circadian rhythms in the endocrine system. In: Hormones, Brain and Behavior. Academic Press, pp. 33–91. <https://doi.org/10.1016/B978-012532104-4/50020-2>.
- Lee, H.-J., Wu, Y.-L., 1994. Mating effects on the feeding and locomotion of the German cockroach, *Blattella germanica*. Physiol. Entomol. 19, 39–45. <https://doi.org/10.1111/j.1365-3032.1994.tb01071.x>.
- Lin, T.M., Lee, H.J., 1998. Parallel control mechanisms underlying locomotor activity and sexual receptivity of the female German cockroach, *Blattella germanica* (L.). J. Insect Physiol. 44, 1039–1051. [https://doi.org/10.1016/S0022-1910\(98\)00069-9](https://doi.org/10.1016/S0022-1910(98)00069-9).
- Melo, P.R., Belsio, A.S., Menezes, A.A.L., Azevedo, C.V.M., 2010. Influence of seasonality on circadian motor activity rhythm in common marmosets during puberty. Chronobiol. Int. 27, 1420–1437. <https://doi.org/10.3109/07420528.2010.501416>.
- Michener, C.D., 1974. The social behavior of the bees. In: The Social Behavior of the Bees. Harvard University Press, Cambridge, MA, pp. 1–418.
- Model, Z., Butler, M.P., LeSauter, J., Silver, R., 2015. Suprachiasmatic nucleus as the site of androgen action on circadian rhythms. Horm. Behav. 73, 1–7. <https://doi.org/10.1016/j.yhbeh.2015.05.007>.
- Morin, L.P., 1980. Effect of ovarian hormones on synchrony of hamster circadian rhythms. Physiol. Behav. 24, 741–749. [https://doi.org/10.1016/0031-9384\(80\)90406-0](https://doi.org/10.1016/0031-9384(80)90406-0).
- Morin, L.P., Fitzgerald, K.M., Zucker, I., 1977. Estradiol shortens the period of hamster circadian rhythms. Science 196, 305–307. <https://doi.org/10.1126/science.557840>.
- Nagari, M., Gera, A., Jonsson, S., Bloch, G., 2019. Bumble bee workers give up sleep to care for offspring that are not their own. Curr. Biol. 29, 3488–3493. <https://doi.org/10.1016/j.cub.2019.07.091>.
- Nakamura, T.J., Moriya, T., Inoue, S., Shimazoe, T., Watanabe, S., Ebihara, S., Shinohara, K., 2005. Estrogen differentially regulates expression of Per 1 and Per 2 genes between central and peripheral clocks and between reproductive and nonreproductive tissues in female rats. J. Neurosci. Res. 82, 622–630. <https://doi.org/10.1002/jnr.20677>.
- Nakamura, T.J., Sellix, M.T., Menaker, M., Block, G.D., 2008. Estrogen directly modulates circadian rhythms of PER2 expression in the uterus. Am. J. Physiol. Endocrinol. Metab. 295, E1025–E1031. <https://doi.org/10.1152/ajpendo.90392.2008>.
- Neumann, A.M., Schmidt, C.X., Brockmann, R.M., Oster, H., 2019. Circadian regulation of endocrine systems. Auton. Neurosci. Basic Clin. 216, 1–8. <https://doi.org/10.1016/j.autneu.2018.10.001>.
- Pandey, A., Motro, U., Bloch, G., 2020. Juvenile hormone interacts with multiple factors to modulate aggression and dominance in groups of orphan bumble bee (*Bombus terrestris*) workers. Horm. Behav. 117, 104602. <https://doi.org/10.1016/j.yhbeh.2019.104602>.
- Peitsch, D., Fietz, A., Hertel, H., de Souza, J., Ventura, D., Menzel, R., 1992. The spectral input systems of hymenopteran insects and their receptor-based colour vision. J. Comp. Physiol. 170, 23–40.
- Robinson, G.E., Vargo, E.L., 1997. Juvenile hormone in adult eusocial hymenoptera: gonadotropin and behavioral pacemaker. Arch. Insect Biochem. Physiol. 35, 559–583.
- Rosenthal, R., 1978. Combining results of independent studies. Psychol. Bull. 85, 185–193. <https://doi.org/10.1037/0033-2909.85.1.185>.
- Royston, S.E., Bunick, D., Mahoney, M.M., 2016. Oestradiol exposure early in life programs daily and circadian activity rhythms in adult mice. J. Neuroendocrinol. 28. <https://doi.org/10.1111/jne.12335>.
- Sellix, M.T., Egli, M., Henderson, R.P., Freeman, M.E., 2004. Ovarian steroid hormones modulate circadian rhythms of neuroendocrine dopaminergic neuronal activity. Brain Res. 1005, 164–181. <https://doi.org/10.1016/j.brainres.2004.01.049>.
- Sellix, M.T., Menaker, M., 2011. Circadian clocks in mammalian reproductive physiology: effects of the “other” biological clock on fertility. Discov. Med. 11, 273–281.
- Sellix, M.T., Murphy, Z.C., Menaker, M., 2013. Excess androgen during puberty disrupts circadian organization in female rats. Endocrinology 154, 1636–1647. <https://doi.org/10.1210/en.2012-2066>.
- Shaw, P.J., Cirelli, L., Greensoan, R.J., Tononi, G., 2000. Correlates of sleep and waking in *Drosophila melanogaster*. Science 287, 1834–1837. <https://doi.org/10.1126/science.287.5459.1834>.
- Shemesh, Y., Cohen, M., Bloch, G., 2007. Natural plasticity in circadian rhythms is mediated by reorganization in the molecular clockwork in honeybees. FASEB J. 21, 2304–2311. <https://doi.org/10.1096/fj.06-8032com>.
- Shepard, M., Keeley, L.L., 1972. Circadian rhythmicity and capacity for enforced activity in the cockroach, *Blaberus discoidalis*, after cardiectomy-allatectomy. J. Insect Physiol. 18, 595–601. [https://doi.org/10.1016/0022-1910\(72\)90089-3](https://doi.org/10.1016/0022-1910(72)90089-3).
- Shpigler, H., Amsalem, E., Huang, Z.Y., Cohen, M., Siegel, A.J., Hefetz, A., Bloch, G., 2014. Gonadotropic and physiological functions of juvenile hormone in bumblebee (*Bombus terrestris*) workers. PLoS One 9, e100650. <https://doi.org/10.1371/journal.pone.0100650>.
- Shpigler, H.Y., Siegel, A.J., Huang, Z.Y., Bloch, G., 2016. No effect of juvenile hormone on task performance in a bumblebee (*Bombus terrestris*) supports an evolutionary link between endocrine signaling and social complexity. Horm. Behav. 85, 67–75. <https://doi.org/10.1016/j.yhbeh.2016.08.004>.
- Sláma, K., Ilona, K.S., 1979. Hypermetabolic response induced by juvenile hormone analogues in an insect. Zeitschrift für Naturforsch. - Sect. C J. Biosci. 34, 599–607. <https://doi.org/10.1515/znc-1979-7-818>.
- Sullivan, J.P., Fahrback, S.E., Harrison, J.F., Capaldi, E.A., Fewell, J.H., Robinson, G.E., 2003. Juvenile hormone and division of labor in honey bee colonies: effects of allatectomy on flight behavior and metabolism. J. Exp. Biol. 206, 2287–2296. <https://doi.org/10.1242/jeb.00432>.
- Sullivan, J.P., Jassim, O., Fahrback, S.E., Robinson, G.E., 2000. Juvenile hormone paces behavioral development in the adult worker honey bee. Horm. Behav. 37, 1–14. <https://doi.org/10.1006/hbeh.1999.1552>.
- Tsang, A.H., Barclay, J.L., Oster, H., 2013. Interactions between endocrine and circadian systems. J. Mol. Endocrinol. 52, 1–16. <https://doi.org/10.1530/JME-13-0118>.
- van Honk, C.G.J., Röseler, P.F., Velthuis, H.H.W., Hoogeveen, J.C., 1981. Factors influencing the egg laying of workers in a captive *Bombus terrestris* colony. Behav. Ecol. Sociobiol. 9, 9–14. <https://doi.org/10.1007/BF00299847>.
- Walker, W.F., 1977. Juvenile hormone: evidence for a role in the feeding rhythm of *Oncopeltus fasciatus* (Dallas). Experientia 33, 1539–1541. <https://doi.org/10.1007/BF01918859>.
- Woodard, H.D., Rankin, M.A., 1980. Effect of precocene II on circadian rhythms of feeding and mating behavior in the milkweed bug, *Oncopeltus fasciatus*. Experientia 36, 554–555. <https://doi.org/10.1007/BF01965796>.

- Wu, B., Ma, L., Zhang, E., Du, J., Liu, S., Price, J., Li, S., Zhao, Z., 2018. Sexual dimorphism of sleep regulated by juvenile hormone signaling in *Drosophila*. *PLoS Genet.* 14, e1007318 <https://doi.org/10.1371/journal.pgen.1007318>.
- Yerushalmi, S., Bodenheimer, S., Bloch, G., 2006. Developmentally determined attenuation in circadian rhythms links chronobiology to social organization in bees. *J. Exp. Biol.* 209, 1044–1051. <https://doi.org/10.1242/jeb.02125>.
- Zhao, Y., Zhang, K., Fent, K., 2018. Regulation of zebrafish (*Danio rerio*) locomotor behavior and circadian rhythm network by environmental steroid hormones. *Environ. Pollut.* 232, 422–429. <https://doi.org/10.1016/j.envpol.2017.09.057>.