

OPEN

Treatment Effect of Drug-Coated Balloons Is Durable to 3 Years in the Femoropopliteal Arteries Long-Term Results of the IN.PACT SFA Randomized Trial

Peter A. Schneider, MD; John R. Laird, MD; Gunnar Tepe, MD; Marianne Brodmann, MD; Thomas Zeller, MD; Dierk Scheinert, MD; Christopher Metzger, MD; Antonio Micari, MD; Ravish Sachar, MD; Michael R. Jaff, DO; Hong Wang, MD, MH; Melissa S. Hasenbank, PhD; Prakash Krishnan, MD; for the IN.PACT SFA Trial Investigators

Background—Randomized controlled trials have reported favorable 1-year outcomes with drug-coated balloons (DCBs) for the treatment of symptomatic peripheral arterial disease when compared with standard percutaneous transluminal angioplasty (PTA). Evidence remains limited on the durability of the treatment effect with DCBs in the longer term.

Methods and Results—IN.PACT SFA is a single-blind, randomized trial (Randomized Trial of IN.PACT Admiral Paclitaxel-Coated Percutaneous Transluminal Angioplasty [PTA] Balloon Catheter vs Standard PTA for the Treatment of Atherosclerotic Lesions in the Superficial Femoral Artery [SFA] and/or Proximal Popliteal Artery [PPA]) that enrolled 331 patients with symptomatic (Rutherford 2–4) femoropopliteal lesions up to 18 cm in length. Patients were randomized 2:1 to receive treatment with DCB or PTA. The 36-month assessments included primary patency, freedom from clinically driven target lesion revascularization, major adverse events, and functional outcomes. At 36 months, primary patency remained significantly higher among patients treated with DCB compared with PTA (69.5% versus 45.1%; log rank $P<0.001$). The rates of clinically driven target lesion revascularization were 15.2% and 31.1% ($P=0.002$) for the DCB and PTA groups, respectively. Functional outcomes were similarly improved between treatment groups even though subjects in the DCB group required significantly fewer reinterventions versus those in the PTA group ($P<0.001$ for target lesion revascularization, $P=0.001$ for target vessel revascularization). There were no device- or procedure-related deaths as adjudicated by an independent Clinical Events Committee.

Conclusions—Three-year results demonstrate a durable and superior treatment effect among patients treated with DCB versus standard PTA, with significantly higher primary patency and lower clinically driven target lesion revascularization, resulting in similar functional improvements with reduced need for repeat interventions.

Clinical Trial Registration—URL: <http://www.clinicaltrials.gov>. Unique identifiers: NCT01175850 for IN.PACT SFA phase I in the European Union and NCT01566461 for IN.PACT SFA phase II in the United States. (*Circ Cardiovasc Interv.* 2018;11:e005891. DOI: 10.1161/CIRCINTERVENTIONS.117.005891.)

Key Words: angioplasty ■ peripheral arterial disease ■ target lesion revascularization

Endovascular intervention has become the primary mode of revascularization for patients with symptomatic femoropopliteal peripheral artery disease. Multiple modalities of treatment exist; however, the mainstay is percutaneous transluminal angioplasty (PTA) and implantation of a bare metal stent (BMS).¹ Angioplasty, though effective in luminal gain, has been associated with restenosis rate of up to 60% at 12 months.^{2,3} Although implantation of a BMS has been shown

to reduce this restenosis rate by nearly half,²⁻⁷ BMSs are associated with inherent problems, including in-stent restenosis, thrombosis, and stent fracture.⁸⁻¹⁰

See Editorial by Sethi and Parikh

To overcome the limitations of standard interventions such as PTA or BMSs, drug-coated balloons (DCBs) were developed in hopes of improved patency over the long term,

Received September 6, 2017; accepted November 29, 2017.

From the Hawaii Permanente Medical Group, Kaiser Foundation Hospital, Honolulu, HI (P.A.S.); Adventist Heart and Vascular Institute, St Helena Hospital, CA (J.R.L.); Department of Diagnostic and Interventional Radiology, RoMed Klinikum, Rosenheim, Germany (G.T.); Department of Internal Medicine, Division of Angiology, Medical University, Graz, Austria (M.B.); Angiology Division, Universitäts-Herzzentrum Freiburg–Bad Krozingen, Germany (T.Z.); Department of Angiology, University Hospital Leipzig, Germany (D.S.); Wellmont Holston Valley Medical Center, Kingsport, TN (C.M.); GVM Care and Research, Maria Cecilia Hospital, Cotignola, Italy (A.M.); North Carolina Heart and Vascular, UNC-REX Health Care, Raleigh, NC (R.S.); Newton-Wellesley Hospital, MA (M.R.J.); Medtronic, Santa Rosa, CA (H.W., M.S.H.); and Icahn School of Medicine, Mount Sinai Medical Center, New York, NY (P.K.).

The Data Supplement is available at <http://circinterventions.ahajournals.org/lookup/suppl/doi:10.1161/CIRCINTERVENTIONS.117.005891/-/DC1>.

Correspondence to Peter A. Schneider, MD, Hawaii Permanente Medical Group, Kaiser Foundation Hospital, 3288 Moanalua Rd, Honolulu, HI 96819. E-mail peterschneidermd@aol.com

© 2018 The Authors. *Circulation: Cardiovascular Interventions* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution Non-Commercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited, the use is noncommercial, and no modifications or adaptations are made.

Circ Cardiovasc Interv is available at <http://circinterventions.ahajournals.org>

DOI: 10.1161/CIRCINTERVENTIONS.117.005891

WHAT IS KNOWN

- There are limited long-term data on the durability of drug-coated balloon treatment effect in patients with symptomatic femoropopliteal peripheral artery disease.

WHAT THE STUDY ADDS

- Three-year results from the IN.PACT Global Study (Randomized Trial of IN.PACT Admiral Paclitaxel-Coated Percutaneous Transluminal Angioplasty [PTA] Balloon Catheter vs Standard PTA for the Treatment of Atherosclerotic Lesions in the Superficial Femoral Artery [SFA] and/or Proximal Popliteal Artery [PPA]) show that the effectiveness of paclitaxel drug-coated balloon treatment is durable up to 3 years and is associated with significantly higher primary patency and a lower rate of repeat interventions compared with percutaneous transluminal angioplasty.

without the disadvantages of a major metal implant such as a stent. DCBs are covered with an antiproliferative agent, paclitaxel, using urea as an excipient to facilitate the transfer of paclitaxel to the inner vessel surface on balloon inflation. After angioplasty, paclitaxel can persist in the vessel wall for up to 180 days in experimental models.¹¹ The sustained presence of paclitaxel provides continuous antiproliferative activity that inhibits neointimal hyperplasia, which is a major contributing factor to restenosis after angioplasty.

Clinical studies have demonstrated the short-term safety and effectiveness of paclitaxel-coated DCBs for the treatment of femoropopliteal peripheral artery lesions,^{12–19} but there are limited long-term data beyond 2 years on the durability of treatment effect. This is the first report of a large randomized clinical trial to evaluate the long-term safety and effectiveness of treatment with a paclitaxel DCB in femoropopliteal lesions.

Methods

The data, analytic methods, and study materials may be made available to other researchers on request from the sponsor.

Study Design

The IN.PACT SFA trial (Randomized Trial of IN.PACT Admiral Paclitaxel-Coated Percutaneous Transluminal Angioplasty [PTA] Balloon Catheter vs Standard PTA for the Treatment of Atherosclerotic Lesions in the Superficial Femoral Artery [SFA] and/or Proximal Popliteal Artery [PPA]) was a prospective, multicenter, multinational, randomized, single-blind trial that was designed to evaluate the safety and effectiveness of a paclitaxel DCB (IN.PACT Admiral, Medtronic Inc, Dublin, Ireland) versus standard PTA balloons in the treatment of subjects with symptomatic superficial femoral artery and proximal popliteal artery disease. The IN.PACT SFA trial was conducted in 2 consecutive phases (IN.PACT SFA phase I in the European Union and IN.PACT SFA phase II in the United States). A list of investigators and clinical sites is given in Table I in the [Data Supplement](#). The details of IN.PACT SFA trial design have been previously reported, as have outcomes through 2 years.^{18,19}

Table 1. Baseline Subject and Procedural Characteristics*

| Characteristic | DCB (Subjects=220/ Lesions=221) | PTA (Subjects=111/ Lesions=113) | P Value |
|------------------------------|---------------------------------------|---------------------------------------|---------|
| Age, y | 67.5±9.5 | 68.0±9.2 | 0.612 |
| Male | 65.0 (143/220) | 67.6 (75/111) | 0.713 |
| Diabetes mellitus | 40.5 (89/220) | 48.6 (54/111) | 0.161 |
| Hypertension | 91.4 (201/220) | 88.3 (98/111) | 0.431 |
| Hyperlipidemia | 84.5 (186/220) | 82.0 (91/111) | 0.637 |
| Current smoker | 38.6 (85/220) | 36.0 (40/111) | 0.719 |
| ABI/TBI† | 0.769±0.228 | 0.744±0.189 | 0.308 |
| Rutherford Clinical Category | | | 0.898 |
| 2 | 37.7 (83/220) | 37.8 (42/111) | |
| 3 | 57.3 (126/220) | 55.9 (62/111) | |
| 4 | 5.0 (11/220) | 5.4 (6/111) | |
| 5 | 0.0 (0/220) | 0.9 (1/111)‡ | |
| Lesion length, cm | 8.94±4.89 | 8.81±5.12 | 0.815 |
| Total occlusions | 25.8 (57/221) | 19.5% (22/113) | 0.222 |
| Calcification | 59.3 (131/221) | 58.4 (66/113) | 0.907 |
| Severe calcification | 8.1 (18/1221) | 6.2 (7/113) | 0.662 |
| Dissections | | | 0.360 |
| 0 | 36.2 (80/221) | 38.9 (44/113) | |
| A–C | 63.8 (141/221) | 60.2 (68/113) | |
| D–F | 0.0 (0/221) | 0.9 (1/113) | |
| Provisional stenting | 7.3 (16/220) | 12.6 (14/111) | 0.110 |

ABI indicates ankle-brachial index; DCB, drug-coated balloon; PTA, percutaneous transluminal angioplasty; and TBI, toe-brachial index.

*Values are mean±SD or % (n/N).

†TBI allowed in cases of incompressible vessels in IN.PACT SFA phase II.

‡One subject in the PTA group was Rutherford Clinical Category 5 despite application of Rutherford Category 2 to 4 as inclusion criteria for enrollment.

Subjects were randomized in a 2:1 fashion into a DCB or PTA group. Eligible subjects had moderate to severe intermittent claudication or ischemic rest pain (Rutherford Clinical Category 2–4); stenosis of 70% to 99% with lesion lengths between 4 and 18 cm, or a complete occlusion with lengths of ≤10 cm involving the superficial femoral and proximal popliteal arteries and were required to have successful predilatation of the lesion before enrollment. Details on the use of preprocedure and periprocedure anticoagulant and antiplatelet medications are given in Table II in the [Data Supplement](#).

Informed consent was obtained from all subjects before enrollment. An institutional review board or ethics committee approved all protocols at each trial site. The trial was conducted in accordance with the Declaration of Helsinki, good clinical practice guidelines, and applicable laws as specified by all relevant governmental bodies.

A Clinical Events Committee reviewed and adjudicated all major adverse events through the 36-month follow-up period. The Clinical Events Committee included interventional and noninterventional clinicians with pertinent expertise who were not participants in the study and did not have conflicts of interest. The Clinical Events Committee used a set of criteria that was specifically designed to classify major adverse events in the study. Independent core laboratories (duplex ultrasonography: VasCore, Boston, MA; angiography: SynvaCor, Springfield, IL) analyzed all procedural and follow-up images through 36 months. The independent core laboratories and Clinical Events Committee were blinded to treatment assignments.




Figure 1. Subject flow in the IN.PACT SFA trial through 36 months. Three hundred thirty-one subjects were randomized 2:1 into groups that received percutaneous transluminal angioplasty (PTA) with a paclitaxel drug-coated balloon (DCB) or a standard uncoated balloon (PTA). Subjects are being followed for 5 years, and the results of intent-to-treat analyses have been previously reported for 12 months¹⁸ and 24 months.¹⁹

End Point Definitions

Primary patency was defined as freedom from clinically driven target lesion revascularization (CD-TLR) and freedom from restenosis (duplex ultrasonography peak systolic velocity ratio ≤ 2.4) and was analyzed through 36 months. CD-TLR was defined as reintervention at the target lesion because of symptoms or a decrease in ankle-brachial index by $\geq 20\%$ or >0.15 when compared with postprocedure baseline ankle-brachial index. The primary composite safety end point was freedom from device- and procedure-related death through 30 days and freedom from target limb major amputation and clinically driven (CD) target vessel revascularization through 36 months.

The rate of major adverse events (death from any cause, CD target vessel revascularization, target limb major amputation, and thrombosis) was evaluated at 36 months. Thrombosis was defined as occlusion because of thrombus formation, confirmed by sudden onset of symptoms and documented by duplex ultrasonography and angiography. Additional end points included the rate of each individual component of the major adverse event composite, primary sustained clinical improvement (defined as freedom from target limb major amputation, target vessel revascularization, and increase in Rutherford Clinical Category), all TLR, and ankle-brachial index (toe-brachial index allowed in cases of incompressible vessels in IN.PACT SFA phase II) at 36 months. Functional assessments included evaluation of walking capacity with the Walking Impairment Questionnaire (WIQ) and the 6-minute walk test (IN.PACT SFA phase II only).

Statistical Analysis

Analyses were based on the intent-to-treat principle. Baseline demographics and clinical characteristics were summarized on a subject basis; lesion characteristics were summarized on a lesion basis. For baseline characteristics, continuous variables were described as mean \pm SD and were compared by Student *t* tests; dichotomous and categorical variables were described as counts and proportions and were compared by the Fisher exact test or Cochran–Mantel–Haenszel

modified ridit scores, respectively. Outcome analyses were performed at a subject level. The Kaplan–Meier method was used to evaluate time to event data for primary patency and CD-TLR over the 36-month follow-up period. The difference in the survival curves between treatment groups was assessed using the log-rank test. For other outcomes, the Fisher exact test was used for binary outcomes, and the Wilcoxon rank-sum test or Student *t* test was used for continuous outcomes. The level of statistical significance was set at $P < 0.05$ with no correction for multiple comparisons. For event rates that were expressed as a proportion, the number of subjects with an event within 1080 days was the numerator and the number of subjects with an event or at least 1050 days of clinical follow-up was the denominator. For assessment of clinical characteristics at 36 months, subjects were required to have data at both baseline and 36 months to assess any changes from baseline. Statistical analyses were performed using SAS (SAS Institute, NC) version 9.4.

Results

Three hundred thirty-one subjects were randomized to receive treatment with DCB ($n=220$) or PTA ($n=111$). Demographic, clinical, and lesion characteristics were well matched between treatment groups at baseline (Table 1). The mean lesion length was 8.94 ± 4.89 cm in the DCB group and 8.81 ± 5.12 cm in the PTA group ($P=0.815$). Provisional stents were implanted in 7.3% of subjects in the DCB group and 12.6% of subjects in the PTA group ($P=0.110$; Table 1). Details on the reasons for provisional stenting are reported in Table III in the [Data Supplement](#). Through 36 months of follow-up, 11.4% of DCB subjects and 9.0% of PTA subjects withdrew from the trial, and 10.7% of DCB subjects and 1.9% of PTA subjects died. Of the remaining 173 DCB subjects eligible for 36-month evaluation,




Figure 2. Durability of effect after treatment with a paclitaxel drug-coated balloon (DCB) for femoropopliteal lesions: primary patency and freedom from clinically driven target lesion revascularization (CD-TLR) at 36 months. **Top.** Primary patency by Kaplan–Meier estimate was significantly higher in the DCB group compared with the percutaneous transluminal angioplasty (PTA) group (log-rank test, $P < 0.001$). **Bottom.** Freedom from CD-TLR by Kaplan–Meier estimate was significantly higher in the DCB group compared with the PTA group (log-rank test, $P < 0.001$). **Top, Bottom.** Bars represent 95% confidence intervals. The number of subjects at risk represents the number of evaluable subjects at the beginning of each 90-day interval. An independent and blinded Clinical Events Committee adjudicated all target lesion revascularization events, and independent and blinded core laboratories reviewed all ultrasound and angiographic images.

93.1% had a completed 36-month follow-up visit (Figure 1). Of the remaining 99 PTA subjects eligible for 36-month evaluation, 92.9% had a completed 36-month follow-up visit.

Effectiveness Outcomes

Primary patency by Kaplan–Meier estimate was significantly higher in the DCB group compared with the PTA group at 36 months (Figure 2; 69.5% for DCB versus 45.1% for PTA; log rank $P < 0.001$). Rates of CD-TLR were lower in the DCB group compared with the PTA group (15.2% for DCB versus 31.1% for PTA; $P = 0.002$; Table 2). The time to first CD-TLR was longer in the DCB group compared with the PTA group (543±278 days for DCB versus 303±213 days for PTA; $P < 0.001$). Freedom from CD-TLR by Kaplan–Meier estimate is shown in Figure 2. Primary sustained clinical improvement was achieved by a greater percentage of subjects in the DCB group compared with the PTA group (68.7% for DCB versus 52.6% for PTA; $P = 0.012$).

A post hoc analysis was performed to evaluate the rates of CD-TLR among nonstented subjects at 36 months. The rate of CD-TLR was lower among nonstented subjects in the DCB group (14.4%) compared with those in the PTA

group (32.2%; $P = 0.001$). CD-TLR rates were not analyzed for stented subjects because of the small numbers of subjects that received stents in each group (16 subjects in DCB group and 14 in PTA group).

Safety Outcomes

Safety outcomes through 36 months are reported in Table 3. The primary composite safety end point was achieved in 81.2% of subjects in the DCB group and 64.1% in the PTA group ($P = 0.002$). Rates of CD target vessel revascularization were significantly lower in the DCB group compared with the PTA group (18.8% for DCB versus 35.9% for PTA; $P = 0.002$). Six deaths occurred between the 24- and 36-month follow-up periods: 5 in the DCB group (septic shock at 756 days, hemorrhagic stroke at 788 days, cardiac arrest at 907 days, acute respiratory failure at 1051 days, and gastric perforation at 1055 days) and 1 in the PTA group (worsening preexisting carcinoma at 877 days). Independent adjudication by the Clinical Events Committee determined that none of the deaths were related to the study device or procedure. The median time to death was 610 days in the DCB group and 637 days in the PTA group.

Table 2. Effectiveness Outcomes at 36 Months*

| Characteristic | DCB (Subjects=220) | PTA (Subjects=111) | <i>P</i> Value† |
|---|-----------------------|-----------------------|-----------------|
| Primary patency‡ | 69.5 (59) | 45.1 (59) | <0.001§ |
| CD-TLRI | 15.2 (30/197) | 31.1 (32/103) | 0.002 |
| Time to first CD-TLR, d | | | |
| Mean±SD | 542.9±278.2 | 302.9±213.0 | <0.001¶ |
| Median | 464.0 | 243.5 | |
| Interquartile range (Q3–Q1) | 370.0 | 186.0 | |
| Min, Max | 1, 1080 | 1, 1045 | |
| All TLR# | 16.2 (32/197) | 34.0 (35/103) | <0.001 |
| Primary sustained clinical improvement** | 68.7 (114/166) | 52.6 (51/97) | 0.012 |
| ABI/TBI†† | 0.917±0.231 | 0.894±0.194 | 0.429 |

ABI indicates ankle-brachial index; CD-TLR, clinically driven target lesion revascularization; DCB, drug-coated balloon; PTA, percutaneous transluminal angioplasty; and TBI, toe-brachial index.

*Values are % (n), % (n/N), or mean±SD.

†Unless otherwise specified, all tests were for superiority using the Fisher exact test for binary variables and Student *t* test for continuous variables.

‡Defined as freedom from CD-TLR or freedom from restenosis as determined by duplex ultrasonography peak systolic velocity ratio ≤2.4 within 36 months. The 36-month primary patency was calculated based on Kaplan–Meier estimate, and the number of primary patency failure subjects is displayed in the parentheses.

§Log rank *P* value.

¶Defined as any reintervention at the target lesion because of symptoms or drop of ABI of ≥20% or >0.15 when compared with postprocedure baseline ABI/TBI.

¶¶Wilcoxon rank-sum test *P* value.

#Includes clinically-driven and incidental or duplex-driven TLR.

**Defined as freedom from target limb major amputation, target vessel revascularization, and increase in Rutherford class.

††TBI allowed in cases of incompressible vessels in IN.PACT SFA phase II.

Functional Outcomes

There was sustained improvement in functional outcomes in both treatment groups (Table 4) through 3 years. Subjects in the DCB group achieved the same level of functional improvement despite 48% fewer reinterventions than those required in the PTA group. Six-minute walk test distance was similar between groups at 36 months (283±130 m for DCB versus 293±124 m for PTA; *P*=0.583). Total WIQ score was also similar between groups at 36 months (72±34 for DCB versus 75±29 for PTA; *P*=0.785).

Discussion

The prospective, multicenter, multinational, randomized IN.PACT SFA trial demonstrated that treatment with a paclitaxel-coated balloon is superior to standard PTA through 3 years in subjects with symptomatic femoropopliteal artery disease. The study design included adjudication of all major adverse events by an independent and blinded Clinical Events Committee and all angiography and duplex ultrasonography by independent and blinded core laboratories.

DCB angioplasty is intended to be a durable stand-alone therapy that, especially in cases of noncomplex lesions, obviates the need for permanent implants such as BMSs, stent grafts, and

Table 3. Safety Outcomes at 36 Months*

| Characteristic | DCB (Subjects=220) | PTA (Subjects=111) | <i>P</i> Value† |
|--|-----------------------|-----------------------|--------------------|
| Primary composite safety‡ | 81.2 (160/197) | 64.1 (66/103) | 0.002 |
| Major adverse events§ | 27.9 (55/197) | 37.9 (39/103) | 0.089 |
| All-cause deaths | 10.7 (21/197) | 1.9 (2/103) | 0.006 |
| Device- or procedure-related deaths | 0.0 (0/197) | 0.0 (0/103) | N/A |
| Clinically driven TVR | 18.8 (37/197) | 35.9 (37/103) | 0.002 |
| Target limb major amputation | 0.0 (0/197) | 0.0 (0/103) | N/A |
| Thrombosis | 2.0 (4/197) | 4.9 (5/103) | 0.283 |

DCB indicates drug-coated balloon; PTA, percutaneous transluminal angioplasty; and TVR, target vessel revascularization.

*Values are % (n/N).

†*P* values are based on Fisher exact test for superiority with significance level of 0.05.

‡Defined as 30-day freedom from device- and procedure-related death and target limb major amputation and 36-month freedom from clinically driven TVR.

§Composite of death, clinically driven TVR, target limb major amputation, and thrombosis.

||No deaths were adjudicated as device- or procedure-related by the Clinical Events Committee. Median postindex days to death was 610 days (DCB) and 637 days (PTA).

drug-eluting stents (DES). There are inherent difficulties with comparing outcomes across trials with 3-year findings, given differences in end point definitions and study populations. Nevertheless, the primary patency and CD-TLR rates were significantly better in the DCB group compared with the PTA group over this longer-term follow-up period and compare favorably to outcomes reported for other endovascular interventions in the same vascular bed at 3 years. Primary patency was 69.5% at 36 months for DCBs in this study, compared with 60.0% to 71.5% for BMS, DES, and stent grafts.^{7,20,21} Freedom from CD-TLR was 84.5% at 36 months for DCBs in this study, compared with previous reports of 69.7% to 75.5% freedom from TLR at 36 months for BMS^{5,7} and 83.6% freedom from TLR for a DES.²⁰ The 36-month CD-TLR rate of 15.2% for DCBs is less than half of what has been reported for BMS (31.1% CD-TLR at 3 years)⁷ and stent grafts (34.7% TLR at 3 years).²¹ We observed these results despite the fact that the mean lesion length in the DCB arm of the IN.PACT SFA trial was at the upper margin of lesion length reported in other studies. The mean lesion length in the DCB arm of the IN.PACT SFA trial was 8.94 cm compared with 6.64 cm in a randomized controlled trial of a DES²⁰ and 7.00 to 8.91 cm in the major BMS trials.^{5,7} The mean lesion length in the stent graft trial was longer (19 cm).²¹ These data suggest that treatment with DCBs is associated with results at least as favorable as those obtained with stenting but without the risks associated with an endoprosthesis.

This long-term treatment effect is supported by the persistence of paclitaxel in the vessel up to 180 days after application in experimental models.¹¹ The mechanical force of balloon angioplasty triggers inflammatory pathways that stimulate neointimal hyperplasia and restenosis.²² The sustained presence of paclitaxel is thought to provide antiproliferative activity over an extended period of time that is sufficient to inhibit the neointimal hyperplasia that causes restenosis.

Table 4. Functional Outcomes at 36 Months*

| Outcomes | DCB | | PTA | | P Value |
|-------------------------|---------------|---------------|---------------|---------------|---------|
| | Baseline | 36 mo | Baseline | 36 mo | |
| 6MWT† | | | | | |
| Mean distance, m | | | | | 0.988‡ |
| | | | | | 0.583§ |
| N | 119 | 55 | 60 | 29 | |
| Mean±SD | 253.2±123.0 | 282.9±129.7 | 256.0±114.7 | 292.6±124.0 | |
| Median [IQR] | 274.3 [208.8] | 289.6 [219.5] | 278.6 [159.9] | 304.8 [156.4] | |
| Change from baseline, m | | | | | 0.117§ |
| N | ... | 54 | ... | 29 | |
| Mean±SD | ... | 9.0±119.1 | ... | 56.0±101.4 | |
| Median [IQR] | ... | 25.1 [149.5] | ... | 61.0 [83.2] | |
| WIQ, % | | | | | |
| Walking impairment | | | | | 0.799‡ |
| | | | | | 0.785§ |
| N | 214 | 158 | 109 | 91 | |
| Mean±SD | 42.1±28.9 | 71.8±34.2 | 41.3±29.9 | 74.7±29.2 | |
| Median [IQR] | 50.0 [25.0] | 100.0 [50.0] | 50.0 [25.0] | 75.0 [50.0] | |
| Walking distance | | | | | 0.861‡ |
| | | | | | 0.680§ |
| N | 177 | 95 | 83 | 46 | |
| Mean±SD | 32.3±27.7 | 67.4±37.8 | 30.4±24.2 | 65.0±37.9 | |
| Median [IQR] | 26.5 [41.2] | 89.3 [65.5] | 27.6 [36.2] | 79.8 [72.4] | |
| Walking speed | | | | | 0.847‡ |
| | | | | | 0.351§ |
| N | 177 | 96 | 82 | 46 | |
| Mean±SD | 31.8±23.5 | 52.4±31.6 | 29.3±17.1 | 47.1±28.4 | |
| Median [IQR] | 26.1 [29.3] | 56.5 [51.1] | 27.2 [26.1] | 46.7 [38.0] | |
| Stair climbing | | | | | 0.802‡ |
| | | | | | 0.786§ |
| N | 175 | 96 | 83 | 46 | |
| Mean±SD | 42.5±31.3 | 66.4±36.8 | 40.7±29.0 | 68.2±35.9 | |
| Median [IQR] | 41.7 [50.0] | 75.0 [62.5] | 37.5 [50.0] | 83.3 [67.7] | |

6MWT indicates 6-minute walk test; DCB, drug-coated balloon; IQR, interquartile range; PTA, percutaneous transluminal angioplasty; and WIQ, Walking Impairment Questionnaire.

*Values are mean±SD (n). The number of subjects evaluated at each interval is reported in parentheses. Based on number of subjects with available data. Site-reported data.

†Data collected in IN.PACT SFA phase II only.

‡Wilcoxon test for comparison of treatment groups at baseline.

§Wilcoxon test for comparison of treatment groups at 36 months.

The rate of all-cause death was significantly higher in the DCB group compared with the PTA group. However, the site investigator and blinded Clinical Events Committee reviewed each event, and none of the deaths were determined related to the balloon treatment (DCB or PTA group). All events occurred late after therapy. The death rate in the DCB group (10.7%) was consistent with reported death rates in other similar peripheral artery disease studies,^{5,7,21,23} whereas the death

rate in the PTA group (1.9%) was significantly lower than in previous trials.

The WIQ is the most commonly used self-report tool to evaluate patient's walking ability.²⁴ Previous studies have reported moderate to strong correlations between scores on the WIQ and walking performance as measured by treadmill tests in patients with intermittent claudication.²⁵⁻²⁷ The WIQ contains 14 items that contribute to 3 subscales (walking distance,

walking speed, and stair-climbing ability) and an overall score. Each score ranges from 0 to 100 with a lower score indicating greater walking impairment (low performance). Validation studies have identified a WIQ overall score of <42.5 as an indicator of low performance.²⁸ The combined walking distance and stair-climbing score has been shown to provide the best indication of high performance with a cutoff of 75.5.²⁸ In the current study, subjects in the DCB and PTA treatment groups had evidence of significant walking impairment at baseline (overall score 42.1 versus 41.3). At 36 months, there was evidence of sustained improvement in walking ability (combined walking distance and stair-climbing score 133.8 versus 133.2) that was similar for both groups. This same level of functional improvement in the DCB group was achieved with 48% fewer reinterventions. The time to first CD-TLR was nearly twice as long in the DCB group (543 days) compared with the PTA group (303 days), and the rate of reintervention over the 3-year follow-up period was relatively low and without evidence of any significant late surge in TLRs or catch-up effect.

Recent analysis of IN.PACT SFA phase II data showed that the cost of DCB or PTA treatment within 2 years of follow-up was nearly identical. The higher initial cost of the DCB was offset by improved clinical outcomes, including a reduced need for repeat reinterventions, supporting the ultimate conclusion that treatment with DCB is cost-effective, and an economically dominant option compared with standard PTA.²⁹ A similar study from the United Kingdom found that DCBs, especially paclitaxel-coated DCBs with a urea excipient, had the greatest economic value compared with PTA, BMSs, and DES.³⁰ These studies suggest that the sustained longer-term maintenance of patency and avoidance of TLR should produce additional cost benefits over time.

There are several minimally invasive options for revascularization in patients with symptomatic peripheral artery disease, and there is a need for level-one evidence on long-term outcomes after treatment with a DCB for femoropopliteal lesions. The IN.PACT SFA is the first large, randomized trial to show sustained effect of a DCB through 3 years after treatment, with significantly higher primary patency and a lower rate of repeat interventions compared with standard PTA.

Limitations

Physicians involved in follow-up evaluations were aware of ultrasound and angiographic results to ensure optimal decision-making in each case. However, the subject and study sponsor were blinded through the 12-month follow-up period, and the Clinical Events Committee and core laboratories were blinded throughout the entire 3-year follow-up period.

As in any clinical trial, the number of subjects that are available for evaluation is gradually reduced as the length of the follow-up period increases. Despite the >90% 36-month follow-up compliance rates, because of the 2:1 randomization, the number of subjects in the PTA arm at 3 years was relatively low.

Conclusions

The 3-year outcomes from the prospective, multicenter, multinational, randomized IN.PACT SFA trial demonstrate the durable effectiveness after treatment with a paclitaxel DCB

compared with a standard PTA balloon. Treatment with a DCB was associated with significantly higher primary patency and a lower rate of repeat interventions compared with PTA. The results of this analysis show sustained clinical benefit up to 3 years after treatment with a DCB and support use of the paclitaxel-coated DCB as an intervention for patients with symptomatic femoropopliteal disease.

Acknowledgments

We thank Eric Fernandez, MD, and Azah Tabah, PhD, for technical review of the article, and Paula Soto, PhD, and Zachary Harrelson, PhD, of Meridius Health Communications Inc, San Diego, CA, for providing medical writing support, which was funded by Medtronic Inc, Minneapolis, MN, in accordance with Good Publication Practice (GPP3) guidelines.

Sources of Funding

This work was supported by Medtronic.

Disclosures

Dr Schneider is Chief Medical Officer for Intact Vascular; has received modest royalties from Cook Medical; and has served as a non-compensated advisor for Medtronic, Abbott Vascular, and Cardinal. Dr Laird has received research grants from W.L. Gore and Medtronic and has served as a compensated consultant/advisory board member for Abbott Vascular, Bard Peripheral Vascular, Boston Scientific, and Medtronic. Dr Tepe has received research grants from and is a compensated advisory board member for Medtronic and B. Braun and has received support from Biotronik. Dr Brodmann has received speaking honoraria from Bard Peripheral Vascular, Biotronik, Medtronic, Spectranetics, and VIVA Physicians; and is a consultant for Medtronic and Spectranetics. Dr Zeller has received speaking honoraria from Abbott Vascular, Bard Peripheral Vascular, Biotronik, Boston Scientific Corp, Cook Medical, Cordis Corp, GLG, Gore & Associates, Medtronic, Philips, Spectranetics, Straub Medical, TriReme, Veryan, and VIVA Physicians; is a consultant for Abbott Vascular, Bard Peripheral Vascular, Boston Scientific Corp, Cook Medical, Gore & Associates, Medtronic, and Spectranetics; and his clinic has received study funds or funds for research or clinical trials from 480 Biomedical, Abbott Vascular, B. Braun, Bard Peripheral Vascular, Bayer Pharma, Biotronik, Caveo Med, Contego Medical, Cook Medical, CSI, Gore & Associates, Innora, Intact Vascular, Medtronic, Mercator, Philips, Pluristem, Shockwave, Spectranetics, Terumo, TriReme, and Veryan. Dr Scheinert has served as a compensated consultant for Abbott Vascular, Biotronik, Boston Scientific, Cook Medical, Cordis, CR Bard, Gardia Medical, Hemoteq, Medtronic/Covidien, Ostial Inc, TriReme Medical, TriVascular, and Upstream Peripheral Technologies. Dr Metzger has received hands-on proctor fees from Abbott Vascular and TriVascular; has served as a consultant for Abbott Vascular, Bard Peripheral Vascular, Boston Scientific, and TriVascular; and has received symposium honoraria from Abbott Vascular, Bard, Boston Scientific, Medtronic, and CSI. Dr Micari has served as a compensated consultant for Medtronic, Boston Scientific, CR Bard, and Terumo. Dr Sachar has served as a compensated consultant and advisory board member for Abbott Vascular, Boston Scientific, Cordis Corp, and Medtronic/Covidien; has received compensation for educational programs from Abbott Vascular, Boston Scientific, Cordis Corp, and Medtronic/Covidien; has received funds for research or clinical trials from Abbott Vascular, Bard Peripheral Vascular, CardioMEMS, Cook Medical, Cordis Corp, CSI, W.L. Gore & Associates, Medtronic, Spectranetics, and Veryan; and is a major shareholder of Contego Medical. M.R. Jaff has served as a noncompensated advisor for Medtronic; has served as a compensated board member of VIVA Physicians, a 501(c)(3) not-for-profit education and research organization and is an equity investor of PQ Bypass and Vascular Therapies. Dr Wang is a full-time salaried employee of Medtronic. Dr Hasenbank is a full-time salaried employee

of Medtronic. Dr Krishnan has served as a compensated consultant for Abbott Vascular, Bard Peripheral Vascular, and Medtronic.

References

- Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FG; TASC II Working Group. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *J Vasc Surg.* 2007;45(suppl S):S5–S67. doi: 10.1016/j.jvs.2006.12.037.
- Dick P, Wallner H, Sabeti S, Loewe C, Mlekusch W, Lammer J, Koppenssteiner R, Minar E, Schillinger M. Balloon angioplasty versus stenting with nitinol stents in intermediate length superficial femoral artery lesions. *Catheter Cardiovasc Interv.* 2009;74:1090–1095. doi: 10.1002/ccd.22128.
- Schillinger M, Sabeti S, Loewe C, Dick P, Amighi J, Mlekusch W, Schlager O, Cejna M, Lammer J, Minar E. Balloon angioplasty versus implantation of nitinol stents in the superficial femoral artery. *N Engl J Med.* 2006;354:1879–1888. doi: 10.1056/NEJMoa051303.
- Laird JR, Katzen BT, Scheinert D, Lammer J, Carpenter J, Buchbinder M, Dave R, Ansel G, Lansky A, Cristea E, Collins TJ, Goldstein J, Jaff MR; RESILIENT Investigators. Nitinol stent implantation versus balloon angioplasty for lesions in the superficial femoral artery and proximal popliteal artery: twelve-month results from the RESILIENT randomized trial. *Circ Cardiovasc Interv.* 2010;3:267–276. doi: 10.1161/CIRCINTERVENTIONS.109.903468.
- Laird JR, Katzen BT, Scheinert D, Lammer J, Carpenter J, Buchbinder M, Dave R, Ansel G, Lansky A, Cristea E, Collins TJ, Goldstein J, Cao AY, Jaff MR; RESILIENT Investigators. Nitinol stent implantation vs. balloon angioplasty for lesions in the superficial femoral and proximal popliteal arteries of patients with claudication: three-year follow-up from the RESILIENT randomized trial. *J Endovasc Ther.* 2012;19:1–9. doi: 10.1583/11-3627.1.
- Matsumura JS, Yamanouchi D, Goldstein JA, Pollock CW, Bosiers M, Schultz GA, Scheinert D, Rocha-Singh KJ. The United States Study for evaluating endovascular treatments of lesions in the superficial femoral artery and proximal popliteal by using the Protégé EverFlex Nitinol STent SYstem II (DURABILITY II). *J Vasc Surg.* 2013;58:73.e1–83.e1. doi: 10.1016/j.jvs.2012.12.066.
- Rocha-Singh KJ, Bosiers M, Schultz G, Jaff MR, Mehta M, Matsumura JS; Durability II Investigators. A single stent strategy in patients with lifestyle limiting claudication: 3-year results from the Durability II trial. *Catheter Cardiovasc Interv.* 2015;86:164–170. doi: 10.1002/ccd.25895.
- Scheinert D, Scheinert S, Sax J, Piorowski C, Bränlich S, Ulrich M, Biamino G, Schmidt A. Prevalence and clinical impact of stent fractures after femoropopliteal stenting. *J Am Coll Cardiol.* 2005;45:312–315. doi: 10.1016/j.jacc.2004.11.026.
- Iida O, Uematsu M, Soga Y, Hirano K, Suzuki K, Yokoi H, Muramatsu T, Inoue N, Nanto S, Nagata S. Timing of the restenosis following nitinol stenting in the superficial femoral artery and the factors associated with early and late restenoses. *Catheter Cardiovasc Interv.* 2011;78:611–617. doi: 10.1002/ccd.23064.
- Banerjee S, Sarode K, Mohammad A, Gigliotti O, Baig MS, Tsai S, Shammam NW, Prasad A, Abu-Fadel M, Klein A, Armstrong EJ, Jeon-Slaughter H, Brilakis ES, Bhatt DL. Femoropopliteal artery stent thrombosis: report from the excellence in peripheral artery disease registry. *Circ Cardiovasc Interv.* 2016;9:e002730. doi: 10.1161/CIRCINTERVENTIONS.115.002730.
- Speck U, Cremers B, Kelsch B, Biedermann M, Clever YP, Schaffner S, Mahnkopf D, Hanisch U, Böhm M, Scheller B. Do pharmacokinetics explain persistent restenosis inhibition by a single dose of paclitaxel? *Circ Cardiovasc Interv.* 2012;5:392–400. doi: 10.1161/CIRCINTERVENTIONS.111.967794.
- Tepe G, Zeller T, Albrecht T, Heller S, Schwarzwälder U, Beregi JP, Claussen CD, Oldenburg A, Scheller B, Speck U. Local delivery of paclitaxel to inhibit restenosis during angioplasty of the leg. *N Engl J Med.* 2008;358:689–699. doi: 10.1056/NEJMoa0706356.
- Werk M, Langner S, Reinkensmeyer B, Boettcher HF, Tepe G, Dietz U, Hosten N, Hamm B, Speck U, Rieke J. Inhibition of restenosis in femoropopliteal arteries: paclitaxel-coated versus uncoated balloon: femoral paclitaxel randomized pilot trial. *Circulation.* 2008;118:1358–1365. doi: 10.1161/CIRCULATIONAHA.107.735985.
- Werk M, Albrecht T, Meyer DR, Ahmed MN, Behne A, Dietz U, Eschenbach G, Hartmann H, Lange C, Schnorr B, Stiepani H, Zoccai GB, Hänninen EL. Paclitaxel-coated balloons reduce restenosis after femoro-popliteal angioplasty: evidence from the randomized PACIFIER trial. *Circ Cardiovasc Interv.* 2012;5:831–840. doi: 10.1161/CIRCINTERVENTIONS.112.971630.
- Scheinert D, Duda S, Zeller T, Krankenberg H, Rieke J, Bosiers M, Tepe G, Naisbitt S, Rosenfield K. The LEVANT I (Lutonix paclitaxel-coated balloon for the prevention of femoropopliteal restenosis) trial for femoropopliteal revascularization: first-in-human randomized trial of low-dose drug-coated balloon versus uncoated balloon angioplasty. *JACC Cardiovasc Interv.* 2014;7:10–19. doi: 10.1016/j.jcin.2013.05.022.
- Rosenfield K, Jaff MR, White CJ, Rocha-Singh K, Mena-Hurtado C, Metzger DC, Brodmann M, Pilger E, Zeller T, Krishnan P, Gammon R, Muller-Hulsbeck S, Nehler MR, Benenati JF, Scheinert D; for the LEVANT 2 Investigators. Trial of a paclitaxel-coated balloon for femoropopliteal artery disease. *N Engl J Med.* 2015;373:145–153. doi: 10.1056/NEJMoa1406235.
- Schroeder H, Meyer DR, Lux B, Ruecker F, Martorana M, Duda S. Two-year results of a low-dose drug-coated balloon for revascularization of the femoropopliteal artery: outcomes from the ILLUMENATE first-in-human study. *Catheter Cardiovasc Interv.* 2015;86:278–286. doi: 10.1002/ccd.25900.
- Tepe G, Laird J, Schneider P, Brodmann M, Krishnan P, Micari A, Metzger C, Scheinert D, Zeller T, Cohen DJ, Snead DB, Alexander B, Landini M, Jaff MR; IN.PACT SFA Trial Investigators. Drug-coated balloon versus standard percutaneous transluminal angioplasty for the treatment of superficial femoral and popliteal peripheral artery disease: 12-month results from the IN.PACT SFA randomized trial. *Circulation.* 2015;131:495–502. doi: 10.1161/CIRCULATIONAHA.114.011004.
- Laird JR, Schneider PA, Tepe G, Brodmann M, Zeller T, Metzger C, Krishnan P, Scheinert D, Micari A, Cohen DJ, Wang H, Hasenbank MS, Jaff MR; IN.PACT SFA Trial Investigators. Durability of treatment effect using a drug-coated balloon for femoropopliteal lesions: 24-month results of IN.PACT SFA. *J Am Coll Cardiol.* 2015;66:2329–2338. doi: 10.1016/j.jacc.2015.09.063.
- Dake MD, Ansel GM, Jaff MR, Ohki T, Saxon RR, Smouse HB, Machan LS, Snyder SA, O'Leary EE, Ragheb AO, Zeller T; Zilver PTX Investigators. Durable clinical effectiveness with paclitaxel-eluting stents in the femoropopliteal artery: 5-year results of the Zilver PTX randomized trial. *Circulation.* 2016;133:1472–1483; discussion 1483. doi: 10.1161/CIRCULATIONAHA.115.016900.
- Geraghty PJ, Mewissen MW, Jaff MR, Ansel GM; VIBRANT Investigators. Three-year results of the VIBRANT trial of VIABAHN endoprosthesis versus bare nitinol stent implantation for complex superficial femoral artery occlusive disease. *J Vasc Surg.* 2013;58:386.e4–395.e4. doi: 10.1016/j.jvs.2013.01.050.
- Schillinger M, Minar E. Restenosis after percutaneous angioplasty: the role of vascular inflammation. *Vasc Health Risk Manag.* 2005;1:73–78.
- Dake MD, Ansel GM, Jaff MR, Ohki T, Saxon RR, Smouse HB, Snyder SA, O'Leary EE, Tepe G, Scheinert D, Zeller T; Zilver PTX Investigators. Sustained safety and effectiveness of paclitaxel-eluting stents for femoropopliteal lesions: 2-year follow-up from the Zilver PTX randomized and single-arm clinical studies. *J Am Coll Cardiol.* 2013;61:2417–2427. doi: 10.1016/j.jacc.2013.03.034.
- Regensteiner JG, Steiner JF, Panzer RJ, Hiatt WR. Evaluation of walking impairment by questionnaire in patients with peripheral arterial disease. *J Vasc Med Biol.* 1990;2:142–158.
- McDermott MM, Liu K, Guralnik JM, Martin GJ, Criqui MH, Greenland P. Measurement of walking endurance and walking velocity with questionnaire: validation of the Walking Impairment Questionnaire in men and women with peripheral arterial disease. *J Vasc Surg.* 1998;28:1072–1081.
- Myers SA, Johanning JM, Stergiou N, Lynch TG, Longo GM, Pipinos II. Claudication distances and the Walking Impairment Questionnaire best describe the ambulatory limitations in patients with symptomatic peripheral arterial disease. *J Vasc Surg.* 2008;47:550–555. doi: 10.1016/j.jvs.2007.10.052.
- Nicolaï SP, Kruidenier LM, Rouwet EV, Graffius K, Prins MH, Teijink JA. The Walking Impairment Questionnaire: an effective tool to assess the effect of treatment in patients with intermittent claudication. *J Vasc Surg.* 2009;50:89–94. doi: 10.1016/j.jvs.2008.12.073.
- Sagar SP, Brown PM, Zelt DT, Pickett WL, Tranmer JE. Further clinical validation of the walking impairment questionnaire for classification of walking performance in patients with peripheral artery disease. *Int J Vasc Med.* 2012;2012:190641. doi: 10.1155/2012/190641.
- Scheinert D. Drug coated balloon treatment for patients with intermittent claudication: new insights from the IN.PACT Global Study long lesion (≥15 cm) imaging cohort. Paper presented at: EuroPCR 2015, Meeting of the European Association of Percutaneous Cardiovascular Interventions; May 19–22, 2015; Paris, France.
- Katsanos K, Geisler BP, Garner AM, Zayed H, Cleveland T, Pietzsch JB. Economic analysis of endovascular drug-eluting treatments for femoropopliteal artery disease in the UK. *BMJ Open.* 2016;6:e011245. doi: 10.1136/bmjopen-2016-011245.