



Cite this: *RSC Adv.*, 2018, 8, 23199

Combining $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ to identify the distribution and the potential sources of nitrate in human-impacted watersheds, Shandong, China†

Baoshan Yang,^{ab} Hui Wang,^{*ab} Yingkui Jiang,^{ab} Fang Dong,^{ab} Xinhua He^c and Xiaoshuang Lai^a

Identifying the anthropogenic and natural sources of nitrate emissions contributing to surface water continues to be an enormous challenge. It is necessary to control the water quality in the watershed impacted by human disturbance. In this study, water chemical parameters including nitrate (NO_3^-) concentrations, $\delta^{15}\text{N}-\text{NO}_3^-$, $\delta^{18}\text{O}-\text{NO}_3^-$, and $\delta^{18}\text{O}-\text{H}_2\text{O}$ were analyzed to investigate the contamination and sources of NO_3^- in two watershed rivers (Jinyun, JYN and Jinyang, JYA), Jinan, Shandong, China. Results indicated NO_3^- concentrations in the JYN were significantly higher than those in the JYA ($P < 0.05$), probably because of high N input of the extensive farmlands or orchards in the drainage basin. $\delta^{15}\text{N}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{NO}_3^-$, associated with Cl^- , indicated that nitrate-nitrogen (NO_3^--N) was not derived from atmospheric deposition but came principally from manure/sewage and soil organic matter in these two watersheds. The microbial nitrification took place in the nitrate of manure/sewage and soil nitrate. The combination of NO_3^- concentration and nitrogen and oxygen isotope suggested that NO_3^- had undergone microbial denitrification after entering the rivers. Furthermore, NO_3^- concentrations had significant temporal and spatial variation highlighting differential sources and fates. These results expand our understanding of mechanisms driving NO_3^- retention and transport and provide strategies in managing NO_3^- contamination in different land use watersheds around the world.

Received 22nd May 2018
Accepted 16th June 2018

DOI: 10.1039/c8ra04364g

rsc.li/rsc-advances

1. Introduction

Nitrate (NO_3^-) is a widespread contamination source which often poses potential adverse health effects to humans through surface and underground water systems.^{1,2} Higher concentrations have occurred due to mixed land use and anthropogenic activities, resulting in acidification, eutrophication, drinking water degradation and human health risks.³⁻⁶ There are complicated sources of NO_3^- under different watershed conditions. Anthropogenic N is becoming the dominant contributor to the water quality impairments with potential sources of NO_3^- contamination from N deposition, septic decomposition, animal waste, synthetic fertilizer, and decaying organic matter.^{7,8} A worldwide analysis suggests that NO_3^- exports from watersheds are highly variable, and that a broader array of studies is more important for global-scale analyses.⁹⁻¹¹ Identifying the distribution and sources of NO_3^- in rivers across

land use and hydrologic duration will be critical in prioritizing effective NO_3^- reduction methods and providing an effective watershed management strategy.

Although the concentration of NO_3^- and the N-isotopic signature in NO_3^- have been successfully applied in various case studies of surface and ground waters in widespread watersheds around the world to identify the sources of nitrate-nitrogen (NO_3^--N), the wide range and overlapping values for some sources make it difficult to identify the source and transformation of NO_3^- in surface water and groundwater.¹²⁻¹⁵ It is promising to employ $\delta^{18}\text{O}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{H}_2\text{O}$ associated with $\delta^{15}\text{N}-\text{NO}_3^-$ to differentiate the sources according to the well-established linear model between $\delta^{18}\text{O}-\text{NO}_3^-$ ranges and microbial nitrification/denitrification.¹⁶⁻¹⁸

The rivers flow through different human disturbed watersheds, which could result in the different nitrate level. In this research, the stable isotopes of ^{15}N and ^{18}O in NO_3^- and $\delta^{18}\text{O}-\text{H}_2\text{O}$, in conjunction with land use information and water quality under different land use and anthropogenic activities, were used to characterize the source and transformation of NO_3^- that delivered to the one important reservoir from two rivers in pairwisd watersheds (Jinyun and Jinyang watersheds). The objectives of this research were to determine: (1) if the dissolved NO_3^- displays significantly seasonal and spatial differences in these two different watersheds; (2) if the water

^aSchool of Water Conservancy and Environment, University of Jinan, Jinan 250022, China. E-mail: hwang_118@163.com; Fax: +86 053182769233; Tel: +86 82767237

^bKey Laboratory of Water Resources and Environmental Engineering in Universities of Shandong, Jinan 250022, China

^cSchool of Plant Biology, University of Western Australia, Crawley, WA 6009, Australia

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8ra04364g



quality and isotopic data could indicate the NO_3^- sources, *i.e.*, if NO_3^- originates from fertilizer or manure in the JYN watershed or from organism decomposition or sewage in the JYA watershed; and (3) is there different occurrence of NO_3^- transformation with the microbial nitrification or denitrification in these two rivers. The possible source of NO_3^- from fertilizer, manure, septic system, or sewage explanation could be identified due to different land use in these two watersheds. The expected results could provide important base for the risk assessment and management of the nitrate contaminant from the comparable watersheds with different land use practices.

2. Materials and methods

2.1 Study area description

The study area (237.1 km²) is within the Jinyun and Jinyang watershed (117°01'–117°14' E, 36°19'–36°28' N; elevation range of 150–175 m, with respective 55.2 km² and 181.9 km² basin areas). These two watersheds have the Cambrian gneisses and Ordovician limestone landscape. The research region has a warm temperate continental monsoon climate with four distinct seasons. The annual mean precipitation is 710 mm (~65% between June and September). The annual mean temperature is 14.7 °C (−0.4 °C in January and +27.5 °C in July) (Chinese Meteorological Administration Public Service Center, website: <http://www.weather.com.cn>). The soil types are primary the calcareous cinnamon soil and brown soil, which promote water loss and soil erosion.¹⁹

Jinyun river (JYN) and Jinyang river (JYA), locating in southern Jinan city, Shandong, China, are typical watersheds, with a total 55.2 km² and 181.9 km² basin areas, respectively. Both rivers together contribute about half of water to the Wohushan reservoir, which supplies the primary drinking water source for 300 000 residents in the south central Jinan city and a variety of industrial and agricultural activities of Jinan city (Fig. S1†).²⁰ The rivers' source originates from the northern slope of Mountain Tai at >1500 m above sea level, and the rivers traverse the southern part of Jinan city before flowing into the Wohushan reservoir (Fig. S1†). Both the JYN and JYA are fed by both precipitation and ground water throughout the year. The input of these two rivers is mainly dominated by surface runoff from local precipitation during the summer rainy season (June to September), whilst, the rivers are sustained by precipitation and seepage of ground water during the remainder of the year. In general, the daily discharge shows a distinct seasonal variation, ranging from 0.67 to 1.26 m³ s^{−1} in January (winter, lowest) and 4.15 to 7.82 m³ s^{−1} in August (summer, highest) for both the JYN and JYA (the office of Wohushan reservoir Administration, see website: <http://www.whssk.com>). The water contribution to the Wohushan reservoir from the JYN watershed is similar to the JYA watershed, whilst their land use and anthropogenic activities are very contrasting. For instance, high N input from chemical fertilizer and manure applications for intensive agricultural activities and orchard plantations are common in the JYN watershed. The NO_3^- contribution is thus from runoff of precipitation in this watershed. In contrast, more natural tour sites and residential properties are along the JYA

watershed. This JYA watershed is thus associated with septic systems and potential point sources of contamination.²¹

2.2 Water sampling

To investigate seasonal and spatial distribution pattern of the nitrate concentration and isotopic composition in these two watersheds, water samples from 22 sites were collected four times in January (winter, low-water stage), May (spring, rising-water stage), August (summer, high-water stage) and October (fall, falling-water stage) 2011 and 2014 (Fig. 1S†). Four sites were distributed at the upstream Jinyang river (JYA-U), the middle reach of Jinyang river (JYA-M), and the downstream Jinyang river (JYA-D), respectively. Three samples were collected and the samples were combined into one sample at each site. The total samples were four at upstream, middle stream, and downstream Jinyang river, respectively. Four of the sites were selected at the upstream and downstream Jinyun river (JYN-U and JYN-D), respectively. Two sites were from the integrated channel because they are shorter than the Jinyang river. The total number of samples is 8 for Jinyun river. Three samples were collected from integrated channel. The samples were from combined two sites and each of two. In 2014, the samples were only collected from the sites where the data were not obtained well. These sampling sites (Fig. S1†) were selected to avoid fixed point-source contamination. At each site, a depth-integrated (0.5–1.5 m) mixed sample was taken from the center of the river and its temperature, pH and electrical conductivity (EC) were immediately measured. Samples for chemical and isotopic analyses were passed through a 0.45 μm membrane filter and kept at 4 °C until analyses within 2 weeks. In order to reduce the interference of Cl^- , the pretreatment was implemented with AG/H SPE column. Anions were analyzed by ion chromatography (Dionex ICS-1500, USA) at Shandong Analysis and Test Center, Jinan, China. Ammonium N (NH_4^+ -N) concentrations were concluded by the indophenol blue method (Ministry of Environment Protection of PRC, 1989).

2.3 Analytical methods

Three collected water samples from upstream, middle stream and downstream river were selected for isotope analysis. The water was filtered with 0.45 μm cellulose acetate membrane. Before measurement, the water samples were stored in refrigerator at 4 °C. In order to prevent biological activity, the samples were preserved with HCl for isotope analysis within one month. The refrigerated samples were measured for $\delta^{15}\text{N}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{NO}_3^-$ using the denitrifier method at the State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, Guizhou, China. Briefly, the denitrifying bacterium *Pseudomonas aureofaciens* converts NO_3^- to gaseous nitrous oxide (N_2O) for isotopic analysis (Casciotti *et al.*, 2002; Kaushal *et al.*, 2011). With a minimum of 50 nmol NO_3^- requirement to a continuous flow Trace Gas Pre-concentrator-IsoPrime isotope ratio mass spectrometer (Isoprime 100, UK), the $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ values (‰) in the NO_3^- samples were corrected using international reference standards USGS32, USGS34, IAEA-N3 and values are reported in

parts per thousand (‰) relative to atmospheric N₂ and Vienna Standard Mean Ocean Water (V-MOW) for δ¹⁵N and δ¹⁸O, respectively.²² The oxygen-18 (¹⁸O) isotope of water was analyzed through equilibrating with carbon dioxide gas at 25.0 ± 0.1 °C (Lee *et al.*, 2008).⁸ The carbon dioxide gas was then extracted and cryogenically purified. The δ¹⁸O values in the water were determined using a Sercon Integra stable isotope ratio mass spectrometer (Sercon 20, UK) at Nanjing Wanso Testing Services, China.

The following equation was used to calculate the δ¹⁵N and δ¹⁸O values:

$$\delta (\text{‰}) = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 1000 \quad (1)$$

where *R* denote the ratio of ¹⁵N/¹⁴N or ¹⁸O/¹⁶O. The standard deviation of laboratory analyses was ± 0.1‰ for δ¹⁸O–H₂O, ± 0.3‰ for both δ¹⁵N–NO₃[−] and δ¹⁸O–NO₃[−], respectively.

In order to quantify the measurement, data presentation, calculation and statistical analysis were performed using Microsoft office Excel and SPSS 16.0 (SPSS Inc., USA). The paired *t*-test and one-way ANOVA were performed at a significant level of *P* = 0.05.

3. Results and discussion

3.1 Seasonal and spatial distribution pattern of nitrate

The averaged NO₃[−] concentrations were not significantly different between the upstream and downstream site (*P* > 0.05) in the whole hydrological year of the JYN and JYA (Table 1). However, the average NO₃[−] was higher in the JYN than in the JYA, especially in spring and winter (Table 1), which probably resulted from more extensive distributions of agricultural farmlands and orchards in the JYN drainage basin compared with the JYA drainage basin. The NO₃[−] concentrations varied with seasons in both the JYN and JYA and were lower in spring (rising-water stage) and summer (high-water stage) both in the JYN and JYA (Table 1). Although the NO₃[−] concentrations of integrated channel showed a mixed effect, the same temporal patterns were found (Table 1). The possible reasons were the hydrological change, runoff of precipitation, and the transformation between organic N and inorganic N.²³ Both NO₃[−] and Cl[−] concentrations in the JYN and JYA showed little spatial variation (Table 2 and Fig. 1). All of the Cl[−] values in the sites were out of the compositional range of precipitation.²⁴ No rock salts or evaporation sediments were found in the two watersheds.²⁰ The elevated Cl[−] concentration in the rivers of south Jinan indicated that a potential large contribution of anthropogenic sources such as fertilizer, manure and sewage may dominate the Cl[−] concentrations of these two rivers. This also suggested that the NO₃[−] contamination in these two rivers may be from anthropogenic sources rather than from atmospheric deposition.

The nitrate concentration in JYA is lower than JYN during winter, spring and fall, but higher than JYN during summer. This nitrate seasonal variation corresponds to the seasonal variation of nitrogen isotope and oxygen isotope, which displayed high nitrate concentration with high nitrogen isotopes and low oxygen isotope. The results showed that the possible

nitrate sources is from manure or sewage waste. The possible reasons are that there are more agricultural lands distributed in the subwatershed of JYN. The application of manures and runoff in the summer may result in the drastic disturbance.

3.2 Nitrate sources

Both δ¹⁵N–NO₃[−] and δ¹⁸O–NO₃[−] can be a good indicator to quantify the different contribution of nitrate sources.^{2,16} Values of δ¹⁵N–NO₃[−] and δ¹⁸O–NO₃[−] were not significantly different between the JYN and JYA across the whole hydrological years (Table 2). The δ¹⁵N–NO₃[−] values were slightly higher from JYA than from JYN (Table 2). The δ¹⁸O–NO₃[−] values showed contrary pattern compared with the δ¹⁵N–NO₃[−] in these two rivers. Both the δ¹⁵N–NO₃[−] and δ¹⁸O–NO₃[−] presented the different seasonal patterns in the two rivers.

The Cl[−] concentrations in JYN and JYA were distinctly higher than in the precipitation (Fig. 1). It was primarily concluded that NO₃[−] were not only from precipitation. δ¹⁵N–NO₃[−] values varied between +4.61 and +9.39‰ (averaged 8.54 ± 0.92‰, *n* = 56, Tables 1 and 2) and were lower in the upstream (JYN-U, 8.01 ± 1.20‰) than in the downstream (JYN-D, 8.07 ± 1.15‰) of the Jinyun river (Fig. 2A). δ¹⁵N–NO₃[−] values were not significantly different among the upstream of the Jinyang river (JYA-U, averaged 7.33‰), the middle reach of the Jinyang river (JYA-M, averaged 7.08‰), and the downstream of the Jinyang river (JYA-D, averaged 8.02‰); they varied between +8.07–9.68‰ (Fig. 2A). Across these spatial river sites, δ¹⁵N–NO₃[−] values were significantly higher in spring than other seasons (Fig. 2B). The addition of NO₃[−] from manure or sewage might produce elevations of both the NO₃[−] concentration and the δ¹⁵N–NO₃[−] signature.⁷ In this study, although the runoff of precipitation is more drastic in summer than other season, δ¹⁵N–NO₃[−] was not higher in the summer. This indicated that NO₃[−] from discharged sewage may be the main sources.

The combination of analyzing the concentration and the isotopic composition of NO₃[−] is a useful tool to identify the NO₃[−] source and the denitrification process in a watershed.^{17,18,25} Values of NO₃[−] and δ¹⁵N–NO₃[−] were widely scattered in both the JYN and JYA (Fig. 2A). Values of δ¹⁵N–NO₃[−] for these two rivers were not significantly different by paired *t*-test (*P* = 0.22). The integrated channel showed similar patterns and the values were between the JYN and JYA rivers, reflecting the mixing of these two rivers (Table 1).

The ranges of δ¹⁵N–NO₃[−] indicated the NO₃[−] originated primarily from the manure or sewage, with only a little from NO₃[−] fertilizer in the JYN. However, the NO₃[−] concentrations did not increase with δ¹⁵N–NO₃[−] (Fig. 2A and B). Therefore, it was likely that the state of NO₃[−] from manure or sewage changed with nitrification and denitrification controlled by microbes.

3.3 Nitrification and denitrification

δ¹⁸O–NO₃[−] values in the two watershed rivers ranged from +2.89 to +9.21‰, indicating that neither atmospheric NO₃[−] deposition (+25–75‰) nor NO₃[−]-containing fertilizer (+17–25‰) was a dominant source of riverine NO₃[−] in these two rivers (Fig. 3). Values of δ¹⁵N–NO₃[−] (+6.68 to +9.68‰) and δ¹⁸O–

Table 1 Base properties, oxygen isotopic compositions of H₂O, and nitrogen and oxygen isotopic compositions of NO₃⁻ in the Jinyun river (JYN), Jinyang river (JYA), and integrated channel (IntC)

| Sample sites | Winter | | | Spring | | | Summer | | | Fall | | |
|--|---------------------------|---------------|--------------|--------------|---------------|--------------|--------------|---------------|--------------|---------------|---------------|--------------|
| | Upstream | Middle stream | Downstream | Upstream | Middle stream | Downstream | Upstream | Middle stream | Downstream | Upstream | Middle stream | Downstream |
| T (°C) | JYN 0.6(0.3) ^a | 0.8(0.2) | 0.5(0.1)a | 22.5(1.9)a | 23.6(0.4) | 22.5(1.2)a | 28.5(2.0)a | 26.8(0.3) | 27.2(1.7)a | 16.6(1.5)a | 13.7(0.3) | 13.9(2.1)a |
| | JYA 0.5(0.1)a | | 0.5(0.1)a | 22.8(0.4)a | | 24.0(0.1)a | 26.5(0.2)a | | 27.0(0.1)a | 13.5(0.1)a | | 13.8(0.2)a |
| | IntC 0.6(0.1)a | | 0.5(0.1)a | 23.1(0.3)a | | 22.5(0.1)a | 26.8(1.3)a | | 27.4(0.1)a | 14.5(0.2)a | | 15.2(0.1)a |
| pH | JYN 7.7(0.1)a | | 8.1(0.3)a | 8.1(0.1)a | | 8.0(0.1)a | 8.0(0.1)a | | 7.7(0.2)a | 7.7(0.1)a | | 8.1(0.1)a |
| | JYA 8.0(0.01)a | 8.1(0.04) | 8.0(0.02)a | 8.2(0.08)a | 8.0(0.05) | 8.2(0.02)a | 8.0(0.03)a | 7.9(0.02) | 8.0(0.04)a | 7.9(0.03)a | 8.0(0.05) | 8.0(0.02)a |
| | IntC 8.0(0.01)a | | 8.0(0.01)a | 8.1(0.03)a | | 8.1(0.02)a | 7.9(0.02)a | | 8.0(0.01)a | 8.1(0.01)a | | 7.9(0.02)a |
| DO (mg L ⁻¹) | JYN 14.6(1.5)a | | 12.3(1.1)a | 9.7(1.5)a | | 8.9(1.2)a | 11.2(1.5)a | | 7.0(1.0)a | 10.2(0.9)a | | 9.8(1.0)a |
| | JYA 12.8(1.2)b | 11.5(1.1) | 13.7(1.4)b | 10.2(1.4)b | 8.8(1.3) | 13.4(1.2)b | 9.4(0.4)b | 8.7(0.8) | 8.2(0.4)a | 9.8(0.3)b | 10.7(1.0) | 10.2(0.2)a |
| | IntC 12.7(1.4)ab | | 13.2(1.2)a | 9.6(0.8)a | | 8.8(1.1)a | 8.0(1.3)b | | 9.2(1.0)a | 9.1(2.2)b | | 9.3(1.9)a |
| Cl ⁻ (mg L ⁻¹) | JYN 27.6(3.2)a | 26.2(0.1) | 23.4(2.8)a | 23.9(3.0)a | | 24.1(3.4)a | 19.1(1.7)a | 16.4(0.3) | 21.0(3.5)a | 19.4(3.2)a | | 21.7(2.5)a |
| | JYA 25.4(9.2)a | | 27.6(0.03)a | 23.6(0.03)a | 23.0(0.1) | 21.8(0.03)a | 18.1(0.4)a | | 13.6(0.1)a | 20.7(0.1)a | 19.9(0.03) | 20.2(0.5)a |
| | IntC 26.4(5.1)a | | 25.8(3.5)a | 23.8(2.8)a | | 23.7(4.2)a | 19.2(5.0)a | | 18.8(2.7)a | 20.9(3.5)a | | 21.2(4.4)a |
| NH ₄ ⁺ (mg L ⁻¹) | JYN 0.19(0.02)a | 0.12(0.03) | 0.13(0.01)a | 0.14(0.01)a | | 0.15(0.03)a | 0.25(0.02)a | 0.13(0.01)b | 0.18(0.01)b | 0.15(0.01)b | 0.44(0.01) | 0.53(0.03)b |
| | JYA 0.09(0.01)b | | 0.15(0.01)a | 0.25(0.01)b | 0.12(0.01) | 0.23(0.02)a | 0.13(0.01)b | | 0.46(0.03)a | 0.38(0.01)a | | 0.39(0.11)a |
| | IntC 0.98(0.01)c | | 0.12(0.02)a | 0.15(0.09)b | | 0.15(0.01)a | 0.37(0.03)a | | 4.47(0.15)a | 8.31(0.02)a | | 5.76(0.01)a |
| NO ₃ ⁻ (mg L ⁻¹) | JYN 11.47(0.07)a | 8.92(0.04) | 11.00(0.09)a | 6.27(0.05)a | 3.92(0.01) | 7.08(0.05)a | 5.56(0.13)a | 5.43(0.16) | 4.47(0.15)a | 8.31(0.02)a | 6.05(0.01) | 7.70(0.01)a |
| | JYA 8.49(0.43)b | | 9.98(0.02)b | 4.73(0.01)b | | 2.48(0.01)b | 6.32(0.16)b | | 5.36(0.09)b | 7.25(0.22)b | | 7.19(1.03)b |
| | IntC 9.87(0.33)b | | 10.05(0.29)b | 5.46(0.41)a | | 5.21(0.38)b | 5.70(0.22)a | | 5.59(0.25)b | 7.21(1.29)b | | 7.19(1.03)b |
| δ ¹⁸ O-H ₂ O (‰) | JYN -7.33(1.42)a | -8.15(0.39) | -7.92(1.38)a | -9.31(0.38)a | -9.24(0.43) | -8.45(0.36)a | -8.34(0.35)a | -10.51(0.33) | -7.68(0.33)a | -11.07(2.27)a | -9.45(0.15) | -9.22(1.90)a |
| | JYA -9.60(0.42)a | | -8.72(0.73)a | -8.73(0.25)a | | -9.08(0.37)a | -9.20(0.23)a | | -8.73(0.51)a | -8.74(0.22)b | | -7.94(0.26)b |
| | IntC -9.20(0.65)a | | -7.98(1.14)a | -9.52(0.57)a | | -8.19(0.95)a | -8.85(0.48)a | | -9.52(0.36)b | -8.76(2.60)b | | -9.13(1.58)a |
| δ ¹⁵ N-NO ₃ ⁻ (‰) | JYN 6.68(0.09)a | 8.07(0.32) | 6.81(0.07)a | 9.39(0.05)a | 9.35(0.19) | 9.32(0.04)a | 8.55(0.13)a | 8.07(0.17) | 8.73(0.19)a | 7.41(0.01)a | 8.41(0.23) | 4.61(0.02)a |
| | JYA 8.26(0.29)b | | 8.24(0.27)b | 9.68(0.24)b | | 9.47(0.26)b | 8.25(0.13)b | | 8.15(0.20)b | 8.72(0.22)a | | 8.52(0.20)b |
| | IntC 8.16(0.07)b | | 7.86(0.19)a | 9.65(0.08)b | | 9.54(0.22)a | 8.24(0.08)b | | 8.76(0.17)a | 8.21(0.11)a | | 8.35(0.30)b |
| δ ¹⁸ O-NO ₃ ⁻ (‰) | JYN 7.31(1.42)a | 3.46(0.42) | 7.30(1.40)a | 6.14(0.36)a | 5.65(0.57) | 6.65(0.39)a | 8.22(0.33)a | 5.84(0.30) | 8.68(0.29)a | 7.43(0.35)a | 5.71(0.28) | 5.11(0.28)a |
| | JYA 2.89(0.40)b | | 3.25(0.38)b | 6.41(0.54)a | | 6.18(0.52)b | 6.29(0.32)b | | 6.13(0.34)b | 6.13(0.30)b | | 6.09(0.31)a |
| | IntC 5.46(0.52)a | | 6.13(0.44)a | 6.52(0.37)a | | 7.03(0.55)c | 7.84(0.62)a | | 9.21(0.49)b | 6.54(0.35)b | | 5.98(0.43)a |

^a Values reported are means ($n = 4$) and standard errors in the brackets. The different lowercase letters of a and b in each column of upstream and downstream indicate significant differences between rivers in the same sampling position.

Table 2 Spatial pattern comparison of main anion concentration and isotopic fractions among the rivers across all the seasons

| | Cl ⁻ (mg L ⁻¹) | NH ₄ ⁺ (μg L ⁻¹) | NO ₃ ⁻ (mg L ⁻¹) | δ ¹⁸ O-H ₂ O (‰) | δ ¹⁵ N-NO ₃ ⁻ (‰) | δ ¹⁸ O-NO ₃ ⁻ (‰) |
|--------------------|---------------------------------------|--|--|--|--|--|
| Spatial site | AVR ± SD | AVR ± SD | AVR ± SD | AVR ± SD | AVR ± SD | AVR ± SD |
| JYN-U ^a | 22.49 ± 4.08a | 94.84 ± 9.00a | 7.90 ± 2.65a | -9.01 ± 1.59a | 8.01 ± 1.20a | 6.57 ± 1.56a |
| JYN-D | 22.58 ± 1.44a | 84.13 ± 9.32b | 7.07 ± 2.82b | -8.32 ± 0.68a | 8.07 ± 1.15a | 6.94 ± 1.48a |
| JYA-U | 21.94 ± 3.24a | 85.84 ± 5.33a | 6.70 ± 1.58b | -9.07 ± 0.42a | 8.73 ± 0.67a | 5.43 ± 1.70a |
| JYA-M | 21.38 ± 4.21a | 88.08 ± 11.56b | 6.08 ± 2.09b | -9.34 ± 0.97a | 8.46 ± 0.61a | 5.17 ± 1.14b |
| JYA-D | 20.78 ± 5.75a | 84.41 ± 16.50b | 6.38 ± 3.21a | -8.62 ± 0.48a | 8.60 ± 0.60a | 5.41 ± 1.44a |
| Integrated channel | 22.46 ± 2.90a | 90.36 ± 9.31b | 7.24 ± 1.96c | -8.89 ± 0.57a | 8.60 ± 0.67 ab | 6.84 ± 1.20a |

^a JYU-U, the upstream Jiyun river; JYU-D, the downstream Jiyun river; JYA-U, the upstream Jiyang river; JYA-M, the middle reach of Jiyang river; JYA-D, the downstream Jiyang river. Values reported are means ($n \geq 12$) ± standard errors. The different lowercase letters of a and b in each column indicate significant differences among the sampling positions.

NO₃⁻ (+2.89 to +9.21‰) were distributed within the ranges of manure/sewage and the microbial nitrification (Fig. 3). Instead, the data indicated that most of the NO₃⁻ were derived from manure/sewage and microbial nitrification (Fig. 3)

Although all δ¹⁸O-NO₃⁻ values fell in the range typically cited as representing ammonium fertilizer sources (-8‰ < δ¹⁸O-NO₃⁻ < +14‰), whilst, δ¹⁵N-NO₃⁻ values were out of the range for ammonium fertilizer sources (-7‰ < δ¹⁵N-NO₃⁻ < +5‰).^{26,27} The δ¹⁸O-NO₃⁻ values produced by microbial nitrification in watersheds range from -10 to +10‰ (Kendall *et al.*, 2007).¹⁶ All of these sample values had some overlap range of typical soil nitrate (+3‰ < δ¹⁵N-NO₃⁻ < +8‰, -8‰ < δ¹⁸O-NO₃⁻ < +14‰). Soil NO₃⁻ is a product of bacterial decomposition of organic N that results from the decomposition of plants and animals and their products (*i.e.* organic waste). Inorganic or organic N based fertilizers are a primary source of nitrate in agricultural areas, while septic waste plays a significant role in residential areas.^{18,28} The results indicated that the NO₃⁻ for these two rivers could be derived from the nitrification of manure/sewage other than ammonium fertilizers.²⁶⁻²⁸ Furthermore, it did support our hypothesis that different

land uses could result in different NO₃⁻ sources in these two rivers. However, there was no information available on the rates of various fractionation processes, the usage of manure in these two watersheds, or the amount of sewage discharging to the JYN and JYA, and therefore soil NO₃⁻ could not be separated from other NO₃⁻ sources derived from based on δ¹⁵N-NO₃⁻ values alone.




Fig. 1 Plots of NO₃⁻ and Cl⁻ in the Jinyun river and Jiyang river and integrated channel of Jinan, China.




Fig. 2 NO₃⁻ versus δ¹⁵N-NO₃⁻ in spatial sites of these two watershed rivers (A) and in different seasons across the watersheds (B). The ranges in A are from Isotope Hydrology (Joel R Gat, 2010).




Fig. 3 Patterns of $\delta^{18}\text{O}-\text{H}_2\text{O}$ and $\delta^{18}\text{O}-\text{NO}_3^-$ for the Jinyun river and Jinyang river in different seasons.

$\delta^{18}\text{O}-\text{NO}_3^-$ values in these two rivers were within -10 to $+10\text{‰}$ (Fig. 2B), possibly suggesting that most of NO_3^- contributed by soil nitrate could be from microbial nitrification before flowing into the JYN and JYA. Nitrogen-fixing bacteria account for two-thirds of the oxygen atoms in NO_3^- from soil water and one-third from atmospheric oxygen.^{29,30} As shown in Fig. 4, the $\delta^{18}\text{O}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{H}_2\text{O}$ values in these two rivers were highly and positively correlated ($r^2 = 0.393$, $P = 0.04$), further suggesting that the $\delta^{18}\text{O}-\text{NO}_3^-$ values were largely controlled by the $\delta^{18}\text{O}-\text{H}_2\text{O}$ in the river during the microbial nitrification. However, there was no regular seasonal distribution of $\delta^{18}\text{O}-\text{H}_2\text{O}$ in these two rivers compared with the oxygen isotopic composition of dissolved NO_3^- , indicating that the

nitrification capacity from microbial activities might not be always consistent with the seasons change (Fig. 4). The correlations between the $\delta^{18}\text{O}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{H}_2\text{O}$ in the spring and autumn are better than the summer and winter. The possible reasons are the temperatures and water quantities in these two seasons are apt to the microbial nitrification in the river.

Denitrification can occur in anaerobic pockets within a water body and can cause a characteristic increase of both the $\delta^{15}\text{N}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{NO}_3^-$ values in the remaining NO_3^- .^{16,31} The process of ammonia evaporation causes an enrichment of heavier N isotope and hence increases the $\delta^{15}\text{N}$ values.^{7,16} Indeed, these changes of isotopic compositions would correlate with decreasing NO_3^- concentrations as nitrate is being transformed to nitrogen gas and then out of the system. It is important to validate whether the observed changes in the isotopic composition of the remaining NO_3^- correspond to a decrease of the total dissolved NO_3^- in the water body. This can be done by plotting $\delta^{15}\text{N}$ against a concentration scale.³² Although the ratio of changes in $\delta^{15}\text{N}-\text{NO}_3^-$ and $\delta^{18}\text{O}-\text{NO}_3^-$ is not typically close to $1 : 2$, the $\delta^{15}\text{N}-\text{NO}_3^-$ values across these two river water negatively correlated with NO_3^- in the winter in this study (Fig. 2B). The variation in the negative correlation between $\delta^{15}\text{N}-\text{NO}_3^-$ and NO_3^- concentrations showed that the denitrification had happened although the dissolved oxygen is higher in the winter river (Kendall, 1998).²⁶




Fig. 4 Patterns of $\delta^{18}\text{O}-\text{H}_2\text{O}$ and $\delta^{18}\text{O}-\text{NO}_3^-$ for the Jinyun river and Jinyang river in different seasons.

4. Conclusion

Results in this study showed that NO_3^- concentrations had distinct seasonal patterns. They were lower in spring and summer either in the Jinyun river or Jinyang river and were higher in the Jinyun river than the Jinyang river, especially in

spring and winter. Both river waters indicated that NO_3^- concentration deviated from manure/sewage and anthropogenic contamination might be a dominant NO_3^- origination over the watersheds. The sources of NO_3^- could be from microbial nitrification and manure/sewage in the Jinyun river and Jinyang river. The nitrification would have taken place in manure/sewage and soil, but not in the ammonium fertilizer. This study provides a reference in identifying the sources of nitrate in the similar multipurpose watershed around the world. It is necessary to control or reduce nitrate contamination in the waters.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The study was jointly supported by Natural Science Foundation of Shandong Province, China (ZR2017MD022 and ZR2018MD002) and Natural Science Foundation of China (31270586). We thank Bonnie Chapman Beers, working in the SUNY Upstate Medical University, for polishing the language. We are appreciated with Qian Li and Fujun Yue for chemical analyses.

References

- 1 K. S. Lee, Y. S. Bong, D. Lee, Y. Kim and K. Kim, *Sci. Total Environ.*, 2008, **395**, 117–124.
- 2 E. Pastén-Zapata, R. Ledesma-Ruiz, T. Harter, A. Ramírez and J. Mahlknecht, *Total Environ.*, 2014, **470–471**, 855–864.
- 3 C. Liu, S. Li, Y. Lang and H. Xiao, *Environ. Sci. Technol.*, 2006, **40**, 6928–6933.
- 4 S. S. Kaushal, P. M. Groffman, L. E. Band, E. M. Elliott, C. A. Shields and C. Kendall, *Environ. Sci. Technol.*, 2011, **45**, 8225–8232.
- 5 A. Menció, M. Boy and J. Mas-Pla, *Sci. Total Environ.*, 2011, **409**, 3049–3058.
- 6 M. Itoh, Y. Takemon, A. Makabe, C. Yoshimizu, A. Kohzu, N. Ohte, D. Tumurskh, I. ayasu, N. Yoshida and T. Nagata, *Sci. Total Environ.*, 2011, **409**, 1530–1538.
- 7 M. Zhang, Y. Zhi, J. Shi and L. Wu, *Sci. Total Environ.*, 2018, **639**, 1175–1187.
- 8 K. S. Lee, Y. S. Bong, D. Lee, Y. Kim and K. Kim, *Sci. Total Environ.*, 2008, **395**, 117–124.
- 9 M. Alvarez-Cobelas, D. G. Angeler and S. Sanchez-Carrillo, Export of nitrogen from catchments: a worldwide analysis, *Environ. Pollut.*, 2008, **156**, 261–269.
- 10 S. Li, C. Liu, J. Li, X. Liu, B. Chetelat, B. Wang and F. Wang, *Environ. Sci. Technol.*, 2010, **44**, 1753–1758.
- 11 N. Chen and H. Hong, *Biogeochemistry*, 2011, **106**, 311–321.
- 12 L. M. Kellman and C. Hillaire-Marcel, *Agric., Ecosyst. Environ.*, 2003, **95**, 87–102.
- 13 M. L. Cole, K. D. Kroeger, J. W. McClelland and I. Valiela, *Biogeochemistry*, 2006, **77**, 199–215.
- 14 C. C. Y. Chang, C. Kendall, S. R. Silva, W. A. Battaglin and D. H. Campbell, *J. Fish. Aquat. Sci.*, 2002, **59**, 1874–1885.
- 15 M. M. Savard, A. Smirnov, D. Paradis, E. Van Bochove and S. Liao, *J. Hydrol.*, 2010, **381**, 134–141.
- 16 C. Kendall, E. M. Elliott and S. D. Wankel, *Stable isotopes in ecology and environmental science*, 2007, pp. 375–449.
- 17 D. A. Burns and C. Kendall, *Water Resour. Res.*, 2002, **38**, 1051–1061.
- 18 S. Danielescu and K. T. MacQuarrie, *Biogeochemistry*, 2013, **115**, 111–127.
- 19 L. Wang and Y. Qin, *J. Soil Water Conserv.*, 2005, **25**, 87–91.
- 20 M. Liu, Q. Wang, M. Li and D. Liu, *J. Shandong Norm. Univ., Nat. Sci.*, 2000, **15**, 238–240.
- 21 M. Niu, K. Kong and Z. Xu, *J. Jinan Univ., Sci. Technol.*, 2013, **27**, 418–422.
- 22 F. Yue, S. Li, C. Liu, N. An and H. Cai, *Chin. J. Ecol.*, 2012, **31**, 1–6.
- 23 S. Zhang, H. Wang, Q. Chen, B. Yang and L. Yang, *J. Soil Water Conserv.*, 2012, **26**, 169–200.
- 24 W. Wang, S. Qu, Q. Ye and X. Sun, *J. Hydraul. Eng.*, 2011, **42**, 477–489.
- 25 Y. Xia, Y. Li, X. Zhang and X. Yan, *J. Geophys. Res.: Biogeosci.*, 2017, **122**(1), 2–14.
- 26 E. Minet, R. Goodhue, W. Meier-Augenstein, R. Kalin, O. Fenton, K. Richards and C. Coxon, *Water Res.*, 2017, **124**, 85–96.
- 27 C. Kendall and R. Aravena, *Environmental tracers in subsurface hydrology*, 2000, pp. 261–297.
- 28 Z. Jin, Q. Zheng, C. Zhu, Y. Wang, J. Cen and F. Li, *Appl. Geochem.*, 2018, **93**, 10–19.
- 29 D. Marconi, PhD thesis, ProQuest Dissertations Publishing, 2017.
- 30 L. Bristow, M. Altabet, I. Gregory-Eaves and R. Maranger, *Biogeochemistry*, 2017, **135**(3), 221–237.
- 31 G. Anornu and D. Adomako, *Sci. Total Environ.*, 2017, **603–604**, 687–698.
- 32 C. Leibundgut, P. Maloszewski and C. Külls, *Tracers in Hydrology*, 2009, 50–65.