

Src42A modulates tumor invasion and cell death via Ben/dUev1a-mediated JNK activation in *Drosophila*

X Ma^{1,2}, Y Shao^{1,2}, H Zheng^{1,2}, M Li¹, W Li^{*1} and L Xue^{*1}

Loss of the cell polarity gene could cooperate with oncogenic Ras to drive tumor growth and invasion, which critically depends on the c-Jun N-terminal Kinase (JNK) signaling pathway in *Drosophila*. By performing a genetic screen, we have identified Src42A, the ortholog of mammalian Src, as a key modulator of both *Ras*^{V12}/*Igl*^{-/-} triggered tumor invasion and loss of cell polarity gene-induced cell migration. Our genetic study further demonstrated that the Bendless (Ben)/dUev1a ubiquitin E2 complex is an essential regulator of Src42A-induced, JNK-mediated cell migration. Furthermore, we showed that ectopic Ben/dUev1a expression induced invasive cell migration along with increased MMP1 production in wing disc epithelia. Moreover, Ben/dUev1a could cooperate with *Ras*^{V12} to promote tumor overgrowth and invasion. In addition, we found that the Ben/dUev1a complex is required for ectopic Src42A-triggered cell death and endogenous Src42A-dependent thorax closure. Our data not only provide a mechanistic insight into the role of Src in development and disease but also propose a potential oncogenic function for Ubc13 and Uev1a, the mammalian homologs of Ben and dUev1a.

Cell Death and Disease (2013) 4, e864; doi:10.1038/cddis.2013.392; published online 17 October 2013

Subject Category: Cancer

Cancer is the leading cause of death worldwide, causing an estimated total of 577 190 deaths in United States alone in 2012.¹ Cancer-related mortalities are mainly caused by metastasis, rather than the primary tumor that arises from a malignant lesion.² Although significant progress has been made towards the understanding of cancer development, the molecular and genetic mechanisms of tumor metastasis have remained poorly understood. With reduced genome redundancy and amenable genetic tools, *Drosophila melanogaster* has become an excellent model system to investigate the genetic mechanism of cancer biology over the past decades.^{3–6} Several invasion and metastasis models have been established in both adult flies and developing larvae.^{4,7,8} In the eye–antennal discs, expression of oncogenic mutants, such as *Ras*^{V12}, can cooperate with mutants that disrupt cell polarity, such as *scrib* or *lethal giant larvae (Igl)*, to induce invasive tumors into the ventral nerve cord (VNC).^{9–11} In the epithelia of the developing wing disc, loss of the cell polarity gene (*scrib*) or C-terminal SRC kinase (*Csk*) along the anterior/posterior (A/P) boundary using *patched*-Gal4 driver produces an invasive migration phenotype,^{12–14} which has been used to model cell migration *in vivo*.³

The c-Jun N-terminal Kinase (JNK) pathway is evolutionarily conserved from fly to human and has an essential role in regulating a wide range of cellular activities including proliferation, differentiation, migration and apoptosis.¹⁵

Recently, JNK signaling has been shown to have an important role in modulating *Ras*^{V12}/*scrib*^{-/-}-triggered tumor growth and metastasis, as well as *Raf*^{GOF}/*RhoGEF2*-induced tumorigenesis.^{14,16,17}

Src was the first discovered oncogene encoding a non-receptor membrane-associated tyrosine kinase.^{18–20} Nine Src family members have been identified in mammals, whereas the *Drosophila* genome encodes only two Src homologs, Src42A and Src64B.^{21–23} Src42A and Src64B have a redundant role in regulating dorsal closure and are both required for tracheal cell morphogenesis.^{24,25} Recent studies found that Src64B could induce JNK-dependent overgrowth and expression of Yorkie's (Yki) target genes when cell death was blocked by expressing p35.²⁶ When Src64B was overexpressed in a clone context, it induced non-autonomous tumor overgrowth through JNK-dependent propagation of Yki activity.²⁷ Consistent with the well-documented role of Src family kinases in promoting mammalian tumor invasion,²⁸ Cagan and colleagues found that inhibition of Csk, a negative regulator of Src family kinases, triggered JNK signaling-mediated cell invasion in *Drosophila* wing discs.¹³ However, a direct involvement of Src in tumor metastasis and cell invasion, and its underlying mechanisms, remain largely elusive.

We have performed a genetic screen and identified Src42A as a crucial regulator of *Ras*^{V12}/*Igl*^{-/-}-induced tumor

¹Department of Interventional Radiology, Shanghai Key Laboratory of Signaling and Disease Research, Shanghai 10th People's Hospital, School of Life Science and Technology, Tongji University, Shanghai 200092, China

*Corresponding author: L Xue or W Li, Department of Interventional Radiology, Shanghai Key Laboratory of Signaling and Disease Research, School of Life Science and Technology, Shanghai 10th People's Hospital, Tongji University, 1239 Siping Road, Shanghai 200092, China. Tel: +86 21 65985407; Fax: +86 21 65985407; E-mails: lwz@tongji.edu.cn or lei.xue@tongji.edu.cn

²These authors contribute equally to this work.

Keywords: Src42A; invasion; cell death; JNK; Bendless; dUev1a

Abbreviations: VNC, ventral nerve cord; JNK, c-Jun N-terminal Kinase; MMP1, matrix metalloproteinase 1; AEL, after egg laying; A/P, anterior/posterior; AO, acridine orange; MARCM, mosaic analysis with a repressible cell marker; Ey, eyeclose

Received 16.7.13; revised 24.8.13; accepted 10.9.13; Edited by M Agostini




Figure 1 Src42A is required for *Ras-IgI* oncogenic cooperation and loss of *scrib*-induced invasion. (a–c) GFP-labeled clones of cells with indicated genotypes are created in the developing eye–antennal discs. *Ras^{V12}/Igl^{-/-}*-induced tumor growth (a) and invasion of the VNC (a') were dramatically suppressed by expressing Puc (b–b'). The invasion behavior was also strongly suppressed by expressing an *src42A* RNAi (c'), whereas the tumor overgrowth was not significantly changed (c). Scale bars, 200 μ m in a–c and 100 μ m in a'–c'. (d–f) Fluorescence micrographs of wing discs are shown. Compared with the control (d–d''), loss of *scrib*-induced cell invasion and MMP1 expression (e–e'') was suppressed by the expression of an *src42A* RNAi (f–f'). Scale bars, 20 μ m

invasion, as well as the loss of *scrib*-triggered cell migration in development. Our genetic epistasis analysis demonstrated that Src42A modulates JNK signaling-mediated cell invasion, cell death and thorax closure upstream of Bendless (Ben) and dUev1a. Furthermore, we showed that ectopic expression of Ben/dUev1a not only induces cell migration and matrix

metalloproteinase 1 (MMP1) upregulation but also cooperates with *Ras^{V12}* to promote tumor growth and invasive behavior. Together, these data highlight the importance of Src42A-Ben/dUev1a-JNK signaling in regulating cell invasion and extend our knowledge towards the underlying mechanism of Src42A in tumor progression.

Results and discussion

Src42A is required for tumor invasion and cell migration.

As previously described, co-expression of oncogenic Ras (*Ras^{V12}*) in *Igl*-mutant cells in eye–antenna discs using the *ey-FLP/MARCM* (mosaic analysis with a repressible cell marker) system induces strong tumor-like growths (Figure 1a), with invasive migration into the VNC 8 days after egg laying (AEL) (Figure 1a').^{10,16,29} Such animals kept growing as oversized larvae carrying huge tumors in their head and died before pupation (Supplementary Figure S1A). Blocking JNK signaling by expressing the JNK phosphatase Puc dramatically suppressed the growth and invasion behaviors of *Ras^{V12}/Igl^{-/-}* tumors (Figures 1b and b').^{12,16} To identify additional genes required for tumor growth and invasion, we performed a genetic screen for dominant suppressors of the *Ras^{V12}/Igl^{-/-}* tumor progression phenotype and identified dUev1a as a crucial regulator.¹² From this screen, we also found that the loss of *src42A* dramatically suppressed tumor cell invasion into the VNC (Figure 1c') and enabled the animals to survive to the pupa stage (Supplementary Figure S1B), whereas the tumor size remained largely unaffected (compare Figures 1a and c). Loss of *src64B* produced a similar suppression effect (Supplementary Figure S2), consistent with previous finding that the two *Drosophila* Src proteins share redundant functions in development.²⁴

To examine whether Src42A modulates cell invasion in other contexts, we turned to another well-characterized model in the larval wing imaginal disc.^{3,12,13,30–32} Compared with the controls (Figures 1d–d''), RNAi downregulation of the cell polarity gene *scrib* driven by *ptc*-Gal4 triggered strong cell migration phenotype towards the posterior part of the discs (Figure 1e'), along with JNK-dependent upregulation of matrix metalloproteinase1 (MMP1, Figure 1e''),^{12,14} a protein that is essential for basement membrane degradation.^{29,33} We found that the loss of *src42A* dramatically suppressed the epithelial migration phenotype as well as the upregulation of MMP1 (Figures 1f–f''). Taken together, these data indicate that Src42A is required for *Ras^{V12}/Igl^{-/-}*-triggered tumor invasion in the eye discs as well as the loss of cell-polarity-induced cell migration and MMP1 expression in the developing wing.

Src42A induce JNK-mediated cell migration through Ben/dUev1a.

Ectopic expression of Src42A along the A/P boundary of wing imaginal discs driven by *ptc*-Gal4 (*ptc*>Src42A) resulted in strong cell migration phenotype along with the upregulation of MMP1 (Figures 2b–b'''). To examine whether cell invasion is a primary consequence of Src42A activation, or a secondary effect of cell death triggered by elevated Src activity,^{34–36} we expressed the caspase inhibitor p35 to block cell death. Co-expression of




Figure 2 Src42A induces JNK-mediated cell invasion in developing wings. Fluorescence micrographs of wing discs are shown. Compared with the controls (a–a'''), ectopic Src42A expression-induced MMP1 secretion (b') and cell invasion phenotype (b''') were completely suppressed by co-expressing of Bsk^{DN} (c–c'''). Scale bars, 100 μm in a–c and 20 μm in a'''–c'''

p35 with Src42A resulted in a widening of the *ptc* domain and enhanced the production of MMP1 (Supplementary Figure S3). Importantly, these 'undead cells'^{37,38} can still migrate to the posterior part (Supplementary Figure S3), indicating that cell migration is a primary event induced by Src42A expression.

JNK signaling is misregulated in many human cancers and contributes to invasion in different *Drosophila* models.^{7,13,29,31} We found that *ptc*>Src42A-induced cell migration and MMP1 induction (Figure 2b) were significantly suppressed by the expression of a dominant-negative form of Basket (Bsk) (Figure 2c), indicating that JNK signaling is required for Src42A-induced cell migration. We performed genetic epistasis analysis between Src42A and components of JNK signaling and found that *ptc*>Src42A-induced migration phenotype was significantly blocked by the loss of JNK kinase Hep, JNK Kinase Kinase dTAK1 and the E3 ubiquitin ligase dTRAF2 (Figure 3c), suggesting that Src42A induces cell migration upstream of dTRAF2.

Our recent work identified the E2 ubiquitin-conjugating enzyme complex consisting of Bendless (Ben) and dUev1a as the upstream regulators of dTRAF2 in JNK signaling.^{12,39} We explored the genetic interactions between the Src42A and Ben/dUev1a complex and found that *ptc*>Src42A-triggered migration phenotype and MMP1 expression (Figures 3a–a'')

were suppressed strongly by RNAi downregulation of either *Ben* or *dUev1a* alone (Figure 3c), and completely by both (Figures 3 b–b''), suggesting that the Ben/dUev1a complex is necessary for Src42A-induced cell migration. We previously showed that co-expression of Ben and dUev1a resulted in JNK activation;¹² consistently, Ben and dUev1a expression along the A/P boundary induced a mild cell migration phenotype and upregulation of MMP1 (Figures 4a,b and d), which was significantly enhanced by removing one copy of endogenous *puc* (Figure 4c), indicating that Ben/dUev1a expression could induce JNK-dependent cell migration.

Together, the above data suggested that the Ben/dUev1a complex is necessary and sufficient for Src42A-induced JNK-dependent cell migration.

Ben/dUev1a cooperates with *Ras*^{V12} to induce tumor growth and invasion. Gain of function of cell morphology regulators or loss of cell polarity could promote JNK-dependent cell migration and cooperate with *Ras*^{V12} to drive tumor growth and invasion.^{36,40} As Ben/dUev1a expression could also induce JNK-dependent cell migration, we wonder whether Ben/dUev1a could cooperate with *Ras*^{V12} to promote tumor progression. As reported previously, clonal expression of *Ras*^{V12} in eye disc induced benign overgrowth (Figure 4e), with no cells invading into the VNC




Figure 3 Src42A induces cell invasion through Ben/dUev1a. (a–b) Fluorescence micrographs of wing discs are shown. Ectopic Src42A expression-induced cell invasion (a') and MMP1 secretion (a'') were strongly suppressed by co-expressing a *ben* and *dUev1a* RNAi (b–b''). Scale bars, 20 μ m. (c) Quantification of invasion phenotype induced by Src42A expression. Results are shown as mean \pm S.D., $n \geq 30$. Student's *t*-test was used to compute *P*-values, significance is indicated with asterisks (****P* < 0.001, ***P* < 0.01)

(Figure 4e').¹⁶ On the other hand, expression of Ben/dUev1a resulted in no obvious growth or invasion phenotype (Figures 4f and f'). When Ben/dUev1a was simultaneously expressed with *Ras*^{V12}, tumor overgrowth (Figure 4g) and invasion into the VNC region (Figure 4g') were observed in the eye discs 8 days AEL, suggesting that Ben/dUev1a could cooperate with *Ras*^{V12} to promote tumor growth and invasion behavior.

Src42A activates JNK signaling *in vivo*. To investigate whether Src42A activates JNK signaling *in vivo*, we examined JNK phosphorylation and *puc* expression, a read-out of JNK activity,⁴¹ in response to Src42 expression. We found that ectopic expression of Src42A under the *ptc* promoter in the wing disc resulted in strong JNK phosphorylation (compare Figures 5a and b) and expression of a *puc*-LacZ reporter (Figures 5c and d). Furthermore, ectopic




Figure 4 Ben/dUev1a expression promotes tumor growth and metastasis of *Ras*^{V12} cells. (a–d) Fluorescence micrographs of wing discs are shown. Compared with the controls (a), Ben/dUev1a expression induced mild cell invasion phenotype (b) was intensely enhanced by deleting one copy of endogenous *puc* (c). Ben/dUev1a expression under the *ptc* promoter upregulates MMP1 secretion. Crosses were performed at 29 °C. Scale bars, 20 μ m. (e–g) Dorsal views of the cephalic complexes (e–g) and NVCs are shown (e'–g'). *Ras*^{V12}-expressing clones show a moderate growth advantage (e), but never invade into NVC (e'). Clones with elevated Ben/dUev1a activity gave no obvious overgrowth or invasion phenotypes (f–f'). Massive tumor growth (g) and invasion to the VNC (g) were observed by co-expression of *Ras*^{V12} and Ben/dUev1a. Scale bars, 200 μ m in (e–g) and 100 μ m in (e'–g')

Src42A expression driven by *GMR*-Gal4 is sufficient to activate *puc* transcription in developing eye discs (Figures 5e and f). Together, these results indicate that Src42A can activate JNK signaling *in vivo*.

Src42A triggers JNK-dependent cell death in the developing eye. Apart from the role of Src42A in regulating tumor invasion and cell migration showed above, we and others^{35,36} also found that ectopic Src42A expression in developing eyes could induce extensive cell death, as shown with acridine orange (AO) staining (Supplementary Figure S4B) that detects dying cells⁴² and produce a small eye phenotype (Figure 6b). Consistent with the genetic epistasis analysis in cell invasion, the *GMR*>Src42A-triggered cell death and small eye phenotype were dramatically suppressed by inactivation of Ben, dUev1a or downstream components of the JNK signaling (Supplementary Figure S4 and Figures 6c–g). Besides, the ectopic Src42A-caused small eye phenotype was also reverted by co-expressing two




Figure 5 Src42A expression activates JNK *in vivo*. (a–b) Fluorescence micrographs of wing discs are shown. Compared with the controls (a), Src42A expression induced strong JNK phosphorylation (b). Scale bars, 100 μ m. (c–f) X-Gal staining of the *puc-LacZ* reporter gene in the developing wing (c–d) and eye (e–f) are shown. Ectopic Src42A expression under the *ptc* promoter strongly upregulated *puc* expression along the A/P boundary (d). The weak endogenous *puc* expression posterior to the morphogenetic furrow of eye disc (e) was significantly enhanced by expressing a wild-type Src42A (f)

independent RNAi lines against *src42A* (Figure 6h and Supplementary Figure S5), which further confirmed the efficiency of the two RNAi lines.

Ben/dUev1a is physiologically required for Src42A.

Apart from its role in modulating cell migration and cell death, *src42A* also regulates JNK-dependent dorsal closure.²⁴ Consistently, JNK signaling, as revealed by *puc* expression, is activated in the dorsal patch of the third instar larval wing disc (Figure 7a).⁴³ Downregulation of *src42A* under the *pannier* (*pnr*) promoter resulted in reduced *puc* expression in the wing disc (Figure 7b) and produced a cleft phenotype in the adult thorax (Figure 7d). This cleft phenotype resembled that of JNK inactivation²² and could be partially rescued by deleting one copy of endogenous *puc* (Figure 7f), or expression of a wild-type Hep (Hep^{WT}) (Figure 7h), or co-expression of Ben and dUev1a (Figure 7j). Together, these results suggested that the Ben/dUev1a complex is required for the physiological function of Src42A in thorax development.

Conclusion

We have identified Src42A as a key modulator of both *Ras*^{V12}/*Igf*^{-/-}-triggered tumor invasion and loss of cell-polarity-induced cell migration in *Drosophila*. Our genetic evidence established the Ben/dUev1a complex as an essential positive regulator that mediates Src42A-induced, JNK-dependent cell migration and death (Figure 7k). We showed that ectopic Ben/dUev1a expression not only induced MMP1 upregulation and cell migration but also cooperated with *Ras*^{V12} to promote tumor growth and invasion, suggesting that Ben and dUev1a




Figure 6 Src42A induces Ben/dUev1a-mediated cell death in developing eyes. Light micrographs of *Drosophila* adult eyes are shown. Compared with the wild-type eye (a), the *GMR>Src42A* small eye phenotype (b) was dramatically suppressed by the expression of Bsk^{DN} (c), a *hep* RNAi (d), dTAK1^{DN} (e), a *dTRAF2* RNAi (f), reducing *ben* and *dUev1a* activity (g), or a *src42A* RNAi (h)




Figure 7 Src42A modulates endogenous JNK activity upstream of Ben/dUev1a. (a–b) X-Gal staining of the *puc-LacZ* reporter gene in the developing wing are shown. The endogenous *puc* expression pattern in the notum region of wild-type wing disc (a) was reduced by expressing a *src42A* RNAi (b) indicated by the arrow head. (c–j) Light micrographs of *Drosophila* adult thoraxes are shown. Compared with the controls (c), loss of *src42A*-produced strong thorax cleft phenotype (d) was restored by deleting on copy of *puc* (f), overexpression of Hep (h) or overexpression of Ben/dUev1a (j), whereas loss of *puc*, expression of Hep or Ben/dUev1a alone gave no obvious phenotype (e, g, i respectively). (k) A schematic diagram of Ben/dUev1a-JNK signaling in regulating Src42A-induced cell migration and death

are candidate proto-oncogenes. Consistent with our findings, Uev1A, the mammalian homolog of dUev1a, is upregulated in many human cancer cell lines.^{44,45} Our data suggested a link between Src activation and Ben/dUev1a-JNK signaling in regulating cell migration and death in development and disease, which provided beneficial information for mammalian studies and target therapies for cancer.

Materials and Methods

***Drosophila* strains and generation of clones.** All stocks were raised on standard *Drosophila* media, and crosses were performed at 25 °C unless otherwise indicated. Fluorescently labeled invasive tumors were produced in the eye discs as previously described¹⁰ using the following strains: *y, w, ey-Flp; tub-Gal80, FRT40A; Act>y⁺>Gal4, UAS-GFP* (40A tester), *Igf¹ FRT40A UAS-Ras^{V12}* (40A tester) and *ey-Flp, Act>y⁺>Gal4, UAS-GFP*. Additional strains, including *GMR-Gal4, ptc-Gal4, pnr-Gla4, UAS-p35, UAS-GFP* and *UAS-Src42A^{CA}*, were obtained from Bloomington stock center; *UAS-ben-IR* (no. 9413) and *UAS-src42A-IR* (no. 26019) were obtained from the VDRC center; *UAS-dTAK1-IR, UAS-Bsk^{DN}, UAS-hep-IR, UAS-dTRAF2-IR, UAS-Puc, puc^{E69,46} UAS-src42A-IR²⁵ UAS-src64B-IR⁴⁷ UAS-Src42A³⁵ UAS-Ben^{T8}, UAS-dUev1a, UAS-dUev1a-IR* and *UAS-scrib-IR^{12,39}* were previously described. Genotypes of flies used in each figure can be found in online Supplementary Information.

Immunohistochemistry. Antibody staining of imaginal discs was performed as previously described.¹⁶ The following antibodies were used: rabbit anti-phospho-JNK (1:200, Calbiochem, San Diego, CA, USA), mouse anti-MMP1 (1:200, Developmental Studies Hybridoma Bank, Iowa City, IA, USA). Secondary antibodies were anti-rabbit-Alexa (1:1000, Cell Signaling Technology, Danvers, MA, USA) and anti-mouse-Cy3 (1:1000, Jackson Immuno Research, West Grove, PA, USA).

X-gal staining. Eye and wing discs were dissected from the third instar larvae in PBT (1 × PBS pH 7.0, 0.1% Triton X 100) and stained for β-galactosidase activity as described.⁴⁸

AO staining. Eye and wing discs were dissected from the third instar larvae in PBT and incubated in 1 × 10⁻⁵ M AO for 5 min at room temperature prior to imaging.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. We thank Bloomington, VDRC and NIG stock centers for fly stocks, Edward Yao for critical reading of the manuscript, and Yujia Hu for statistical assistance. This work was supported by the National Basic Research Program of China (973 Program) (2010CB944901, 2011CB943903), National Natural Science Foundation of China (31071294, 31171413, 31371490), the Specialized Research Fund for the Doctoral Program of Higher Education of China (20120072110023, 20120072120030) and Shanghai Committee of Science and Technology (09DZ2260100).

1. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012; **62**: 10–29.
2. Valastyan S, Weinberg RA. Tumor metastasis: molecular insights and evolving paradigms. *Cell* 2011; **147**: 275–292.
3. Miles WO, Dyson NJ, Walker JA. Modeling tumor invasion and metastasis in *Drosophila*. *Dis Model Mech* 2011; **4**: 753–761.
4. Gonzalez C. *Drosophila melanogaster*: a model and a tool to investigate malignancy and identify new therapeutics. *Nat Rev Cancer* 2013; **13**: 172–183.

5. Willoughby LF, Schlosser T, Manning SA, Parisot JP, Street IP, Richardson HE *et al*. An *in vivo* large-scale chemical screening platform using *Drosophila* for anti-cancer drug discovery. *Dis Model Mech* 2013; **6**: 521–529.
6. Elsum I, Yates L, Humbert PO, Richardson HE. The Scribble-Dlg-Lgl polarity module in development and cancer: from flies to man. *Essays Biochem* 2012; **53**: 141–168.
7. Rudrapatna VA, Cagan RL, Das TK. *Drosophila* cancer models. *Dev Dyn* 2012; **241**: 107–118.
8. Brumby AM, Richardson HE. Using *Drosophila melanogaster* to map human cancer pathways. *Nat Rev Cancer* 2005; **5**: 626–639.
9. Brumby AM, Richardson HE. scribble mutants cooperate with oncogenic Ras or Notch to cause neoplastic overgrowth in *Drosophila*. *EMBO J* 2003; **22**: 5769–5779.
10. Pagliarini RA, Xu T. A genetic screen in *Drosophila* for metastatic behavior. *Science* 2003; **302**: 1227–1231.
11. Wu M, Pastor-Pareja JC, Xu T. Interaction between Ras(V12) and scribbled clones induces tumour growth and invasion. *Nature* 2010; **463**: 545–548.
12. Ma X, Yang L, Yang Y, Li M, Li W, Xue L. dUev1a modulates TNF-JNK mediated tumor progression and cell death in *Drosophila*. *Dev Biol* 2013; **380**: 211–221.
13. Vidal M, Larson DE, Cagan RL. Csk-deficient boundary cells are eliminated from normal *Drosophila* epithelia by exclusion, migration, and apoptosis. *Dev Cell* 2006; **10**: 33–44.
14. Cordero JB, Macagno JP, Stefanatos RK, Strathdee KE, Cagan RL, Vidal M. Oncogenic Ras diverts a host TNF tumor suppressor activity into tumor promoter. *Dev Cell* 2010; **18**: 999–1011.
15. Weston CR, Davis RJ. The JNK signal transduction pathway. *Curr Opin Cell Biol* 2007; **19**: 142–149.
16. Igaki T, Pagliarini RA, Xu T. Loss of cell polarity drives tumor growth and invasion through JNK activation in *Drosophila*. *Curr Biol* 2006; **16**: 1139–1146.
17. Khoo P, Allan K, Willoughby L, Brumby AM, Richardson HE. In *Drosophila*, RhoGEF2 cooperates with activated Ras in tumorigenesis through a pathway involving Rho1-Rok-Myosin-II and JNK signalling. *Dis Model Mech* 2013; **6**: 661–678.
18. Yeatman TJ. A renaissance for SRC. *Nat Rev Cancer* 2004; **4**: 470–480.
19. Thomas SM, Brugge JS. Cellular functions regulated by Src family kinases. *Annu Rev Cell Dev Biol* 1997; **13**: 513–609.
20. Parsons SJ, Parsons JT. Src family kinases, key regulators of signal transduction. *Oncogene* 2004; **23**: 7906–7909.
21. Simon MA, Drees B, Kornberg T, Bishop JM. The nucleotide sequence and the tissue-specific expression of *Drosophila* c-src. *Cell* 1985; **42**: 831–840.
22. Takahashi F, Endo S, Kojima T, Saigo K. Regulation of cell-cell contacts in developing *Drosophila* eyes by Dsrc41, a new, close relative of vertebrate c-src. *Genes Dev* 1996; **10**: 1645–1656.
23. Dodson GS, Guarneri DJ, Simon MA. Src64 is required for ovarian ring canal morphogenesis during *Drosophila* oogenesis. *Development* 1998; **125**: 2883–2892.
24. Tateno M, Nishida Y, Adachi-Yamada T. Regulation of JNK by Src during *Drosophila* development. *Science* 2000; **287**: 324–327.
25. Shindo M, Wada H, Kaido M, Tateno M, Aigaki T, Tsuda L *et al*. Dual function of Src in the maintenance of adherens junctions during tracheal epithelial morphogenesis. *Development* 2008; **135**: 1355–1364.
26. Fernandez BG, Jezowska B, Janody F. *Drosophila* actin-Capping Protein limits JNK activation by the Src proto-oncogene. *Oncogene* 2013; e-pub ahead of print 3 May 2013; doi:10.1038/onc.2013.155.
27. Enomoto M, Igaki T. Src controls tumorigenesis via JNK-dependent regulation of the Hippo pathway in *Drosophila*. *EMBO Rep* 2013; **14**: 65–72.
28. Guarino M. Src signaling in cancer invasion. *J Cell Physiol* 2010; **223**: 14–26.
29. Uhlirova M, Bohmann D. JNK- and Fos-regulated Mmp1 expression cooperates with Ras to induce invasive tumors in *Drosophila*. *EMBO J* 2006; **25**: 5294–5304.
30. Rudrapatna VA, Bangi E, Cagan RL. Caspase signalling in the absence of apoptosis drives Jnk-dependent invasion. *EMBO Rep* 2013; **14**: 172–177.
31. Rudrapatna VA, Bangi E, Cagan RL. A Jnk-Rho-Actin remodeling positive feedback network directs Src-driven invasion. *Oncogene* 2013; e-pub ahead of print 8 July 2013; doi:10.1038/onc.2013.232.
32. Portela M, Richardson HE. Death takes a holiday—non-apoptotic role for caspases in cell migration and invasion. *EMBO Rep* 2013; **14**: 107–108.
33. Srivastava A, Pastor-Pareja JC, Igaki T, Pagliarini R, Xu T. Basement membrane remodeling is essential for *Drosophila* disc eversion and tumor invasion. *Proc Natl Acad Sci USA* 2007; **104**: 2721–2726.
34. Singh J, Aaronson SA, Mlodzik M. *Drosophila* Abelson kinase mediates cell invasion and proliferation through two distinct MAPK pathways. *Oncogene* 2010; **29**: 4033–4045.
35. Pedraza LG, Stewart RA, Li DM, Xu T. *Drosophila* Src-family kinases function with Csk to regulate cell proliferation and apoptosis. *Oncogene* 2004; **23**: 4754–4762.
36. Vidal M, Warner S, Read R, Cagan RL. Differing Src signaling levels have distinct outcomes in *Drosophila*. *Cancer Res* 2007; **67**: 10278–10285.
37. Huh JR, Guo M, Hay BA. Compensatory proliferation induced by cell death in the *Drosophila* wing disc requires activity of the apical cell death caspase Dronc in a nonapoptotic role. *Curr Biol* 2004; **14**: 1262–1266.
38. Ryoo HD, Gorenc T, Steller H. Apoptotic cells can induce compensatory cell proliferation through the JNK and the Wingless signaling pathways. *Dev Cell* 2004; **7**: 491–501.
39. Ma X, Huang J, Yang L, Yang Y, Li W, Xue L. NOPO modulates Egr-induced JNK-independent cell death in *Drosophila*. *Cell Res* 2012; **22**: 425–431.
40. Brumby AM, Goulding KR, Schlosser T, Loi S, Galea R, Khoo P *et al*. Identification of novel Ras-cooperating oncogenes in *Drosophila melanogaster*: a RhoGEF/Rho-family/JNK pathway is a central driver of tumorigenesis. *Genetics* 2011; **188**: 105–125.
41. Agnes F, Suzanne M, Noselli S. The *Drosophila* JNK pathway controls the morphogenesis of imaginal discs during metamorphosis. *Development* 1999; **126**: 5453–5462.
42. Abrams JM, White K, Fessler LI, Steller H. Programmed cell death during *Drosophila* embryogenesis. *Development* 1993; **117**: 29–43.
43. Campuzano S, Modolell J. Patterning of the *Drosophila* nervous system: the achaete-scute gene complex. *Trends Genet* 1992; **8**: 202–208.
44. Xiao W, Lin SL, Broomfield S, Chow BL, Wei YF. The products of the yeast MMS2 and two human homologs (hMMS2 and CROC-1) define a structurally and functionally conserved Ubc-like protein family. *Nucleic Acids Res* 1998; **26**: 3908–3914.
45. Syed NA, Andersen PL, Warrington RC, Xiao W. Uev1A a ubiquitin conjugating enzyme variant, inhibits stress-induced apoptosis through NF-kappaB activation. *Apoptosis* 2006; **11**: 2147–2157.
46. Xue L, Igaki T, Kuranaga E, Kanda H, Miura M, Xu T. Tumor suppressor CYLD regulates JNK-induced cell death in *Drosophila*. *Dev Cell* 2007; **13**: 446–454.
47. Pechkovsky A, Lahav M, Bitman E, Salzberg A, Kleinberger T. E4orf4 induces PP2A- and Src-dependent cell death in *Drosophila melanogaster* and at the same time inhibits classic apoptosis pathways. *Proc Natl Acad Sci USA* 2013; **110**: E1724–E1733.
48. Xue L, Noll M. *Drosophila* female sexual behavior induced by sterile males showing copulation complementation. *Proc Natl Acad Sci USA* 2000; **97**: 3272–3275.



Cell Death and Disease is an open-access journal published by Nature Publishing Group. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies this paper on Cell Death and Disease website (<http://www.nature.com/cddis>)