

Integrating surgical and clinical andrology is essential to improve the quality of care delivered to infertile couples

Sandro C. Esteves¹, Ahmad Majzoub², Ashok Agarwal³

¹ANDROFERT, Andrology and Human Reproduction Clinic, Referral Center for Male Reproduction, Campinas, SP, Brazil; ²Department of Urology, Hamad Medical Corporation, Doha, Qatar; ³American Center for Reproductive Medicine, Cleveland Clinic, Cleveland, OH, USA

Correspondence to: Sandro C. Esteves. Director, ANDROFERT, Andrology and Human Reproduction Clinic, Referral Center for Male Reproduction, Av. Dr. Heitor Penteado 1464, Campinas, SP 13075-460, Brazil. Email: s.esteves@androfert.com.br.

Response to: Yovich JL, Keane KN. Assessing the male in fertility clinics—men undervalued, undermanaged and undertreated. *Transl Androl Urol* 2017;6:S624-8.

Submitted May 10, 2017. Accepted for publication May 11, 2017.

doi: 10.21037/tau.2017.05.37

View this article at: <http://dx.doi.org/10.21037/tau.2017.05.37>

Drs. Yovich and Keane's commentary (1) about the recently published clinical practice guidelines for sperm DNA fragmentation (SDF) testing based on clinical scenarios (2) is an excellent piece of writing. The authors commence by acknowledging the importance of male infertility evaluation and integration of clinical and laboratory andrology in the modern workup of infertile couples. However, they pointed out that current practice values the male factor only if semen analysis is severely abnormal in contrast to the usual comprehensive female infertility evaluation. Given the success of ICSI, male infertility is indeed often neglected. This factor is aggravated by the limitations of semen analysis, to which we concurred and discussed elsewhere (3-6). To illustrate their point, Yovich and Keane discussed the often debated issue of varicocele and male infertility. Despite overwhelming evidence confirming the adverse effect of varicocele on several sperm markers and the benefit of varicocelectomy in selected men (7-12), only recently—and after more than 150 years since the first publication about varicocele—the Cochrane review confirmed that there might be a benefit to performing varicocelectomy in subfertile men (13). Regrettably, many subfertile men could have benefitted from treatment; the benefit of performing varicocele repair using microsurgery techniques has been advocated for a long time by eminent urological microsurgeons (14-16). Along these lines, Dr. Yovich himself contributed significantly to the evolution of laboratory and clinical andrology in his distinguished career in reproductive

medicine for over 35 years (17). His seminal works on human sperm function in the 90's had a tremendous impact on clinical practice and set the path for future research (18-22).

Drs. Yovich and Keane go on by suggesting ways to integrate clinical andrology as an essential element in evaluation of infertile couple. A comprehensive male evaluation including sperm function testing, such as SDF, ultrasound, use of microsurgical techniques for varicocele repair, and incorporation of clinical and surgical andrology are amongst their proposals. Additionally, the authors shared their experience in Australia, where the lack of urologists dedicated to male infertility represents a significant shortcoming to improved clinical care. To overcome this fact that seems to plague many countries alike, the authors propose the integration of both reproductive endocrinology and andrology within the scope of reproductive medicine fellowship programs. Obviously, this proposition needs to be analyzed in the perspective of each country due to the existence of plain boundaries limiting the practice of gynecologists and urologists. But if successful, urologists may become an integral part of reproductive care rather than serving as mere technicians performing sperm retrievals. Lastly, Yovich and Keane stressed the importance of quality management to fertility centers willing to improve their quality of care for both male and female, an element that we also feel to be essential as discussed elsewhere (23).

Yovich and Keane overall supported the use of SDF

testing in all clinical scenarios presented by Agarwal *et al.* (2). However, they have reservations to bypassing the epididymis as this organ is necessary for sperm maturation. The authors' preference is to search for debris-free micro-epididymal sperm aspiration (MESA) samples containing motile spermatozoa in preference to testicular sperm extraction (TESE). We, in contrast, advocate the use of testicular sperm in preference over ejaculated and testicular sperm in men with high SDF in semen undergoing ART, provided all measures to reduce DNA damage have been attained (2). The reason is that SDF rates in testicular sperm are significantly lower than both testicular sperm and epididymal sperm (24) [(reviewed by Esteves *et al.* (25)]. With regards to epididymal sperm, in particular, Steele *et al.*, using the Comet assay, showed that DNA integrity was higher (83.0%±1.2%) in testicular specimens of men with obstructive azoospermia than in proximal epididymal counterparts (75.4%±2.3%; $P<0.05$) (26). Their results were corroborated by Suganuma *et al.*, who used an experimental mice model to demonstrate that the passage of sperm through the epididymis was associated with a loss of sperm DNA integrity and fertilizing capacity (27). The results of these studies and others (28) indicate that SDF may be triggered during sperm transport through the epididymis as a result of excessive oxidative stress. Therefore, the use of testicular sperm for ICSI in preference over epididymal and ejaculated sperm becomes attractive as the probability of selecting spermatozoa free of DNA damage can be increased (25).

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Yovich JL, Keane KN. Assessing the male in fertility clinics—men undervalued, undermanaged and undertreated. *Transl Androl Urol* 2017;6:S624-8.
2. Agarwal A, Majzoub A, Esteves SC, et al. Clinical utility of sperm DNA fragmentation testing: practice recommendations based on clinical scenarios. *Transl Androl Urol* 2016;5:935-50.
3. Esteves SC, Hamada A, Kondray V, et al. What every gynecologist should know about male infertility: an update. *Arch Gynecol Obstet* 2012;286:217-29.
4. Esteves SC. A clinical appraisal of the genetic basis in unexplained male infertility. *J Hum Reprod Sci* 2013;6:176-82.
5. Esteves SC. Clinical relevance of routine semen analysis and controversies surrounding the 2010 World Health Organization criteria for semen examination. *Int Braz J Urol* 2014;40:443-53.
6. Esteves SC, Sharma RK, Gosálvez J, et al. A translational medicine appraisal of specialized andrology testing in unexplained male infertility. *Int Urol Nephrol* 2014;46:1037-52.
7. Esteves SC, Agarwal A. Afterword to varicocele and male infertility: current concepts and future perspectives. *Asian J Androl* 2016;18:319-22.
8. Cho CL, Esteves SC, Agarwal A. Novel insights into the pathophysiology of varicocele and its association with reactive oxygen species and sperm DNA fragmentation. *Asian J Androl* 2016;18:186-93.
9. Majzoub A, Esteves SC, Gosálvez J, et al. Specialized sperm function tests in varicocele and the future of andrology laboratory. *Asian J Androl* 2016;18:205-12.
10. Esteves SC, Gosálvez J, López-Fernández C, et al. Diagnostic accuracy of sperm DNA degradation index (DDSi) as a potential noninvasive biomarker to identify men with varicocele-associated infertility. *Int Urol Nephrol* 2015;47:1471-7.
11. Esteves SC, Roque M, Agarwal A. Outcome of assisted reproductive technology in men with treated and untreated varicocele: systematic review and meta-analysis. *Asian J Androl* 2016;18:254-8.
12. Tiseo BC, Esteves SC, Cocuzza MS. Summary evidence on the effects of varicocele treatment to improve natural fertility in subfertile men. *Asian J Androl* 2016;18:239-45.
13. Kroese AC, de Lange NM, Collins JA, et al. Surgery or embolization for varicocele in subfertile men. *Cochrane database Syst Rev* 2012;10:CD000479.
14. Marmar JL. The evolution and refinements of varicocele surgery. *Asian J Androl* 2016;18:171-8.
15. Schlegel PN, Goldstein M. Alternate indications for varicocele repair: non-obstructive azoospermia, pain, androgen deficiency and progressive testicular dysfunction. *Fertil Steril* 2011;96:1288-93.
16. Mehta A, Goldstein M. Microsurgical varicocelectomy: a review. *Asian J Androl* 2013;15:56-60.
17. Yovich JL. A clinician's personal view of assisted

- reproductive technology over 35 years. *Reprod Biol* 2011;11 Suppl 3:31-42.
18. Yovich JM, Edirisinghe WR, Yovich JL. Use of the acrosome reaction to ionophore challenge test in managing patients in an assisted reproduction program: a prospective, double-blind, randomized controlled study. *Fertil Steril* 1994;61:902-10.
 19. Yovich JL. Pentoxifylline: actions and applications in assisted reproduction. *Hum Reprod* 1993;8:1786-91.
 20. Edirisinghe WR, Junk S, Yovich JM, et al. Sperm stimulants can improve fertilization rates in male-factor cases undergoing IVF to the same extent as micromanipulation by partial zona dissection (PZD) or subzonal sperm insemination (SUZI): a randomized controlled study. *J Assist Reprod Genet* 1995;12:312-8.
 21. Hirsh AV, Mills C, Bekir J, et al. Factors influencing the outcome of in-vitro fertilization with epididymal spermatozoa in irreversible obstructive azoospermia. *Hum Reprod*. 1994;9:1710-6.
 22. Edirisinghe WR, Junk SM, Matson PL, et al. Changes in motility patterns during in-vitro culture of fresh and frozen/thawed testicular and epididymal spermatozoa: implications for planning treatment by intracytoplasmic sperm injection. *Hum Reprod* 1996;11:2474-6.
 23. Bento FC, Esteves SC. Establishing a quality management system in a fertility center: experience with ISO 9001. *MedicalExpress* 2016;3:M160302.
 24. Esteves SC, Sánchez-Martín F, Sánchez-Martín P, et al. Comparison of reproductive outcome in oligozoospermic men with high sperm DNA fragmentation undergoing intracytoplasmic sperm injection with ejaculated and testicular sperm. *Fertil Steril* 2015;104:1398-405.
 25. Esteves SC, Roque M, Garrido N. Use of testicular sperm for intracytoplasmic sperm injection in men with high sperm DNA fragmentation: a SWOT analysis. *Asian J Androl* 2017. [Epub ahead of print].
 26. Steele EK, McClure N, Maxwell RJ, et al. A comparison of DNA damage in testicular and proximal epididymal spermatozoa in obstructive azoospermia. *Mol Hum Reprod* 1999;5:831-5.
 27. Suganuma R, Yanagimachi R, Meistrich ML. Decline in fertility of mouse sperm with abnormal chromatin during epididymal passage as revealed by ICSI. *Hum Reprod* 2005;20:3101-8.
 28. Muratori M, Tamburrino L, Marchiani S, et al. Investigation on the origin of sperm DNA fragmentation: role of apoptosis, immaturity and oxidative stress. *Mol Med* 2015;21:109-22.

Cite this article as: Esteves SC, Majzoub A, Agarwal A. Integrating surgical and clinical andrology is essential to improve the quality of care delivered to infertile couples. *Transl Androl Urol* 2017;6(Suppl 4):S629-S631. doi: 10.21037/tau.2017.05.37