

RESEARCH ARTICLE

Acceleration and sprint profiles of professional male football players in relation to playing position

José M. Oliva-Lozano¹✉*, Víctor Fortes²✉, Peter Krstrup³✉, José M. Muñor^{1,4}✉

1 Health Research Centre, University of Almería, Almería, Spain, **2** UD Almería, Almería, Spain, **3** Department of Sports Science and Clinical Biomechanics, Faculty of Health Sciences, University of Southern Denmark, Odense, Denmark, **4** Laboratory of Kinesiology, Biomechanics and Ergonomics (KIBIOMER Lab.), Research Central Services, University of Almería, Almería, Spain

✉ These authors contributed equally to this work.

✉ Current address: University of Almería, Edificio Central, Almería, Spain

* jol908@ual.es



OPEN ACCESS

Citation: Oliva-Lozano JM, Fortes V, Krstrup P, Muñor JM (2020) Acceleration and sprint profiles of professional male football players in relation to playing position. PLoS ONE 15(8): e0236959. <https://doi.org/10.1371/journal.pone.0236959>

Editor: Chris Harnish, Mary Baldwin University Murphy Deming College of Health Sciences, UNITED STATES

Received: March 18, 2020

Accepted: July 16, 2020

Published: August 6, 2020

Copyright: © 2020 Oliva-Lozano et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: José M. Oliva-Lozano was supported by a scholarship funded by the Spanish Ministry of Science, Innovation and Universities (FPU18/04434). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

The study aims were to describe positional differences in the acceleration and sprint profiles of professional football players in match-play, and analyse start speeds required based on the intensity of accelerations and decelerations. This longitudinal study was conducted over thirteen competitive microcycles in a professional football team from LaLiga 123. Data were collected through electronic performance tracking systems. Every player was categorised based on the playing position: central defender (CD), full-back (FB), forward (FW), midfielder (MF), and wide midfielder (WMF). In respect of acceleration profile, positional differences were found for all variables ($p < 0.05$), except average magnitude of accelerations (ACC_{AVG} , $p = 0.56$) and decelerations (DEC_{AVG} , $p = 0.76$). The sprint profile also showed positional differences for all variables ($p < 0.05$), apart from sprint duration ($p = 0.07$). In addition, although low-intensity accelerations required significantly greater start speeds (V_0) than high-intensity accelerations in WMF (0.4 ± 0.2 km/h; $p < 0.05$) and FW (0.4 ± 0.2 km/h; $p < 0.05$), no significant differences ($p > 0.05$) were found in CD, FB, and MF. However, high-intensity decelerations were performed at significantly higher V_0 than low-intensity decelerations in MF (2.65 ± 0.1 km/h; $p < 0.05$), FW (3.3 ± 0.1 km/h; $p < 0.05$), FB (3.9 ± 0.4 km/h; $p < 0.05$), WMF (4.3 ± 0.3 km/h; $p < 0.05$), and CD (4.1 ± 0.7 km/h; $p < 0.05$). Therefore, positional differences exist for most variables of the acceleration and sprint profiles. In addition, different V_0 were observed between high-intensity and low-intensity accelerations as well as high-intensity and low-intensity decelerations.

Introduction

During the past decade, there has been an increase in the literature related to athlete monitoring [1,2]. The large number of electronic performance tracking systems available on the market [2] has allowed a detailed understanding of football match demands, enabling coaches to

achieve optimal training targets [1]. Football is a team sport that combines intermittent periods of high-intensity activity with longer periods of lower-intensity activity [3,4]. Professional football players cover around 10 km per match [3,5], but only 10% of the total distance is performed at high-intensity [3]. However, these high-intensity periods contribute in particular to neuromuscular fatigue, consequently increasing the risk of injury [6].

High-speed running actions, or sprints, are considered a prerequisite for successful performance in football [7,8]. In fact, sprinting skills are of prime importance in modern football [7,9]. For instance, straight sprints are the actions most frequently performed when scoring a goal [10], evading an opponent, and creating a shot on goal [11]. Thus, selection, testing, and physical conditioning of players should put emphasis on developing sprinting skills [8]. In addition, careful monitoring of these actions is necessary [9,12], taking into consideration different playing positions [13,14]. For example, a wide midfielder (WMF) may cover 294 ± 76 m of sprinting distance per match, whereas a central defender (CD) may cover 123 ± 48 m [13]. However, research on the sprint profile of professional football match-play has so far been limited [13], while different components of the sprint profile, such as sprint duration, start speed of each sprint or distance covered per sprint, have not yet been studied.

In addition, another determinant factor of football performance is the acceleration profile [7,8,13]. The acceleration profile is understood as a group of acceleration-based variables that are physically demanding [15] given the rate of change in velocity performed by the player [16]. High-intensity accelerations and decelerations have a significant impact on football players' mechanical load [12,17] and indicators of muscle damage post-match [18]. Accelerations have a high metabolic cost [19], while decelerations increase the mechanical load [20]. Moreover, these actions are significantly associated with the neuromuscular fatigue [12] and rating of perceived exertion (RPE) [21]. For example, a previous investigation reported that the total of accelerations performed by professional football players in training sessions was significantly correlated with their session RPE [21]. Consequently, previous research suggests that understanding the acceleration profile would help to understand the potential impact that this profile might have on match performance and risk of injury [12].

Some studies have tried to individualise (based on playing position) and contextualise both acceleration [13,15,20,22] and sprint profiles [7,13,14], but most studies have analysed these profiles separately. Also, little is known, for example, about the start speed required to perform a high-intensity acceleration or deceleration [23]. The aims of this study were therefore to: 1) describe the acceleration profile of players and compare it by playing position; 2) describe the sprint profile of players and compare it by playing position; and 3) analyse the start speed (V_0) required based on the intensity of the acceleration and deceleration by playing position. Regarding the first and second aims, we hypothesised that greater positional differences may be found, particularly, between defensive and offensive positions. When it comes to the third aim, we hypothesised that high intensity accelerations and decelerations would elicit greater start speeds than low intensity accelerations or decelerations.

Materials and methods

Study design

The study was conducted over thirteen competitive microcycles in a professional football team from LaLiga 123. The team played one match per microcycle. The match location (home or away) alternated with each microcycle; seven matches were played away and six at home. The playing formation was 4-4-2 for all matches. Data were collected using wearable sensors (Real-Track Systems, Almería, Spain). In addition, every player was categorised based on their

playing position: central defender (CD), full-back (FB), forward (FW), midfielder (MF), and wide midfielder (WMF).

Subjects

Twenty-three professional male football players (age: 26.79 ± 3.78 years; height: 180.81 ± 6.20 cm; weight: 75.71 ± 6.88 kg; professional career: 8.40 ± 2.80 years) voluntarily took part in the study. Each player had a specific playing position: CD ($n = 4$), FW ($n = 5$), FB ($n = 5$), MF ($n = 5$) and WMF ($n = 4$). Given the very different nature of goalkeeping, this position was not included in the study [24]. Additionally, only players who completed the total duration of the match were analysed. Consequently, although a total of 13 matches were analyzed, not all the players from the sample participated in all the matches. The club allowed the research team to access players' data and informed consent was provided. The study was conducted ethically according to Declaration of Helsinki and it was approved by the Bioethics Committee at the University of Almeria.

Procedures

Data were collected using WIMU Pro 10 Hz global positioning system (GPS) devices (RealTrack Systems, Almería, Spain). This device also contains inertial sensors (four 3D accelerometers, three 3D gyroscopes, one 3D magnetometer and one barometer), which collected data at 100Hz. The validity and reliability of this device has been analysed for the collection of time-motion variables and is considered a suitable instrument for this purpose in football [25]. Regarding the validity of the device, the total bias in mean velocity measurement was between 1.18 and 1.32 km/h while the bias in distance was between 2.32 and 4.32 m [25]. In addition, good inter-unit and intra-unit reliability was reported (intraclass correlation coefficients > 0.93) [25]. The devices were calibrated according to the manufacturer's instructions before the start of each match. All the devices were placed in the Smart Station (RealTrack Systems, Almería, Spain). First, the battery of the devices had to be fully charged. Then, a flat surface was found without any nearby magnetic devices in order to turn on the devices. After 60 seconds, the recording button was pressed. Once the calibration procedure was complete, the devices were placed in a vertical position in the back pocket of a chest vest (Rasán, Valencia, Spain).

The devices were placed in the Smart Station (RealTrack Systems, Almería, Spain) at the end of the match in order to transfer the data to the SPro software (RealTrack Systems, Almería, Spain). This software reported a database with the performance variables which were categorised into the acceleration and sprint profile as indicated in Table 1. In addition, the start speed of the action (V_0) for each high or low-intensity acceleration and deceleration was calculated in order to investigate the third aim of this study. This variable (i.e., V_0) was obtained from the "Sprint Extended" section of the "Intervals Pro" report, which was created by SPro (RealTrack Systems, Almería, Spain).

Statistical analysis

First, descriptive statistics were produced for all variables of the acceleration and sprint profiles. Playing position was set as an independent variable. Then, the Shapiro-Wilk test was used to analyse the normality of the variables and Levene's test for homogeneity. Parametric and non-parametric tests were used, since only ACC_{DIS} , ACC_{LOW} , DEC_{LOW} , $DIFF_{ACDC}$, V_{MAX} and sprint time were variables with normal distribution. On the one hand, when comparing the acceleration and sprint profiles relative to playing position, one-way analysis of variance (ANOVA) with Bonferroni post-hoc and Kruskal Wallis tests was used. On the other

Table 1. Description of acceleration and sprint profile variables.

Profile	Variable	Definition
Acceleration	ACC _{DIS}	Total distance covered by accelerations (m)
	DEC _{DIS}	Total distance covered by decelerations (m)
	ACC _{LOW}	Total number of low-intensity accelerations (below 3 m/s ²)
	ACC _{HIGH}	Total number of high-intensity accelerations (above 3 m/s ²)
	DEC _{LOW}	Total number of low-intensity decelerations (above -3 m/s ²)
	DEC _{HIGH}	Total number of high-intensity decelerations (below -3 m/s ²)
	DIFF _{ACDC}	ACC _{HIGH} —DEC _{HIGH}
	ACC _{AVG}	Average magnitude of accelerations (m/s ²)
	DEC _{AVG}	Average magnitude of decelerations (m/s ²)
	ACC _{MAX}	Maximum magnitude of accelerations (m/s ²)
DEC _{MAX}	Maximum magnitude of decelerations (m/s ²)	
Sprint	SPA	Total sprint actions (above 24 km/h)
	SPD	Total distance covered by sprinting (above 24 km/h)
	SPD _{AVG}	Average distance covered per sprint (above 24 km/h)
	V _{MAX}	Maximum speed reached in the match (km/h)
	Sprint time	Duration of sprint (s)

<https://doi.org/10.1371/journal.pone.0236959.t001>

hand, when comparing the intensity of the accelerations (low-intensity accelerations, high-intensity accelerations, low-intensity decelerations, and high-intensity decelerations) based on start speed (V_0), the Mann-Whitney U test was used. The statistical power, which was calculated by G*Power software (Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany), was greater than 0.85 in all the variables that were analyzed with the sample size of this study. The level of significance was set at $p \leq 0.05$ and the statistical analysis was carried out using IBM SPSS Statistics version 25 (SPSS Inc., Chicago, IL, USA). Effect sizes (ES) were also calculated and categorised as trivial (0–0.19), small (0.20–0.49), moderate (0.50–0.79) and large (0.80 or higher) effect [26].

Results

Table 2 shows the descriptive statistics of the data collected for all variables of the acceleration profile and the positional differences. Significant positional differences ($p < 0.05$) were found for all variables, except DIFF_{ACDC} ($F_4 = 1.15$; $p = 0.33$), ACC_{AVG} ($p = 0.56$) and DEC_{AVG} ($p = 0.76$). WMF was the position with the greatest ACC_{DIS} covered (436.5 ± 86.3 m; $F_4 = 13.63$; $p < 0.05$; $ES = 0.9$ – 2.3) and resulted in a greater DEC_{DIS} covered compared to CD (104.4 ± 17.2 m; $p < 0.05$; $ES = 1.7$), FW (75.2 ± 17.2 m; $p < 0.05$; $ES = 1.1$), and MF (105.7 ± 16.9 m; $p < 0.05$; $ES = 1.6$). In addition, WMF performed greater ACC_{HIGH} than CD (8.5 ± 2.1 ; $p < 0.05$; $ES = 1.3$) and MF (7.8 ± 2.0 ; $p < 0.05$; $ES = 1.3$), greater DEC_{HIGH} than CD (13.6 ± 3.6 ; $p < 0.05$; $ES = 1.1$), and greater ACC_{MAX} (0.3 ± 0.1 m/s²; $p < 0.05$; $ES = 0.6$) and DEC_{MAX} (0.4 ± 0.2 m/s²; $p < 0.05$; $ES = 0.5$) than MF.

MF showed significantly greater ACC_{LOW} than WMF (62.1 ± 14.7 ; $F_4 = 5.93$; $p < 0.05$; $ES = 1.1$), FW (54.1 ± 14.7 ; $F_4 = 5.93$; $p < 0.05$; $ES = 0.9$), and CD (51.2 ± 14.7 ; $F_4 = 5.93$; $p < 0.05$; $ES = 0.9$). MF also resulted in greater DEC_{LOW} than FW (46.8 ± 13.9 ; $F_4 = 3.96$; $p < 0.05$; $ES = 0.9$) and WMF (43.9 ± 13.9 ; $F_4 = 3.96$; $p < 0.05$; $ES = 0.8$). However, MF showed lower ACC_{DIS} than WMF (175.8 ± 25.5 m; $F_4 = 13.63$; $p < 0.05$; $ES = 2.3$) and FB (90.5 ± 26.1 m; $p < 0.05$; $ES = 1.1$). In addition, FW showed higher values of DEC_{MAX} compared to CD (0.6 ± 0.2 m/s²; $p < 0.05$; $ES = 0.8$).

Table 2. Acceleration profile of professional football players and differences between playing positions.

Variables	Position					p	ES
	CD (M ± SD)	FB (M ± SD)	MF (M ± SD)	WMF (M ± SD)	FW (M ± SD)		
ACC _{DIS} (m)	290.8 ± 76.0 ^d	351.3 ± 99.3 ^{cd}	260.7 ± 64.1 ^{bd}	436.5 ± 86.3 ^{abce}	333.6 ± 118.1 ^d	0.01	0.17–2.32
DEC _{DIS} (m)	229.9 ± 43.0 ^d	271.2 ± 66.3	228.7 ± 54.5 ^d	334.4 ± 74.5 ^{ace}	259.2 ± 55.9 ^d	0.01	0.03–1.72
ACC _{LOW} (total)	354.7 ± 54.5 ^c	378.1 ± 45.8	405.9 ± 61.9 ^{ade}	343.8 ± 48.0 ^c	351.7 ± 46.2 ^c	0.01	0.06–1.11
ACC _{HIGH} (total)	26.5 ± 6.1 ^d	30.4 ± 7.6	27.1 ± 5.5 ^d	34.9 ± 6.7 ^{ac}	29.9 ± 9.6	0.02	0.06–1.32
DEC _{LOW} (total)	355.9 ± 46.9	371.6 ± 45.9	387.4 ± 59.0 ^{de}	343.5 ± 15.8 ^c	340.6 ± 40.1 ^c	0.01	0.06–0.92
DEC _{HIGH} (total)	50.9 ± 8.6 ^d	54.1 ± 13.3	54.8 ± 12.4	64.5 ± 15.8 ^a	55.2 ± 12.1	0.02	0.04–1.07
DIFF _{ACDC} (total)	-24.4 ± 7.8	-23.7 ± 11.0	-27.7 ± 11.6	-29.5 ± 13.5	-25.3 ± 10.0	0.33	0.07–0.47
ACC _{AVG} (m/s ²)	0.6 ± 0.1	0.6 ± 0.1	0.6 ± 0.1	0.6 ± 0.1	0.5 ± 0.1	0.56	0.03–0.22
DEC _{AVG} (m/s ²)	-0.6 ± 0.1	-0.6 ± 0.1	-0.6 ± 0.1	-0.6 ± 0.1	-0.6 ± 0.1	0.76	0.01–0.26
ACC _{MAX} (m/s ²)	4.5 ± 0.4	4.5 ± 0.3	4.4 ± 0.6 ^d	4.70 ± 0.31 ^c	4.5 ± 0.6	0.01	0.06–0.67
DEC _{MAX} (m/s ²)	-5.7 ± 0.5 ^e	-6.1 ± 0.6	-5.8 ± 0.8 ^d	-6.20 ± 0.9 ^c	-6.3 ± 0.9 ^a	0.04	0.11–0.77

M: mean; SD: standard deviation; ES: effect size.

^aStatistical difference to CD (p < 0.05);

^bStatistical difference to FB (p < 0.05);

^cStatistical difference to MF (p < 0.05);

^dStatistical difference to WMF (p < 0.05);

^eStatistical difference to FW (p < 0.05).

<https://doi.org/10.1371/journal.pone.0236959.t002>

Table 3 gives the descriptive statistics for the sprint profile as well as the significant positional differences observed for all variables, apart from sprint time ($F_4 = 2.18$; $p = 0.07$). WMF reached greater V_{MAX} compared to CD (1.4 ± 0.5 km/h; $F_4 = 14.47$; $p < 0.05$; $ES = 0.9$), MF (3.5 ± 0.5 km/h; $F_4 = 14.47$; $p < 0.05$; $ES = 2.2$), and FW (2.1 ± 0.5 km/h; $F_4 = 14.47$; $p < 0.05$; $ES = 1.2$). WMF also resulted in greater SPA compared to CD (7.3 ± 1.1 ; $p < 0.05$; $ES = 1.7$), MF (11.3 ± 1.1 ; $p < 0.05$; $ES = 2.9$) and FW (7.2 ± 1.1 ; $p < 0.05$; $ES = 1.8$). These differences were also observed for SPD covered, which was greater in WMF than CD (189.3 ± 22.5 m; $p < 0.05$; $ES = 2.1$), MF (250.1 ± 22.1 m; $p < 0.05$; $ES = 3.06$) and FW (172.9 ± 22.5 m; $p < 0.05$; $ES = 2.05$). WMF also showed higher SPD_{AVG} than CD (4.7 ± 1.2 ; $p < 0.05$;

Table 3. Sprint profile of professional football players and differences between playing positions.

Variables	Position					p	ES
	CD (M ± SD)	FB (M ± SD)	MF (M ± SD)	WMF (M ± SD)	FW (M ± SD)		
SPA (total)	8.6 ± 3.7 ^{cd}	11.7 ± 3.9 ^c	4.6 ± 2.9 ^{abde}	15.9 ± 4.7 ^{ace}	8.7 ± 3.1 ^{cd}	0.01	0.04–2.87
SPD (m)	148.9 ± 73.1 ^d	228.3 ± 92.5 ^c	88.2 ± 53.9 ^{bde}	338.2 ± 103.9 ^{ace}	165.3 ± 58.1 ^{cd}	0.01	0.24–3.05
SPD _{AVG} (m)	16.9 ± 2.6 ^d	19.2 ± 3.2	20.1 ± 7.7	21.6 ± 3.8 ^a	19.6 ± 4.8	0.01	0.07–1.45
V_{MAX} (km/h)	30.6 ± 1.4 ^{acd}	30.7 ± 1.6 ^c	28.5 ± 1.7 ^{abde}	32.0 ± 1.6 ^{ace}	29.9 ± 2.0 ^{cd}	0.01	0.07–2.16
Sprint time (s)	2.5 ± 0.3	2.8 ± 0.4	2.8 ± 1.2	3.1 ± 0.5	2.9 ± 0.7	0.07	0.03–1.45

M: mean; SD: standard deviation; ES: effect size.

^aStatistical difference to CD (p < 0.05);

^bStatistical difference to FB (p < 0.05);

^cStatistical difference to MF (p < 0.05);

^dStatistical difference to WMF (p < 0.05);

^eStatistical difference to FW (p < 0.05).

<https://doi.org/10.1371/journal.pone.0236959.t003>

$ES = 1.46$). However, MF was the position with the lowest SPA (4.6 ± 2.9 ; $p < 0.05$; $ES = 1.2-2.03$) and V_{max} (28.5 ± 1.7 km/h; $F_4 = 14.47$; $p < 0.05$; $ES = 0.7-2.1$).

Fig 1 describes the V_0 required to perform low-intensity accelerations, high-intensity accelerations, low-intensity decelerations, and high-intensity decelerations. On the one hand, low-intensity accelerations required significantly greater V_0 than high-intensity accelerations in WMF (0.4 ± 0.2 km/h; $p < 0.05$; $ES = 0.12$) and FW (0.4 ± 0.2 km/h; $p < 0.05$; $ES = 0.13$) (Fig 1a), but no significant differences were found in CD, FB, and MF. On the other hand, high-intensity decelerations were performed at significantly higher V_0 than low-intensity decelerations in MF (2.7 ± 0.1 km/h; $p < 0.05$; $ES = 0.57$), FW (3.3 ± 0.1 km/h; $p < 0.05$; $ES = 0.64$), FB (3.9 ± 0.4 km/h; $p < 0.05$; $ES = 0.71$), WMF (4.3 ± 0.3 km/h; $p < 0.05$; $ES = 0.75$), and CD (4.1 ± 0.7 km/h; $p < 0.05$; $ES = 0.79$) (Fig 1b).

Discussion

The main purpose of this study was to describe the positional differences in the acceleration and sprint profiles of professional football players in match-play and to analyse the start speed (V_0) required based on the intensity of the acceleration and deceleration. This study showed positional differences for most variables of the acceleration and sprint profiles. Also, significant differences were observed in V_0 when comparing high-intensity accelerations and high-intensity decelerations to low-intensity accelerations and low-intensity decelerations.

This study is the first to provide detailed information on the acceleration and sprint profiles of professional football players. Previous investigations [13,20,27] have described positional differences for some of the high-intensity profile variables examined in the present study but conclusions from most studies were limited because only a few variables, which are the most common in the literature, were analyzed. These studies have examined, for instance, professional Norwegian [13,20] and British [27] football teams, and found differences between playing positions for variables of the acceleration profile [13,20,27]. For example, WMF covered significantly greater ACC_{DIS} (559 ± 232 m) and DEC_{DIS} (456 ± 107 m) than MF (ACC_{DIS} : 559 ± 232 m; DEC_{DIS} 360 ± 120 m) [20]. Similarly, our study showed that WMF covered significantly greater ACC_{DIS} (436.5 ± 86.3 m) and DEC_{DIS} (334.4 ± 74.5 m) than MF (ACC_{DIS} : 260.7 ± 64.1 m; DEC_{DIS} 228.7 ± 54.5 m). The same study also found that FB was another position with greater ACC_{DIS} covered (714 ± 298 m) than MF (559 ± 232 m) [20]. This suggests that playing in the lateral side of the pitch in addition to the offensive and defensive roles of FB let this position cover longer ACC_{DIS} compared to central playing positions such as MF [28].

With regard to the frequency of the accelerations, another study clearly showed that the totals for ACC_{HIGH} and DEC_{HIGH} were lower than for ACC_{LOW} and DEC_{LOW} [29], which represent the nature of football as a sport that involves intermittent repeated periods of high-intensity activity [4] and thus, high neuromuscular fatigue may be developed [12]. In addition, our results support previous research reporting that WMF always performed a higher amount of ACC_{HIGH} (35 ± 5 accelerations) and DEC_{HIGH} (62 ± 9 decelerations) than CD (ACC_{HIGH} : 27 ± 7 accelerations; DEC_{HIGH} : 45 ± 8 decelerations) [27]. Consequently, the variable $DIFF_{ACDC}$ also supports the same study, since the differences between playing positions were repeated once again and a higher number of DEC_{HIGH} than ACC_{HIGH} was observed in all positions [27]. However, MF performed significantly greater ACC_{LOW} and DEC_{LOW} than WMF (mean difference, ACC_{LOW} : 62.1 ± 14.7 accelerations; DEC_{LOW} : 43.9 ± 13.9 decelerations) and FW (mean difference, ACC_{LOW} : 54.1 ± 14.7 accelerations; DEC_{LOW} : 46.8 ± 13.9 decelerations) in our study, which may be explained by the fact that density increases (reduced area per player) as the ball is closer to the central zones of the pitch in match play [30].

Although WMF also had the highest $DIFF_{ACDC}$ (-27 ± 4), this study reported that FW had the




Fig 1. Differences on start speed of the action (V_0) based on the intensity of the accelerations/decelerations; a) Differences between low-intensity accelerations and high-intensity accelerations; b) Differences between low-intensity decelerations and high-intensity decelerations.

<https://doi.org/10.1371/journal.pone.0236959.g001>

lowest $\text{DIFF}_{\text{ACDC}}$ (-17 ± 4) [27]. Players, therefore decelerate at high-intensity more than they accelerate at high-intensity, so special focus on mechanical load indicators is recommended for strength and conditioning coaches [20]. When it comes to the magnitude of the accelerations and decelerations, the results cannot be compared to previous studies [31,32], since this is the first study to carry out this analysis in match-play. However, when analysing this variable in training contexts, differences between playing positions remained low for ACC_{MAX} ($0.17 \pm 0.03 \text{ m/s}^2$) and DEC_{MAX} ($0.26 \pm 0.03 \text{ m/s}^2$) [32]. Consequently, this study supports the assertion that the acceleration profile is position-dependent and that different training strategies may be adopted to improve match performance and decrease risk of injury [12].

Positional differences were found in the sprint profile, and similar conclusions were reached by previous studies that analysed some of the variables of this profile [5,15,20,33]. For example, the total of SPA performed by FW (14 ± 6 actions), FB (12 ± 5 actions) and WMF (8 ± 4 actions) was also greater than CD (5 ± 3 actions) in previous research [15]. When it comes to MF, this position showed the lowest amount of SPA not only in our study (5 ± 3 actions) but also in previous research (4 ± 4 actions) [15]. These results may be explained by the fact that this position is limited to reach high-speed actions given its tactical role (e.g., keeping ball possession, passing) and playing area [31,34,35]. Also, other studies [13,33] found that WMF covered the highest SPD ($294 \pm 76 \text{ m}$ and $185 \pm 23 \text{ m}$, respectively) whereas CD covered the lowest ($123 \pm 48 \text{ m}$ and $77 \pm 17 \text{ m}$, respectively). Thus, SPD_{AVG} was also position-dependent in this study. However, no significant differences ($p > 0.05$) were observed in a previous study in European professional players [36]. In this sense, it is interesting to note that data were collected from players who belonged to ten different teams, which might explain differences in the conclusions reached [36]. However, the same study found significant differences between playing positions when comparing V_{MAX} , which is in line with our study [36]. Specifically, the V_{MAX} reached by WMF ($32.0 \pm 1.6 \text{ km/h}$) and FW ($29.9 \pm 2.0 \text{ km/h}$) was significantly higher compared to CD ($30.6 \pm 1.4 \text{ km/h}$) in our study as well as in the above-mentioned study (WMF: $32.9 \pm 2.0 \text{ km/h}$; FW: $33.1 \pm 1.9 \text{ km/h}$; CD: $31.7 \pm 1.8 \text{ km/h}$) [36]. Similarly, MF showed the lowest V_{MAX} ($31.0 \pm 1.7 \text{ km/h}$) [36]. In this regard, the V_{MAX} from WMF and FW may be explained by their greater SPD_{AVG} (WMF: $21.6 \pm 3.8 \text{ m}$; FW: $19.6 \pm 4.8 \text{ m}$), which allow them to maximize their acceleration capacity [36]. On the contrary, MF are limited to the increase in density of players in central zones of the pitch as explained above [30]. The only variable from the sprint profile that did not show significant differences in our study was sprint time, which was in line with previous research, where the largest differences observed between positions ranged from 0.1 to 0.2 seconds [37]. Overall, these findings imply that offensive positions such as WMF and FW are subjected to the highest sprint demands, whereas CD, a defensive position characterised by short runs [13,15,33], is the least demanding position in respect of the sprint profile in professional football.

Finally, another novel finding of this study was that football players required significantly higher V_0 when performing high-intensity decelerations than low-intensity decelerations across all playing positions (mean difference, CD: $4.1 \pm 0.7 \text{ km/h}$; WMF: $4.3 \pm 0.3 \text{ km/h}$; FB: $3.9 \pm 0.4 \text{ km/h}$; FW: $3.3 \pm 0.1 \text{ km/h}$; MF: $2.7 \pm 0.1 \text{ km/h}$). However, V_0 was significantly greater for low-intensity accelerations than high-intensity accelerations, and only in specific positions (WMF and FW: $0.4 \pm 0.2 \text{ km/h}$). These results indicate that the ability to accelerate or decelerate is highly dependent on the V_0 of the player, particularly, when decelerating at high-intensity ($p < 0.01$; $ES \sim 0.69$). In this regard, a recent study, which tested maximal accelerations during sprint actions at different V_0 , observed a linear decrease in the maximal acceleration when V_0 increased [38]. This may be explained by a typical speed-time curve from any sprint test in which the largest increase in speed is at the start of the action and then, the curve is flattened with increasing running speed [38,39]. In consequence, the acceleration capacity

decreases at higher speeds [38]. Training drills could take this into consideration, since it is not only the frequency and intensity of the acceleration that is important, but also the V_0 required to perform the acceleration [23]. However, it is important to understand that players occupying positions such as CD may accelerate or decelerate from lower V_0 than players in other positions [23]. Since football players perform actions which start at different V_0 [15], previous studies assumed that different V_0 may result in different neuromuscular preload, body inclines and, therefore, different muscle group activation [38,40]. Furthermore, the players experience a massive metabolic load every time the acceleration is increased, even when speed is low [41]. Thus, when designing effective match-based drills, these profiles and positional differences may be considered for a more accurate approach to players' performance [23]. In addition, an individualised approach to performance is needed, since these results lead to the hypothesis that, for example, a faster CD may have more ability to perform high-intensity accelerations and high-intensity decelerations than a slower player in the same position.

There are some limitations that need to be considered when interpreting the findings of this research. The data were collected using GPS technology. Only one professional football team was assessed over 13 official matches from a Spanish competition. In addition, not all the players could participate in all the matches. Another limitation of the study is that the effect of different variables such as team formation, competitive standard, or style of play on the acceleration and sprint profiles was not analysed. Future studies may consider these variables since these variables may affect match running performance [42–44]. For example, a previous study showed that teams which were characterised by possession-play covered greater distance in high-intensity running actions [44]. Also, although speed and acceleration bands/thresholds were selected according to previous studies [29,36], it is worth noting that this issue has not yet been standardised in the literature [11].

Conclusions

The findings of this longitudinal study provide meaningful information related to the sprint and acceleration profiles of Spanish professional football match-play. This study found that positional differences exist across playing positions in both profiles, which should be considered by strength and conditioning coaches when designing effective match-based drills in training sessions. Also, special focus should be given to WMF since this position was the most demanding of the acceleration and sprint profiles. In addition, only a few studies have analysed the acceleration-velocity relationship, which is deemed important when designing training drills. Despite strength and conditioning coaches still focusing on training sprint actions starting at zero speed, most of these actions are performed at 5–6 km/h in match-play. In addition, different V_0 were observed between high-intensity and low-intensity accelerations as well as high-intensity and low-intensity decelerations. This may potentially affect the neuromuscular load of the players and coaching strategies are necessary in order to maximize players' performance. Finally, these data could also serve as a comparison source for future researchers or sports scientists and coaches from professional football teams. For example, if players decelerate at high-intensity more than they accelerate at high-intensity, coaches who choose to put more emphasis on acceleration capacity than deceleration capacity in training sessions should understand that match-play may require the opposite.

Supporting information

S1 File.
(PDF)

Acknowledgments

The results of the current study do not constitute endorsement of the product by the authors or the journal.

Author Contributions

Conceptualization: José M. Oliva-Lozano, Víctor Fortes, Peter Krstrup, José M. Muoyor.

Data curation: José M. Oliva-Lozano.

Formal analysis: José M. Oliva-Lozano.

Investigation: José M. Oliva-Lozano, Víctor Fortes, Peter Krstrup, José M. Muoyor.

Methodology: José M. Oliva-Lozano.

Supervision: Víctor Fortes, José M. Muoyor.

Validation: José M. Oliva-Lozano, José M. Muoyor.

Visualization: Peter Krstrup, José M. Muoyor.

Writing – original draft: José M. Oliva-Lozano.

Writing – review & editing: José M. Oliva-Lozano, Víctor Fortes, Peter Krstrup, José M. Muoyor.

References

1. Gabbett TJ, Nassis GP, Oetter E, Pretorius J, Johnston N, Medina D, et al. The athlete monitoring cycle: A practical guide to interpreting and applying training monitoring data. *Br J Sports Med.* 2017; 51: 1451–1452. <https://doi.org/10.1136/bjsports-2016-097298> PMID: 28646100
2. Gómez-Carmona CD, Pino-Ortega J, Sánchez-Ureña B, Ibáñez SJ, Rojas-Valverde D. Accelerometry-based external load Indicators in sport: too many options, same practical outcome? *Int J Environ Res Public Health.* 2019; 16: 1–13. <https://doi.org/10.3390/ijerph16245101> PMID: 31847248
3. Rampinini E, Coutts A, Castagna C, Sassi R, Impellizzeri F. Variation in top level soccer match performance. *Int J Sports Med.* 2007; 28: 1018–1024. <https://doi.org/10.1055/s-2007-965158> PMID: 17497575
4. Oliva-Lozano JM, Gómez-Carmona CD, Pino-Ortega J, Moreno-Pérez V, Rodríguez-Pérez MA. Match and training high intensity activity-demands profile during a competitive mesocycle in youth elite soccer players. *J Hum Kinet.* 2020;Epub Ahead: 1–11.
5. Palucci-Vieira LH, Aquino R, Lago-Peñas C, Munhoz-Martins GH, Puggina EF, Barbieri FA, et al. Running performance in Brazilian professional football players during a congested match schedule. *J Strength Cond Res.* 2018; 32: 313–325. <https://doi.org/10.1519/JSC.0000000000002342> PMID: 29369952
6. Carling C, Gall FL, Reilly TP. Effects of physical efforts on injury in elite soccer. *Int J Sports Med.* 2010; 31: 180–185. <https://doi.org/10.1055/s-0029-1241212> PMID: 20024885
7. Buchheit M, Samozino P, Glynn JA, Michael BS, Al Haddad H, Mendez-Villanueva A, et al. Mechanical determinants of acceleration and maximal sprinting speed in highly trained young soccer players. *J Sports Sci.* 2014; 32: 1906–1913. <https://doi.org/10.1080/02640414.2014.965191> PMID: 25356503
8. Haugen TA, Tønnessen E, Seiler S. Anaerobic performance testing of professional soccer players 1995–2010. *Int J Sports Physiol Perform.* 2013; 8: 148–156. <https://doi.org/10.1123/ijssp.8.2.148> PMID: 22868347
9. Oliva-Lozano JM, Rojas-Valverde D, Gómez-Carmona CD, Fortes V, Pino-Ortega J. Impact of contextual variables on the representative external load profile of Spanish professional soccer match-play: a full season study. *Eur J Sport Sci.* 2020; 1–22. <https://doi.org/10.1080/17461391.2020.1751305> PMID: 32233969
10. Faude O, Koch T, Meyer T. Straight sprinting is the most frequent action in goal situations in professional football. *J Sports Sci.* 2012; 30: 625–631. <https://doi.org/10.1080/02640414.2012.665940> PMID: 22394328

11. Sweeting AJ, Cormack SJ, Morgan S, Aughey RJ. When is a sprint a sprint? A review of the analysis of team-sport athlete activity profile. *Front Physiol.* 2017; 20: 1–12. <https://doi.org/10.3389/fphys.2017.00432> PMID: 28676767
12. Harper DJ, Carling C, Kiely J. High-intensity acceleration and deceleration demands in elite team sports competitive match play: a systematic review and meta-analysis of observational studies. *Sport Med.* 2019; 49: 1923–1947. <https://doi.org/10.1007/s40279-019-01170-1> PMID: 31506901
13. Ingebrigtsen J, Dalen T, Hjelde GH, Drust B, Wisløff U. Acceleration and sprint profiles of a professional elite football team in match play. *Eur J Sport Sci.* 2015; 15: 101–110. <https://doi.org/10.1080/17461391.2014.933879> PMID: 25005777
14. Miñano-Espin J, Casáis L, Lago-Peñas C, Gómez-Ruano MÁ. High speed running and sprinting profiles of elite soccer players. *J Hum Kinet.* 2017; 58: 169–176. <https://doi.org/10.1515/hukin-2017-0086> PMID: 28828087
15. Varley M, Aughey R. Acceleration profiles in elite Australian soccer. *Int J Sports Med.* 15.08.2012. 2012; 34: 34–39. <https://doi.org/10.1055/s-0032-1316315> PMID: 22895869
16. Little T, Williams AG. Specificity of acceleration, maximum speed, and agility in professional soccer players. *J Strength Cond Res.* 2005; 19: 76–78. PMID: 15705049
17. Vanrenterghem J, Nedergaard NJ, Robinson MA, Drust B. Training load monitoring in team sports: a novel framework separating physiological and biomechanical load-adaptation pathways. *Sport Med.* 2017; 47: 2135–2142. <https://doi.org/10.1007/s40279-017-0714-2> PMID: 28283992
18. De Hoyo M, Cohen DD, Sañudo B, Carrasco L, Álvarez-Mesa A, del Ojo JJ, et al. Influence of football match time–motion parameters on recovery time course of muscle damage and jump ability. *J Sports Sci.* 2016; 34: 1363–1370. <https://doi.org/10.1080/02640414.2016.1150603> PMID: 26930226
19. Hader K, Mendez-Villanueva A, Palazzi D, Ahmadi S, Buchheit M. Metabolic power requirement of change of direction speed in young soccer players: not all is what it seems. McCrory JL, editor. *PLoS One.* 2016; 11: 1–21. <https://doi.org/10.1371/journal.pone.0149839> PMID: 26930649
20. Dalen T, Jørgen I, Gertjan E, Geir Havard H, Ulrik W. Player load, acceleration, and deceleration during forty-five competitive matches of elite soccer. *J Strength Cond Res.* 2016; 30: 351–359. <https://doi.org/10.1519/JSC.0000000000001063> PMID: 26057190
21. Gaudino P, Iaia FM, Strudwick AJ, Hawkins RD, Alberti G, Atkinson G, et al. Factors influencing perception of effort (session rating of perceived exertion) during elite soccer training. *Int J Sports Physiol Perform.* 2015; 10: 860–864. <https://doi.org/10.1123/ijspp.2014-0518> PMID: 25671338
22. De Hoyo M, Sañudo B, Suárez-Arrones L, Carrasco L, Joel T, Domínguez-Cobo S, et al. Analysis of the acceleration profile according to initial speed and positional role in elite professional male soccer players. *J Sports Med Phys Fitness.* 2018; 58: 22. <https://doi.org/10.23736/S0022-4707.17.08003-3> PMID: 29199786
23. De Hoyo M, Sañudo B, Suárez-Arrones L, Carrasco L, Joel T, Domínguez-Cobo S, et al. Analysis of the acceleration profile according to initial speed and positional role in elite professional male soccer players. *J Sports Med Phys Fitness.* 2018; 58: 22. <https://doi.org/10.23736/S0022-4707.17.08003-3> PMID: 29199786
24. White A, Hills SP, Cooke CB, Batten T, Kilduff LP, Cook CJ, et al. Match-play and performance test responses of soccer goalkeepers: a review of current literature. *Sport Med.* 2018; 48: 2497–2516. <https://doi.org/10.1007/s40279-018-0977-2> PMID: 30144021
25. Bastida-Castillo A, Gómez-Carmona CD, De la Cruz E, Pino-Ortega J, Bastida Castillo A, Gómez Carmona CD, et al. Accuracy, intra- and inter-unit reliability, and comparison between GPS and UWB-based position-tracking systems used for time–motion analyses in soccer. *Eur J Sport Sci.* 2018; 18: 1–8.
26. Cohen J. *Statistical power analysis for the behavioral sciences.* 2nd ed. Hillsdale N.J.: L. Erlbaum Associates; 1988. <https://www.worldcat.org/title/statistical-power-analysis-for-the-behavioral-sciences/oclc/17877467>
27. Tierney PJ, Young A, Clarke ND, Duncan MJ. Match play demands of 11 versus 11 professional football using Global Positioning System tracking: variations across common playing formations. *Hum Mov Sci.* 2016; 49: 1–8. <https://doi.org/10.1016/j.humov.2016.05.007> PMID: 27269201
28. Konefał M, Chmura P, Andrzejewski M, Puksza D, Chmura J. Analysis of match performance of full-backs from selected European soccer leagues. *Cent Eur J Sport Sci Med.* 2015; 11: 45–53. <https://doi.org/10.18276/cej.2015.3-05>
29. Russell M, Sparkes W, Northeast J, Cook CJ, Love TD, Bracken RM, et al. Changes in acceleration and deceleration capacity throughout professional soccer match-play. *J Strength Cond Res.* 2016; 30: 2839–2844. <https://doi.org/10.1519/JSC.0000000000000805> PMID: 25474342
30. Fradua L, Zubillaga A, Caro O, Iván Fernández-García A, Ruiz-Ruiz C, Tenga A. Designing small-sided games for training tactical aspects in soccer: extrapolating pitch sizes from full-size professional

- matches. *J Sports Sci.* 2013; 31: 573–581. <https://doi.org/10.1080/02640414.2012.746722> PMID: 23186250
31. Gaudino P, Alberti G, Iaia FM. Estimated metabolic and mechanical demands during different small-sided games in elite soccer players. *Hum Mov Sci.* 2014; 36: 123–133. <https://doi.org/10.1016/j.humov.2014.05.006> PMID: 24968370
 32. Rago V, Silva JR, Mohr M, Barreira D, Krstrup P, Rebelo AN. Variability of activity profile during medium-sided games in professional soccer. *J Sports Med Phys Fitness.* 2019; 59. <https://doi.org/10.23736/S0022-4707.18.08376-7> PMID: 29697215
 33. Vigh-Larsen JF, Dalgas U, Andersen TB. Position-specific acceleration and deceleration profiles in elite youth and senior soccer players. *J Strength Cond Res.* 2018; 32: 1114–1122. <https://doi.org/10.1519/JSC.0000000000001918> PMID: 28699924
 34. Carling C. Analysis of physical activity profiles when running with the ball in a professional soccer team. *J Sports Sci.* 2010; 28: 319–326. <https://doi.org/10.1080/02640410903473851> PMID: 20077273
 35. Bradley PS, Carling C, Gomez Diaz A, Hood P, Barnes C, Ade J, et al. Match performance and physical capacity of players in the top three competitive standards of English professional soccer. *Hum Mov Sci.* 2013; 32: 808–821. <https://doi.org/10.1016/j.humov.2013.06.002> PMID: 23978417
 36. Andrzejewski M, Chmura J, Pluta B, Konarski JM. Sprinting activities and distance covered by top level Europa League soccer players. *Int J Sports Sci Coach.* 2015; 10: 39–50. <https://doi.org/10.1260/1747-9541.10.1.39>
 37. Suárez L, Torreno N, Requena B, De Villarreal ES, Casamichana D, Barbero-Álvarez JC, et al. Match-play activity profile in professional soccer players during official games and the relationship between external and internal load. *J Sport Med Phys Fit.* 2014; 55: 1417–1422.
 38. Sonderegger K, Tschopp M, Taube W. The challenge of evaluating the intensity of short actions in soccer: a new methodological approach using percentage acceleration. Lucia A, editor. *PLoS One.* 2016; 11: 1–10. <https://doi.org/10.1371/journal.pone.0166534> PMID: 27846308
 39. Morin J-B, Sève P. Sprint running performance: comparison between treadmill and field conditions. *Eur J Appl Physiol.* 2011; 111: 1695–1703. <https://doi.org/10.1007/s00421-010-1804-0> PMID: 21210279
 40. Young W, Benton D, Pryor J. Resistance training for short sprints and maximum-speed sprints. *Strength Cond J.* 2001; 23: 1–7. <https://doi.org/10.1519/00126548-200104000-00001>
 41. Osgnach C, Poser S, Bernardini R, Rinaldo R, Di Prampero PE. Energy cost and metabolic power in elite soccer. *Med Sci Sport Exerc.* 2010; 42: 170–178. <https://doi.org/10.1249/MSS.0b013e3181ae5cfd> PMID: 20010116
 42. Aquino R, Vieira LHP, Carling C, Martins GHM, Alves IS, Puggina EF. Effects of competitive standard, team formation and playing position on match running performance of Brazilian professional soccer players. *Int J Perform Anal Sport.* 2017; 17: 695–705. <https://doi.org/10.1080/24748668.2017.1384976>
 43. Bradley PS, Carling C, Archer D, Roberts J, Dodds A, Di Mascio M, et al. The effect of playing formation on high-intensity running and technical profiles in English FA Premier League soccer matches. *J Sports Sci.* 2011; 29: 821–830. <https://doi.org/10.1080/02640414.2011.561868> PMID: 21512949
 44. Yi Q, Gómez MA, Wang L, Huang G, Zhang H, Liu H. Technical and physical match performance of teams in the 2018 FIFA World Cup: effects of two different playing styles. *J Sports Sci.* 2019; 37: 2569–2577. <https://doi.org/10.1080/02640414.2019.1648120> PMID: 31354060