


Differential Recognition of *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa* by Human Monocyte-Derived Macrophages and Dendritic Cells

Manuela Gómez-Gaviria¹, Iván Martínez-Duncker², Laura C García-Carnero¹, Héctor M Mora-Montes¹ 

¹Departamento de Biología, División de Ciencias Naturales y Exactas, Campus Guanajuato, Universidad de Guanajuato, Guanajuato, Guanajuato México; ²Laboratorio de Glicobiología Humana y Diagnóstico Molecular; Centro de Investigación en Dinámica Celular, Instituto de Investigación en Ciencias Básicas y Aplicadas, Universidad Autónoma del Estado de Morelos, Cuernavaca, Morelos, México

Correspondence: Héctor M Mora-Montes, Departamento de Biología, División de Ciencias Naturales y Exactas, Campus Guanajuato, Universidad de Guanajuato, Noria Alta s/n, Col. Noria Alta, C.P. Guanajuato, Guanajuato, 36050, México, Tel +52 473-7320006 Ext 8193, Fax +52 473-7320006 Ext 8153, Email hmora@ugto.mx

Background: Sporotrichosis is a mycosis frequently caused by *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*. The cell wall is a species-specific fungal structure with a direct role in activating the host's immune response. The current knowledge about anti-*Sporothrix* immunity comes from studies using *S. schenckii* or *S. brasiliensis* and murine cells. Macrophages and dendritic cells detect and eliminate pathogens, and although the function of these cells links innate with adaptive immunity, little is known about their interaction with *Sporothrix* spp.

Methods: *S. schenckii*, *S. brasiliensis*, and *S. globosa* conidia or yeast-like cells were co-incubated with human monocyte-derived macrophages or dendritic cells, and the phagocytosis and cytokine stimulation were assessed. These interactions were also performed in the presence of specific blocking agents of immune receptors or fungal cells with altered walls to analyze the contribution of these molecules to the immune cell-fungus interaction.

Results: Both types of immune cells phagocytosed *S. globosa* conidia and yeast-like cells to a greater extent, followed by *S. brasiliensis* and *S. schenckii*. Furthermore, when the wall internal components were exposed, the phagocytosis level increased for *S. schenckii* and *S. brasiliensis*, in contrast to *S. globosa*. Thus, the cell wall components have different functions during the interaction with macrophages and dendritic cells. *S. globosa* stimulated an increased proinflammatory response when compared to the other species. In macrophages, this was a dectin-1-, mannose receptor-, and TLR2-dependent response, but dectin-1- and TLR2-dependent stimulation in dendritic cells. For *S. schenckii* and *S. brasiliensis*, cytokine production was dependent on the activation of TLR4, CR3, and DC-SIGN.

Conclusion: The results of this study indicate that these species are recognized by immune cells differently and that this may depend on both the structure and cell wall organization of the different morphologies.

Keywords: cell wall, cytokine production, fungal morphologies, immune response, phagocytosis

Introduction

Sporotrichosis is a subcutaneous mycosis that affects humans and other mammals. Its etiological agents are members of the *Sporothrix* pathogenic clade, and currently, the infection is mainly caused by *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*.^{1–3} This disease has global distribution; however, most of the animal and human cases are concentrated in countries located between the Tropics of Cancer and Capricorn, with exceptions reported in South Africa, China, Japan, and the anecdotic cases documented in European countries and the United States of America.^{3–10} The lymphocutaneous infection is the most frequent clinical form of sporotrichosis, which rarely becomes a life-threatening disseminated disease.^{11–13} However, when there is temporal or permanent immunosuppression in the

host, the subcutaneous infection can develop into a systemic disease with high mortality rates in both humans and animals.^{5,10,14,15} Historically, *S. schenckii* was the first fungal species associated with sporotrichosis, and as a consequence, it is the *Sporothrix* species most thoroughly studied at both basic and clinical levels.^{3,16} *S. globosa*, like *S. schenckii*, is distributed worldwide,⁴ however, within the pathogenic clade is considered the least virulent species due to its limited growth at high temperatures.^{17–19} *S. brasiliensis* has been responsible for infectious outbreaks in Brazil and is considered the most virulent species.^{18–21} These three fungal species can grow as conidium-producer mycelia in the environment and undergo dimorphism to yeast-like cells when colonizing and damaging the host tissues. Thus, conidia and hyphae are the first fungal morphologies to establish contact with the host. However, the zoonotic and epizootic *S. brasiliensis* outbreaks have changed this paradigm, and currently, not only conidia but also yeast-like cells can infect host tissues.^{3,5,21,22}

The study of antifungal immunity is nowadays an active research area because it is thought that the knowledge accumulated in this subject could be translated to immunotherapeutic approaches to help in the treatment of mycoses caused by etiological agents that commonly develop antifungal drug resistance; and also because it is clear that most of the severe presentations of mycosis and those associated with high mortality rates occur in individuals without an intact immunity, stressing the role of the immune response in controlling most of the aggressive manifestations of fungi when interacting with the host.^{23,24} Since the cell wall is the outermost *Sporothrix* structure and thus is the primary point of contact with components of the host immunity it has been thoroughly studied in recent years.^{19,22,25–32} Thus far, it is known that the *S. schenckii* and *S. brasiliensis* cell wall contains two layers. Chitin, β -1,6-, β -1,4-, and β -1,3-glucans are the main components of the inner layer;^{3,26,28} while *N*-linked and *O*-linked glycoproteins and peptidorhamnomannan are the main components of the outermost layer.^{5,28,33,34}

Thus far, most of our knowledge about anti-*Sporothrix* immunity comes from the study of the interaction of *S. schenckii* with murine immune cells, and to a lesser extent the study of *S. brasiliensis* using the same experimental model.³⁵ Cellular components of innate immunity, such as granulocytes, monocytes, macrophages, and dendritic cells are capable of detecting and kill fungal pathogens, recruit additional immune effectors, and to link innate and adaptive immunities.³⁶ Currently, there are relatively few reports dealing with the *Sporothrix*-macrophage interaction.^{37–41} Using murine macrophages, it has been demonstrated that *S. schenckii* chitin-rich heteroglycan modulates cytokine production and promoted *S. schenckii* phagocytosis.⁴² Regarding immune receptors involved in this interaction, it was found that dectin-1 expression was increased in peritoneal murine macrophages after infection with *S. schenckii*, and there were nitric oxide and cytokines productions driven by this immune receptor,⁴³ while TLR2 is required for fungal phagocytosis and the stimulation of IL-1 β , IL-12, and TNF- α .⁴⁴ TLR4 has also been involved in the sensing of *S. schenckii* by murine macrophages, and the absence of this receptor led to an increment in TGF- β production⁴⁵ and a reduction of pro and anti-inflammatory cytokines.⁴⁶ In the same line, TLR4 is required for phagocytosis of *S. brasiliensis* and stimulation of proinflammatory cytokines.⁴⁷ Interestingly, TLR2 seems that does not contribute to cytokine stimulation when *S. brasiliensis* interacts with murine macrophages, but it is required for phagocytosis.⁴⁸

The interaction of *Sporothrix* cells with human macrophages has also been studied. Thus far, it is known that *S. schenckii* *N*-linked and *O*-linked glycans promote uptake by human monocyte-derived macrophages in a dectin-1-, mannose receptor- (MR), complement receptor 3- (CR3), and TLR4-dependent mechanism;^{29,31} while peptidorhamnomannan seems to be the CR3 ligand for stimulation of IL-1 β by both *S. schenckii* and *S. brasiliensis* cells.⁴⁹ In addition, *S. schenckii* rhamnose-containing oligosaccharides are required for fungal uptake by human monocyte-derived macrophages in a TLR4-dependent pathway.²² The *S. globosa*-macrophage interaction has been lesser studied than that described for the other two species. Using murine macrophages, it has been reported that *S. globosa* melanin inhibits antigen presentation in macrophages, which could be an immunoevasion mechanism.⁵⁰ When interacting with human macrophages, *S. globosa* melanin inhibited fungal phagocytosis, protected from radical oxygen and nitrogen species, negatively affected the production of proinflammatory cytokines, and modulated TLR2 and TLR4 expression.⁵¹

The information related to the *Sporothrix*-dendritic cell interactions is even more scarce. Immature human monocyte-derived dendritic cells were stimulated with *S. schenckii* yeast-like cells, and was found that fungal isolates from cutaneous origins were more potent to stimulate a strong Th1 response than strains from the visceral origin.⁵² Murine bone marrow-dendritic cells stimulated with *S. schenckii* cell wall proteins showed higher expression of CD80, CD86,

and CD40, but not TLR-4, and Th1-prone cytokine profile with IL-6, IL-17A, and TNF α production.⁵³ Similarly, *S. schenckii* yeast-like cells stimulated the production of a Th1-prone cytokine profile and phagocytosis when interacting with these immune cells.^{54,55}

Despite all this information, the interaction between macrophages or dendritic cells with different *Sporothrix* species is poorly studied. Therefore, here we assessed the relevance of fungal morphology, cell wall components, and immune receptors on the interaction of different *Sporothrix* species when interacting with either human monocyte-derived macrophages or monocyte-derived dendritic cells.

Materials and Methods

Strains and Culturing Conditions

The following strains were used in this study: *S. schenckii* ATCC MYA-4821, *S. brasiliensis* ATCC MYA 4823, and *S. globosa* FMR 9624. All are clinical isolates characterized by molecular techniques elsewhere.^{32,56} These strains of *S. schenckii* and *S. brasiliensis* are considered model organisms of these species, and their genomes and transcriptome have been previously sequenced.^{57–59} Cells were grown in YPD medium, pH 4.5 (1% [w/v] yeast extract, 2% [w/v] gelatin peptone, and 3% [w/v] dextrose) at 28°C, and 2% [w/v] agar was added when the solid medium was required. For conidia propagation, cells were grown on YPD, pH 4.5 plates at 28°C and after seven days of incubation, 10 mL of deionized water was added to the surface of the plates; this was gently dispersed with a plastic spreader, and the conidia suspension was collected.⁶⁰ Yeast-like cells were obtained by inoculating conidia in YPD, pH 7.8 broth, and incubating at 37°C and 120 rpm for four days.²⁶ For both morphotypes, cells were thoroughly washed with deionized water and kept at –20°C until used. To inactivate cells by heating (heat-killed cells, HK), these were incubated at 60°C for 2 h, and the loss of viability was confirmed by incubation in YPD plates, pH 4.5, at 28°C for 5 days.²⁶ The *N*-linked and *O*-linked glycans were removed from cell walls by incubating with endoglycosidase H and β -elimination, respectively, as reported.^{22,29}

Ethics Statement

This study was conducted following the Declaration of Helsinki. In agreement, cells were obtained from healthy adult volunteers after information about the study was disclosed and written informed consent was signed. The use of human cells was approved by the Ethics Committee from Universidad de Guanajuato (reference CIBIUG-P36-2019).

Differentiation of Primary Human Cells to Macrophages and Dendritic Cells

Human PBMCs were isolated from venous blood using Histopaque-1077 (Sigma-Aldrich, Saint Louis, MO, USA), as described.⁶¹ For differentiation to macrophages, 1 mL aliquots containing 2×10^7 PBMCs in RPMI supplemented with 1% (v/v) penicillin-streptomycin solution (PS, Sigma-Aldrich) were seeded in flat bottom 12-well plates and incubated 1.5 h at 37°C, 5% (v/v) CO₂. Non-adherent cells were discarded, and the remaining adherent cells were washed with PBS at 37°C, and 1 mL of X–VIVO serum-free medium (Lonza, Basel, Switzerland) supplemented with 1% (v/v) PS and 10 ng mL⁻¹ recombinant human granulocyte-macrophage colony-stimulating factor (Sigma-Aldrich) were included per well.⁶² For differentiation to immature dendritic cells (DCs), adherent cells were added 1 mL X–VIVO serum-free medium supplemented with 1% (v/v) PS, 500 U mL⁻¹ recombinant human IL-4 (Sigma-Aldrich), and 800 U mL⁻¹ recombinant human granulocyte-macrophage colony-stimulating factor (Sigma-Aldrich) for 6–7 days.^{63,64} In both cases, plates were incubated for 6–7 days at 37 °C, 5% (v/v) CO₂, replacing the medium every 2 days. Differentiation to DC cells was confirmed by flow cytometry, detecting DC-SIGN with FITC-conjugated mAb anti-human DC-SIGN (Thermo-Fisher Scientific, Waltham, MA, USA), as described elsewhere.⁶³

Phagocytosis Assays

For fungal labeling, cells were incubated with 1 mg mL⁻¹ Acridine Orange (Sigma-Aldrich) for 30 min at room temperature, the excess dye was washed with PBS, and cell concentration was adjusted at 3×10^7 cells mL⁻¹.⁶⁵ Six-well plates were used to perform the interactions at an immune cell: fungus ratio of 1:6 in 800 μ L DMEM medium (Sigma-Aldrich). The plates were incubated for 2 h at 37°C and 5% (v/v) CO₂,^{29,66} immune cells were detached from

plates with chilled PBS and incubated with 1.25 mg mL^{-1} Trypan Blue.⁶⁷ The phagocytic event was analyzed by cytometry using a FACSCanto II system (Becton Dickinson, Franklin Lakes, NJ, USA). Fifty thousand events were collected per sample through the FL1 and FL2 channels, which were previously calibrated with non-labeled immune cells.^{22,29,65,67} The following compounds were used in preincubation experiments: $200 \text{ } \mu\text{g mL}^{-1}$ laminarin (Sigma-Aldrich),⁶⁸ $10 \text{ } \mu\text{g mL}^{-1}$ anti-MR antibody (Thermo-Fisher Scientific, MA5-44033), $10 \text{ } \mu\text{g mL}^{-1}$ anti-TLR4 antibody (Santa Cruz Biotechnology, Dallas, TX, sc-293072), $10 \text{ } \mu\text{g mL}^{-1}$ anti-TLR2 antibody (Thermo-Fisher Scientific, 16-9922-82), $10 \text{ } \mu\text{g mL}^{-1}$ anti-CD209 antibody (DC-SIGN; Thermo Fisher Scientific, 14-2099-82), $10 \text{ } \mu\text{g mL}^{-1}$ anti-CD11b antibody (CR3, Thermo Fisher Scientific, MA5-16528), Isotype-matched, irrelevant IgG1 antibody ($10 \text{ } \mu\text{g mL}^{-1}$, Santa Cruz Biotechnology, Cat. No. sc-52003, used as a control in experiments where MR and TLR4 were blocked), $10 \text{ } \mu\text{g mL}^{-1}$ IgG2 α antibody (Thermo-Fisher Scientific, 14-4724-85, to control TLR2 and CD209 blocking experiments), and $10 \text{ } \mu\text{g mL}^{-1}$ IgG2 antibody (R&D, Minneapolis, MN, USA, Cat. No. MAB9794, to control CD11b blocking assays).^{26,49,69} Immune cells were preincubated for 1 h at 37°C and 5% (v/v) CO_2 with any of these compounds before interacting with fungal cells. Even though the above-listed reagents were negative for bacterial LPS, the preincubations were performed in the presence of $5 \text{ } \mu\text{g mL}^{-1}$ polymyxin B (Sigma-Aldrich).^{70,71} Cells detected with the FL1 channel were considered in the early stage of phagocytosis, cells detected by both FL1 and FL2 channels were regarded as in the intermediate stage of phagocytosis, and those detected only with the FL2 channel were considered in the late stage of phagocytosis.^{22,29,65}

Cytokine Stimulation

The cell-cell interactions were performed in 96-well microplates. Each well contained $100 \text{ } \mu\text{L}$ of immune cells in RPMI 1640 Dutch modification (Sigma-Aldrich) at $1 \times 10^6 \text{ cells mL}^{-1}$ and $100 \text{ } \mu\text{L}$ of fungal cells at $1 \times 10^6 \text{ mL}^{-1}$. Plates were incubated for 24 h at 37°C with 5% (v/v) CO_2 . Similar to the experiments described in the previous section, in some cases, immune cells were pre-incubated for 60 min at 37°C with 5% (v/v) CO_2 with one of the antibodies already described or laminarin and the interactions were performed in the presence of $5 \text{ } \mu\text{g mL}^{-1}$ polymyxin B (Sigma-Aldrich).^{70,71} After incubation, plates were centrifuged for 10 min at $1800 \times g$ at 4°C , supernatants were collected and used for cytokine quantification by ELISA. The Standard ABTS ELISA Development kits (Peprotech, Cranbury, NJ, USA) were used for quantification of tumor necrosis factor- α (TNF α), interleukin 6 (IL-6), and interleukin 10 (IL-10); while DuoSet ELISA from R&D was used for interleukin 1 β (IL-1 β) and Interleukin 12 p70 (IL-12 p70) quantification. Immune cells suspended in RPMI 1640 Dutch modification were included in each plate as controls. In all cases, these control wells gave threshold levels that were subtracted from all the cytokine determinations.

Statistical Analysis

The Mann–Whitney U and Kruskal–Wallis tests from the GraphPad Prism 6 software were used for data analysis, establishing significance at $P < 0.05$. Since the Kruskal–Wallis analysis indicated that data had no normal distribution ($P < 0.05$), non-parametric tests were applied to the data. All experiments were carried out in duplicate with samples from eight healthy donors. Results are shown as the mean and standard deviation.

Results

Differential Phagocytosis of *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa* by Human Monocyte-Derived Macrophages

To compare the ability of primary human phagocytic cells to uptake fungi from the three most frequently isolated *Sporothrix* species from medical and veterinary samples, named *S. schenckii*, *S. brasiliensis*, and *S. globosa*, conidia or yeast-like cells were grown and co-incubated with the human cells. Since the phagocytic process was analyzed by flow cytometry, phagocytosis of germling could not be analyzed using this strategy, as hyphae can clothe the instrument pipes.⁷² When conidia were used in the interactions, this tended to be more phagocytosed than yeast-like cells in *S. schenckii* and *S. brasiliensis*, but for *S. globosa*, conidia, and yeast-like cells were similarly phagocytosed (Figure 1). When the phagocytosis of the three species was compared among themselves, *S. schenckii* cells were the less

phagocytosed, followed by *S. brasiliensis*, and the most phagocytosed species was *S. globosa*. This observation applied to the two analyzed morphologies (Figure 1). In addition, these differences were observed only in the intermediate and late stages of phagocytosis (Figure 1). There were no significant differences in the early stage of the immune process when *S. brasiliensis* cells were compared with *S. globosa* cells (Figure 1). Control wells containing only the human cells showed threshold uptake levels (Figure 1).

Next, we assessed the contribution of immune receptors on the uptake of either conidia or yeast-like cells from the three species under study. We focused on MR, dectin-1, CD11b (Complement receptor-3; CR3) TLR2, and TLR4, as these receptors have been previously involved in the recognition of *S. schenckii* and *S. brasiliensis* cells.^{26,43–46,49} For this, the immune cells were preincubated with monoclonal antibodies against particular receptors or with laminarin, an antagonist of dectin-1,^{62,69} before being challenged with the fungal cells. Since most of the fungal cells interacting with monocyte-derived macrophages are in the late stage of the phagocytic process, under our experimental conditions, we focused only on the analysis of this subset. For the case of *S. schenckii* conidia and yeast-like cells, the five receptors under analysis were required for fungal uptake, as the blocking of any of them significantly reduced the fungal uptake (Figure 2A). However, the contribution of receptors varied in conidia and yeast-like cells, being MR more relevant for yeast-like cell uptake than in conidia and dectin-1 and TLR2 for conidia uptake (Figure 2A). For the case of *S. brasiliensis* cells, the five receptors were required for proper yeast-like cells uptake, but MR was dispensable for conidia phagocytosis (Figure 2A). Like the observation in *S. schenckii*, the receptor relevance varied depending on the fungal morphology, being dectin-1, and TLR2 more relevant for conidia uptake and CR3 and TLR4 for yeast-like cells phagocytosis (Figure 2A). Interestingly, *S. globosa* conidia and yeast-like cells were phagocytosed via dectin-1 and TLR2, whilst MR, CR3, and TLR4 were dispensable for fungal uptake (Figure 2A). Isotype-matched irrelevant antibodies were used to preincubate human cells and did not affect the fungal phagocytosis of any of the morphologies or species under study (data not shown).

The contribution of cell wall components to fungal uptake by human monocyte-derived macrophages was also analyzed. For this, cells were inactivated by heat, as this treatment artifactually exposes internal cell wall components, such as β -1,3-glucan, and chitin at the cell wall surface.^{26,62,69} Moreover, cell wall *N*-linked or *O*-linked glycans were removed by treatment with endoglycosidase H or β -elimination, respectively.^{22,29,31} The *S. schenckii* HK conidia and yeast-like cells were more phagocytosed than live cells, but for the case of *S. brasiliensis*, this treatment has only a positive effect on the phagocytosis of yeast-like cells and had no effect on the *S. globosa* uptake (Figure 2B). Removal of *N*-linked glycans, *O*-linked glycans, or both kinds of glycans negatively affected the phagocytosis of the *S. schenckii* and *S. brasiliensis* conidia and yeast-like cells, but once again, phagocytosis of *S. globosa* was not affected by trimming of these cell wall components (Figure 2B). Collectively, these results




Figure 1 Phagocytosis of *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa* by human monocyte-derived macrophages. Interactions of human monocyte-derived macrophages and *Sporothrix* cells at a ratio 1:6 were incubated for 2 h at 37°C, and 5% (v/v) CO₂. Macrophages were analyzed by flow cytometry, collecting 50,000 events, which were defined as a human cell interacting with at least one fluorescent fungal cell. Control, human cells were incubated only with PBS. **P* < 0.05 when compared to conidia from the same species. †*P* < 0.05 when compared with *S. schenckii* cells of the same morphology. Cell numbers obtained with *S. brasiliensis* conidia and yeast-like cells at the intermediate and late stages were significantly different from those obtained with *S. globosa* cells (*P* < 0.05). Data are shown as means ± SD from eight donors analyzed by duplicate.




Figure 2 Contribution of pattern recognition receptors and cell wall components on the phagocytosis of *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa* by human monocyte-derived macrophages. In (A), Human monocyte-derived macrophages were preincubated with 200 $\mu\text{g mL}^{-1}$ laminarin or 10 $\mu\text{g mL}^{-1}$ of any of the following antibodies: anti-mannose receptor (MR), anti-CR3, anti-TLR2, or anti-TLR4. Then, cells were coincubated with Acridine Orange-labeled conidia or yeast-like cells at a macrophage-fungus ratio of 1:6, for 2 h at 37°C and 5% (v/v) CO_2 . Human cells were analyzed by flow cytometry, collecting 50,000 events, which were defined as a human cell interacting with at least one fungal cell. All the interactions were performed in the presence of 5 $\mu\text{g mL}^{-1}$ polymyxin B. No treatment, cells preincubated with PBS. Results correspond to cells in the late stage of phagocytosis. For all cases, 100% corresponds to human cells preincubated with PBS, and the absolute values were similar to those shown in Figure 1 or (B) of this figure. CR3, complement receptor 3. * $P < 0.05$ when compared to the no-treatment condition of the same strain. † $P < 0.05$ when compared to conidia from the same species. In (B), Similar experiments as described in (A), but human cells were not preincubated with any blocking agent. Before coincubation with human monocyte-derived macrophages, conidia, and yeast-like cells were inactivated by heat (HK), treated with endoglycosidase H (Endo-H), β -eliminated to remove O-linked glycans, or both treated with endoglycosidase H and β -elimination. No treatment refers to live cells without any treatment. * $P < 0.05$ when compared to the no-treatment condition of the same strain. For both panels, data were shown as means \pm SD from eight donors analyzed by duplicate.

suggest that conidia and yeast-like cells from the three species under analysis are differentially recognized by human monocyte-derived macrophages, and the pattern recognition receptors and cell wall components have species- and morphology-specific relevance.

Cytokine Production by Human Monocyte-Derived Macrophages Stimulated with *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*

In addition to the study of the phagocytic process, we also assessed the ability of human monocyte-derived macrophages to produce cytokines after stimulation by the species under study. The TNF α and IL-6 levels stimulated by *S. schenckii* yeast-like cells were significantly lower when compared to the levels stimulated by *S. schenckii* conidia; whilst IL-1 β and

IL-10 levels were similar when cells were stimulated by any of these to *S. schenckii* morphologies (Figure 3). For *S. brasiliensis*, the four cytokines analyzed were stimulated at lower levels by yeast-like cells, when compared with conidia (Figure 3). However, when cells were stimulated with *S. globosa* conidia or yeast-like cells these produced similar levels of TNF α , IL-6, IL-1 β , and IL-10 (Figure 3). When comparing the cytokine levels stimulated among the three different species, *S. schenckii* conidia stimulated the lowest TNF α , IL-6, IL-1 β , and IL-10 levels, and conidia from *S. brasiliensis* and *S. globosa* stimulated similar levels of the four cytokines (Figure 3). Similarly, *S. schenckii* yeast-like cells stimulated the lowest cytokines levels, while *S. brasiliensis* intermediate and *S. globosa* yeast-like cells had the highest cytokine levels (Figure 3).

The TNF α and IL-10 production were taken as readouts of the host-fungus interaction and we also explored the contribution of different PRRs to the stimulation of both cytokines. For the case of TNF α , its stimulation by *S. schenckii* conidia was negatively influenced when MR, TLR4, or CR3 were blocked, suggesting a role of these receptors in the induction of this proinflammatory cytokine (Figure 4A). However, for *S. schenckii* yeast-like cells only dectin-1 and TLR2 were required for proper cytokine production, being dispensable the rest of the receptors analyzed (Figure 4A). *S. brasiliensis* conidia and yeast-like cells stimulated TNF α via TLR4 and CR3, but in the latter, there was also a contribution of MR (Figure 4A); while in *S. globosa*, both conidia and yeast-like cells have a similar dependence in dectin-1, MR, and TLR2 for cytokine induction (Figure 4A). When a similar approach was used to analyze the relevance of receptors for IL-10 production, we found that *S. schenckii* conidia and yeast-like cells stimulated this cytokine via dectin-1 and MR, but in case of conidia, TLR2 was also involved (Figure 4B). Dectin-1, TLR2, TLR4, and CR3 were required for proper IL-10 stimulation by *S. brasiliensis* conidia, an observation that contrasted with the sole dependency on MR to IL-10 stimulation by *S. brasiliensis* yeast-like cells (Figure 4B). Finally, both morphologies of *S. globosa* did not show significant differences between them, both stimulating IL-10 via dectin-1 and TLR2 (Figure 4B). Isotype-matched irrelevant antibodies were used to preincubate human cells and did not affect the fungal phagocytosis of any of the morphologies or species under study (data not shown).

We also assessed the contribution of cell wall components to TNF α and IL-10 stimulation by monocyte-derived macrophages. Both, *S. schenckii* HK conidia and yeast-like cells stimulated higher TNF α levels than live cells, but this cytokine production was significantly reduced when *N*-linked or *O*-linked glycans or the combination of both were removed from the wall (Figure 5A). Similarly, both, *N*-linked and *O*-linked glycans were required for proper TNF α stimulation by *S. brasiliensis* conidia or yeast-like cells, but only in the case of HK yeast-like cells was observed a stronger cytokine production when compared with live cells (Figure 5A). None of the cell wall treatments affected the ability of *S. globosa* conidia or yeast-like cells to stimulate TNF α (Figure 5A). Similar results were observed for IL-10




Figure 3 Cytokine production by human monocyte-derived macrophages stimulated with *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*. Human and fungal cells were cocultured for 24 h, and the supernatants were collected and used for cytokine quantification by ELISA. * $P < 0.05$, when compared with the cytokine level stimulated by conidia from the same species. † $P < 0.05$, when compared with the same morphology of the other two fungal species analyzed. Results are shown as mean \pm standard deviation from data generated with samples from eight donors analyzed by duplicate.




Figure 4 Contribution of pattern recognition receptors to cytokine production by human monocyte-derived macrophages stimulated with *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*. Human cells were preincubated with 200 $\mu\text{g mL}^{-1}$ laminarin or 10 $\mu\text{g mL}^{-1}$ of any of the following antibodies: anti-mannose receptor (MR), anti-complement receptor 3 (CR3), anti-TLR2, or anti-TLR4. Then, cells were cocultured with either conidia or yeast-like cells of the fungal species for 24 h at 37°C and 5% (v/v) CO_2 . Supernatants were collected and used to measure the levels of TNF α (A) or IL-10 (B) by ELISA. No treatment, cells preincubated with PBS. In all cases, 100% corresponds to the system with no treatment, and the absolute values were similar to those shown in Figure 3. * $P < 0.05$ when compared to the no-treatment condition of the same strain. † $P < 0.05$ when compared to conidia from the same species. Results are shown as mean \pm standard deviation from data generated with samples from eight donors analyzed by duplicate.

stimulation by *S. globosa*, but in the case of *S. schenckii* both conidia and yeast-like cells positively stimulated IL-10 production when HK cells were used for the interaction or when *N*-linked and *O*-linked glycans were removed from the wall (Figure 5B). For the case of *S. brasiliensis* cells, conidia did not affect the ability to stimulate IL-10 after modification of the cell wall, but yeast-like cells increased the ability to stimulate this cytokine when cells were heat-inactivated or *N*-linked and *O*-linked glycans were trimmed off from walls (Figure 5B). Collectively, these data indicate that monocyte-derived macrophages are capable to discriminate between cell morphologies and the different *Sporothrix* species, using different PRRs and having a differential response against these cells.




Figure 5 Contribution of cell wall components on cytokine stimulation by *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa* interacting with human monocyte-derived macrophages. Macrophages were incubated with conidia or yeast-like cells at a macrophage-fungus ratio of 1:6, for 24 h at 37°C and 5% (v/v) CO₂. Plates were centrifuged and supernatants were used to quantify TNFα (A) or IL-10 (B) by ELISA. Before coinubation with human monocyte-derived macrophages, conidia, and yeast-like cells were inactivated by heat (HK) treated with endoglycosidase H (Endo-H), β-eliminated to remove O-linked glycans, or both treated with endoglycosidase H and β-elimination. No treatment refers to live cells without any treatment. *P < 0.05 when compared to the no-treatment condition of the same strain. Results are shown as mean ± standard deviation from data generated with samples from eight donors analyzed by duplicate.

Human Monocyte-Derived Dendritic Cells Differentially Phagocytose *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*

Next, we performed similar experiments but using monocyte-derived DCs. Using the same strategy to assess fungal uptake by macrophages, we found that *S. schenckii* and *S. brasiliensis* conidia were hardly phagocytosed by monocyte-derived DCs, but this was not the case for *S. globosa*, which were readily phagocytosed, in a similar rate than yeast-like cells (Figure 6). Both types of fungal cells generated the highest phagocytosis levels in the early, intermediate, and late stages (Figure 6). *S. schenckii* yeast-like cells were modestly phagocytosed by these immune cells but the levels were significantly different from those observed on *S. brasiliensis* yeast-like cells, which were the fungal cells with the lowest ability to be phagocytosed (Figure 6). Control reaction with immune cells incubated with buffer gave threshold detections (Figure 6).

When the relevance of immune receptors for DCs uptake was analyzed, we found that *S. schenckii* conidia uptake was dependent on MR, CR3, DC-SIGN, and TLR4 (Figure 7). Similarly, yeast-like cells depended on the same receptor




Figure 6 Phagocytosis of *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa* by human monocyte-derived dendritic cells. Fungal cells were labeled with Acridine Orange and incubated with human monocyte-derived dendritic cells at an immune cell-fungus ratio of 1:6, for 2 h at 37°C and 5% (v/v) CO₂. Human cells were analyzed by flow cytometry, collecting 50,000 events, which were defined as a human cell interacting with at least one fluorescent fungal cell. Control, human cells were incubated only with PBS. * $P < 0.05$ when compared to conidia from the same species. † $P < 0.05$ when compared with the same morphology in the other two species. Results are shown as mean \pm standard deviation from data generated with samples from eight donors analyzed by duplicate.

subsets, along with dectin-1 (Figure 7). The *S. brasiliensis* conidia and yeast-like cells depended on the engagement of MR, CR3, DC-SIGN, and TLR4 with its ligands, and both *S. globosa* morphologies were phagocytosed via dectin-1 and TLR2 (Figure 7). Isotype-matched irrelevant antibodies were used to preincubate human cells and did not affect the fungal phagocytosis of any of the morphologies or species under study (data not shown). These results indicate the three fungal species are sensed by different subsets of immune receptors found on the monocyte-derived DCs surface.

Then, the relevance of cell wall components during the *Sporothrix*-monocyte-derived DCs interaction was analyzed. Both morphologies of *S. schenckii* were positively affected when HK cells were used in the interaction with immune cells, significantly increasing the number of phagocytosed cells (Figure 7B). However, the removal of *N*-linked, *O*-linked, or both types of glycans from the cell wall negatively affected cell uptake (Figure 7B). Similar results were observed for *S. brasiliensis* yeast-like cells, but no significant changes were observed for *S. brasiliensis* conidia subjected to these treatments (Figure 7B). *S. globosa* conidia and yeast-like cells did not show any effect on the phagocytosis when HK or deglycosylated cells were used to interact with monocyte-derived DCs. Collectively, these results suggest that the cell wall components play different roles during interaction with these immune cells.

Next, to assess the ability of these fungal species to stimulate cytokine production by human monocyte-derived DCs, the proinflammatory cytokines TNF α , IL-6, IL-1 β , and IL-12 p70 were quantified by ELISA, after the immune cell-fungus interaction. *S. schenckii* yeast-like cells stimulated lower TNF α , IL-6, and IL-1 β levels than conidia, whilst both cell morphologies stimulated similar IL-12 p70 levels (Figure 8). For the case of *S. brasiliensis* yeast-like cells, only TNF α and IL-1 β levels were lower when compared to those stimulated by conidia; while both *S. globosa* conidia and yeast-like cells stimulated similar levels of the four cytokines analyzed (Figure 8). When comparing the stimulation profiles among species, *S. globosa* stimulated the highest levels of the four cytokines followed by *S. brasiliensis*, and the lowest cytokine levels were associated with *S. schenckii* cells (Figure 8). For the case of conidia, only TNF α and IL-1 β stimulated by *S. brasiliensis* conidia were higher than those stimulated by *S. schenckii* conidia (Figure 8).

Next, we assessed the contribution of PRRs to TNF α stimulation. CR3 and DC-SIGN were required to induce proper TNF α production by both *S. schenckii* conidia and yeast-like cells, although the latter also required dectin-1 engagement with its ligand and conidia of TLR4 signaling (Figure 9A). Similarly, *S. brasiliensis* conidia and yeast-like cells depended on interaction with CR3 and DC-SIGN for TNF α production but in these cases, TLR4 was also required (Figure 9A). Only *S. brasiliensis* yeast-like cells required dectin-1 signaling for proper TNF α production (Figure 9A). Finally, both *S. globosa* conidia and yeast-like cells stimulated TNF α via dectin-1 and TLR2 (Figure 9A). Isotype-matched irrelevant antibodies were used to preincubate human cells and did not affect the fungal phagocytosis of any of the morphologies or




Figure 7 Contribution of pattern recognition receptors and cell wall components on the phagocytosis of *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa* by human monocyte-derived Dendritic cells. In (A), Immune cells were preincubated with $200 \mu\text{g mL}^{-1}$ laminarin or $10 \mu\text{g mL}^{-1}$ of any of the following antibodies: anti-mannose receptor (MR), anti-CR3, anti-DCSIGN, anti-TLR2, or anti-TLR4. Then, cells were coincubated with Acridine Orange-labeled conidia or yeast-like cells at an immune cell-fungus ratio of 1:6, for 2 h at 37°C and 5% (v/v) CO_2 . Human cells were analyzed by flow cytometry, collecting 50,000 events, which were defined as a human cell interacting with at least one fluorescent fungal cell. All the interactions were performed in the presence of $5 \mu\text{g mL}^{-1}$ polymyxin B. No treatment refers to cells preincubated only with PBS. Results correspond to cells in the late stage of phagocytosis. For all cases, 100% corresponds to the system with no treatment, and the absolute values were similar to those shown in Figure 6 or (B) of this figure. CR3, complement receptor 3. * $P < 0.05$ when compared to the no-treatment condition of the same strain. $^{\dagger}P < 0.05$ when compared to conidia from the same species. In (B), Similar experiments as described in (A), but human cells were not preincubated with any blocking agent. Before coincubation with human monocyte-derived dendritic, conidia and yeast-like cells were inactivated by heat (HK) treated with endoglycosidase H (Endo-H), β -eliminated to remove O-linked glycans, or both treated with endoglycosidase H and β -elimination. No treatment refers to live cells without any treatment. * $P < 0.05$ when compared to the no-treatment condition of the same strain. Results are shown as mean \pm standard deviation from data generated with samples from eight donors analyzed by duplicate.

species under study (data not shown). Regarding the cell wall components involved in this stimulation, *S. schenckii* conidia and yeast-like HK cells stimulated higher cytokine levels than live cells (Figure 9B). When glycans were removed from the wall there was a negative effect on $\text{TNF}\alpha$ production by conidia, but in the case of yeast-like cells, this negative effect was more evident when both N-linked and O-linked glycans were removed from the wall (Figure 9B). For the case of *S. brasiliensis*, the removal of any of the two kinds of glycans or both together had a negative impact on the cytokine levels stimulated by conidia but not for yeast-like cells (Figure 9B). Heat-inactivation had a positive effect only on *S. brasiliensis* yeast-like cells, being live and HK conidia similar in the ability to stimulate $\text{TNF}\alpha$ production (Figure 9B). For the case of *S. globosa*, none of the treatments affected the ability of conidia or yeast-like cells to stimulate $\text{TNF}\alpha$ (Figure 9B).




Figure 8 Cytokine production by human monocyte-derived dendritic cells stimulated with *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*. Human and fungal cells were cocultured for 24 h., and the supernatants were collected and used for cytokines quantification by ELISA. * $P < 0.05$, when compared with the cytokine level stimulated by conidia from the same species. † $P < 0.05$, when compared with the same morphology of the other two fungal species analyzed. ‡ $P < 0.05$, when compared with the same morphology of *S. schenckii*. Results are shown as mean \pm standard deviation from data generated with samples from eight donors analyzed by duplicate.

Discussion

Members of the *Sporothrix* pathogenic clade have different cell wall components that are considered potential antigens.^{57,73,74} Previous studies in *S. schenckii* and *S. brasiliensis* determined that the cell wall composition and organization can change among species and morphologies,^{19,26,28} thus, it is feasible to conceive that immune cells interact differently with each species and morphotype. Thus far, few reports have assessed the interaction between *Sporothrix* and macrophages or DCs.^{42–45} Most of the knowledge about this interaction comes from the study of *S. schenckii* and *S. brasiliensis*,³⁵ and little is known about the interactions of these immune cells with *S. globosa*. This work aimed to compare the importance of fungal morphology, cell wall components, and immune receptors during the interaction of *S. schenckii*, *S. brasiliensis*, and *S. globosa* with macrophages and DCs.

Phagocytosis assay with human monocyte-derived macrophages and DCs showed that *S. globosa* is the most phagocytosed species in both conidial and yeast-like cell morphologies, followed by *S. brasiliensis* and *S. schenckii*. The increased phagocytosis of *S. globosa* could be related to the organization of the cell wall components of this species. Phagocytosis is known to initiate with the recognition of β -1,3-glucan and chitin, through dectin-1, TLR2, and TLR4 receptors.⁷⁵ So, it is likely that there could be an association between the proportion of these polysaccharides and phagocytosis. *S. globosa* has a higher amount of β -1,3-glucans exposed on the surface, which could favor its phagocytosis in both cell morphologies.¹⁹ In *S. schenckii* and *S. brasiliensis*, cells are less phagocytosed, and previous reports have shown that β -1,3-glucans are not exposed on the cell surface, and these are in a lower proportion than those of *S. globosa*, which hinders their recognition by these immune cells.^{19,26} When the contribution of cell wall components, such as β -glucans, *N*-glycans, and *O*-glycans of the three species was evaluated, it was found that HK *S. schenckii* and *S. brasiliensis* showed an increase in macrophage- and DCs-mediated phagocytosis of yeast-like cells, and a decrease in this when *N*-linked and *O*-linked glycans of these two species are removed. This suggests that *N*-glycans and *O*-glycans from *S. schenckii* and *S. brasiliensis* play an important role during the detection of immune cells to initiate phagocytosis. For the case of *S. globosa*, phagocytosis was not affected by any of the fungal treatments, and this could be related to the fact that both cell wall *N*-linked and *O*-linked glycans of this species are found in a lower proportion than in *S. schenckii* and *S. brasiliensis*.^{22,29} Furthermore, the absence of changes in phagocytosis of *S. globosa* could be because all components of the cell wall of this species contribute to fungal uptake by macrophages and DCs, and when altered by the treatments applied to cells the other wall components can compensate for the disturbance.

When the contribution of the different immune receptors in the uptake of conidia and yeasts was analyzed, it was found that all receptors are important in the detection of *S. schenckii* and *S. brasiliensis* by human monocyte-derived macrophages, mainly in yeast-like cells. However, some variations were observed in the case of *S. brasiliensis* conidia,




Figure 9 Contribution of pattern recognition receptors and cell wall components to TNF α production by human monocyte-derived dendritic cells stimulated with *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*. In **A**, Dendritic cells were preincubated with 200 $\mu\text{g mL}^{-1}$ laminarin or 10 $\mu\text{g mL}^{-1}$ of any of the following antibodies: anti-mannose receptor (MR), anti-complement receptor 3 (CR3), anti-DC-SIGN, anti-TLR2, or anti-TLR4. Then, cells were coincubated with either conidia or yeast-like cells of the fungal species for 24 h at 37°C and 5% (v/v) CO $_2$, the supernatants were collected and used to measure the levels of TNF α . No treatment refers to cells preincubated with PBS. In all cases, 100% corresponds to the system with no treatment, and the absolute values were similar to those shown in Figure 8. * $P < 0.05$ when compared to the no-treatment condition of the same strain. † $P < 0.05$ when compared to conidia from the same species. In **B**, dendritic cells were incubated with conidia or yeast-like cells at an immune cell-fungus ratio of 1:6, for 24 h at 37°C and 5% (v/v) CO $_2$ and supernatants were collected and used to quantify TNF α by ELISA. Before coincubation with human monocyte-derived dendritic cells, conidia, and yeast-like cells were inactivated by heat (HK) treated with endoglycosidase H (Endo-H), β -eliminated to remove O-linked glycans, or both treated with endoglycosidase H and β -elimination. No treatment refers to live cells without any treatment. * $P < 0.05$ when compared to the no-treatment condition of the same strain. † $P < 0.05$ when compared to conidia from the same species. Results are shown as mean \pm standard deviation from data generated with samples from eight donors analyzed by duplicate.

where the essential receptors are dectin-1 and TLR2, a result that could be related to the cell wall organization of this morphology (Figure 10).^{26,76}

Based on the results obtained, it can be observed that dectin-1 is a relevant receptor for the recognition of these three species by monocyte-derived macrophages. Most of the β -1,3-glucan required to trigger dectin-1-dependent signaling pathways may be already accessible on the cell surface, as reported with PBMCs in *S. schenckii* and *S. brasiliensis* species.²⁶ Previous work has reported that the interaction of *S. schenckii* and *S. brasiliensis* with monocyte-derived macrophages depends on complement protein C3, which facilitates yeast phagocytosis.⁴⁹ Here, we confirm this observation but interestingly, it does not apply to *S. globosa*, since the uptake of conidia and yeast-like cells was dependent on dectin-1 and TLR2 (Figure 10).




Figure 10 Phagocytosis and receptors involved in the immune recognition of different species of the *Sporothrix* pathogenic clade. Macrophages and dendritic cells can phagocytose yeast-like cells and conidia of *S. schenckii*, *S. brasiliensis*, and *S. globosa*. *S. globosa* is the species most phagocytosed by both types of cells, followed by *S. brasiliensis* and, finally, *S. schenckii*. The receptors involved in such recognition vary between species and morphologies. *S. globosa* is recognized by dectin-1 and TLR2 receptors, while *S. brasiliensis* and *S. schenckii* are recognized by dectin-1, TLR2, TLR4, CR3, DC-SIGN, and MR.

When the relevance of these receptors by human monocyte-derived DCs was assessed, it was found that the three species are detected by different sets of receptors. DCs play a key role in regulating the balance between pro- and anti-inflammatory responses, in addition, these cells can translate fungal-associated molecular signatures and coordinate an inflammatory response that protects against infection.^{54,77} For these immune cells, the receptors showed variation in the recognition of conidia and yeast-like cells of the three species, like that found in macrophages. However, it is striking that the dectin-1 receptor was dispensable for the recognition of conidia of *S. schenckii* and *S. brasiliensis* since it is a receptor that is involved in important signaling pathways and has a significant contribution in the recognition of fungal cells.⁷⁸ According to the results obtained in this work, it is possible to hypothesize that the role of the different receptors analyzed may be influenced by the type of immune cell interacting with *Sporothrix*, highlighting the relevance of studying the immune cell-*Sporothrix* interaction with different subtypes of cellular immune effectors.

The interaction of either *S. globosa* conidia or yeast-like cells with human monocyte-derived macrophages and DCs induced a higher production of IL-6 and TNF α when compared to *S. schenckii* and *S. brasiliensis*. It is tempting to speculate that this proinflammatory trend could contribute to the lower ability to colonize tissues and establish the infection of this species. In agreement with this hypothesis, the contrary scenario was observed with the most virulent species of the clade, *S. schenckii* and *S. brasiliensis*, which stimulated lower pro-inflammatory levels (Figure 11).

The analysis of different PRRs and cell wall components during the stimulation of TNF α and IL-10 by human monocyte-derived macrophages and DCs showed that the PRRs analyzed here have a morphology- and species-dependent role in this immunological event. When human-derived macrophages interacted with *S. globosa* conidia or yeast-like cells, dectin-1, MR, and TLR2 were required for TNF α stimulation, while dectin-1 and TLR-2 for IL-10 production. This contrast with the involvement of CR3 in the stimulation of TNF α by *S. schenckii* and *S. brasiliensis* cells, as reported previously and as shown here.⁴⁹ Interestingly, IL-10 stimulation by *S. brasiliensis* yeast-like cells was solely dependent on MR, stressing again that glycan structures are different in these organisms, as reported previously for *S. schenckii* and *S. brasiliensis*.^{27,49} Finally, similar to other fungal species,^{63,64} DC-SIGN was required for stimulation of TNF α by *S. schenckii* and *S. brasiliensis*, but not for *S. globosa*, suggesting once again that β -1,3-glucan sensing by dectin-1 and TLR2 are the main PRRs-PAMPs interactions that these fungal cells establish with DCs.




Figure 11 Cytokine profiles of macrophages and dendritic cells stimulated with *Sporothrix schenckii*, *Sporothrix brasiliensis*, or *Sporothrix globosa*. When human cells interact with *S. globosa* conidia or yeast-like cells, the production of the different cytokines analyzed increases. In the case of *S. schenckii*, there is a low stimulation of these cytokines compared to *S. globosa* and *S. brasiliensis*. These results were obtained from the inter-species comparison.

Conclusion

In conclusion, this study reported that the cell wall components and the different morphologies of *S. schenckii*, *S. brasiliensis*, and *S. globosa* play an important role during recognition by human monocyte-derived macrophages and DCs. The interaction of these immune cells with the different *Sporothrix* species generates specific profiles of cytokines and uptake via different immune receptors. This study contributes to expanding the knowledge of the immune response against different species of the *Sporothrix* pathogenic clade.

Acknowledgments

This work was supported by Consejo Nacional de Ciencia y Tecnología (ref. FC 2015-02-834, Ciencia de Frontera 2019-6380), and Red Temática Glicociencia en Salud (CONACYT-México). The funding sources that supported this work did not have any involvement in the design, acquisition, and analysis of data and writing of the manuscript.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis, and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

The authors declare no conflicts of interest in this work.

References

- de Beer ZW, Duong TA, Wingfield MJ. The divorce of *Sporothrix* and *Ophiostoma*: solution to a problematic relationship. *Stud Mycol.* 2016;83:165–191. doi:10.1016/j.simyco.2016.07.001

2. de Beer ZW, Procter M, Wingfield MJ, Marincowitz S, Duong TA. Generic boundaries in the *Ophiostomatales* reconsidered and revised. *Stud Mycol.* 2022;101:57–120. doi:10.3114/sim.2022.101.02
3. Lopes-Bezerra LM, Mora-Montes HM, Zhang Y, et al. Sporotrichosis between 1898 and 2017: the evolution of knowledge on a changeable disease and on emerging etiological agents. *Med Mycol.* 2018;56(suppl_1):126–143. doi:10.1093/mmy/myx103
4. Chakrabarti A, Bonifaz A, Gutierrez-Galhardo MC, Mochizuki T, Li S. Global epidemiology of sporotrichosis. *Med Mycol.* 2015;53(1):3–14. doi:10.1093/mmy/myu062
5. Lopez-Romero E, Reyes-Montes MR, Perez-Torres A, et al. *Sporothrix schenckii* complex and sporotrichosis, an emerging health problem. *Future Microbiol.* 2011;6(1):85–102. doi:10.2217/fmb.10.157
6. Lv S, Hu X, Liu Z, Lin Y, Wu H, Li F. Clinical epidemiology of sporotrichosis in Jilin province, China (1990–2019): a series of 4969 cases. *Infect Drug Resist.* 2022;15:1753–1765. doi:10.2147/idr.S354380
7. Takenaka M, Sato S, Nishimoto K. Survey of 155 sporotrichosis cases examined in Nagasaki Prefecture from 1951 to 2007. *Nihon Ishinkin Gakkai Zasshi.* 2009;50(2):101–108. doi:10.3314/jjmm.50.101
8. Bongomin F, Adetona Fayemiwo S. Epidemiology of fungal diseases in Africa: a review of diagnostic drivers. *Curr Med Mycol.* 2021;7(1):63–70. doi:10.18502/cmm.7.1.6246
9. Barile F, Mastrodonato M, Loconsole F, Rantuccio F. Cutaneous sporotrichosis in the period 1978–1992 in the province of Bari, Apulia, Southern Italy. *Mycoses.* 1993;36(5–6):181–185. doi:10.1111/j.1439-0507.1993.tb00747.x
10. Gold JA, Derado G, Mody RK, Benedict K. Sporotrichosis-associated hospitalizations, United States, 2000–2013. *Emerg Infect Dis.* 2016;22(10):1817–1820. doi:10.3201/eid2210.160671
11. Lopes-Bezerra LM, Mora-Montes HM, Bonifaz A. *Sporothrix* and Sporotrichosis. In: Mora-Montes HM, Lopes-Bezerra LM, editors. *Current Progress in Medical Mycology*. Cham: Springer; 2017:309–331.
12. Mora-Montes HM, Dantas Ada S, Trujillo-Esquivel E, de Souza Baptista AR, Lopes-Bezerra LM. Current progress in the biology of members of the *Sporothrix schenckii* complex following the genomic era. *FEMS Yeast Res.* 2015;15:6.
13. Orofino-Costa R, Macedo PM, Rodrigues AM, Bernardes-Engemann AR. Sporotrichosis: an update on epidemiology, etiopathogenesis, laboratory and clinical therapeutics. *An Bras Dermatol.* 2017;92(5):606–620. doi:10.1590/abd1806-4841.2017279
14. Queiroz-Telles F, Buccheri R, Benard G. Sporotrichosis in immunocompromised hosts. *J Fungi.* 2019;5(1):8. doi:10.3390/jof5010008
15. de Miranda LHM, Meli M, Conceição-Silva F, et al. Co-infection with feline retrovirus is related to changes in immunological parameters of cats with sporotrichosis. *PLoS One.* 2018;13(11):e0207644. doi:10.1371/journal.pone.0207644
16. Mora-Montes HM. Special Issue “*Sporothrix* and Sporotrichosis 2.0”. *J Fungi.* 2022;8(8):821. doi:10.3390/jof8080821
17. Nava-Pérez N, Neri-García LG, Romero-González OE, Terrones-Cruz JA, García-Carnero LC, Mora-Montes HM. Biological and clinical attributes of *Sporothrix globosa*, a causative agent of sporotrichosis. *Infect Drug Resist.* 2022;15:2067–2090. doi:10.2147/idr.s362099
18. Arrillaga-Moncrieff I, Capilla J, Mayayo E, et al. Different virulence levels of the species of *Sporothrix* in a murine model. *Clin Microbiol Infect.* 2009;15(7):651–655. doi:10.1111/j.1469-0691.2009.02824.x
19. Lozoya-Pérez NE, Clavijo-Giraldo DM, Martínez-Duncker I, et al. Influences of the culturing media in the virulence and cell wall of *Sporothrix schenckii*, *Sporothrix brasiliensis*, and *Sporothrix globosa*. *J Fungi.* 2020;6(4):323. doi:10.3390/jof6040323
20. Rodrigues AM, Gonçalves SS, de Carvalho JA, Borba-Santos LP, Rozental S, Camargo ZP. Current progress on epidemiology, diagnosis, and treatment of sporotrichosis and their future trends. *J Fungi.* 2022;8(8):776. doi:10.3390/jof8080776
21. Etchecopaz A, Toscanini MA, Gisbert A, et al. *Sporothrix brasiliensis*: a review of an emerging South American fungal pathogen, its related disease, presentation and spread in Argentina. *J Fungi.* 2021;7(3):170. doi:10.3390/jof7030170
22. Tamez-Castrellón AK, van der Beek SL, López-Ramírez LA, et al. Disruption of protein rhamnosylation affects the *Sporothrix schenckii*-host interaction. *Cell Surf.* 2021;7:100058. doi:10.1016/j.tscw.2021.100058
23. Lionakis MS, Drummond RA, Hohl TM. Immune responses to human fungal pathogens and therapeutic prospects. *Nat Rev Immunol.* 2023;1–20. doi:10.1038/s41577-022-00826-w
24. Loh JT, Lam KP. Fungal infections: immune defense, immunotherapies and vaccines. *Adv Drug Deliv Rev.* 2023;196:114775. doi:10.1016/j.addr.2023.114775
25. Gómez-Gaviria M, Vargas-Macías AP, García-Carnero LC, Martínez-Duncker I, Mora-Montes HM. Role of protein glycosylation in interactions of medically relevant fungi with the host. *J Fungi.* 2021;7(10):875. doi:10.3390/jof7100875
26. Martínez-Álvarez JA, Pérez-García LA, Mellado-Mojica E, et al. *Sporothrix schenckii sensu stricto* and *Sporothrix brasiliensis* are differentially recognized by human peripheral blood mononuclear cells. *Front Microbiol.* 2017;8:843. doi:10.3389/fmicb.2017.00843
27. Villalobos-Duno HL, Barreto LA, Alvarez-Aular Á, et al. Comparison of cell wall polysaccharide composition and structure between strains of *Sporothrix schenckii* and *Sporothrix brasiliensis*. *Front Microbiol.* 2021;12:726958. doi:10.3389/fmicb.2021.726958
28. Lopes-Bezerra LM, Walker LA, Niño-Vega G, et al. Cell walls of the dimorphic fungal pathogens *Sporothrix schenckii* and *Sporothrix brasiliensis* exhibit bilaminar structures and sloughing of extensive and intact layers. *PLoS Negl Trop Dis.* 2018;12(3):e0006169–e0006169. doi:10.1371/journal.pntd.0006169
29. Lozoya-Pérez NE, Casas-Flores S, de Almeida JRF, et al. Silencing of *OCH1* unveils the role of *Sporothrix schenckii* N-linked glycans during the host-fungus interaction. *Infect Drug Resist.* 2019;12:67–85. doi:10.2147/idr.S185037
30. Bezerra L. *Sporothrix schenckii* cell wall peptidorhamnomannans. Mini Review. *Front Microbiol.* 2011;2:243. doi:10.3389/fmicb.2011.00243
31. López-Ramírez LA, Martínez-Duncker I, Márquez-Márquez A, Vargas-Macías AP, Mora-Montes HM. Silencing of *ROT2*, the encoding gene of the endoplasmic reticulum glucosidase II, affects the cell wall and the *Sporothrix schenckii*-host interaction. *J Fungi.* 2022;8(11):1220. doi:10.3390/jof8111220
32. Castro RA, Kubitschek-Barreira PH, Teixeira PAC, et al. Differences in cell morphometry, cell wall topography and Gp70 expression correlate with the virulence of *Sporothrix brasiliensis* clinical isolates. *PLoS One.* 2013;8(10):e75656. doi:10.1371/journal.pone.0075656
33. Lloyd KO, Bitoon MA. Isolation and purification of a peptido-rhamnomannan from the yeast form of *Sporothrix schenckii*. Structural and immunochemical studies. *J Immunol.* 1971;107(3):663–671. doi:10.4049/jimmunol.107.3.663
34. García-Carnero LC, Salinas-Marín R, Lozoya-Pérez NE, et al. The Heat shock protein 60 and Pap1 participate in the *Sporothrix schenckii*-host interaction. *J Fungi.* 2021;7(11):960. doi:10.3390/jof7110960

35. Vargas-Macías AP, Gómez-Gaviria M, García-Carnero LC, Mora-Montes HM. Current models to study the *Sporothrix*-host interaction. Review. *Front Fungal Biol.* 2022;3:833111. doi:10.3389/ffunb.2022.833111
36. Ward RA, Vyas JM. The first line of defense: effector pathways of anti-fungal innate immunity. *Curr Opin Microbiol.* 2020;58:160–165. doi:10.1016/j.mib.2020.10.003
37. Alegranci P, de Abreu Ribeiro LC, Ferreira LS, et al. The predominance of alternatively activated macrophages following challenge with cell wall peptide-polysaccharide after prior infection with *Sporothrix schenckii*. *Mycopathologia.* 2013;176(1–2):57–65. doi:10.1007/s11046-013-9663-y
38. Franco Dde L, Nascimento RC, Ferreira KS, Almeida SR. Antibodies against *Sporothrix schenckii* enhance TNF- α production and killing by macrophages. *Scand J Immunol.* 2012;75(2):142–146. doi:10.1111/j.1365-3083.2011.02636.x
39. Carlos IZ, Sgarbi DB, Santos GC, Placeres MC. *Sporothrix schenckii* lipid inhibits macrophage phagocytosis: involvement of nitric oxide and tumour necrosis factor- α . *Scand J Immunol.* 2003;57(3):214–220. doi:10.1046/j.1365-3083.2003.01175.x
40. Tachibana T, Matsuyama T, Mitsuyama M. Involvement of CD4+ T cells and macrophages in acquired protection against infection with *Sporothrix schenckii* in mice. *Med Mycol.* 1999;37(6):397–404. doi:10.1046/j.1365-280x.1999.00239.x
41. Oda LM, Kubelka CF, Alviano CS, Travassos LR. Ingestion of yeast forms of *Sporothrix schenckii* by mouse peritoneal macrophages. *Infect Immun.* 1983;39(2):497–504. doi:10.1128/iai.39.2.497-504.1983
42. Huang L, Zhang J, Du W, et al. Chitin-rich heteroglycan from *Sporothrix schenckii sensu stricto* potentiates fungal clearance in a mouse model of sporotrichosis and promotes macrophages phagocytosis. *BMC Microbiol.* 2021;21(1):190. doi:10.1186/s12866-021-02243-w
43. Jellmayer JA, Ferreira LS, Manente FA, et al. Dectin-1 expression by macrophages and related antifungal mechanisms in a murine model of *Sporothrix schenckii sensu stricto* systemic infection. *Microb Pathog.* 2017;110:78–84. doi:10.1016/j.micpath.2017.06.025
44. Negrini Tde C, Ferreira LS, Alegranci P, et al. Role of TLR-2 and fungal surface antigens on innate immune response against *Sporothrix schenckii*. *Immunol Invest.* 2013;42(1):36–48. doi:10.3109/08820139.2012.719982
45. Sassá MF, Ferreira LS, Ribeiro LC, Carlos IZ. Immune response against *Sporothrix schenckii* in TLR-4-deficient mice. *Mycopathologia.* 2012;174(1):21–30. doi:10.1007/s11046-012-9523-1
46. Sassá MF, Sauri AE, Souza LF, Ribeiro LC, Sgarbi DB, Carlos IZ. Response of macrophage Toll-like receptor 4 to a *Sporothrix schenckii* lipid extract during experimental sporotrichosis. *Immunology.* 2009;128(2):301–309. doi:10.1111/j.1365-2567.2009.03118.x
47. Rossato L, Santos SSD, Ferreira LG, de Almeida SR. The importance of Toll-like receptor 4 during experimental *Sporothrix brasiliensis* infection. *Med Mycol.* 2019;57(4):489–495. doi:10.1093/mmy/myy048
48. Rossato L, Silvana Dos Santos S, Ferreira LG, Rogério de Almeida S. The impact of the absence of Toll-like receptor-2 during *Sporothrix brasiliensis* infection. *J Med Microbiol.* 2019;68(1):87–94. doi:10.1099/jmm.0.000876
49. Neves GWP, Wong SSW, Aimaniana V, et al. Complement-mediated differential immune response of human macrophages to *Sporothrix* species through interaction with their cell wall peptidoglycan-mannans. *Front Immunol.* 2021;12:749074. doi:10.3389/fimmu.2021.749074
50. Song Y, Yao L, Zhen Y, et al. *Sporothrix globosa* melanin inhibits antigen presentation by macrophages and enhances deep organ dissemination. *Braz J Microbiol.* 2021;52(1):19–31. doi:10.1007/s42770-020-00345-7
51. Guan MQ, Yao L, Zhen Y, Song Y, Cui Y, Li SS. Melanin of *Sporothrix globosa* affects the function of THP-1 macrophages and modulates the expression of TLR2 and TLR4. *Microb Pathog.* 2021;159:105158. doi:10.1016/j.micpath.2021.105158
52. Uenotsuchi T, Takeuchi S, Matsuda T, et al. Differential induction of Th1-prone immunity by human dendritic cells activated with *Sporothrix schenckii* of cutaneous and visceral origins to determine their different virulence. *Int Immunol.* 2006;18(12):1637–1646. doi:10.1093/intimm/dx1097
53. Quinello C, Souza Ferreira L, Picolli I, et al. *Sporothrix schenckii* cell wall proteins-stimulated BMDCs are able to induce a Th1-prone cytokine profile *in vitro*. *J Fungi.* 2018;4(3):106. doi:10.3390/jof403106
54. Verdan FF, Faleiros JC, Ferreira LS, et al. Dendritic cell are able to differentially recognize *Sporothrix schenckii* antigens and promote Th1/Th17 response *in vitro*. *Immunobiology.* 2012;217(8):788–794. doi:10.1016/j.imbio.2012.04.006
55. Kusuvara M, Qian H, Li X, et al. Mouse bone marrow-derived dendritic cells can phagocytize the *Sporothrix schenckii*, and mature and activate the immune response by secreting interleukin-12 and presenting antigens to T lymphocytes. *J Dermatol.* 2014;41(5):386–392. doi:10.1111/1346-8138.12472
56. Madrid H, Cano J, Gené J, Bonifaz A, Toriello C, Guarro J. *Sporothrix globosa*, a pathogenic fungus with widespread geographical distribution. *Rev Iberoam Micol.* 2009;26(3):218–222. doi:10.1016/j.riam.2009.02.005
57. Teixeira MM, de Almeida LG, Kubitschek-Barreira P, et al. Comparative genomics of the major fungal agents of human and animal sporotrichosis: *Sporothrix schenckii* and *Sporothrix brasiliensis*. *BMC Genomics.* 2014;15:943. doi:10.1186/1471-2164-15-943
58. Giosa D, Felice MR, Giuffrè L, et al. Transcriptome-wide expression profiling of *Sporothrix schenckii* yeast and mycelial forms and the establishment of the *Sporothrix* Genome DataBase. *Microb Genom.* 2020;6(10):mgen000445. doi:10.1099/mgen.0.000445
59. Giosa D, Giuffrè L, Felice MR, et al. P421 Whole-transcriptome analysis of *Sporothrix brasiliensis* grown in mold- and yeast-inducing conditions. *Med Mycol.* 2022;60(Supplement_1):myac072P421. doi:10.1093/mmy/myac072.P421
60. Trujillo-Esquivel E, Martínez-Álvarez JA, Clavijo-Giraldo DM, et al. The *Sporothrix schenckii* gene encoding for the ribosomal protein L6 has constitutive and stable expression and works as an endogenous control in gene expression analysis. *Front Microbiol.* 2017;8:1676. doi:10.3389/fmicb.2017.01676
61. Endres S, Ghorbani R, Lonnemann G, van der Meer JW, Dinarello CA. Measurement of immunoreactive interleukin-1 beta from human mononuclear cells: optimization of recovery, intrasubject consistency, and comparison with interleukin-1 alpha and tumor necrosis factor. *Clin Immunol Immunopathol.* 1988;49(3):424–438. doi:10.1016/0090-1229(88)90130-4
62. Perez-Garcia LA, Csonka K, Flores-Carreón A, et al. Role of protein glycosylation in *Candida parapsilosis* cell wall integrity and host interaction. *Front Microbiol.* 2016;7:306. doi:10.3389/fmicb.2016.00306
63. Cambi A, Gijzen K, de Vries IJM, et al. The C-type lectin DC-SIGN (CD209) is an antigen-uptake receptor for *Candida albicans* on dendritic cells. *Eur J Immunol.* 2003;33(2):532–538. doi:10.1002/immu.200310029
64. Cambi A, Netea MG, Mora-Montes HM, et al. Dendritic cell interaction with *Candida albicans* critically depends on N-linked mannan. *J Biol Chem.* 2008;283(29):20590–20599. doi:10.1074/jbc.M709334200
65. Hernandez-Chavez MJ, Franco B, Clavijo-Giraldo DM, Hernandez NV, Estrada-Mata E, Mora-Montes HM. Role of protein phosphomannosylation in the *Candida tropicalis*-macrophage interaction. *FEMS Yeast Res.* 2018. doi:10.1093/femsyr/foy053

66. Lozoya-Pérez NE, Casas-Flores S, Martínez-Álvarez JA, et al. Generation of *Sporothrix schenckii* mutants expressing the green fluorescent protein suitable for the study of host-fungus interactions. *Fungal Biol.* 2018;122:1023–1030. doi:10.1016/j.funbio.2018.07.004
67. Gonzalez-Hernandez RJ, Jin K, Hernandez-Chavez MJ, et al. Phosphomannosylation and the functional analysis of the extended *Candida albicans* *MNN4*-like gene family. *Front Microbiol.* 2017;8:2156. doi:10.3389/fmicb.2017.02156
68. Mora-Montes HM, Bates S, Netea MG, et al. A multifunctional mannosyltransferase family in *Candida albicans* determines cell wall mannan structure and host-fungus interactions. *J Biol Chem.* 2010;285(16):12087–12095. doi:10.1074/jbc.M109.081513
69. Estrada-Mata E, Navarro-Arias MJ, Perez-Garcia LA, et al. Members of the *Candida parapsilosis* complex and *Candida albicans* are differentially recognized by human peripheral blood mononuclear cells. *Front Microbiol.* 2015;6:1527. doi:10.3389/fmicb.2015.01527
70. Navarro-Arias MJ, Defosse TA, Dementhon K, et al. Disruption of protein mannosylation affects *Candida guilliermondii* cell wall, immune sensing, and virulence. *Front Microbiol.* 2016;7:1951. doi:10.3389/fmicb.2016.01951
71. Schwartz SN, Medoff G, Kobayashi GS, Kwan CN, Schlessinger D. Antifungal properties of polymyxin B and its potentiation of tetracycline as an antifungal agent. *Antimicrob Agents Chemother.* 1972;2(1):36–40. doi:10.1128/AAC.2.1.36
72. Cossarizza A, Chang H-D, Radbruch A, et al. Guidelines for the use of flow cytometry and cell sorting in immunological studies. *Eur J Immunol.* 2021;51(12):2708–3145. doi:10.1002/eji.202170126
73. Tamez-Castrellón AK, Romeo O, García-Carnero LC, Lozoya-Pérez NE, Mora-Montes HM. Virulence factors in *Sporothrix schenckii*, one of the causative agents of sporotrichosis. *Curr Protein Pept Sci.* 2020;21(3):295–312. doi:10.2174/1389203720666191007103004
74. Ruiz-Baca E, Mora-Montes HM, Lopez-Romero E, Toriello C, Mojica-Marin V, Urtiz-Estrada N. 2D-immunoblotting analysis of *Sporothrix schenckii* cell wall. *Mem Inst Oswaldo Cruz.* 2011;106(2):248–250. doi:10.1590/S0074-02762011000200021
75. Thomas CJ, Schroder K. Pattern recognition receptor function in neutrophils. *Trends Immunol.* 2013;34(7):317–328. doi:10.1016/j.it.2013.02.008
76. García-Carnero LC, Martínez-Duncker I, Gómez-Gaviria M, Mora-Montes HM. Differential recognition of clinically relevant *Sporothrix* species by human mononuclear cells. *J Fungi.* 2023;9(4):448. doi:10.3390/jof9040448
77. Bozza S, Perruccio K, Montagnoli C, et al. A dendritic cell vaccine against invasive aspergillosis in allogeneic hematopoietic transplantation. *Blood.* 2003;102(10):3807–3814. doi:10.1182/blood-2003-03-0748
78. Reid DM, Gow NAR, Brown GD. Pattern recognition: recent insights from Dectin-1. *Curr Opin Immunol.* 2009;21(1):30–37. doi:10.1016/j.coi.2009.01.003

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>