

Effect of Low Level Laser Therapy on Pain Reduction After Midpalatal Expansion in Rats

Ali Mohammad-Hoseyni¹, Maryam Pirmoradian-Najafabadi², Sajjad Ashnagar³, Hanieh Nokhbatolfoghahaei⁴, Reza Fekrazad⁵

¹Postgraduate Student, Department of Orthodontics, School of Dentistry, Shahed University, Tehran, Iran

²PhD Candidate, Department of Dental Biomaterials, School of Dentistry, Research Center for Science and Technology in Medicine, Tehran University of Medical Sciences, Tehran, Iran

³Department of Periodontics and Oral Medicine, Michigan School of Dentistry, Mi, USA

⁴Dentist, Laser Research Center of Dentistry, Dental Research Center, Dentistry Research Institute, Tehran University of Medical Sciences, Tehran, Iran

⁵Associate Professor, Department of Periodontics, Dental Faculty and Laser Research Center in Medical Sciences, AJA University of Medical Sciences, Tehran, Iran

Abstract

Objectives: The purpose of this experimental study was to evaluate the efficacy of low level laser therapy (LLLT) for pain reduction after midpalatal expansion in rats.

Materials and Methods: Sixty male Sprague six-week old rats weighing 180±10g were divided into seven groups (two experimental groups of 24 rats and one control group of 12 rats). The experimental groups were subjected to expansion with or without LLLT. The health status of each rat was monitored starting seven days prior to the experiment and evaluated by regular body weight monitoring during the study period. Diode laser with 810nm wavelength and 100 mw output power was used. Laser therapy and body weight monitoring were performed on days 0, 2, 4, 6, 8, 10, 12 and 14. The data were analyzed by One-way repeated measures ANOVA.

Results: The body weight of the experimental groups significantly decreased in the first two days because of the pain and difficult nutrition with the new appliance. Within the next two days, the body weight of all rats increased but this increase was significantly higher in the irradiated compared to the non-irradiated group. This significant improvement continued until day 14 and then between days 14 and 30 the rats gained weight similarly in the irradiated and control groups.

Conclusion: The study results showed that laser irradiated group continued to gain weight easier than the control group. This may be due to more efficient pain control due to laser irradiation after midpalatal expansion.

Key words: Laser Therapy, Low-Level; Palatal Expansion Technique; Rats

(*Journal of Dentistry, Tehran University of Medical Sciences, Tehran, Iran (2015; Vol. 12, No. 9)*)

Corresponding author:

H. Nokhbatolfoghahaei, Laser Research Center of Dentistry, Dental Research Center, Dentistry Research Institute, Tehran University of Medical Sciences, Tehran, Iran

h-nokhbeh@farabi.tums.ac.ir

Received: 16 February 2015

Accepted: 27 March 2015

INTRODUCTION

Pain is inevitable during orthodontic treatment. Orthodontists have always tried to minimize pain and discomfort. Due to fear of pain after placement of appliances, many patients may prefer other therapeutic approaches or even

reject treatment [1-6]. Previous studies indicate that 90-95% of orthodontic patients report pain during treatment period [2,7]. Orthodontic pain is due to forces applied on the teeth, which cause edema in periodontal tissues, resulting in release of inflammatory mediators [1,8].

Orthodontists use a variety of methods to alleviate orthodontic pain, including cognitive behavioral therapy (CBT) [2], use of non-steroidal anti-inflammatory drugs (NSAIDs) [1-3,7,9], anesthetic gels [1,2], chewing gums and wafer bytes [1,2], electrical nerve stimulation and vibratory stimulation [1]. All the afore-mentioned methods have merits and demerits. Although NSAIDs reduce orthodontic pain, they have side effects such as allergy, peptic ulcer, congestive heart problems and most importantly, they may negatively affect the process of tooth movement by inhibition of bone resorption. These side effects question the use of these drugs [1,2]. Low level laser therapy has been used in dentistry since 1970 and there is evidence of its pain controlling potential and anti-inflammatory properties due to its regenerative effect on neurons [1-3,6]. It is defined as laser treatment with low output power causing insignificant thermal changes and biomodulatory effects [1,2,4,5]. It has been demonstrated that LLLT can postpone the onset and reduce the severity and duration of pain; also it has very few side effects [1]. However, a number of reports have shown that laser therapy has no effect on the onset or severity of pain compared to the placebo group [3]. In order to standardize pain measurement, many studies have used visual analog scale (VAS), which is a psychometric response scale. When responding to a VAS, respondents specify their level of pain by indicating a position along a continuous line between two endpoints, which are no pain as number 0 and worst possible pain as number 10. However, studies on animal models for evaluation of pain and tooth movement cannot use VAS and thus, pain measurement is done through observing animal behavior [4,10,11]. The purpose of this study was to evaluate the efficacy of LLLT for pain reduction after midpalatal expansion in rats by monitoring the body weight of animals during the treatment period.

MATERIALS AND METHODS

In this animal study, 60 six-week old Sprague rats weighing 180 ± 10 g (Razi Vaccine and Serum Research Institute, Tehran, Iran) were used. To standardize the conditions, they were kept in Razi Institute for one week and then moved to the animal house of the Veterinary Faculty of Tehran University (survey site) under similar conditions in terms of water and food supply at 23°C [12]. Rats were monitored after their transfer and were randomly divided into three groups of two experimental ($n=24$) and one control ($n=12$) group as follows:

A: Control group: Without expansion or laser therapy

B: Experimental groups:

(a) Expansion, without laser therapy for 14 days

(b) Expansion, with laser therapy for 14 days

Orthodontic intervention:

All rats except for the control group were deeply sedated by intramuscular injection of 10% animal ketamine and 2% xylazine in order to place orthodontic appliances for maxillary expansion. The protocol by Saito and Shimizu [13] was slightly modified and followed for opening the midpalatal suture. This method involved using a 1.5 mm thick stainless steel ring (made of 0.5 mm thick orthodontic wire by an orthodontic technician)(Fig. 1). An initial space, smaller than ring diameter, was created with an osteotome between the incisors.

Then the ring was pushed with a slight pressure into the created space for expansion. A hole was created distal to both central incisors along the gingival papillae, in order to keep the ring tight in place, using a straight handpiece with a maximum speed of 35,000 rpm and a $\frac{1}{4}$ round bur. After ring placement, it was fixed using a brass wire with a round cross-section and a diameter of 0.5 mm through the created holes. The aim of this operation was to achieve 1.5mm expansion of midpalatal suture, equal to the



Fig. 1. A 1.5 mm diameter stainless steel ring made of 0.5 mm orthodontic wire with two additional arms for better handling.



Fig. 2. Expansion by 1.5mm between the upper incisors.




Fig. 3. Low-level GaAlAs laser device used in this study

size of the ring used in between the anterior teeth.

At the end, the fixing wire was bent so that it did not irritate the soft palatal tissue of rats.

According to Saito and Shimizu [13] describing the appliance placement in between incisors of rats, initial holes were created in proximal surfaces of incisors and then a 1.5 mm appliance was placed between the teeth (Fig. 2). Pilot assessments revealed that this process can highly make the teeth susceptible to fracture, and this was due to the location of holes adjacent to gingival papilla.

Hence, a slight change was made to make the process safer.

Primarily, the required space was provided by putting the appliance in between the incisors; afterwards, two holes along the long axis of the appliances were created on either side of the incisors. These changes minimized the risk of tooth fracture.

Laser therapy:

In this study, Photon-Lase III (DMC, Sao Paulo, Brazil) was used (Fig. 3). The rats received Gallium-Aluminum-Arsenide (GaAlAs) diode laser irradiation with 810 nm wavelength and output power of 100 mW and energy density of 4 J/cm², considering the safety regulations, on days 0, 2, 4, 6, 8, 10, 12 and 14 on four points, with the 0.6 mm optical fiber in contact with the mucosa. Irradiation was performed under general anesthesia. Irradiated points were: A buccal point 0.5 mm apical to the bone crest and three points in the palatal at 1, 2 and 3 mm apical to the bone crest. The non-lased group was generally anesthetized in the same periods of time as the laser treated group in order to have similar anesthetic intervention in both groups. Statistical analysis was performed using one-way repeated measures ANOVA. The results were obtained by weighing the rats daily.




Fig. 4. The differences in weight of the three groups: control, non-lased and lased. Horizontal axis: days of examination. Vertical axis: weight.

RESULTS

The mean (\pm standard deviation) weight in all groups before and after treatment is summarized in Tables 1 and 2 and Fig. 4.

The groups had significant differences in terms of weight ($P=0.008$). The difference in weight of the control and intervention groups was significant ($P\leq 0.05$).

But there was no difference in weight change between the two intervention groups ($P=0.461$). All experimental groups wearing the appliance exhibited a weight loss within the first two days.

The weight loss in the laser-treated group was significantly less than the value in the non-lased group.

Table 1. The mean weight (gr) of groups during the weighing sessions

| Mean weight (gr) | Day 0 | Day 2 | Day 4 | Day 6 | Day 8 | Day 10 | Day 12 | Day 14 |
|-------------------------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Control | 170.2±32.5 | 174.2±33.1 | 177±35.7 | 178.8±33.5 | 181.2±31.6 | 183.2±20.1 | 187.6±28.7 | 191±28.4 |
| Experimental non-lased | 203.3±28.5 | 192.7±28.3 | 191.4±28.0 | 194.2±27.7 | 200.4±28.3 | 207.8±28 | 210.7±27.5 | 213.3±27.2 |
| Experimental lased | 180.5±32.2 | 172.2±31.8 | 173.6±31.5 | 176±31 | 180.0±30.8 | 182.0±31.2 | 184.6±30.9 | 185.5±30.5 |

Table 2. The mean weight changes (gr) among the groups during the weighing sessions

| Change in weight (gr) | Days 0-2 | Days 2-4 | Days 4-6 | Days 6-8 | Days 8-10 | Days 10-12 | Days 12-14 |
|-------------------------------|-----------|----------|----------|----------|-----------|------------|------------|
| Control | +4±0.8 | +2.8±0.8 | +1.8±0.5 | +2.3±0.6 | +2±0.6 | +4.4±0.6 | +3.4±0.6 |
| Experimental non-lased | -10.5±1.5 | -1.3±1 | +2.8±0.8 | +6.2±0.8 | +7.3±0.8 | +2.9±0.4 | +2.5±0.4 |
| Experimental lased | -8.2±1.3 | +1.4±0.9 | +2.3±0.8 | +4.0±0.8 | +2±0.7 | +2.5±0.7 | +0.9±0.2 |

Weight loss in laser-treated group turned into weight gain after the second day. However, in the non-lased experimental group weight loss continued until day four. Despite the fact that the non-lased experimental group started to compensate the weight loss after day four, the laser-treated group still showed a significant difference in weight gain in the same period of time, compared to the non-lased experimental group. At the end of day eight, all the rats in laser treated group fully compensated their weight loss.

However, this process took 10 days for the non-lased experimental group. Weight change after day eight in the laser treated group was greater than that in the non-lased experimental group, but not significantly (Table 1).

DISCUSSION

Pain is one of the most important reasons, which discourages patients from seeking orthodontic treatment. It is believed that pain caused by appliance placement in orthodontic treatment cannot be accurately and reliably measured. In 1962, Burstone [14] introduced a scale for measuring pain; however due to physical and mental differences among people and their impact on pain processing and perception, these scales are not much reliable [15].

However, the effect of pain on life habits such as chewing and eating food is obvious and measurable. One method for measuring the quantity and quality of eating is regular weight measurement, which was used in the current study. Sprague rats were used in the current study, which are a common breed similar to Wistar in animal studies. Only male rats were chosen in order to eliminate the possible confounding effect of hormonal changes on the results. In our study, weight loss was seen during the first two days in both intervention groups. This weight loss, attributed to postoperative pain, could also be due to difficulty eating dry food while wearing the appliance.

However, continuous weight loss in the non-lased experimental group indicates presence of a factor not seen in the laser-treated group. Considering the effects of LLLT on pain reduction through various means, pain is thought to be the distinct factor preventing the rats in the non-lased experimental group from eating properly. Although the reason for pain encountered during rapid palatal expansion is not fully understood, various concepts have been discussed. In this study a special method for rapid midpalatal expansion was used which includes involvement of periodontal ligament (PDL) of both maxillary incisors and the midpalatal suture. Thus, the pain caused by the appliance insertion has both periodontal and osseous origins.

Furstman and Bernick [16] in 1972 suggested that periodontal pain is caused by pressure ischemia, inflammation and edema. Burstone [14] identified immediate and delayed pain response, the former being related to the initial compression of the periodontal ligament immediately after placement of the appliance and the latter, which started a few hours later, was termed hyperalgesia of the PDL. Prostaglandins have been shown to cause hyperalgesia, which is increased sensitivity to noxious agents such as histamine, bradykinin, serotonin and substance P. Perception of pain is due to changes in blood flow of the PDL and is correlated with the presence of substances such as prostaglandins and substance P. Factors involved in pain development in PDL can be somehow attributed to bone pain and include factors such as protons, kinins, histamine and serotonin [17]. However, LLLT can affect cell membrane permeability to calcium, sodium and potassium ions leading to events that ultimately result in pain control. These events include degradation of bradykinin [18], increased activity of cellular receptors responsible for inducing production of endorphins [19], decreased activity of C fibers and increased action potential of neurons [18]. This is why various studies have reported LLLT to have

pain relieving and anti-inflammatory properties [18, 20, 21].

Weight loss of rats in the initial days after placing the appliance is probably due to pain from their rapid maxillary expansion and as mentioned earlier, laser irradiation decreases inflammation, edema and pain. Lasers cause such effects through several mechanisms such as increasing the levels of certain prostaglandins such as PGI₂, which have anti-inflammatory effects, increasing the levels of immunoglobulins and lymphokines, which play a role in the immune system and increasing beta-endorphins that are involved in analgesia, inhibition of bradykinin synthesis, doubling the lymphatic drainage and balancing osmotic and oncotic pressure [22]. Significant difference in weight changes within the first few days after insertion of appliance among the groups confirms it. However, as time passed and the critical phase of inflammation caused by the appliance subsided, the level of inflammatory cytokines decreased. Hence, within the second week after orthodontic treatment we noticed regain of weight in the non-lased experimental group rats.

Analgesic effects of lasers have been confirmed in several clinical studies. Lim et al, [23] in 1995, Trotamano et al, [5] in 2009 and Artes-Ribas et al, [9] in 2013 have shown that irradiation of low level laser can reduce pain in patients receiving orthodontic treatment. In the current study, early relief of pain and lower pain peak in days one and two were noted in the laser-treated group. It is worth bearing in mind that in all the afore-mentioned studies, lasers with 830 nm wavelength were used; but in our study, 810 nm wavelength was utilized. It must be underlined that Esper et al, [7] in a study on the effect of LLLT and LED irradiation on pain following orthodontic treatment found that LED was the best choice for this purpose (superior to LLLT). They used 660nm InGaAlP laser for LLLT and 640nm LED. The results can be related to the wavelength used in their

study, which affects the superficial parts of tissues.

CONCLUSION

These results show that irradiated group continued to gain weight easier than the control group. This result can be due to more efficient pain control after midpalatal expansion in the irradiated group. After day 14, the conditions of both groups were similar.

REFERENCES

- 1- He WL, Li CJ, Liu ZP, Sun JF, Hu ZA, Yin X, et al. Efficacy of low-level laser therapy in the management of orthodontic pain: a systematic review and meta-analysis. *Lasers Med Sci*. 2013 Nov;28(6):1581-9.
- 2- Eslamian L, Borzabadi-Farahani A, Hassanzadeh-Azhiri A, Badiie MR, Fekrazad R. The effect of 810-nm low-level laser therapy on pain caused by orthodontic elastomeric separators. *Lasers Med Sci*. 2014 Mar;29(2):559-64.
- 3- Kim WT, Bayome M, Park JB, Park JH, Baek SH, Kook YA. Effect of frequent laser irradiation on orthodontic pain. A single-blind randomized clinical trial. *Angle Orthod*. 2013 Jul;83(4):611-6.
- 4- Xiaoting L, Yin T, Yangxi C. Interventions for pain during fixed orthodontic appliance therapy. A systematic review. *Angle Orthod*. 2010 Sep;80(5):925-32.
- 5- Tortamano A, Lenzi DC, Haddad AC, Bottino MC, Dominguez GC, Vigorito JW. Low-level laser therapy for pain caused by placement of the first orthodontic archwire: a randomized clinical trial. *Am J Orthod Dentofacial Orthop*. 2009 Nov;136(5):662-7.
- 6- Turhani D, Scheriau M, Kapral D, Benesch T, Jonke E, Bantleon HP. Pain relief by single low-level laser irradiation in orthodontic patients undergoing fixed appliance therapy. *Am J Orthod Dentofacial Orthop*. 2006 Sep;130(3):371-7.
- 7- Esper MA, Nicolau RA, Arisawa EA. The

- effect of two phototherapy protocols on pain control in orthodontic procedure--a preliminary clinical study. *Lasers Med Sci.* 2011 Sep;26(5):657-63.
- 8- Stanfeld J, Jones J, Laster L, Davidovitch Z. Biochemical aspects of orthodontic tooth movement. I. Cyclic nucleotide and prostaglandin concentrations in tissues surrounding orthodontically treated teeth in vivo. *Am J Orthod Dentofacial Orthop.* 1986 Aug;90(2):139-48.
- 9- Artes-Ribas M, Arnabat-Dominguez J, Puigdollers A. Analgesic effect of a low-level laser therapy (830 nm) in early orthodontic treatment. *Lasers Med Sci.* 2013 Jan;28(1):335-41.
- 10- Mogil JS. Animal models of pain: progress and challenges. *Nat Rev Neurosci.* 2009 Apr;10(4):283-94.
- 11- Molony V, Kent JE. Assessment of acute pain in farm animals using behavioral and physiological measurements. *J Anim Sci.* 1997 Jan;75(1):266-72.
- 12- ACLAM Task Force Members, Kohn DF, Martin TE, Foley PL, Morris TH, Swindle MM, Vogler GA, Wixson SK. Public statement: guidelines for the assessment and management of pain in rodents and rabbits. *J Am Assoc Lab Anim Sci.* 2007 Mar;46(2):97-108.
- 13- Saito S, Shimizu N. Stimulatory effects of low-power laser irradiation on bone regeneration in midpalatal suture during expansion in the rat. *Am J Orthod Dentofacial Orthop.* 1997 May;111(5):525-32.
- 14- Burstone C. Biomechanics of tooth movement. In: Kraus BS, Riedel RA (eds.) *Vistas in orthodontics* Lea & Febiger Philadelphia:1962:197-213.
- 15- Shavit Y, Fish G, Wolf G, Mayburd E, Meerson Y, Yirmiya R, et al. The effects of perioperative pain management techniques on food consumption and body weight after laparotomy in rats. *Anesth Analg.* 2005 Oct;101(4):1112-6.
- 16- Furstman L, Bernick S. Clinical considerations of the periodontium. *Am J Orthod.* 1972 Feb;61(2):138-55.
- 17- Krishnan V, Davidovitch Z. Cellular, molecular, and tissue-level reactions to orthodontic force. *Am J Orthod Dentofacial Orthop.* 2006 Apr;129(4):469.e1-32.
- 18- Pozza DH1, Fregapani PW, Weber JB, de Oliveira MG, de Oliveira MA, Ribeiro Neto N, et al. Analgesic action of laser therapy (LLLT) in an animal model. *Med Oral Patol Oral Cir Bucal.* 2008 Oct 1;13(10):E648-52.
- 19- Hagiwara S, Iwasaka H, Okuda K, Noguchi T. GaAlAs (830 nm) low-level laser enhances peripheral endogenous opioid analgesia in rats. *Lasers Surg Med.* 2007 Dec;39(10):797-802.
- 20- Albertini R, Villaverde AB, Aimbire F, Salgado MA, Bjordal JM, Alves LP, et al. Anti-inflammatory effects of low-level laser therapy (LLLT) with two different red wavelengths (660nm and 684nm) in carrageenan-induced rat paw edema. *J Photochem Photobiol B.* 2007 Nov 12;89(1): 50-5.
- 21- Boschi ES, Leite CE, Saciura VC, Caberlon E, Lunardelli A, Bitencourt S, et al. Anti-inflammatory effects of low-level laser therapy (660 nm) in the early phase in carrageenan-induced pleurisy in rat. *Lasers Surg Med.* 2008 Sep;40(7):500-8.
- 22- Gama SK, Habib FA, Monteiro JS, Paraguassú GM, Araújo TM, Cangussú MC, et al. Tooth movement after infrared laser phototherapy: clinical study in rodents. *Photomed Laser Surg.* 2010 Oct;28 Suppl2: S79-83.
- 23- Lim HM, Lew KK, Tay DK. A clinical investigation of the efficacy of low level laser therapy in reducing orthodontic postadjustment pain. *Am J Orthod Dentofacial Orthop.* 1995 Dec;108(6):614-22.