







Article

Synthesis of Novel Crown Ether-Squaramides and Their Application as Phase-Transfer Catalysts

Zsuzsanna Fehér¹, Dóra Richter¹, Sándor Nagy¹ , Péter Bagi¹ , Zsolt Rapi¹ , András Simon²,
László Drahos³ , Péter Huszthy¹ , Péter Bakó¹ and József Kupai^{1,*} 

- ¹ Department of Organic Chemistry and Technology, Budapest University of Technology and Economics, Szent Gellért tér 4, H-1111 Budapest, Hungary; zsuzsanna.feher@edu.bme.hu (Z.F.); richterdora98@gmail.com (D.R.); nagy.sandor@mail.bme.hu (S.N.); bagi.peter@vbk.bme.hu (P.B.); rapi.zsolt@vbk.bme.hu (Z.R.); huszthy.peter@vbk.bme.hu (P.H.); bako.peter@vbk.bme.hu (P.B.)
- ² Department of Inorganic and Analytical Chemistry, Budapest University of Technology and Economics, Szent Gellért tér 4, H-1111 Budapest, Hungary; simon.andras@vbk.bme.hu
- ³ MS Proteomics Research Group, Research Centre for Natural Sciences, Hungarian Academy of Sciences, Magyar Tudósok körútja 2, H-1117 Budapest, Hungary; drahos.laszlo@ttk.hu
- * Correspondence: kupai.jozsef@vbk.bme.hu; Tel.: +36-14-632-369

Abstract: This work presents the synthesis of six new phase-transfer organocatalysts in which the squaramide unit is directly linked to the nitrogen atom of an aza-crown ether. Four chiral skeletons, namely hydroquinine, quinine, cinchonine (cinchonas), and α -D-glucopyranoside were responsible for the asymmetric construction of an all-carbon quaternary stereogenic center in α -alkylation and Michael addition reactions of malonic esters. We investigated the effects of these different chiral units and that of crown ethers with different sizes on catalytic activity and enantioselectivity. During extensive parameter investigations, both conventional and emerging green solvents were screened, providing valuable α,α -disubstituted malonic ester derivatives with excellent yields (up to 98%).

Keywords: asymmetric catalysis; phase-transfer catalysis; enantioselectivity; allylation; crown compounds; carbohydrates; amino acids



Citation: Fehér, Z.; Richter, D.; Nagy, S.; Bagi, P.; Rapi, Z.; Simon, A.; Drahos, L.; Huszthy, P.; Bakó, P.; Kupai, J. Synthesis of Novel Crown Ether-Squaramides and Their Application as Phase-Transfer Catalysts. *Molecules* **2021**, *26*, 6542. <https://doi.org/10.3390/molecules26216542>

Academic Editors: Dimitrios Bikiaris and Maurizio Benaglia

Received: 15 September 2021
Accepted: 27 October 2021
Published: 29 October 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Investigating the creation of quaternary stereogenic centers is essential in organic synthesis as it poses a challenge to organic chemists due to the possible steric repulsion between the groups around the stereocenter. In the last decade, the synthesis of chiral α,α -disubstituted malonates has gained considerable attention [1,2]. These compounds can be applied in the preparation of α,α -disubstituted amino acids, as they can undergo chemoselective transformations if their carboxylic acid moieties are protected with different groups, e.g., through Curtius rearrangement [3]. A decade ago, a new method involving enantioselective phase-transfer catalytic double α -alkylation of malonates was developed for the construction of chiral quaternary carbon centers, applying *tert*-butyl diphenylmethyl α -alkylmalonates as starting materials [1]. This method was later extended in numerous publications by examining a broad substrate scope [3–6]. Moreover, similar phase-transfer catalytic reactions, such as the α -benzoyloxylation of *tert*-butyl methyl α -alkylmalonates [7] and the Michael addition reaction of *tert*-butyl methyl α -benzylmalonate to acrylates, have also been elaborated [2]. In all of these papers, quaternary ammonium salts were applied as phase-transfer catalysts (PTCs) in the enantioselective α -alkylation of malonates, while chiral macrocycle PTCs remain a rather neglected tool for these asymmetric reactions.

Phase-transfer catalysis is one of the most efficient asymmetric synthetic methods, which is also inexpensive and sustainable as it involves simple procedures and mild reaction conditions. Thanks to these advantages, phase-transfer catalysis is particularly suitable for industrial applications [8].

The most common types of asymmetric phase-transfer catalysts are chiral quaternary ammonium and phosphonium salts, but chiral crown ether derivatives and other macrocycles emerge as alternatives despite their cumbersome and costly synthesis [9–11]. Crown ethers are neutral ligands that can complex and transport alkali metal cations into the organic phase. This crown ether–metal cation complex plays the same role as the quaternary onium cation, but crown ethers work with a different mechanism, called cation-binding catalysis, during which the entire reacting ion pair is transported into the organic phase, not just the anion [12]. The most conspicuous benefits that help crown ether derivatives to stand out from other catalysts are as follows: they are usually more resistant to strong bases, and the cation is usually more accessible than the positively charged nitrogen or phosphorus atom in ammonium or phosphonium salts; consequently, a stronger interaction can take place with the reactive anion. Furthermore, crown ethers are more effective in extracting inorganic salts from their solid form [13].

In asymmetric phase-transfer catalysis, introducing hydrogen bond donor units into catalysts has recently attained broad application [14]. The most common of those units include the hydroxyl group, amide, (thio)urea and squaramide. In the case of quaternary onium salts, there are many examples for the installation of these units into catalyst scaffold [15–20]. However, in crown ether derivatives, only hydroxyl groups are frequently used as ancillary components capable of forming hydrogen bonds.

Squaramides are highly effective double hydrogen bond donor units due to their rigid, aromatic four-membered ring [21]. There are a few examples in which a squaramide unit was connected indirectly to crown ether derivatives [22–27]. However, these ligands have only been applied for ion pair transport so far.


Herein, we present the synthesis of novel chiral crown ether-squaramide phase-transfer organocatalysts, and the catalytic performance of these new derivatives were tested in the synthesis of α,α -disubstituted malonates. The effects of different chiral units and various cavity sizes of crown ethers on catalytic activity and enantioselectivity were investigated. In these catalysts, the squaramide unit is directly linked to the nitrogen atom of an aza-crown ether; thus, one amino group of the squaramide unit is tertiary. This means that the squaramide unit serves as a linker and a single hydrogen bond donor unit [28,29]. By applying this catalyst design, we have anticipated that one NH group and the cation complexed by the crown ether can form secondary interactions of sufficient strength during the catalysis.

2. Results and Discussion

2.1. Synthesis of Crown Ether-Squaramide Phase-Transfer Catalysts

We have prepared a series of crown ethers with a squaramide hydrogen bond donor unit (Scheme 1) for catalyzing the α -alkylation reaction of malonates, which may be an important method for obtaining α,α -disubstituted α -amino acid intermediates. Different cinchona alkaloids [quinine (**Q**), hydroquinine (**HQ**) and cinchonine (**C**)] and a D-glucose derivative (**G**) were chosen as chiral starting materials for the syntheses of the catalysts. Previously, we have shown that the substituent on the nitrogen atom of glucose-based aza-crown ether **G** influences the catalytic effect [30].

The cinchona alkaloid-based catalysts were prepared in 3 steps. The C9 hydroxyl group of the commercially available starting materials (**Q**, **HQ**, **C**) was converted to an amino group as reported earlier [31]. In the next step, half squaramide derivatives quinine-half squaramide (**Q-HSQ**), hydroquinine-half squaramide (**HQ-HSQ**) and cinchonine-half squaramide (**C-HSQ**) were obtained by the addition of these amines to dimethyl squarate (**DMSQ**) [32,33]. Finally, the addition of 1-aza-15-crown-5 or 1-aza-18-crown-6 ethers to the previously mentioned half squaramides afforded the corresponding cinchona-crown ether-squaramide derivatives (**Q5**, **Q6**, **C5**, **C6**, **HQ5**).



Scheme 1. Synthesis of cinchona alkaloid-based (**Q5**, **Q6**, **HQ5**, **C5**, **C6**) and D-glucose-based (**G5**) crown ether-squaramide phase-transfer catalysts. DIAD: diisopropyl azodicarboxylate, DPPA: diphenylphosphoryl azide, DCM: dichloromethane.


The glucose-based catalyst was prepared in a two-step-synthesis from glucose-aza-crown ether derivative **G**, which can be prepared in 5 steps from D-glucose [34,35]. First, dimethyl squarate (**DMSQ**) was reacted with 3,5-bis(trifluoromethyl)aniline to obtain a half squaramide derivative **HSQ** [36]; then, the treatment of aza-crown ether **G** with this half squaramide (**HSQ**) led to the appropriate glucose-crown ether-squaramide derivative (**G5**).

During the synthetic procedures, all compounds were characterized by well-established methods, including HRMS, IR, ^1H and ^{13}C NMR spectroscopies.

2.2. Application of the Catalysts

The catalysts were applied in the asymmetric α -alkylation of *tert*-butyl methyl α -benzylmalonate [3] (**1**) under phase-transfer conditions (Scheme 2).

First, we compared the activities and enantioselectivities of the catalysts; these results are shown in Table 1. The catalysts provided good yields except catalysts bearing a 1-aza-18-crown-6 ether macroring (**Q6** and **C6**, Table 1, Entries 5–6), and only poor or no enantioselectivity. The highest enantiomeric excess of (*R*)-**3** was obtained with catalyst **C5**, which was still as low as 9% (Table 1, Entry 3). In the absence of a base while applying **C5** as catalyst, no product was observed according to thin-layer chromatography (TLC) analysis. When 50% aq. NaOH was applied as a base in the absence of a phase-transfer catalyst, the yield was only 33%.



Scheme 2. Synthesis of α,α -disubstituted malonate (*R*)-3 by asymmetric α -alkylation of *tert*-butyl methyl α -benzylmalonate (**1**) with allyl halides **2** under phase-transfer conditions.

Table 1. Comparison of the catalysts in the α -alkylation of malonate **1**^a.

Entry	Catalyst	Base ^b	Yield ^c (%)	ee ^d (%)
1	Q5	50% aq. NaOH	98	5
2	HQ5	50% aq. NaOH	93	<5
3	C5	50% aq. NaOH	73	9
4	G5	50% aq. NaOH	86	5
5	Q6	50% aq. KOH	59	<5
6	C6	50% aq. KOH	37	<5

^a Reaction conditions: *tert*-butyl methyl α -benzylmalonate (1 eq), allyl bromide reagent (1 eq), catalyst (10 mol%), base (50 eq), 2.4 mL dichloromethane (DCM) solvent, 25 °C, 24 h reaction time. ^b Applied base chosen considering the phase-transfer catalyst (PTC) crown ether cavity size. ^c Yield of isolated product purified by preparative thin-layer chromatography (TLC). ^d Determined by chiral high-performance liquid chromatography (HPLC).

Next, we conducted a parameter study with catalyst **C5** to maximize the enantioselectivity. First, we examined the effect of the solvent while applying 50% aq. NaOH or solid NaOH as base (Table 2). In the case of aqueous NaOH, six different solvents were tested (Table 2, Entries 1–6); however, only the most suitable solvent (DCM: dichloromethane; Table 2, Entry 7) and two other polar aprotic solvents (MeCN; THF: tetrahydrofuran; Table 2, Entries 8–9) were used with solid NaOH. Dichloromethane proved to be the best solvent in terms of yield and enantiomeric excess by applying 50% aq. NaOH as base (Table 2, Entry 1, 73% yield, 9% ee).

Table 2. Examination of the effect of the solvent in the presence of catalyst **C5**^a.

Entry	Solvent	Base	Base Quantity (eq)	Yield ^b (%)	ee ^c (%)
1	DCM	50% aq. NaOH	50	73	9
2	toluene	50% aq. NaOH	50	60	7
3	EtOAc	50% aq. NaOH	50	65	5
4	MTBE	50% aq. NaOH	50	51	<5
5	CPME	50% aq. NaOH	50	49	<5
6	2-Me-THF	50% aq. NaOH	50	51	<5
7	DCM	solid NaOH	1	38	10
8	MeCN	solid NaOH	1	49	<5
9	THF	solid NaOH	1	38	<5

^a Reaction conditions: *tert*-butyl methyl α -benzylmalonate (1 eq), allyl bromide reagent (1 eq), **C5** catalyst (10 mol%), base, 2.4 mL solvent, 25 °C, 24 h reaction time. ^b Yield of isolated product purified by preparative TLC. ^c Determined by chiral HPLC. ee: enantiomeric excess, DCM: dichloromethane, MTBE: *tert*-butyl methyl ether, CPME: cyclopentyl methyl ether, 2-Me-THF: 2-methyltetrahydrofuran, THF: tetrahydrofuran.

Then, the effect of the quantity of the applied base or its concentration was investigated. By decreasing the quantity of the 50% aq. NaOH from 50 to 25 eq, the yield slightly increased (from 73% to 85%), but the enantiomeric excess did not change. By using 1 eq of

50% aq. NaOH or 50 eq of 30% aq. NaOH, both yield and enantioselectivity declined (from 73% yield and 9% ee to 26–28% yield and <5% ee).

In the next step of the parameter study, we examined the effect of the catalyst quantity (10 or 20 mol%), the reagent quantity (1 or 1.2 eq) and the solvent volume (1.2 or 2.4 mL) (Table 3). This set of experiments indicated that the application of 10 mol% catalyst, 1.2 eq of allyl bromide and 2.4 mL DCM is the most beneficial as it gave (*R*)-3 in an increased yield (83%) with a slightly better enantiomeric excess (11%, Table 3, Entry 3).

Table 3. Examination of the effect of catalyst quantity, reagent quantity and solvent volume in the presence of catalyst C5 ^a.

Entry	Catalyst Quantity (mol%)	Allyl Bromide Quantity (eq)	Solvent Volume (mL)	Yield ^b (%)	ee ^c (%)
1	10	1	2.4	73	9
2	20	1	2.4	79	8
3	10	1.2	2.4	83	11
4	10	1	1.2	92	6

^a Reaction conditions: *tert*-butyl methyl α -benzylmalonate (1 eq), allyl bromide reagent, C5 catalyst, 50% aq. NaOH base (50 eq), DCM solvent, 25 °C, 24 h reaction time. ^b Yield of isolated product purified by preparative TLC. ^c Determined by chiral HPLC. ee: enantiomeric excess.

The reaction was also carried out using allyl iodide instead of allyl bromide as a reagent under the same conditions as at the beginning of the study (1 eq reagent, 10 mol% C5 catalyst, 50 eq 50% aq. NaOH base, DCM solvent, 25 °C, 24 h). In this case, a better yield (98% instead of 73%) and a slightly higher, but still low, enantiomeric excess (12% instead of 9%) were obtained. Consequently, we continued the study by applying allyl iodide.

Next, the reaction was carried out at 0 °C as an endeavor to improve enantioselectivity. As increasing the quantity of the reagent to 1.2 eq and decreasing the quantity of the base (50% aq. NaOH) to 25 eq seemed to be able to ameliorate yield and enantiomeric excess, we also varied these conditions in the following experiments (Table 4). By setting a lower reaction temperature, the enantiomeric excess did not improve significantly in any case. Based on these experiments, the yield was found to deteriorate when only 25 eq base is applied, while it does not depend on the temperature and the reagent quantity in the applied ranges. We also conducted the reaction at −78 °C to examine whether the enantioselectivity could be improved by setting a drastically lower reaction temperature, but no product (3) was formed based on TLC analysis.

Table 4. Examination of the effect of temperature and quantities of reagent and base in the presence of catalyst C5 ^a.


Entry	Temperature (°C)	Reagent Quantity (eq)	Base Quantity (eq)	Yield ^b (%)	ee ^c (%)
1	25	1	50	98	12
2	0	1	50	94	15
3	0	1	25	79	15
4	0	1.2	50	97	15
5	0	1.2	25	74	15

^a Reaction conditions: *tert*-butyl methyl α -benzylmalonate (1 eq), allyl iodide reagent, C5 catalyst (10 mol%), 50% aq. NaOH base, 2.4 mL DCM solvent, 24 h reaction time. ^b Yield of isolated product purified by preparative TLC. ^c Determined by chiral HPLC. ee: enantiomeric excess.

Furthermore, the use of other bases did not improve the outcome: with 2 eq Na₂CO₃ in THF or MeCN solvent, there was no product at 0 °C, and by applying 50% aq. NaOH base in the cases of Q6 and C6, and 50% aq. KOH in the cases of Q5, HQ5, C5 and G5 catalysts (to examine the effect of both bases with each catalyst), enantioselective alkylation occurred only with C5 catalyst at 0 °C (17% ee, but only 47% yield, with 50% aq. KOH).

As another attempt to increase enantioselectivity, we also investigated the activity and enantioselectivity of catalyst C5 in Michael addition reaction of *tert*-butyl methyl


α -benzylmalonate (**1**) to benzyl acrylate (Scheme 3). We anticipated higher enantioselectivity than in the alkylation reaction as benzyl acrylate has a hydrogen bond acceptor carbonyl oxygen, which could form stronger interactions with the hydrogen bond donor groups of the catalyst as opposed to the allyl halide reagents in the alkylation reaction. However, there was practically no enantioselectivity (2% ee) in this reaction, although product **5** was obtained in a good yield (74%).



Scheme 3. Synthesis of α,α -disubstituted malonate **5** by Michael addition of *tert*-butyl methyl α -benzylmalonate (**1**) to benzyl acrylate (**4**) (Reaction conditions: *tert*-butyl methyl α -benzylmalonate (1 eq), benzyl acrylate reagent (1.5 eq), **C5** catalyst (10 mol%), 50% aq. NaOH base (6.5 eq), DCM solvent (4 mL), 0 °C, 24 h reaction time).

2.3. Proposed Mechanism

Based on our previous research, we propose a mechanism for the α -alkylation reaction of *tert*-butyl methyl α -benzylmalonate (**1**) in the presence of our crown ether-squaramide catalysts. According to our preceding study, the Na^+ ion of multiple monosaccharide-based aza-crown ethers' Na^+ complexes can be coordinated by a substrate's carbonyl group [37]. In our other work, we have found that it is possible that only one NH group of a cinchona-based organocatalyst's squaramide unit forms a hydrogen bond with a substrate's carbonyl group [29]. Extending these results to our catalysts, we suggest that after the deprotonation of malonate **1** by the hydroxide ion, the phase-transfer catalyst coordinates the malonate anion in a way that one carbonyl of the malonate binds to the Na^+ ion and the other to the NH group of the squaramide (Scheme 4). Then, the nucleophilic α -carbon of the malonate attacks the allyl halide, either directly on the electrophilic saturated carbon or in a conjugate addition on the unsaturated carbon, resulting in allylic rearrangement. Naturally, in the case of the applied unsubstituted allyl halides, the product is the same in both cases. Thus, the halide anion leaves, and an allyl group is built into the substrate.



Scheme 4. Proposed transition state for the α -alkylation reaction of *tert*-butyl methyl α -benzylmalonate (**1**) with catalyst **C5**. R^1 , R^2 = *tert*-butyl or methyl.

3. Materials and Methods

3.1. General

Starting materials were purchased from commercially available sources (Sigma-Aldrich, Merck, and Alfa Aesar). Infrared spectra were recorded on a Bruker Alpha-T FT-IR spectrometer (Bruker, Ettlingen, Germany). Optical rotations were measured on a Perkin-Elmer 241 polarimeter (Perkin-Elmer, Waltham, MA, USA) that was calibrated by measuring the optical rotations of both enantiomers of menthol. Silica gel 60 F₂₅₄ (Merck) plates were used for TLC. Silica gel 60 (70–230 mesh, Merck) was used for column chromatography. Ratios of solvents for the eluents are given in volumes (mL mL⁻¹). Melting points were taken on a Boetius micro-melting point apparatus (VEB Dresden Analytik, Dresden, Germany), and they were uncorrected.

NMR spectra were recorded on a Bruker Avance 500 MHz spectrometer (Bruker, Ettlingen, Germany), (500.13 MHz for ¹H, 125.76 MHz for ¹³C and 50.68 MHz for ¹⁵N) in DMSO-d₆ and were referenced to residual solvent proton signals ($\delta_{\text{H}} = 2.50$) and solvent carbon signals ($\delta_{\text{C}} = 39.51$). The approximate ¹⁵N chemical shifts were detected by ¹H,¹⁵N-gs-HMBC method optimized for $J(\text{N,H}) = 5$ Hz long-range couplings, and ¹⁵N chemical shifts are given on NH₃ scale. The pulse programs of one-dimensional (¹H, ¹³C and DEPTQ) and two-dimensional (¹H,¹H-gs-COSY, ¹H,¹³C-gs-HSQC, ¹H,¹³C-gs-HMQC, ¹H,¹³C-gs-HMBC, ¹H,¹⁵N-gs-HMBC and ¹H,¹H-gs-ROESY) measurements were utilized. All chemical shifts are reported in parts per million (ppm). Abbreviations used in the description of resonances are: a (methylene hydrogen with higher ¹H chemical shift), b (methylene hydrogen with lower ¹H chemical shift), o (overlapping signal), s (singlet), d (doublet), t (triplet), q (quartet), br (broad), m (multiplet) and nd: not detectable. Coupling constants (J) are given in Hz.

The broad signals, the non-integer integral values in the ¹H NMR spectra (seen in Supplementary Materials) of compounds and the duplication of some signals in the ¹H and ¹³C NMR spectra (e.g., ¹H and ¹³C NMR signals of *N*-methylene groups in aza-crown ether units) of compounds indicate inhibited motions in the molecule. Also, the conformational properties of the molecules resulted in many cases in which not all signals appeared in the NMR spectra due to signal broadening. In these cases, the approximate chemical shift of the above-mentioned signals could be determined based on their HSQC or HMBC correlations. Previous ¹H NMR measurements of similar compounds at higher temperatures showed that not all conformational motion inhibitions were removed.

In the process of structure elucidation, we assigned the ¹H and ¹³C NMR signals of the main conformer with the exception of compound **G5**, in which the chemical shifts of several minor signals were also determined.

The starting points of signal assignment were easily identifiable units of molecules: the methyl and methoxy groups, the terminal olefinic methylene unit, the position 2' in the quinoline ring and the methylidene groups next to nitrogen. The remainder of the molecular structures was elucidated using the same NMR methods mentioned above.

HPLC-MS was performed using a Shimadzu LCMS-2020 (Shimadzu Corp., Kyoto, Japan) device, equipped with a Reprospher (Altmann Analytik Corp., München, Germany) 100 C18 (5 μm ; 100 \times 3 mm) column and a positive/negative double ion source (DUIS \pm) with a quadrupole MS analyzer in a range of 50–1000 m/z . The samples were eluted with gradient elution, using eluent A (0.1% HCOOH in H₂O) and eluent B (0.1% HCOOH in MeCN). The flow rate was set to 1.5 mL min⁻¹. The initial condition was 5% eluent B, followed by a linear gradient to 100% eluent B by 1.5 min; from 1.5 to 4.0 min, 100% eluent B was retained; and from 4 to 4.5 min, it went back by a linear gradient to 5% eluent B, which was retained from 4.5 to 5 min. The column temperature was kept at room temperature, and the injection volume was 1 μL . The purity of the compounds was assessed by HPLC with UV detection at 215 and 254 nm. High-resolution mass measurements were performed on a Q-TOF Premier mass spectrometer (Waters Corp., Milford, MA, USA). The ionization method was electrospray ionization (ESI) operated in positive ion mode. The enantiomeric excess (ee) values were determined by chiral HPLC

with a PerkinElmer Series 200 instrument (PerkinElmer, Waltham, MA, USA). The applied column, eluent, flow rate, column temperature, and detector wavelength are indicated at the corresponding procedure.

3.2. Procedures

3.2.1. 3-(+)-Methoxy-4-(((R)-quinolin-4-yl((1S,2R,4S,5R)-5-vinylquinuclidin-2-yl)methyl)amino)cyclobut-3-ene-1,2-dione (**C-HSQ**)

A solution of **C-N** (535 mg, 1.55 mmol) in DCM (2 mL) was added to a solution of dimethyl squarate (**DMSQ**) (242 mg, 1.70 mmol) in DCM (2 mL) under argon atmosphere. This mixture was stirred for 3 h at room temperature. The solvent was removed under reduced pressure, then the crude product was purified by column chromatography (SiO₂; DCM:methanol = 10:1) to obtain **C-HSQ** as pale-yellow crystals (494 mg, 79% yield). TLC (SiO₂ TLC; DCM:methanol = 5:1, *R_f* = 0.62). M.p. 122 °C (decomposed). $[\alpha]_D^{25}$: +220.0 (*c* = 1.0, chloroform).

IR (KBr): 3233, 3075, 2938, 2869, 1803, 1706, 1606, 1510, 1460, 1393, 1375, 1263, 1240, 1174, 1079, 1053, 927, 851, 826, 768 cm⁻¹.

¹H NMR (500.13 MHz, DMSO-d₆): δ = 9.20 (br m, 1H, NH), 8.94 (d, *J* = 3.8 Hz, 1H, H-2'), 8.35 (d, *J* = 7.9 Hz, 1H, H-5'), 8.07 (d, *J* = 8.2 Hz, 1H, H-8'), 7.80 (br m, 1H, H-7'), 7.72 (o m, 1H, H-6'), 7.71 (o m, 1H, H-3'), 6.08 (br m, 1H, H-9), 5.80 (br m, 1H, H-10), 5.13 (d, *J* = 17.1 Hz, 1H, H-11a), 5.08 (d, *J* = 10.5 Hz, 1H, H-11b), 4.25 (s, 3H, H-5''), 3.28 (br m, 1H, H-8), 2.99 (o m, 1H, H-2a), 2.90 (o m, 1H, H-6a), 2.84 (o m, 1H, H-6b), 2.80 (o m, 1H, H-2b), 2.21 (br m, 1H, H-3), 1.51 (o m, 1H, H-4), 1.49 (o m, 2H, H-5ab), 0.91 (o m, 1H, H-7a), 0.84 (o m, 1H, H-7b).

¹³C NMR (125.76 MHz, DMSO-d₆): δ = 189.8 (C-1'' or C-2''), 181.6 (C-1'' or C-2''), 177.5 (C-3''), 171.2 (C-4''), 150.4 (C-2'), 148.0 (C-8a'), 144.7 (C-4'), 140.6 (C-10), 130.1 (C-8'), 129.6 (C-7'), 127.3 (C-6'), 126.3 (C-4a'), 122.9 (C-5'), 119.8 (C-3'), 114.6 (C-11), 60.1 (C-5''), 59.2 (C-8), 52.2 (C-9), 48.8 (C-6), 45.9 (C-2), 38.6 (C-3), 27.3 (C-4), 26.0 (C-5), 24.7 (C-7).

HRMS (ESI⁺): *m/z* [M + H]⁺ calcd for C₂₄H₂₅N₃O₃: 404.1974; found: 404.1974.

To the best of our knowledge, the synthesis of **C-HSQ** has not been reported so far.

3.2.2. General Procedure for the Preparation of Cinchona Alkaloid-Based Catalysts

To a solution of cinchona half squaramide (**Q-HSQ** or **HQ-HSQ** or **C-HSQ**, 0.482 mmol) in methanol (2.1 mL), a solution of aza-crown ether (1-aza-15-crown-5 ether or 1-aza-18-crown-6 ether, 0.439 mmol) in methanol (2.1 mL) was added under argon atmosphere. The resulting mixture was warmed up to 60 °C, then it was stirred for 6 h at this temperature. The solvent was removed under reduced pressure, then the crude product was purified by column chromatography (SiO₂; DCM:methanol = 10:1) to obtain **Q5** or **Q6** or **HQ5** or **C5** or **C6**.

3.2.3. 3-(+)-(1,4,7,10-Tetraoxa-13-azacyclopentadecan-13-yl)-4-(((S)-(6-methoxyquinolin-4-yl)((1S,2S,4S,5R)-5-vinylquinuclidin-2-yl)methyl)amino)cyclobut-3-ene-1,2-dione (**Q5**)

Yellowish white crystals (93 mg, 34% yield). TLC (SiO₂ TLC; DCM:methanol = 10:1, *R_f* = 0.46). M.p. 97–100 °C. $[\alpha]_D^{25}$: +53.1 (*c* = 1.0, chloroform).

IR (KBr): 3323, 2934, 2863, 1786, 1664, 1621, 1516, 1471, 1432, 1354, 1313, 1298, 1245, 1230, 1176, 1124, 1082, 1057, 1028, 851 cm⁻¹.

¹H NMR (500.13 MHz, DMSO-d₆): δ = 8.77 (d, *J* = 4.3 Hz, 1H, H-2'), 7.95 (d, *J* = 9.2 Hz, 1H, H-8'), 7.89 (br m, 1H, H-5'), 7.69 (d, *J* = 4.3 Hz, 1H, H-3'), 7.64 (br m, 1H, NH), 6.26 (br m, 1H, H-9), 7.42 (dd, *J* = 9.2, and 2.3 Hz, 1H, H-7'), 5.93 (ddd, *J* = 17.2, 10.3, and 7.6 Hz, 1H, H-10), 5.04 (d, *J* = 17.2 Hz, 1H, H-11a), 4.99 (d, *J* = 10.3 Hz, 1H, H-11b), 3.98 (br m, 1H, H-2'''a or H-11'''a), 3.93 (s, 3H, H-9'), 3.65 (br m, 2H, H-2''' and/or H-11'''), 3.00–3.60 (o m, 16H, H-3''', H-4''', H-5''', H-6''', H-7''', H-8''', H-9''', and H-10'''), 3.47 (o m, 1H, H-2'''b or H-11'''b), 3.50 (o m, 1H, H-8), 3.33 (o m, 1H, H-6a), 3.17 (o m, 1H, H-2a), 2.72 (dm, *J* = 13.3 Hz, 1H, H-2b), 2.62 (br m, 1H, H-6b), 2.27 (br m, 1H, H-3), 1.57 (br m, 1H,

H-4), 1.49 (o m, 2H, H-5ab), 1.42 (tm, $J = 12.2$ Hz, 1H, H-7a), 0.55 (dm, $J = 13.2$, and 7.1 Hz, 1H, H-7b).

^{13}C NMR (125.76 MHz, DMSO- d_6): $\delta = 182.2$ (C-1'' or C-2''), 181.8 (C-1'' or C-2''), 168.3 (C-3'' or C-4''), 166.5 (C-3'' or C-4''), 157.8 (C-6'), 144.2 (C-4' or C-8a'), 144.0 (C-4' or C-8a'), 147.8 (C-2'), 142.2 (C-10), 131.4 (C-8'), 127.7 (C-4a'), 121.9 (C-7'), 119.9 (C-3'), 114.4 (C-11), 101.8 (C-5'), 68.5–70.5 (C-3''', C-4''', C-5''', C-6''', C-7''', C-8''', C-9''', and C-10'''), 58.6 (C-8), 55.7 (C-9'), 55.5 (C-2), 52.8 (C-9), 51.8 (C-2''' or C-13'''), 51.4 (C-2''' or C-13'''), 40.1 (C-6), 39.2 (C-3), 27.4 (C-5), 27.3 (C-4), 26.1 (C-7).

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₃₄H₄₄N₄O₇: 621.3288; found: 621.3285.

To the best of our knowledge, the synthesis of **Q5** has not been reported so far.

3.2.4. 3-(+)-(1,4,7,10,13-Pentaoxa-16-azacyclooctadecan-16-yl)-4-(((S)-(6-methoxyquinolin-4-yl)((1S,2S,4S,5R)-5-vinylquinuclidin-2-yl)methyl)amino)cyclobut-3-ene-1,2-dione (**Q6**)

Pale-brown oil (64 mg, 22% yield). TLC (SiO₂ TLC; DCM:methanol = 10:1, $R_f = 0.19$). $[\alpha]_D^{25}$: +64.5 ($c = 1.0$, chloroform).

IR (KBr): 3311, 2909, 2867, 1784, 1666, 1622, 1574, 1521, 1473, 1431, 1347, 1315, 1303, 1248, 1231, 1134, 1107, 1028, 980, 833 cm⁻¹.

^1H NMR (500.13 MHz, DMSO- d_6): $\delta = 8.78$ (d, $J = 4.3$ Hz, 1H, H-2'), 7.96 (d, $J = 9.1$ Hz, 1H, H-8'), 7.90 (br m, 1H, H-5'), 7.71 (br m, 1H, H-3'), 7.66 (br m, 1H, NH), 6.32 (br m, 1H, H-9), 7.42 (dd, $J = 9.1$, and 2.2 Hz, 1H, H-7'), 5.95 (ddd, $J = 17.3$, 10.5, and 6.7 Hz, 1H, H-10), 5.08 (d, $J = 17.3$ Hz, 1H, H-11a), 5.03 (d, $J = 10.5$ Hz, 1H, H-11b), 3.99 (br m, 1H, H-2'''a or H-11'''a), 3.98 (br m, 1H, H-8), 3.93 (s, 3H, H-9'), 3.71 (br m, 2H, H-2''' and/or H-11'''), 3.00–3.64 (o m, 18H, H-4''', H-5''', H-6''', H-7''', H-8''', H-9''', H-10''', H-11''', and H-12'''), 3.52 (o m, 1H, H-2'''b or H-11'''b), 3.38 (o m, 1H, H-6a), 3.27 (o m, 1H, H-2a), 2.78 (br m, 1H, H-2b), 2.69 (br m, 1H, H-6b), 2.34 (br m, 1H, H-3), 1.62 (o m, 1H, H-4), 1.50 (o m, 1H, H-7a), 0.60 (br m, 1H, H-7b), and (2H, H-5).

^{13}C NMR (125.76 MHz, DMSO- d_6): $\delta = 182.1$ (C-1'' or C-2''), 181.9 (C-1'' or C-2''), 168.2 (C-3'' or C-4''), 166.2 (C-3'' or C-4''), 157.9 (C-6'), 144.2 (C-8a'), 147.7 (C-2'), 142.0 (C-10), 131.4 (C-8'), 127.7 (C-4a'), 121.9 (C-7'), 120.0 (C-3'), 114.7 (C-11), 101.7 (C-5'), 68.5–70.5 (C-3''', C-4''', C-5''', C-6''', C-7''', C-8''', C-9''', C-10''', C-11''', and C-12'''), 55.7 (C-8, C-9'), 55.3 (C-2), 52.6 (C-9), 50.6 (C-2''' or C-13'''), 49.7 (C-2''' or C-13'''), 40.2 (C-6), 38.9 (C-3), 27.7 (C-4), 25.8 (C-7), and (C-5, C-4').

^{15}N NMR (50.68 MHz, DMSO- d_6): 312.3 (N-1'').

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₃₆H₄₈N₄O₈: 665.3550; found: 665.3550.

To the best of our knowledge, the synthesis of **Q6** has not been reported so far.

3.2.5. 3-(+)-(1,4,7,10-Tetraoxa-13-azacyclopentadecan-13-yl)-4-(((S)-((1S,2S,4S,5R)-5-ethylquinuclidin-2-yl)(6-methoxyquinolin-4-yl)methyl)amino)cyclobut-3-ene-1,2-dione (**HQ5**)

White powder (208 mg, 76% yield). TLC (SiO₂ TLC; DCM:methanol = 10:1, $R_f = 0.19$). M.p. 107 °C. $[\alpha]_D^{25}$: +50.8 ($c = 1.0$, chloroform).

IR (KBr): 3271, 2930, 2862, 1789, 1672, 1622, 1579, 1519, 1475, 1432, 1353, 1297, 1257, 1241, 1230, 1122, 1029, 939, 854, 831 cm⁻¹.

^1H NMR (500.13 MHz, DMSO- d_6): $\delta = 8.78$ (d, $J = 4.3$ Hz, 1H, H-2'), 7.95 (d, $J = 9.2$ Hz, 1H, H-8'), 7.89 (br, 1H, H-5'), 7.73 (br m, 1H, H-3'), 7.67 (br m, 1H, NH), 6.28 (br m, 1H, H-9), 7.42 (dd, $J = 9.2$, and 2.3 Hz, 1H, H-7'), 4.02 (br m, 1H, H-2'''a or H-11'''a), 3.93 (s, 3H, H-9'), 3.65 (br m, 2H, H-2''' and/or H-11'''), 3.10–3.62 (o m, 14H, H-4''', H-5''', H-6''', H-7''', H-8''', H-9''', and H-10'''), 3.57 (o m, 1H, H-8), 3.45 (o m, 1H, H-2'''b or H-11'''b), 3.36 (o m, 1H, H-6a), 3.20 (o m, 1H, H-2a), 2.67 (br m, 1H, H-6b), 2.51 (o m, 1H, H-2b), 1.58 (o m, 1H, H-4), 1.56 (o m, 1H, H-5a), 1.45 (o m, 1H, H-5b), 1.44 (o m, 1H, H-3), 1.41 (o m, 1H, H-7a), 1.34 (o m, 2H, H-10ab), 0.81 (t, $J = 7.0$ Hz, 3H, H-11), 0.58 (br m, 1H, H-7b).

^{13}C NMR (125.76 MHz, DMSO- d_6): $\delta = 182.2$ (C-1'' or C-2''), 181.8 (C-1'' or C-2''), 168.4 (C-3'' or C-4''), 166.4 (C-3'' or C-4''), 157.8 (C-6'), 144.2 (C-8a'), 147.7 (C-2'), 131.4 (C-8'), 127.6 (C-4a'), 121.9 (C-7'), 119.9 (C-3'), 101.7 (C-5'), 69.0–70.0 (C-3''', C-4''', C-5''', C-6''', C-7''', C-8''', C-9''', and C-10'''), 58.5 (C-8), 56.7 (C-2), 55.7 (C-9'), 52.4 (C-9), 51.8

(C-2''' or C-11'''), 51.4 (C-2''' or C-11'''), 39.8 (C-6), 36.5 (C-3), 27.6 (C-5), 27.0 (C-10), 25.5 (C-7), 24.8 (C-4), 11.9 (C-11), and (C-4').

¹⁵N NMR (50.68 MHz, DMSO-d₆): 312.1 (N-1'').

HRMS (ESI⁺): *m/z* [M + H]⁺ calcd for C₃₄H₄₆N₄O₇: 623.3445; found: 623.3450.

To the best of our knowledge, the synthesis of **HQ5** has not been reported so far.

3.2.6. 3-(+)-(1,4,7,10-Tetraoxa-13-azacyclopentadecan-13-yl)-4-(((R)-quinolin-4-yl((1S,2R,4S,5R)-5-vinylquinuclidin-2-yl)methyl)amino)cyclobut-3-ene-1,2-dione (**C5**)

White powder (122 mg, 47% yield). TLC (SiO₂ TLC; DCM:methanol = 10:1, *R_f* = 0.46). M.p. 204 °C. [α]_D²⁵: +45.2 (*c* = 1.0, chloroform).

IR (KBr): 3308, 2934, 2895, 2856, 1790, 1664, 1575, 1526, 1475, 1433, 1384, 1352, 1317, 1265, 1249, 1124, 1080, 1058, 770 cm⁻¹.

¹H NMR (500.13 MHz, DMSO-d₆): δ = 8.94 (d, *J* = 4.2 Hz, 1H, H-2'), 8.51 (d, *J* = 8.5 Hz, 1H, H-5'), 8.05 (d, *J* = 8.3 Hz, 1H, H-8'), 7.79 (o m, 1H, H-3'), 7.78 (o m, 1H, H-7'), 7.67 (o m, 1H, NH), 7.70 (o m, 1H, H-6'), 6.36 (br m, 1H, H-9), 5.81 (ddd, *J* = 17.3, 10.5, and 6.7 Hz, 1H, H-10), 5.17 (d, *J* = 17.3 Hz, 1H, H-11a), 5.08 (d, *J* = 10.5 Hz, 1H, H-11b), 3.93 (br m, 1H, H-2'''a or H-11'''a), 3.67 (br m, 2H, H-2''' and/or H-11'''), 3.26–3.62 (o m, 14H, H-4''' H-5''' H-6''' H-7''' H-8''' H-9''' and H-10'''), 3.58 (o m, 4H, H-3''' and H-10'''), 3.52 (br m, 1H, H-2'''b or H-11'''b), 3.40 (o m, 1H, H-8), 3.14 (br m, 1H, H-2a), 2.96 (br m, 1H, H-6a), 2.88 (o m, 1H, H-6b), 2.82 (br m, 1H, H-2b), 2.23 (br m, 1H, H-3), 1.52 (o m, 1H, H-4), 1.51 (o m, 2H, H-5ab), 0.95 (br m, 1H, H-7a), 0.88 (br m, 1H, H-7b).

¹³C NMR (125.76 MHz, DMSO-d₆): δ = 182.2 (C-1'' or C-2''), 182.1 (C-1'' or C-2''), 168.2 (C-3'' or C-4''), 166.8 (C-3'' or C-4''), 150.3 (C-2'), 148.0 (C-8a'), 145.7 (C-4'), 140.5 (C-10), 129.9 (C-8'), 129.4 (C-7'), 126.9 (C-6'), 126.6 (C-4a'), 123.5 (C-5'), 119.7 (C-3'), 114.6 (C-11), 69.0–70.0 (C-3''', C-4''', C-5''', C-6''', C-7''', C-8''', C-9''', and C-10'''), 59.0 (C-8), 51.8 (C-9 and C-2''' or C-11'''), 51.4 (C-2''' or C-11'''), 48.8 (C-6), 45.8 (C-2), 38.6 (C-3), 27.3 (C-4), 25.9 (C-5), 24.9 (C-7).

¹⁵N NMR (50.68 MHz, DMSO-d₆): 312.4 (N-1''), 105.8 (9-NH, ¹*J* = 95 Hz).

HRMS (ESI⁺): *m/z* [M + H]⁺ calcd for C₃₃H₄₂N₄O₆: 591.3183; found: 591.3179.

To the best of our knowledge, the synthesis of **C5** has not been reported so far.

3.2.7. 3-(+)-(1,4,7,10,13-Pentaoxa-16-azacyclooctadecan-16-yl)-4-(((R)-quinolin-4-yl((1S,2R,4S,5R)-5-vinylquinuclidin-2-yl)methyl)amino)cyclobut-3-ene-1,2-dione (**C6**)

Pale-brown oil (137 mg, 49% yield). TLC (SiO₂ TLC; DCM:methanol = 5:1, *R_f* = 0.62). [α]_D²⁵: +25.2 (*c* = 1.0, chloroform).

IR (KBr): 3275, 2935, 2867, 1789, 1671, 1637, 1577, 1520, 1477, 1431, 1350, 1297, 1259, 1204, 1116, 987, 916, 852, 828, 771 cm⁻¹.

¹H NMR (500.13 MHz, DMSO-d₆): δ = 8.94 (d, *J* = 4.0 Hz, 1H, H-2'), 8.52 (d, *J* = 8.3 Hz, 1H, H-5'), 8.06 (d, *J* = 8.3 Hz, 1H, H-8'), 7.80 (br m, 1H, H-3'), 7.78 (t, *J* = 7.7 Hz, 1H, H-7'), 7.68 (t, *J* = 7.4 Hz, 1H, H-6'), 7.65 (o m, 1H, NH), 6.41 (br m, 1H, H-9), 5.80 (ddd, *J* = 17.3, 10.4, and 6.3 Hz, 1H, H-10), 5.18 (d, *J* = 17.3 Hz, 1H, H-11a), 5.08 (d, *J* = 10.4 Hz, 1H, H-11b), 3.96 (br m, 1H, H-2'''a or H-11'''a), 3.71 (br m, 2H, H-2''' and/or H-11'''), 3.25–3.64 (o m, 18H, H-4''' H-5''' H-6''' H-7''' H-8''' H-9''' H-10''' H-11''' and H-12'''), 3.57 (o m, 1H, H-2'''b or H-11'''b), 3.47 (br m, 1H, H-8), 3.17 (o m, 1H, H-2a), 2.98 (o m, 1H, H-6a), 2.91 (o m, 1H, H-6b), 2.83 (o m, 1H, H-2b), 2.26 (br m, 1H, H-3), 1.56 (o m, 3H, H-4, H-5ab), 1.01 (br m, 1H, H-7a), 0.89 (br m, 1H, H-7b).

¹³C NMR (125.76 MHz, DMSO-d₆): δ = 182.13 (C-1'' or C-2''), 182.11 (C-1'' or C-2''), 168.0 (C-3'' or C-4''), 166.6 (C-3'' or C-4''), 150.4 (C-2'), 145.5 (C-4'), 148.0 (C-8a'), 140.4 (C-10), 129.9 (C-8'), 129.5 (C-7'), 127.1 (C-6'), 126.6 (C-4a'), 123.6 (C-5'), 119.9 (C-3'), 114.7 (C-11), 68.5–71.0 (C-3''', C-4''', C-5''', C-6''', C-7''', C-8''', C-9''', C-10''', C-11''', and C-12'''), 58.9 (C-8), 52.0 (C-9), 51.6 (C-2''' or C-13'''), 49.7 (C-2''' or C-13'''), 48.8 (C-6), 45.8 (C-2), 38.6 (C-3), 27.2 (C-4), 25.8 (C-5), 25.0 (C-7).

HRMS (ESI⁺): *m/z* [M + H]⁺ calcd for C₃₅H₄₆N₄O₇: 635.3445; found: 635.3459.

To the best of our knowledge, the synthesis of **C6** has not been reported so far.

3.2.8. Procedure for the Preparation of the Glucose-Based Catalyst:

3-(+)-((3,5-bis(trifluoromethyl)phenyl)amino)-4-((2R,4aR,6S,6aR,19aS,19bR)-6-methoxy-2-phenyltetradecahydro-13H-[1,3]dioxino[4',5':5,6]pyrano[3,4-e][1,4,7,10]tetraoxa[13]azacyclopentadecin-13-yl)cyclobut-3-ene-1,2-dione (**G5**)

To a solution of **HSQ** (119 mg, 0.350 mmol) in methanol (2.5 mL), a solution of crown ether derivative **G** (140 mg, 0.319 mmol) in methanol (2.5 mL) was added under argon atmosphere. The resulting mixture was warmed up to 60 °C, then stirred for 6 h at this temperature. The solvent was removed under reduced pressure, then the crude product was purified by column chromatography (SiO₂; DCM:methanol = 40:1) to obtain **G5** as a yellowish-white powder (131 mg, 55% yield). TLC (SiO₂ TLC; DCM:methanol = 40:1, $R_f = 0.19$). M.p. 209–211 °C. $[\alpha]_D^{25}$: +32.5 ($c = 1.0$, chloroform).

IR (KBr): 3262, 3080, 2931, 2868, 1785, 1687, 1593, 1562, 1475, 1458, 1427, 1383, 1278, 1174, 1093, 1059, 1018, 992, 885, 766 cm⁻¹.

¹H NMR (500.13 MHz, DMSO-d₆):

Major signals: $\delta = 9.55$ (br s, 0.6H, NH), 7.92 (s, 2H, H-2''' and H-6'''), 7.56 (br m, 0.55H, H-4'''), 7.41 (o m, 2H, H-2'' and H-6''), 7.37 (o m, 2H, H-3'' and H-5''), 7.36 (o m, 1H, H-4''), 5.61 (s, 0.65H, H-2), 4.92 (br m, 0.55H, H-6), 4.16 (o m, 2H, H-4ab), 3.30–3.90 (o m, 12H, H-8, H-9, H-10, H-15, H-17 and H-18), 3.55 (o m, 2H, H-4''' and H-5'''), 3.52 (o m, 1H, H-19a), 3.35 (o m, 1H, H-6a), 3.31 (o s, 3H, CH₃).

Minor signals: $\delta = 7.65$ (br m, 0.45H, H-4'''), 7.38 (o m, H-3'' and H-5''), 7.35 (o m, 1H, H-4''), 7.32 (o m, H-3'' and H-5''), 7.27 (o m, H-2'' and H-6''), 5.45 (s, 0.5H, H-2), 3.24 (s, 1.7H, CH₃), 3.19 (s, 0.5H, CH₃)

¹³C NMR (125.76 MHz, DMSO-d₆):

Major signals: $\delta = 185.7$ (C-1' or C-2'), 180.9 (C-1' or C-2'), 171.1 (C-3' or C-4'), 162.4 (C-3' or C-4'), 141.0 (C-1'''), 137.6 (C-1''), 130.8 (q, $J = 32.8$ Hz, C-3''' and C-5'''), 128.8 (C-4''), 128.0 (C-3'' and C-5''), 126.1 (C-2'' and C-6''), 123.3 (q, $J = 272.5$ Hz, CF₃), 119.1 (C-2''' and C-6'''), 114.6 (C-4'''), 100.5 (C-2), 97.3 (C-6), 81.1 (C-19b), 78.7 (C-6a), 77.8 (C-19a), 68.3–72.0 (C-8, C-9, C-11, C-15, C-17 and C-18), 68.1 (C-4), 62.0 (C-4a), 54.6 (CH₃), 52.0 (C-2'' or C-11''), 51.7 (C-12 or C-14).

Minor signals: $\delta = 137.8$ (C-1''), 137.7 (C-1''), 128.7 (C-4''), 128.15 (C-3'' and C-5''), 128.12 (C-3'' and C-5''), 125.9 (C-2'' and C-6''), 119.1 (C-2''' and C-6'''), 114.6 (C-4'''), 100.5 (C-2), 98.4 (C-6), 97.8 (C-6), 80.8 (C-19b), 78.8 (C-6a), 62.2 (C-4a), 58.1 (CH₃), 58.0 (CH₃), 54.7 (CH₃).

HRMS (ESI⁺): m/z [M + H]⁺ calcd for C₃₄H₃₆F₆N₂O₁₀: 747.2352; found: 747.2355.

To the best of our knowledge, the synthesis of **G5** has not been reported so far.

3.2.9. General Procedure for the α -alkylation of tert-butyl methyl α -benzylmalonate with allyl halides (**3**)

To a solution of *tert*-butyl methyl α -benzylmalonate (**1**, 26 mg, 0.1 mmol) in the indicated solvent, catalyst **Q5** or **Q6** or **HQ5** or **C5** or **C6** or **G5**, and a base were added (exact reaction conditions are shown in Tables 1–4). Then, the allyl halide (**2**, allyl bromide or allyl iodide) was added, and the resulting mixture was stirred vigorously at the temperature shown in Tables 1–4. After 24 h, water (1 mL) was added to the reaction mixture, and it was extracted with DCM (2 × 1 mL). The combined organic phase was dried over MgSO₄ and filtered (except in the cases of MeCN and THF solvents, when the mixture was only dried over MgSO₄, then filtered). The volatile components were removed under reduced pressure. The residue was purified by preparative TLC (SiO₂; hexane:EtOAc = 4:1) to obtain product **3** as a colorless oil. Yields and enantiomeric excess (ee) values can be seen in Tables 1–4. Spectroscopic data are fully consistent with those reported in the literature [3]. TLC (SiO₂ TLC; hexane:EtOAc = 10:1, $R_f = 0.47$). $[\alpha]_D^{25}$: -1.7 ($c = 1.0$, chloroform, 15% ee, (*R*) abs. config.) (lit. $[\alpha]_D^{25}$: -7.6 ($c = 1.0$, chloroform, 94% ee, (*R*) abs. config.). The enantiomeric excess values were determined by chiral HPLC using a Phenomenex Lux[®] 5- μ m Cellulose-2 column (250 × 4.6 mm ID), and a 95:5 mixture of hexane/ethanol was

used as the eluent with a flow rate of 0.8 mL min⁻¹. The column temperature was 20 °C. UV detector λ = 254 nm. Retention time for (S)-3: 5.0 min, for (R)-3: 5.4 min.

MS (ESI⁺): m/z (%) = 249 (100) [M-*t*-Bu + 2H]⁺.

3.2.10. General Procedure for the Michael Addition of *tert*-butyl methyl α -benzylmalonate to benzyl acrylate (5)

To a solution of *tert*-butyl methyl α -benzylmalonate (**1**, 26 mg, 0.1 mmol) in DCM (4 mL), catalyst **C5** (6 mg, 0.01 mmol) and aq. 50% NaOH (35 μ L, 0.65 mmol) were added. Then, benzyl acrylate (**4**, 23 μ L, 24 mg, 0.15 mmol) was added, and the resulting mixture was stirred vigorously at 0 °C. After 24 h, water (1 mL) was added to the reaction mixture, then it was extracted with DCM (2 \times 1 mL), dried over MgSO₄, and filtered. The volatile components were removed under reduced pressure. The residue was purified by preparative TLC (SiO₂; hexane:EtOAc = 10:1) to obtain product **5** (32 mg, 74% yield, 2% ee) as a colorless oil. Spectroscopic data are fully consistent with those reported in the literature [2]. TLC (SiO₂ TLC; hexane:EtOAc = 10:1, R_f = 0.23). In this case, the enantiomeric excess values were determined by chiral HPLC using a Kromasil[®] 5- μ m AmyCoat[®] column (250 \times 4.6 mm ID), an 85:15 mixture of hexane/ethanol was used as the eluent with a flow rate of 0.8 mL min⁻¹. The column temperature was 20 °C. UV detector λ = 254 nm. Retention times: 6.8 min and 8.7 min.

MS (ESI⁺): m/z (%) = 371 (100) [M-*t*-Bu + 2H]⁺.

4. Conclusions

In conclusion, we have described the synthesis of six new crown ether-squaramide phase-transfer organocatalysts bearing four different chiral units. To the best of our knowledge, this is the first successful direct coupling of squaramide and aza-crown-ether units and the first application of crown ether-squaramides as phase-transfer catalysts. We have tested their performance in the asymmetric α -alkylation of *tert*-butyl methyl α -benzylmalonate with extensive parameter investigation, after which reaction of the products could be converted to α,α -disubstituted amino acids through Curtius rearrangement. The alkylation reactions afforded the products in excellent yields, but with only low enantiomeric excess values. Thus, despite the low enantioselectivity, the new crown ether-based catalysts can catalyze the often-difficult construction of quaternary carbon centers. With the catalyst having the best enantioselectivity, a Michael addition reaction of *tert*-butyl methyl α -benzylmalonate was also conducted; however, no enantioselectivity was observed. Based on our results, we anticipate that a linker between the crown ether and the squaramide unit may potentially increase the enantioselectivity of this new type of phase-transfer catalyst, as two NH groups may form stronger hydrogen bonds during catalysis.

Supplementary Materials: The following are available online: NMR spectra of the new compounds and HPLC chromatograms of products **3** and **5**.

Author Contributions: Conceptualization, Z.F., S.N. and J.K.; methodology, Z.F., S.N. and J.K.; synthesis of compounds, Z.F., D.R. and Z.R.; performing NMR experiments, data analysis, A.S.; determination of ee values by chiral HPLC measurements, P.B. (Péter Bagi); performing HRMS experiments, data analysis, L.D.; writing—original draft preparation, Z.F.; writing—review and editing, J.K., P.H., P.B. (Péter Bakó), P.B. (Péter Bagi), Z.F. and Z.R.; project administration, J.K.; funding acquisition, P.H. and J.K.; resources, P.H. and J.K.; supervision, J.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the New National Excellence Program of the Ministry of Human Capacities, grant numbers ÚNKP-20-3-II-BME-325 (S.N.), ÚNKP-21-2-I-BME-209 (D.R.) and ÚNKP-20-5-BME-322 (J.K.), and the János Bolyai Research Scholarship of the Hungarian Academy of Science (J.K.). It was also supported by the National Research, Development, and Innovation Office (grant numbers FK138037 and K128473), the Servier–Beregi PhD Research Fellowship (S.N.), and the Gedeon Richter's Talentum Foundation (Z.F.).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors are grateful to Hajnalka Szabó-Szentjóni from Budapest University of Technology and Economics for her assistance with the synthesis of crown ether derivatives, and Róbert Ritter from Budapest University of Technology and Economics for his assistance with the NMR measurements.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the compounds are available from the authors.

References

1. Hong, S.; Lee, J.; Kim, M.; Park, Y.; Park, C.; Kim, M.; Jew, S.; Park, H. Highly Enantioselective Synthesis of α,α -Dialkylmalonates by Phase-Transfer Catalytic Desymmetrization. *J. Am. Chem. Soc.* **2011**, *133*, 4924–4929. [[CrossRef](#)]
2. Odanaka, Y.; Kanemitsu, T.; Iwasaki, K.; Mochizuki, Y.; Miyazaki, M.; Nagata, K.; Kato, M.; Itoh, T. Asymmetric Michael addition of malonic diesters to acrylates by phase-transfer catalysis toward the construction of quaternary stereogenic α -carbons. *Tetrahedron* **2019**, *75*, 209–219. [[CrossRef](#)]
3. Kanemitsu, T.; Koga, S.; Nagano, D.; Miyazaki, M.; Nagata, K.; Itoh, T. Asymmetric Alkylation of Malonic Diester Under Phase-Transfer Conditions. *ACS Catal.* **2011**, *1*, 1331–1335. [[CrossRef](#)]
4. Ha, M.W.; Lee, H.; Yi, H.Y.; Park, Y.; Kim, S.; Hong, S.; Lee, M.; Kim, M.; Kim, T.; Park, H. Enantioselective Phase-Transfer Catalytic α -Benzoylation and α -Allylation of α -tert-Butoxycarbonyllactones. *Adv. Synth. Catal.* **2013**, *355*, 637–642. [[CrossRef](#)]
5. Ha, M.W.; Hong, S.; Park, C.; Park, Y.; Lee, J.; Kim, M.; Lee, J.; Park, H. Enantioselective phase-transfer catalytic α -alkylation of 2-methylbenzyl tert-butyl malonates. *Org. Biomol. Chem.* **2013**, *11*, 4030–4039. [[CrossRef](#)] [[PubMed](#)]
6. Park, C.; Ha, M.W.; Kim, B.; Hong, S.; Kim, D.; Park, Y.; Kim, M.; Lee, J.K.; Lee, J.; Park, H. Enantioselective α -Alkylation of Benzylideneamino tert-Butyl Malonates by Phase-Transfer Catalysis. *Adv. Synth. Catal.* **2015**, *357*, 2841–2848. [[CrossRef](#)]
7. Kanemitsu, T.; Sato, M.; Yoshida, M.; Ozasa, E.; Miyazaki, M.; Odanaka, Y.; Nagata, K.; Itoh, T. Enantioselective α -Benzoyloxylation of Malonic Diesters by Phase-Transfer Catalysis. *Org. Lett.* **2016**, *18*, 5484–5487. [[CrossRef](#)] [[PubMed](#)]
8. Tan, J.; Yasuda, N. Contemporary Asymmetric Phase Transfer Catalysis: Large-Scale Industrial Applications. *Org. Process Res. Dev.* **2015**, *19*, 1731–1746. [[CrossRef](#)]
9. Hashimoto, T.; Maruoka, K. Recent Development and Application of Chiral Phase-Transfer Catalysts. *Chem. Rev.* **2007**, *107*, 5656–5682. [[CrossRef](#)] [[PubMed](#)]
10. Shirakawa, S.; Maruoka, K. Recent Developments in Asymmetric Phase-Transfer Reactions. *Angew. Chem. Int. Ed.* **2013**, *52*, 4312–4348. [[CrossRef](#)]
11. Majdecki, M.; Niedbala, P.; Tyszcza-Gumkowska, A.; Jurczak, J. Assisted by Hydrogen-Bond Donors: Cinchona Quaternary Salts as Privileged Chiral Catalysts for Phase-Transfer Reactions. *Synthesis* **2021**, *53*, 2777–2786. [[CrossRef](#)]
12. Brak, K.; Jacobsen, E.N. Asymmetric Ion-Pairing Catalysis. *Angew. Chem. Int. Ed.* **2013**, *52*, 534–561. [[CrossRef](#)]
13. Schettini, R.; Sicignano, M.; De Riccardis, F.; Izzo, I.; Della Sala, G. Macrocyclic Hosts in Asymmetric Phase-Transfer Catalyzed Reactions. *Synthesis* **2018**, *50*, 4777–4795. [[CrossRef](#)]
14. Wang, H. Chiral Phase-Transfer Catalysts with Hydrogen Bond: A Powerful Tool in the Asymmetric Synthesis. *Catalysts* **2019**, *9*, 244. [[CrossRef](#)]
15. Majdecki, M.; Niedbala, P.; Jurczak, J. Amide-Based Cinchona Alkaloids as Phase-Transfer Catalysts: Synthesis and Potential Application. *Org. Lett.* **2019**, *21*, 8085–8090. [[CrossRef](#)] [[PubMed](#)]
16. Majdecki, M.; Grodek, P.; Jurczak, J. Stereoselective α -Chlorination of β -Keto Esters in the Presence of Hybrid Amide-Based Cinchona Alkaloids as Catalysts. *J. Org. Chem.* **2021**, *86*, 995–1001. [[CrossRef](#)]
17. Wang, B.; Liu, Y.; Sun, C.; Wei, Z.; Cao, J.; Liang, D.; Lin, Y.; Duan, H. Asymmetric Phase-Transfer Catalysts Bearing Multiple Hydrogen-Bonding Donors: Highly Efficient Catalysts for Enantio- and Diastereoselective Nitro-Mannich Reaction of Amidosulfones. *Org. Lett.* **2014**, *16*, 6432–6435. [[CrossRef](#)] [[PubMed](#)]
18. Craig, R.; Litvajova, M.; Cronin, S.A.; Connon, S.J. Enantioselective acyl-transfer catalysis by fluoride ions. *Chem. Commun.* **2018**, *54*, 10108–10111. [[CrossRef](#)]
19. Lu, N.; Fang, Y.; Gao, Y.; Wei, Z.; Cao, J.; Liang, D.; Lin, Y.; Duan, H. Bifunctional Thiourea–Ammonium Salt Catalysts Derived from Cinchona Alkaloids: Cooperative Phase-Transfer Catalysts in the Enantioselective Aza-Henry Reaction of Ketimines. *J. Org. Chem.* **2018**, *83*, 1486–1492. [[CrossRef](#)]
20. Cui, D.; Li, Y.; Huang, P.; Tian, Z.; Jia, Y.; Wang, P. Bifunctional phase-transfer catalysts for synthesis of 2-oxazolidinones from isocyanates and epoxides. *RSC Adv.* **2020**, *10*, 12360–12364. [[CrossRef](#)]
21. Alemán, J.; Parra, A.; Jiang, H.; Jørgensen, K.A. Squaramides: Bridging from Molecular Recognition to Bifunctional Organocatalysis. *Chem. Eur. J.* **2011**, *17*, 6890–6899. [[CrossRef](#)]
22. Frontera, A.; Orell, M.; Garau, C.; Quiñonero, D.; Molins, E.; Mata, I.; Morey, J. Preparation, Solid-State Characterization, and Computational Study of a Crown Ether Attached to a Squaramide. *Org. Lett.* **2005**, *7*, 1437–1440. [[CrossRef](#)]

23. Zdanowski, S.; Piątek, P.; Romański, J. An ion pair receptor facilitating the extraction of chloride salt from the aqueous to the organic phase. *New J. Chem.* **2016**, *40*, 7190–7196. [[CrossRef](#)]
24. Jaglenc, D.; Siennicka, S.; Dobrzycki, Ł.; Karbarz, M.; Romański, J. Recognition and Extraction of Sodium Chloride by a Squaramide-Based Ion Pair Receptor. *Inorg. Chem.* **2018**, *57*, 12941–12952. [[CrossRef](#)] [[PubMed](#)]
25. Jaglenc, D.; Dobrzycki, Ł.; Karbarz, M.; Romański, J. Ion-pair induced supramolecular assembly formation for selective extraction and sensing of potassium sulfate. *Chem. Sci.* **2019**, *10*, 9542–9547. [[CrossRef](#)]
26. Yu, X.-H.; Cai, X.-J.; Hong, X.-Q.; Tam, K.Y.; Zhang, K.; Chen, W.-H. Synthesis and biological evaluation of aza-crown ether-squaramide conjugates as anion/cation symporters. *Future Med. Chem.* **2019**, *11*, 1091–1106. [[CrossRef](#)]
27. Zaleskaya, M.; Jaglenc, D.; Karbarz, M.; Dobrzycki, Ł.; Romański, J. Squaramide based ion pair receptors possessing ferrocene as a signaling unit. *Inorg. Chem. Front.* **2020**, *7*, 972–983. [[CrossRef](#)]
28. Yan, L.; Huang, G.; Wang, H.; Xiong, F.; Peng, H.; Chen, F. Squaramide-Linked Chloramphenicol Base Hybrid Catalysts for the Asymmetric Michael Addition of 2,3-Dihydrobenzofuran-2-carboxylates to Nitroolefins. *Eur. J. Org. Chem.* **2018**, *1*, 99–103. [[CrossRef](#)]
29. Kisszekelyi, P.; Alammari, A.; Kupai, J.; Huszthy, P.; Barabas, J.; Holtzl, T.; Szente, L.; Bawn, C.; Adams, R.; Szekely, G. Asymmetric synthesis with cinchona-decorated cyclodextrin in a continuous-flow membrane reactor. *J. Catal.* **2019**, *371*, 255–261. [[CrossRef](#)]
30. Rapi, Z.; Bakó, P.; Drahos, L.; Keglevich, G. Side-Arm Effect of a Methyl α -D-Glucopyranoside Based Lariat Ether Catalysts in Asymmetric Syntheses. *Heteroatom Chem.* **2015**, *26*, 63–71. [[CrossRef](#)]
31. McCooney, S.H.; Connon, S.J. Readily Accessible 9-epi-amino Cinchona Alkaloid Derivatives Promote Efficient, Highly Enantioselective Additions of Aldehydes and Ketones to Nitroolefins. *Org. Lett.* **2007**, *9*, 599–602. [[CrossRef](#)]
32. Nagy, S.; Dargó, G.; Kisszekelyi, P.; Fehér, Z.; Simon, A.; Barabás, J.; Höltzl, T.; Mátravölgyi, B.; Kárpáti, L.; Drahos, L.; et al. New enantiopure binaphthyl-cinchona thiosquaramides: Synthesis and application for enantioselective organocatalysis. *New J. Chem.* **2019**, *43*, 5948–5959. [[CrossRef](#)]
33. Nagy, S.; Fehér, Z.; Dargó, G.; Barabás, J.; Garádi, Z.; Mátravölgyi, B.; Kisszekelyi, P.; Dargó, G.; Huszthy, P.; Höltzl, T.; et al. Comparison of Cinchona Catalysts Containing Ethyl or Vinyl or Ethynyl Group at Their Quinuclidine Ring. *Materials* **2019**, *12*, 3034. [[CrossRef](#)] [[PubMed](#)]
34. Bakó, P.; Tóke, L.J. Novel monoaza- and diazacrown ethers incorporating sugar units and their extraction ability towards picrate salts. *Inclusion Phenom. Mol. Recognit. Chem.* **1995**, *23*, 195–201. [[CrossRef](#)]
35. Bakó, P.; Kiss, T.; Tóke, L. Chiral azacrown ethers derived from D-glucose as catalysts for enantioselective Michael addition. *Tetrahedron Lett.* **1997**, *38*, 7259–7262. [[CrossRef](#)]
36. Yang, W.; Du, D.-M. Highly Enantioselective Michael Addition of Nitroalkanes to Chalcones Using Chiral Squaramides as Hydrogen Bonding Organocatalysts. *Org. Lett.* **2010**, *12*, 5450–5453. [[CrossRef](#)] [[PubMed](#)]
37. Makó, A.; Menyhárd, D.K.; Bakó, P.; Keglevich, G.; Tóke, L. Theoretical study of the asymmetric phase-transfer mediated epoxidation of chalcone catalyzed by chiral crown ethers derived from monosaccharides. *J. Mol. Struct.* **2008**, *892*, 336–342. [[CrossRef](#)]