



Right Heart Function in Cardiorenal Syndrome

Tilman Kramer^{1,2} · Paul Brinkkoetter^{2,3,4} · Stephan Rosenkranz^{1,2,4}

Accepted: 12 August 2022 / Published online: 27 September 2022
© The Author(s) 2022

Abstract

Purpose of Review Since CRS is critically dependent on right heart function and involved in interorgan crosstalk, assessment and monitoring of both right heart and kidney function are of utmost importance for clinical outcomes. This systematic review aims to comprehensively report on novel diagnostic and therapeutic paradigms that are gaining importance for the clinical management of the growing heart failure population suffering from CRS.

Recent Findings Cardiorenal syndrome (CRS) in patients with heart failure is associated with poor outcome. Although systemic venous congestion and elevated central venous pressure have been recognized as main contributors to CRS, they are often neglected in clinical practice. The delicate hemodynamic balance in CRS is particularly determined by the respective status of the right heart.

Summary The consideration of hemodynamic and CRS profiles is advantageous in tailoring treatment for better preservation of renal function. Assessment and monitoring of right heart and renal function by known and emerging tools like renal Doppler ultrasonography or new biomarkers may have direct clinical implications.

Keywords Cardiorenal syndrome · Right heart function · Right heart failure · Pulmonary hypertension · Pulmonary arterial hypertension · Venous congestion · Renal dysfunction · Biomarkers · Treatment approaches

Introduction

Cardiorenal syndrome (CRS) is commonly diagnosed in patients with heart failure (HF) and concomitant chronic kidney disease (CKD). Based on a cardiopulmonary-renal cross talk [1•, 2], acute or chronic dysfunction of one organ impairs the function of the respective other organ [3, 4]. While renal dysfunction in HF has traditionally been considered to result from decreased renal perfusion and associated neural and hormonal changes, recent evidence suggests

that rather persistent venous congestion represents a major contributor [5, 6, 7•]. Right ventricular (RV) function plays a key role in preventing CRS in HF and pulmonary hypertension (PH) as the heart aims at compensating the respective state by balancing pre- and afterload [1•]. In right heart failure (RHF), elevated central venous pressure (CVP) leading to venous congestion by backward transmission, was identified to initiate a vicious cycle of hormonal and endothelial activation, hepatic dysfunction, ascites, increased intra-abdominal pressure (IAP), intestinal mucosal ischemia, inflammation, oxidative stress, excessive renal tubular sodium reabsorption, and volume overload, leading to further RV stress [1•, 2, 7•, 8]. Renal congestion caused by HF- or PH-mediated RHF results in renal edema, increased interstitial pressure, tubular compression, and intracapsular tamponade, which may further aggravate back pressure and thus decrease renal blood flow (RBF) and glomerular filtration rate (GFR) [1•, 1, 10, 11]. In situation of decreased renal perfusion, the amount of glomerular blood filtered increases (filtration fraction = GFR/RBF) to maintain GFR. Increase in filtration fraction leads to proximal nephron sodium retention. Additionally, proximal nephron sodium retention leads to a lower fraction of sodium and chloride in

This article is part of the Topical Collection on *Cardio-renal Interactions in Heart Failure*

✉ Tilman Kramer
tilman.kramer@uk-koeln.de

¹ Klinik III Für Innere Medizin, Herzzentrum Der Universität Zu Köln, Köln, Germany

² Cologne Cardiovascular Research Center (CCRC), Klinikum Der Universität Zu Köln, Köln, Germany

³ Klinik II Für Innere Medizin, Nephrologie, Universität Zu Köln, Köln, Germany

⁴ Center for Molecular Medicine Cologne (CMMC), Universität Zu Köln, Köln, Germany

the tubular ultrafiltrate at the level of macula densa, leading to neurohormonal activation and worsening renal function (WRF) [4, 12]. The pathophysiology of CRS as a consequence of right heart dysfunction remain poorly understood, which creates an urgent need for biomarkers, diagnostic tools, and interventions to improve renal outcomes [13].

Study Selection

The aim of this systematic review article was to review the available literature on the meaning of right heart function in CRS “type 1” and “type 2” with a special focus on implications for current diagnostic and therapeutic management. Additionally, we provide an update on clinical and pathophysiological findings regarding CRS and right heart function. Therefore, we performed a comprehensive computerized literature search through multiple Medline searches on the PubMed database using MeSH terms and keywords. Searched terms included a combination of either “cardiorenal syndrome” and/or “pulmonary hypertension” and/or “pulmonary arterial hypertension” and/or “heart failure” and/or “decompensated heart failure” plus each of the following: “biomarker,” “central venous pressure,” “chronic kidney disease,” “classification,” “definition,” “diagnosis,” “heart failure,” “hemodynamics,” “interactions,” “kidney dysfunction,” “kidney failure,” “kidney injury,” “left heart failure,” “management,” “mortality,” “nephropathy,” “outcome,” “pathophysiology,” “prognosis,” “renal dysfunction,” “renal failure,” “renal function,” “renal insufficiency,” “right heart failure,” “right heart function,” “right-sided heart failure,” “survival,” “therapy,” “treatment,” “venous congestion” for articles published before January 2022. In addition to this search, reference lists of review articles were manually searched to identify other possible eligible references. Studies considered for our review included randomized controlled trials, prospective studies, retrospective studies,

review articles, and case reports. Studies with unavailable full text or inaccurate data extraction were excluded.

Classification of Cardiorenal Syndrome

While cardiac dysfunction impairs renal function, renal dysfunction can also lead to numerous harmful effects on the heart [14], and improvement in renal function can lead to cardiac reverse remodeling [15]. Thus, direct and indirect effects of the respective diseased organ cause a combined disorder of both organ systems by a complex combination of neurohormonal feedback mechanisms [16]. For this reason, the current definition by the “Acute Dialysis Quality Initiative” phenotyped CRS into 2 major groups: CRS and reno-cardiac syndrome, depending on the respective primary origin of the disease process. This classification distinguishes 5 subtypes according to disease severity and sequential organ involvement to enable a more precise and logical approach [3, 16, 17•]. The CRS definition includes acute or chronic conditions, whereby the primary failing organ can be either the heart or the kidney [3, 14, 16]. The exception is subtype 5, which represents a systemic condition affecting both organs simultaneously (Table 1).

In contrast, the previous terminology did not allow to identify and fully characterize the chronology of the pathophysiological interactions that characterize a particular type of combined cardiac/renal disease [16].

The current definition was established to facilitate a precise clinical characterization of CRS and to provide a basis for the development of new diagnostic and treatment approaches [17•]. Since the different subtypes of the current definition often overlap and may change dynamically during disease progression, deriving therapeutic recommendations remain clinically challenging. Functional assessments of cardiac and renal function for exact assignment to the different subtypes are still scarce [4, 17•].

Table 1 Classification of cardiorenal syndrome based on the “Consensus Conference of the Acute Dialysis Quality Initiative” [3, 16, 17•]

CRS phenotype	Nomenclature	Description	Clinical examples
Type 1	Acute	HF resulting in AKI	ACS resulting in cardiogenic shock and AKI, AHF resulting in AKI
Type 2	Chronic	Chronic HF resulting in CKD	Chronic HF
Type 3	Acute renocardiac syndrome	AKI resulting in AHF	HF in the setting of AKI from volume overload, inflammatory surge, and metabolic disturbances in uremia
Type 4	Chronic renocardiac syndrome	CKD resulting in chronic HF	LVH and HF from CKD-associated cardiomyopathy
Type 5	Secondary	Systemic process resulting in HF and renal failure	Amyloidosis, sepsis, cirrhosis

ACS, acute coronary syndrome; AHF, acute heart failure; AKI, acute kidney injury; CKD, chronic kidney disease; CRS, cardiorenal syndrome; HF, heart failure; LVH, left ventricular hypertrophy

Pathophysiological Evolution from Low Perfusion to Renal Congestion

It has been traditionally postulated that reduced cardiac output (CO) leads to diminished renal perfusion and neurohormonal activation representing the main drivers for the development of CRS in HF with reduced ejection fraction (HFrEF) [4, 5, 6, 7•, 18, 19]. The right heart function is primarily important to maintain sufficient RV filling by adequate preload [4]. It has been assumed that a cardiac index (CI) of at least 1.5 l/m²min is necessary to maintain sufficient forward transmission for adequate renal perfusion [4, 19]. The impairment of renal arterial blood flow by either low CO and/or increased peripheral vascular resistance is considered to activate the renin–angiotensin–aldosterone system (RAAS), sympathetic nervous system, and release of arginine vasopressin [4, 18]. Activation of these neurohormones causes systemic and renal vasoconstriction and increased sodium and water retention in the kidneys to compensate arterial underfilling and reduced preload, but at the expense of increased systemic vascular resistance and higher plasma volume, thus leading to worsening HF [13, 18].

Recently, venous congestion has been recognized as important contributor to renal dysfunction in CRS [7•]. Regardless of whether left or right HF is present, venous congestion, increased CVP, and increased renal venous pressure appear to be the major hemodynamic contributors to CRS [1•, 5, 7•, 20•, 21]. The importance of systemic venous congestion in terms of WRF is supported by the finding that deterioration of kidney function occurs more frequently in HF with preserved ejection fraction (HFpEF) than in HFrEF [4, 22].

These findings place the pathophysiological focus of CRS more on right heart function. RV dysfunction, which is also frequently present in patients with left-sided heart disease, leads to backward transmission of elevated filling pressures, resulting in increased CVP and renal venous congestion. Due to complex RV/LV interactions, RHF results in underfilling of the left ventricle (LV) and systemic low-output and in advanced cases to cardiogenic shock [4]. Thus, right heart dysfunction plays a pivotal role for secondary organ damage. Acute RHF may occur because of rapidly increased RV afterload through pulmonary embolism, decompensated RHF/LHF, or hypoxia [4]. Another cause of acute RHF represents impaired RV contractility in consequence of ischemia, myocarditis, or postcardiotomy shock [4]. Chronic RHF is mainly due to chronically elevated pulmonary artery pressure (PAP) and/or pulmonary vascular resistance (PVR), i.e., increased RV afterload in PH. The downstream disturbances in cardiorenal interactions are likely to be similar regardless of whether the primary cause was right-sided or left-sided HF [4].

PH is a hemodynamic condition defined by an elevated mean PAP to ≥ 25 mmHg at rest, measured invasively by right heart catheterization (RHC) [23]. A revised definition was proposed at the 6th World Symposium on Pulmonary Hypertension (WSPH) in 2018, suggesting a redefined threshold of > 20 mmHg at rest [24].

Any kind of PH can contribute to RHF [13, 25] and thus to the development of CRS [4, 13, 26]. PH due to left heart disease represents the most common cause of RHF, and the pathophysiology of CRS in this group is comparable to that in left heart failure (LHF) [13]. Data on the prevalence of CRS in patients with PH and isolated RHF remain scarce [13]. However, PH is associated with impaired kidney function leading to poorer outcome [1•, 20•, 26–30]. Similar to LHF, isolated RHF in PH often results in renal venous congestion and arterial hypoperfusion with subsequent loss of GFR [13, 20•]. Conversely, renal dysfunction may also contribute to aggravation of PH, since cardiopulmonary hemodynamics are crucially dependent on the regulation of fluid homeostasis. Elevated CVP leads to release of inflammatory mediators, neurohormones, and activation of endothelial cells, which may further deteriorate heart and kidney function [13, 21, 31]. Circulatory factors excreted in the kidney are involved not only in the pathogenesis of local structural glomerular and interstitial damage but also in pulmonary inflammation following renal injury [1•, 2, 32]. In PH, the right heart function is elementary to cope with elevated PVR by balancing pre- and afterload resulting in water and salt retention and venous congestion [1•]. LV output is potentially also reduced in PH, as reduced RV output and dilatation frequently lead to leftward shift of the interventricular septum and altered LV geometry, resulting in impaired filling and output despite preserved LVEF [1•, 4, 13, 33–35]. Consequently, forward transmission can also be diminished in PH, leading to reduced renal arterial pressure and thus contributing to the development of CRS [4, 6, 33, 36]. In PAH, reduced CI rather than increased RAP was an independent predictor for WRF [1•, 20•], while in a study on RHF due to miscellaneous PH classes with preserved LVEF elevated RAP rather than CI independently predicted GFR [13, 37]. These results indicate that in PAH, reduced CI and subsequent renal hypoperfusion might be the key hemodynamic drivers of GFR loss and may precede deleterious effects on kidney function of augmented RAP and renal venous congestion during worsening right heart function [20•]. Patients with PH due to left heart disease might be more prone to renal venous congestion similar to the LHF phenotype [13]. Differentiation between hemodynamic profiles may be advantageous in tailoring treatment for preservation of GFR in HF and various forms of PH.

Elevated CVP initially leads to a slightly increased GFR through increased glomerular hydrostatic pressure due to elevated proximal peritubular capillary pressure, resulting

in increased efferent arteriolar pressure [6, 7•, 13, 38, 39]. After glomerular hyperfiltration is exhausted, GFR progressively worsens [7•] due to renal edema, increasing interstitial pressure, tubular compression, and intracapsular tamponade, which may further aggravate back pressure and thus decrease renal perfusion pressure and GFR [1•]. The term “congestive nephropathy” was suggested for a potentially reversible subtype of renal dysfunction caused by reduced renal venous outflow and increased renal interstitial pressure [7•]. Based on increased CVP, another contributor to WRF is ascites and increased IAP [9]. Immediate reduction of IAP by large volume removal improves GFR in decompensated HF [40], which is why ascites and IAP assessment could be integrated into the

diagnostic and therapeutic workflow. Pathophysiological interactions are depicted in Fig. 1.

Diagnosis of Right Heart Dysfunction and Cardiorenal Syndrome

Hemodynamics

RHF can occur in progressive or acutely decompensated PH, leading to chronic or acute kidney damage and thus contributing to CRS. In combined RHF with concomitant acute renal failure, deterioration of either organ function may converge in a vicious circle leading to refractory




Fig. 1 Hemodynamic cardiorenal interactions in left and right heart failure. **Right side:** traditional “low flow” hypothesis of the development of worsening renal function in heart failure due to renal vasoconstriction and hypoperfusion leading to tubular hypoxia and necrosis. **Left side:** concept of “congestive nephropathy” primarily leading worsening renal function in heart failure according to current expert opinion. Heart failure-induced backward transmission results in increased central venous pressure and renal venous congestion with subsequent impaired renal function. CO, cardiac output; CVP, central

venous pressure; GFR, glomerular filtration rate; HF, heart failure; LHF, left heart failure; LV, left ventricle; RAAS, renin–angiotensin–aldosterone system; RHF, right heart failure; RV, right ventricle; WRF, worsening renal function. Adapted from Schefold, J. C. et al. (2016) Heart failure and kidney dysfunction: epidemiology, mechanisms and management. *Nat. Rev. Nephrol.* 10.1038/nrneph.2016.113 and from Rosenkranz S, Howard LS, Gomberg-Maitland M, Hoepfer MM. Systemic Consequences of Pulmonary Hypertension and Right-Sided Heart Failure. *Circulation* 2020;141:678–693

congestive RHF [1•]. In CRS prevention, precise measurements of right heart function are gaining importance. The most accurate way to determine right heart function is RHC [23, 24, 41]. A cutoff value of 15 mmHg for pulmonary arterial Wedge pressure (PAWP) distinguishes between precapillary (≤ 15 mmHg) and postcapillary (> 15 mmHg) PH. This distinction is important since postcapillary PH often occurs in LHF and primarily requires optimized therapy of the left heart condition, while in precapillary PH such as PAH or CTEPH, special treatments are available to improve RHF [23]. Precise hemodynamic assessment is of utmost importance for correct diagnosis and therapeutic management potentially affecting CRS. In experienced hands, RHC is safe and could be warranted in selected difficult-to-treat CRS patients to detect and treat subclinical congestion while avoiding intravascular underfilling [42].

Transthoracic echocardiography represents a more convenient and noninvasive assessment tool of right heart function, which can be performed in nearly every clinical situation. Specific echocardiographic measurements provide information about right heart function in CRS and helps to identify those patients who might require RHC [23]. Echocardiographic signs of right heart strain increase the risk of RHF and systemic venous congestion contributing to CRS. Therefore, control of congestion has been proposed as a central goal in HF treatment [5, 6, 43•]. Assessment of the diameter of the inferior vena cava and its respiratory variability is of particular importance as it correlates positively with CVP and RAP [44, 45].

Renal Doppler ultrasonography represents a relatively new approach to evaluate renal congestion and to guide therapy in patients with left- or right-sided HF [43•, 46•]. Based on Doppler renal venous flow, identification of altered intrarenal venous flow (IRVF) patterns is used to predict adverse outcome and to monitor diuretic response in HF [43•, 47, 48•, 49]. IRVF patterns depend on RAP and the mean circulatory filling pressures [4, 47, 48•]. Backward transmission in RHF results in elevated RAP, which is transmitted into the renal vein with consequently increased pulsatility and change in IRVF patterns, reflecting the renal vasculature's response to elevated intrarenal pressure within the rigid renal capsule [4, 50]. Compared to renal resistance index, the IRVF pattern demonstrates higher prognostic impact [47], confirming renal venous congestion rather than hypoperfusion as the predominant component in CRS development [1•, 4]. Renal venous congestion index (RVSI) was described as a dimensionless continuous ratio that reflects the complete continuum of renal venous congestion by indicating the fraction of the cardiac cycle during which there is no renal venous outflow [43•, 46•]. In PH, RVSI correlates with right heart and renal function [46•]. Its measurement represents a simple and noninvasive approach for the assessment of renal congestion [43•, 46•] and provides additional

prognostic information to stratify the risk of RHF in PH [46•], which in turn can result in CRS [17•].

Biomarkers

Various biomarkers reflecting neurohormonal disorders, myocardial stress/injury, inflammation, oxidative stress, and renal clearance/injury have been considered for their diagnostic and prognostic value in HF and CRS [13, 17•, 51]. Since progressive deterioration in right heart function is associated with slow progressive degenerative processes potentially resulting in CKD, sensitive biomarkers are needed [1•, 52].

Cardiac Biomarkers

Brain Natriuretic Peptide and N-Terminal Pro-brain Natriuretic Peptide

Measurement of brain natriuretic peptide (BNP) or N-terminal pro-BNP (NT-proBNP), which is directly associated with congestion, is supported by current HF and PH guidelines [23, 53] and plays a central role for diagnosis and prognosis of CRS in HF [17•]. BNP/NT-proBNP serum levels in HF with CRS are higher compared to patients with preserved kidney function [4, 54, 55•], which may be explained by several mechanisms including impaired renal excretion and volume overload [4, 56–58]. In patients with $\text{GFR} < 60$ ml/min/1.73m², BNP/NT-proBNP serum concentrations should be interpreted with caution and is only of limited use in diagnosing HF and monitoring treatment response especially when HF and volume overload are present [59]. Nevertheless, elevation of BNP/NT-proBNP in HF with already existing renal dysfunction is also associated with poorer prognosis [54, 55•]. Currently, BNP/NT-proBNP measurement is not recommended for prevention or treatment of acute kidney injury (AKI) [45].

Cardiac Troponin T and Galectin-3

Cardiac troponin T and galectin-3 represent useful cardiac biomarkers which are elevated in CRS [4, 58, 60, 61] and prognostically relevant in HF [4, 60, 61]. However, elevated troponin T serum levels are unspecific for renal damage and occur in most patients with advanced renal dysfunction [54, 55•] and thus must be interpreted with caution when diagnosing acute coronary syndrome in these patients [59]. Determination of troponin T is only recommended on admission to exclude acute myocardial injury; albeit, it is elevated in most acute HF patients [53].

Expression of galectin-3 is linked to fibrosis [55•, 62, 63], including renal [51, 64, 65] and cardiac remodeling and progression of HF [55•, 63, 66–69]. Thus, galectin-3

potentially provides information about the pathophysiology of the underlying renal dysfunction and its progression in HF [55•, 70–72]. High galectin-3 plasma levels are associated with increased risk of renal dysfunction [55•, 73, 74]. In HF, elevated galectin-3 is also linked to impaired kidney function and higher mortality [51, 70]. Through its ability in early detection of WRF, it may be used for risk estimation of CRS progression and potentially as therapeutic target [55•, 75].

Renal Biomarkers

Cystatin C

Cystatin C (CysC) is a protease occurring in all nucleated cells [51, 55•]. In contrast to creatinine, it is filtered freely and then reabsorbed but not secreted in renal tubules [51, 55•]. It was assumed that assessment of CysC could be superior to serum creatinine as it is independently excreted from the respective muscle mass [76]. CysC represents a promising alternative endogenous filtration biomarker for monitoring renal function [51, 59, 76, 77] and a prognostic indicator in HF with normal renal function [51, 55•, 78–80]. However, assessment of CysC provides no information about the different pathomechanisms of CRS [51].

Albuminuria

Albuminuria is associated with increased cardiovascular disease (CVD) risk and is prognostically relevant for CKD progression [59]. In HF, albuminuria is associated with poorer outcome [4, 81, 82]. Albuminuria indicates damage to the glomerular filter, primarily due to dysfunction of podocytes, which are crucial for maintaining glomerular filter barrier [83]. In addition, endothelial dysfunction, inflammation, elevated glomerular pressure, or atherosclerosis contribute to increased albumin excretion [55•, 84, 85]. Albuminuria also occurs in anomalous renal microcirculation [55•, 86] and can reflect impaired renal hemodynamics, such as renal venous congestion caused by RHF and increased CVP [6, 55•]. Accordingly, albuminuria provides information about the pathophysiology leading to WRF [55•, 84]. The assessment of albuminuria integrates urinary creatinine levels to calculate the ratio between urinary albumin and creatinine (UACR) [55•]. Microalbuminuria, defined by UACR between 30 and 300 mg/g [85], is common in HF and associated with worse prognosis [55•, 81, 82, 87].

Neutrophil Gelatinase–Associated Lipocalin

Neutrophil gelatinase–associated lipocalin (NGAL) is a protein that is freely filtered through the glomerulus and completely reabsorbed in the proximal part of the tubule [55•]. Under normal conditions, its concentration in urine

and serum is very low [51, 55•, 88]. In proximal tubular injury, NGAL urinary level rises above the normal range [4, 89], because it cannot be completely reabsorbed due to tubular damage [55•]. High NGAL levels occur in HF with and without renal dysfunction [4, 51, 58, 90] and predict WRF, particularly in acute decompensated HF [51], and adverse clinical outcome [55•, 91, 92], also in CRS patients [4, 93]. However, the diagnostic accuracy of NGAL is not without controversy and may be affected by confounding factors such as sepsis, inflammation, anemia, hypertension, hypoxemia, and cancer [55•, 88].

N-Acetyl Beta Glucosaminidase

N-Acetyl beta glucosaminidase (NAG) is a lysosomal enzyme of the proximal tubule cells, which is excreted into urine if tubular damage with subsequent disruption of lysosomal integrity occurs [51, 55•, 89, 94, 95]. NAG serum levels are gradually elevated in HF with preserved or reduced renal function and are associated with poorer prognosis, independently from GFR but dependent on LVEF [51, 55•, 96–98]. Increase in NAG is predictive for AKI in patients with normal and worsening kidney function and for impaired survival in patients with preexisting AKI [51, 95, 99]. Despite limited specificity [51, 96], NAG is a promising prognostic biomarker for CRS and could also represent a potential therapeutic target since it decreases in response to diuretic-induced decongestion similar to BNP/NT-proBNP [51, 100], which is particularly interesting since venous congestion is the major pathophysiological driver of CRS [7•, 7, 51].

Kidney Injury Molecule-1

Kidney injury molecule-1 (KIM1) is a glycoprotein which is expressed in proximal tubule cells and excreted in the urine after tubular injury [55•, 95]. High urinary KIM1 concentrations predict poorer prognosis including WRF in HF independent of GFR [55•, 96–99] but dependent on LVEF in terms of quantitative characteristics [51, 98]. KIM1 levels correlate with BNP/NT-proBNP and are sensitive to volume fluctuations, reflecting their dependence on congestive clinical states and making it an excellent target for diuretic management of CRS and a possible biomarker for CRS phenotyping [51, 98, 100].

Dickkopf-3

Dickkopf-3 (DKK3) is a stress-induced, tubular epithelial-derived profibrotic glycoprotein which predicts WRF [101–103]. Increased urinary DKK3 levels identify patients at high risk for short-term GFR decline regardless of the respective cause of kidney injury and are associated with

tubulointerstitial fibrosis [102–104]. Although data on the meaning of urinary DKK3 levels in HF are currently pending, DKK3 might represent a promising future biomarker in CRS. Potentially, elevated DKK3 urinary levels are indicative for active renal fibrosis, contributing to poorer outcome in CRS. Hence, urinary DKK3 might serve as a biomarker to monitor CKD progression which may be useful for clinicians to monitor treatment effects and guide therapeutic adjustments [102–104].

Together, there are numerous promising novel biomarkers to monitor kidney function in CRS. However, none of these is specific to impaired right heart function or CRS, making it difficult to identify impaired right heart function as the main contributor to CRS [4]. Therefore, CRS phenotyping must integrate information from biomarkers, hemodynamics, and imaging modalities and should always be interpreted in the clinical context.

Therapeutic Approaches

Diuretics, Volume Optimization, and Ultrafiltration

Since elevated CVP and venous congestion were recognized as main drivers for CRS, diuretics represent the initial drug of choice in decompensated HF associated with inadequate fluid retention to decrease volume overload and to improve cardiorenal hemodynamics [42, 53, 105–107]. Reduction of RV overload, CVP, and renal venous pressure lead to increased renal perfusion and to improved RV/LV interaction, cardiac and kidney function [4, 107]. Loop diuretics are frequently used for fast natriuresis with subsequent extracellular volume reduction [107–109]. There are no differences in outcome, symptom relief, or changes in renal function when loop diuretics are administered as bolus or continuous therapy [4, 42, 106]. High-dose compared to low-dose administrations resulted in a faster relief of congestion with a transient reduction in GFR [42, 106]. The addition of non-loop diuretics might be reasonable to maintain natriuresis without compromising GFR [4, 110, 111]. The level of care for HF patients with CKD should be the same as for those without CKD, but any escalation of therapy and/or clinical deterioration should prompt monitoring of GFR and serum potassium concentration [61]. Acetazolamide as a potent inhibitor of proximal tubular sodium reabsorption could represent another interesting option in decongestive treatment strategies, as targeting sodium reabsorption in the proximal tubules implies potential benefits in HF. The ADVOR trial is currently investigating whether acetazolamide in combination with loop diuretic therapy can improve outcome and decongestion in acute HF with fluid overload [112, 113]. Acetazolamide may be considered if loop diuretic response remains insufficient [53].

During diuretic treatment in acute congestive HF with WRF, intensive volume depletion initially results in increases in serum creatinine and biomarkers of tubular injury (NAG, KIM1, NGAL), while renal function improves over time, suggesting that benefits of decongestion may outweigh transient increases in serum creatinine or tubular injury markers at treatment start [1•, 114]. Caution is advised in decompensated HF without congestion and excessive diuresis, both of which may be associated with reduced RV preload and hence impairment of CO, thus resulting in intravascular hypovolemia, hypotension, and decreased diuresis and natriuresis [4]. When the RV is collapsed, careful volume loading can be beneficial until RV is adequately filled. Further volume expansion can induce adverse effects and should be avoided, especially in patients with mean arterial pressure less than 60 mmHg [13]. Nephrotoxic agents should be temporary avoided [59], regardless of whether decompensated HF is congestive or non-congestive [13]. Since ascites contributes to WRF through increased IAP, paracentesis potentially represents a viable treatment option for acute hemodynamic improvement. Ultrafiltration is another decongestive strategy in decompensated HF and concomitant renal dysfunction, although it does not lead to better outcome or renal function compared to pharmaceutical care only [42, 111]. Ultrafiltration is associated with a higher rate of adverse events [111] and is not an effective therapy in CRS [108]. Future studies on individually titrated ultrafiltration patients are warranted [45, 115], especially in HF and CRS.

Diuretic Resistance

Since decongestion is the key treatment strategy to reduce venous congestion in decompensated HF-associated CRS [4, 42, 116–118], special attention should be paid to the potential development of diuretic resistance, which is predictive for poor outcome in CRS and HF [4, 51, 119–121]. Diuretic resistance is defined as reduced diuretic response resulting in inadequate relief of edema, thereby missing the therapeutic target [4, 122]. Decongestion strategies in CRS patients with diuretic resistance remain clinically challenging, as this is often associated with significant reduction of GFR and CO, which results in diminished delivery of diuretics to the tubules [4, 51, 107, 119]. Another potential contributor to impaired delivery of protein-bound loop diuretics is hypoalbuminemia which is frequent in advanced HF [4, 107]. Further causes are insufficient dosage of diuretics or inadequate substrates (sodium and chloride) at the renal tubules [4, 107, 119]. In addition, activation of the RAAS by diuretic-inherent effects results in reduced renal arterial flow and tubular secretion [51, 123]. Further contributors to diuretic resistance are nephron remodeling due to prolonged use of loop diuretics [4, 124] and diuretic braking phenomena by hemodynamic and neurohormonal mechanisms including RAAS activation and afferent

arteriolar vasoconstriction [4, 51, 125, 126], which lead to diminished natriuresis to preserve intravascular volume in response to increased, diuretics-induced sodium excretion [4, 51, 107, 127]. In CRS, renal dysfunction leads to impaired release of diuretics into the tubular lumen, and sodium excretion is diminished because of reduced filtration [4, 128, 129]. Diuretic resistance is indicative of HF-induced renal dysfunction and less dependent on GFR, suggesting that determination of diuretic resistance is helpful to identify CRS patients [51]. The diuretic dose–response curve in HF patients typically has a sigmoid shape and demonstrates a rightward and downward shift [130]. Given the dependency on RBF, higher doses of loop diuretics might be necessary in CRS. The soon to be published ADVOR trial with acetazolamide as a combinatorial diuretic treatment will clarify whether this strategy is superior to conventional loop diuretic treatment in acute HF [112, 113]. The forthcoming CLOROTIC trial evaluates whether the addition of hydrochlorothiazide to a loop diuretic represents an effective strategy for decongestion in HF [131], since this is recommended for nephrotic patients with diuretic resistance [115]. In two recent trials, the sodium-glucose cotransporter 2 (SGLT2) empagliflozin increased diuresis and demonstrated beneficial effects in acute decompensated HF patients without impairing kidney function [132, 133].

Renin–Angiotensin–Aldosterone System Inhibitors

RAAS activation is crucially involved in pathophysiological changes contributing to WRF. In HFrEF, reno-protective RAAS inhibitors belong to the standard of care and have demonstrated beneficial effects on cardiovascular outcome [53, 59]. Because CRS and hyperkalemia are common in HFrEF [7•, 134–136], particular attention should be paid to RAAS inhibitors and novel treatment agents [53] for possible interactions with renal function. Indeed, the occurrence of renal dysfunction and hyperkalemia are challenges for RAAS inhibitor therapy in clinical practice [59, 135, 136•]. In HFrEF therapy, RAAS inhibition is linked to WRF, which is associated with poorer outcomes compared to no WRF [137]. However, the use of RAAS inhibitors leads to a reduction of all-cause mortality, which is significantly more pronounced in the presence of WRF than in the group without WRF [137]. This striking difference could be explained by the fact that in a state of reduced kidney function there is more intensive RAAS stimulation [137–139], which in turn causes the outstanding potential for improvement with sufficient RAAS blockade [137].

Permissive Acute Kidney Injury

In HFrEF, RAAS inhibitors and SGLT2 inhibitors can reduce renal perfusion, possibly followed by an acute decrease in

GFR [53]. However, this should not lead to immediate discontinuation of these beneficial treatments [53, 137, 140]. Acute declines in GFR should not be misinterpreted as AKI but rather be understood as “permissive AKI” [140], since the use of the respective agents contributes to better preservation of kidney function longer term and reduced all-cause mortality [137]. The optimal therapeutic strategy includes assessment of the clinical setting in which GFR loss occurs [140]. In the absence of alternative treatable causes (e.g., infections, nephrotoxic co-medication, hypotension), a decline in GFR of 30–40%, e.g., under RAAS inhibitor therapy, should be tolerated and not lead to discontinuation of this outcome-modifying therapy [137, 140].

Soluble Guanylate Cyclase Stimulators

Vericiguat, a soluble guanylate cyclase stimulator, represents a new treatment option in HFrEF [53] that improved outcome, irrespective of baseline GFR or WRF [136•, 141–143]. Vericiguat may be considered in addition to standard therapy in HFrEF to reduce risk of cardiovascular mortality and hospitalizations for HF [53]. Vericiguat demonstrated no negative effects on renal function, and, thus, there is no need to down-titrate or interrupt therapy if WRF and/or hyperkalemia occur [136•]. Vericiguat appears to be a potential candidate for the prevention of CRS in HFrEF patients, although the exact mechanisms of action on renal function remain speculative and require further research. On the one hand, vericiguat could improve RBF by its positive effects on cardiac and endothelial function, on the other hand, mild blood pressure lowering effects may potentially impair renal perfusion [136•].

Sodium-Glucose Cotransporter 2 Inhibitors

SGLT2 inhibitors are recommended for the treatment of HFrEF patients with and without type 2 diabetes mellitus [53]. Regardless of the presence of diabetes, SGLT2 inhibitors showed beneficial effects on cardiorenal outcome in HFrEF and HFpEF [108, 144–147]. These cardio-reno-protective benefits render SGLT2 inhibitors a promising drug in CRS prevention and treatment [148]. Selective blockade of the SGLT2 transporter increases renal excretion of glucose and sodium by inhibiting their reabsorption in the renal proximal tubules [149–151]. Reno-protective effects are mainly mediated by reduction in albuminuria, inflammation, hypoxic stress, renal artery stiffness, and restoration of tubuloglomerular feedback [108, 148]. Reinforced diuresis may further contribute to advantageous effects of SGLT2 inhibitors on kidney function [108, 152]. Overall, the safety profile of SGLT2 inhibitors is compelling, with mild genital mycotic infections being the most common but rare adverse

event [153]. Euglycemic diabetic ketoacidosis is a rare but serious side effect of SGLT2 inhibitor treatment, occurring mainly in diabetes with insulin deprivation, after surgery [153] or under infectious conditions [154].

Vasopressin V2 Receptor Antagonists

In HF, RAAS activation leads to elevated vasopressin levels with subsequent impairment of cardiac function, peripheral vasoconstriction, and increased afterload [108]. There is a vasopressin V2 receptor-mediated water retention and, thus, an increase in preload [108, 155, 156]. Tolvaptan is a highly selective vasopressin V2 receptor antagonist with a convincing safety profile that has demonstrated beneficial effects in HF contributing to reduced volume overload, improved symptoms, increased urinary output, and corrected sodium levels without affecting renal function and serum electrolytes through its action on neurohormonal signaling in CRS [108, 155, 157]. Although tolvaptan failed to improve outcome in HF [53, 108, 158], data suggest that it may serve as a potential drug for decongestion in CRS [108, 159], thus promoting renal function by maintaining renal perfusion and avoiding intravascular volume depletion [108, 160]. Tolvaptan can be considered to increase serum sodium and urinary output in patients with persistent hyponatremia and congestion [53].

Treatment of Pulmonary Hypertension

In PH, the use of diuretics is recommended in patients with fluid retention associated with RHF [23], whereas the initiation of targeted therapies requires precise hemodynamic diagnosis and classification [23]. Targeted treatment results in improved cardiopulmonary hemodynamics and RV function without significant changes in GFR [20•], although most medications were shown to have nephroprotective potential in preclinical or clinical settings [26]. Treatment of PAH with the phosphodiesterase type 5 inhibitor (PDE5i) sildenafil was associated with improved kidney function [1•, 161]. Changes in kidney function could be due to several influencing factors, such as progressive PAH, the more intensive use of diuretics or polypharmaceutical effects. Since there are no dedicated clinical trials that have investigated the impact of PAH-targeted therapy on kidney function, further research is warranted [26]. Despite limited data, treatment with pulmonary vasodilators (e.g., inhaled nitric oxide, prostacyclin, and iloprost) and inotropes resulted in beneficial effects on CO and venous congestion [4, 108, 162].

In LHF, backward transmission of elevated left-sided filling pressure into the lung leads to postcapillary PH

[163–166]. The primary strategy to improve cardiopulmonary hemodynamics is optimization of volume status and filling pressures. Although PH is common in HFrEF and HFpEF [167] and is associated with unfavorable cardiorenal outcomes, there is currently no recommendation for targeted PH therapy in both of these entities [23, 168, 169]. Preliminary data suggests that treatment with PDE5i might have advantageous effects in HFpEF and combined post- and precapillary PH [43, 163, 170, 171]. Further studies are needed since there are currently no data on the impact on CRS.

Conclusions

Systemic venous congestion and elevated CVP in the context of RV dysfunction are major contributors in CRS. Although the crucial contribution of congestion to CRS and its impact on outcomes have been well documented, they often remain neglected in clinical practice. Thus, accurate assessment and the recognition of right heart function in CRS gain in importance. CRS in RHF should be suspected when WRF occurs either in cases of AKI without preexisting renal damage or in acute-on-chronic settings. The delicate hemodynamic balance is crucially affected by the respective status of the right heart. RHF patients are potentially threatened by acute decompensation, which must be prevented with close-meshed and strict volume management aiming at preserving or improving renal function and outcome. Assessment and monitoring of right heart and renal function by known and emerging tools like renal Doppler ultrasonography or new biomarkers may have direct clinical implications. The monitoring and differentiation between hemodynamic and CRS profiles may be advantageous in tailoring treatment for preservation of renal function.

Funding Open Access funding enabled and organized by Projekt DEAL.

Declarations

Conflict of Interest TK: Remunerations for lectures from Janssen/Actelion.

PB: Speaker honoraria and consultant fees from Sanofi-Genzyme, AstraZeneca, Alexion, Bayer, Travere, Pfizer, Novartis, Roche; research funding from the German Research Foundation BR-2955/8.

SR: Remunerations for lectures and/or consultancy from Abbot, Janssen/Actelion, Arena, Bayer, BMS, GSK, MSD, Novartis, Pfizer, and United Therapeutics and grants to institution from Actelion, Bayer, Novartis, and United Therapeutics.

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Papers of particular interest, published recently, have been highlighted as:
- Of importance
1. • Rosenkranz S, Howard LS, Gomberg-Maitland M, Hoepfer MM. Systemic consequences of pulmonary hypertension and right-sided heart failure. *Circulation* 2020;141:678–693. **In this article, the author describes systemic consequences of pulmonary hypertension on secondary organ systems. Based on pulmonary hypertension induced right heart failure, pathomechanistic insights of worsening renal function are demonstrated.**
 2. Husain-Syed F, McCullough PA, Birk HW, et al. Cardio-pulmonary-renal interactions: a multidisciplinary approach. *J Am Coll Cardiol*. 2015;65:2433–48.
 3. Ronco C, McCullough P, Anker SD, et al. Cardio-renal syndromes: report from the consensus conference of the acute dialysis quality initiative. *Eur Heart J*. 2010;31:703–11.
 4. Tabucanon T, Tang WHW. Right Heart Failure and Cardiorenal Syndrome. *Cardiol Clin*. 2020;38:185–202.
 5. Mullens W, Abrahams Z, Francis GS, et al. Importance of venous congestion for worsening of renal function in advanced decompensated heart failure. *J Am Coll Cardiol*. 2009;53:589–96.
 6. Damman K, van Deursen VM, Navis G, Voors AA, van Veldhuisen DJ, Hillege HL. Increased central venous pressure is associated with impaired renal function and mortality in a broad spectrum of patients with cardiovascular disease. *J Am Coll Cardiol*. 2009;53:582–8.
 7. • Husain-Syed F, Gröne HJ, Assmus B et al. Congestive nephropathy: a neglected entity? Proposal for diagnostic criteria and future perspectives. *ESC Heart Fail* 2021;8:183–203. **In this article, congestive nephropathy is proposed as neglected clinical entity. Congestive nephropathy is a potentially reversible subtype of renal dysfunction associated with declining renal venous outflow and progressively increasing renal interstitial pressure.**
 8. Verbrugge FH, Dupont M, Steels P, et al. Abdominal contributions to cardiorenal dysfunction in congestive heart failure. *J Am Coll Cardiol*. 2013;62:485–95.
 9. Mullens W, Abrahams Z, Skouri HN, et al. Elevated intra-abdominal pressure in acute decompensated heart failure: a potential contributor to worsening renal function? *J Am Coll Cardiol*. 2008;51:300–6.
 10. Le Clef N, Verhulst A, D'Haese PC, Vervaeke BA. Unilateral renal ischemia-reperfusion as a robust model for acute to chronic kidney injury in mice. *PLoS One*. 2016;11:e0152153.
 11. Hemmi S, Matsumoto N, Jike T, et al. Proximal tubule morphology in rats with renal congestion: a study involving the in vivo cryotechnique. *Med Mol Morphol*. 2015;48:92–103.
 12. Mullens W, Martens P, Testani JM, et al. Renal effects of guideline-directed medical therapies in heart failure: a consensus document from the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail*. 2022;24:603–19.
 13. Bansal S, Prasad A, Linas S. Right heart failure-unrecognized cause of cardiorenal syndrome. *J Am Soc Nephrol*. 2018;29:1795–8.
 14. Berl T, Henrich W. Kidney-heart interactions: epidemiology, pathogenesis, and treatment. *Clin J Am Soc Nephrol*. 2006;1:8–18.
 15. Hawwa N, Shrestha K, Hammadah M, Yeo PSD, Fatica R, Tang WHW. Reverse remodeling and prognosis following kidney transplantation in contemporary patients with cardiac dysfunction. *J Am Coll Cardiol*. 2015;66:1779–87.
 16. Ronco C, Haapio M, House Andrew A, Anavekar N, Bellomo R. Cardiorenal syndrome. *J Am Coll Cardiol*. 2008;52:1527–39.
 17. • Rangaswami J, Bhalla V, Blair JEA et al. Cardiorenal syndrome: classification, pathophysiology, diagnosis, and treatment strategies: a scientific statement from the American heart association. *Circulation* 2019;139:e840–e878. **This work is related to identifying the factors contributing to diuretic resistance representing a key step in optimizing decongestion in cardiorenal syndrome. Additionally, the authors describe biomarkers of cardiac and kidney injury which represent a new dimension in the diagnostic algorithm in evaluating heart failure with impaired kidney function and offer prognostic value in acute and chronic cardiorenal syndrome.**
 18. Schrier RW, Abraham WT. Hormones and hemodynamics in heart failure. *N Engl J Med*. 1999;341:577–85.
 19. Ljungman S, Laragh JH, Cody RJ. Role of the kidney in congestive heart failure. Relationship of cardiac index to kidney function. *Drugs* 1990;39 Suppl 4:10–21; discussion 22–4.
 20. • Bitker L, Sens F, Payet C et al. Presence of kidney disease as an outcome predictor in patients with pulmonary arterial hypertension. *Am J Nephrol* 2018;47:134–143. **The author demonstrates that chronic kidney disease is frequent at diagnosis of pulmonary arterial hypertension and is independently associated with increased mortality. Right heart failure may induce renal hypoperfusion and congestion, and is associated with eGFR decrease.**
 21. Ross EA. Congestive renal failure: the pathophysiology and treatment of renal venous hypertension. *J Card Fail*. 2012;18:930–8.
 22. Sweitzer NK, Lopatin M, Yancy CW, Mills RM, Stevenson LW. Comparison of clinical features and outcomes of patients hospitalized with heart failure and normal ejection fraction (> or =55%) versus those with mildly reduced (40% to 55%) and moderately to severely reduced (<40%) fractions. *Am J Cardiol*. 2008;101:1151–6.
 23. Galie N, Humbert M, Vachiery JL, et al. 2015 ESC/ERS guidelines for the diagnosis and treatment of pulmonary hypertension: the joint task force for the diagnosis and treatment of pulmonary hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS). Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). *Eur Heart J*. 2016;37:67–119.
 24. Simonneau G, Montani D, Celermajer DS et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J* 2019;53.
 25. Arrigo M, Huber LC, Winnik S, et al. Right ventricular failure: pathophysiology, diagnosis and treatment. *Card Fail Rev*. 2019;5:140–6.

26. Nickel NP, O'Leary JM, Brittain EL, et al. Kidney dysfunction in patients with pulmonary arterial hypertension. *Pulm Circ*. 2017;7:38–54.
27. Benza RL, Miller DP, Gomberg-Maitland M, et al. Predicting survival in pulmonary arterial hypertension: insights from the registry to evaluate early and long-term pulmonary arterial hypertension disease management (REVEAL). *Circulation*. 2010;122:164–72.
28. Shah SJ, Thenappan T, Rich S, Tian L, Archer SL, Gomberg-Maitland M. Association of serum creatinine with abnormal hemodynamics and mortality in pulmonary arterial hypertension. *Circulation*. 2008;117:2475–83.
29. O'Leary JM, Assad TR, Xu M, et al. Pulmonary hypertension in patients with chronic kidney disease: invasive hemodynamic etiology and outcomes. *Pulm Circ*. 2017;7:674–83.
30. Haddad F, Fuh E, Peterson T, et al. Incidence, correlates, and consequences of acute kidney injury in patients with pulmonary arterial hypertension hospitalized with acute right-side heart failure. *J Card Fail*. 2011;17:533–9.
31. Colombo PC, Doran AC, Onat D, et al. Venous congestion, endothelial and neurohormonal activation in acute decompensated heart failure: cause or effect? *Curr Heart Fail Rep*. 2015;12:215–22.
32. Scheffold JC, Filipatos G, Hasenfuss G, Anker SD, von Haehling S. Heart failure and kidney dysfunction: epidemiology, mechanisms and management. *Nat Rev Nephrol*. 2016;12:610–23.
33. Konstam MA, Kiernan MS, Bernstein D, et al. Evaluation and Management of Right-Sided Heart Failure: A Scientific Statement From the American Heart Association. *Circulation*. 2018;137:e578–622.
34. Gan C, Lankhaar JW, Marcus JT, et al. Impaired left ventricular filling due to right-to-left ventricular interaction in patients with pulmonary arterial hypertension. *Am J Physiol Heart Circ Physiol*. 2006;290:H1528–33.
35. Marcus JT, Vonk Noordegraaf A, Roeleveld RJ, et al. Impaired left ventricular filling due to right ventricular pressure overload in primary pulmonary hypertension: noninvasive monitoring using MRI. *Chest*. 2001;119:1761–5.
36. Dini FL, Demmer RT, Simioniu A, et al. Right ventricular dysfunction is associated with chronic kidney disease and predicts survival in patients with chronic systolic heart failure. *Eur J Heart Fail*. 2012;14:287–94.
37. Damman K, Navis G, Smilde TD, et al. Decreased cardiac output, venous congestion and the association with renal impairment in patients with cardiac dysfunction. *Eur J Heart Fail*. 2007;9:872–8.
38. Firth JD, Raine AE, Ledingham JG. Raised venous pressure: a direct cause of renal sodium retention in oedema? *Lancet*. 1988;1:1033–5.
39. Doty JM, Saggi BH, Blocher CR, et al. Effects of increased renal parenchymal pressure on renal function. *J Trauma*. 2000;48:874–7.
40. Mullens W, Abrahams Z, Francis GS, Taylor DO, Starling RC, Tang WHW. Prompt reduction in intra-abdominal pressure following large-volume mechanical fluid removal improves renal insufficiency in refractory decompensated heart failure. *J Cardiac Fail*. 2008;14:508–14.
41. Frost A, Badesch D, Gibbs JSR et al. Diagnosis of pulmonary hypertension. *Eur Respir J* 2019;53.
42. Verbrugge FH, Grieten L, Mullens W. Management of the cardiorenal syndrome in decompensated heart failure. *Cardiorenal Med*. 2014;4:176–88.
- 43.● Husain-Syed F, Birk HW, Tello K et al. Alterations in Doppler-derived renal venous stasis index during recompensation of right heart failure and fluid overload in a patient with pulmonary hypertension. *Rev Cardiovasc Med* 2019;20:263–266. **This case report demonstrates the feasibility of using renal Doppler ultrasonography to evaluate the treatment response and to guide therapy in patients with right heart failure. In particular, this case highlights the potential role of renal venous stasis index as a simple, non-invasive, and integrative Doppler-derived measure of renal congestion.**
44. Ciozda W, Kedan I, Kehl DW, Zimmer R, Khandwalla R, Kimchi A. The efficacy of sonographic measurement of inferior vena cava diameter as an estimate of central venous pressure. *Cardiovasc Ultrasound*. 2016;14:33.
45. Kellum JA, Lameire N, Aspelin P et al. Kidney disease: improving global outcomes (KDIGO) acute kidney injury work group. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl* 2012;2:1–138.
- 46.● Husain-Syed F, Birk HW, Ronco C et al. Doppler-derived renal venous stasis index in the prognosis of right heart failure. *J Am Heart Assoc* 2019;8:e013584. **The authors propose renal venous stasis index as a conceptually new and integrative Doppler index of renal congestion. Renal venous stasis index provides additional prognostic information to stratify pulmonary hypertension for the propensity to develop right heart failure.**
47. Iida N, Seo Y, Sai S, et al. Clinical implications of intrarenal hemodynamic evaluation by doppler ultrasonography in heart failure. *JACC Heart Fail*. 2016;4:674–82.
- 48.● Nijst P, Martens P, Dupont M, Tang WHW, Mullens W. Intrarenal flow alterations during transition from euvolemia to intravascular volume expansion in heart failure patients. *JACC Heart Fail* 2017;5:672–681. **In this study, patients with a swan ganz catheter received IV fluids to increase circulatory volume. Even before RAP increases, the renal venous outflow becomes discontinuous. The mean circulatory filling pressure (MCFP) is a combination of stressed and unstressed venous blood that generates a venous pressure that also drives venous return to the right heart. The MCFP needs to overcome resistance on its way and is always higher as RAP (flow to right heart = (MCFP-RAP / resistance). This study illustrates that the kidney responses to MCFP even before RAP is elevated. The fact that MCFP can already be elevated emphasizes the unique significance of renal vein ultrasound.**
49. Puzzovivo A, Monitillo F, Guida P et al. Renal venous pattern: a New parameter for predicting prognosis in heart failure outpatients. *J Cardiovasc Dev Dis* 2018;5.
50. Tang WH, Kitai T. Intrarenal venous flow: a window into the congestive kidney failure phenotype of heart failure? *JACC Heart Fail*. 2016;4:683–6.
51. Brisco MA, Testani JM. Novel renal biomarkers to assess cardiorenal syndrome. *Curr Heart Fail Rep*. 2014;11:485–99.
52. Haase M, Kellum JA, Ronco C. Subclinical AKI—an emerging syndrome with important consequences. *Nat Rev Nephrol*. 2012;8:735–9.
53. McDonagh TA, Metra M, Adamo M, et al. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*. 2021;42:3599–726.
54. Maisel AS, Katz N, Hillege HL, et al. Biomarkers in kidney and heart disease. *Nephrol Dial Transplant*. 2011;26:62–74.
- 55.● Goffredo G, Barone R, Di Terlizzi V, Correale M, Brunetti ND, Iacoviello M. Biomarkers in Cardiorenal Syndrome. *J Clin Med* 2021;10:3433. **The author explains that patients with heart failure, chronic kidney disease is highly prevalent and associated with poor outcome. Estimation of renal function by the use of creatinine is limited due to several confounders. The author describes new biomarkers for better assessment of renal dysfunction and to detect patients at higher risk of renal function worsening as well as to more accurately identify patients prone to developing acute kidney injury.**

56. McCullough PA, Duc P, Omland T, et al. B-type natriuretic peptide and renal function in the diagnosis of heart failure: an analysis from the Breathing Not Properly Multinational Study. *Am J Kidney Dis.* 2003;41:571–9.
57. Jiang K, Shah K, Daniels L, Maisel AS. Review on natriuretic peptides: where we are, where we are going. *Expert Opin Med Diagn.* 2008;2:1137–53.
58. Palazzuoli A, Ruocco G, Pellegrini M, et al. Patients with cardiorenal syndrome revealed increased neurohormonal activity, tubular and myocardial damage compared to heart failure patients with preserved renal function. *Cardiorenal Med.* 2014;4:257–68.
59. Evaluation and Management of Chronic Kidney Disease. Synopsis of the Kidney Disease: Improving Global Outcomes 2012 Clinical Practice Guideline. *Ann Intern Med.* 2013;158:825–30.
60. Colbert G, Jain N, de Lemos JA, Hedayati SS. Utility of traditional circulating and imaging-based cardiac biomarkers in patients with predialysis CKD. *Clin J Am Soc Nephrol.* 2015;10:515–29.
61. Tang WH, Shrestha K, Shao Z, et al. Usefulness of plasma galectin-3 levels in systolic heart failure to predict renal insufficiency and survival. *Am J Cardiol.* 2011;108:385–90.
62. Rubinstein N, Ilarregui JM, Toscano MA, Rabinovich GA. The role of galectins in the initiation, amplification and resolution of the inflammatory response. *Tissue Antigens.* 2004;64:1–12.
63. Sharma UC, Pokharel S, van Brakel TJ, et al. Galectin-3 marks activated macrophages in failure-prone hypertrophied hearts and contributes to cardiac dysfunction. *Circulation.* 2004;110:3121–8.
64. Henderson NC, Mackinnon AC, Farnworth SL, et al. Galectin-3 expression and secretion links macrophages to the promotion of renal fibrosis. *Am J Pathol.* 2008;172:288–98.
65. Okamura DM, Pasichnyk K, Lopez-Guisa JM, et al. Galectin-3 preserves renal tubules and modulates extracellular matrix remodeling in progressive fibrosis. *Am J Physiol Renal Physiol.* 2011;300:F245–53.
66. Vergaro G, Del Franco A, Giannoni A, et al. Galectin-3 and myocardial fibrosis in nonischemic dilated cardiomyopathy. *Int J Cardiol.* 2015;184:96–100.
67. de Boer RA, Lok DJ, Jaarsma T, et al. Predictive value of plasma galectin-3 levels in heart failure with reduced and preserved ejection fraction. *Ann Med.* 2011;43:60–8.
68. McEvoy JW, Chen Y, Halushka MK et al. Galectin-3 and Risk of Heart Failure and Death in Blacks and Whites. *J Am Heart Assoc* 2016;5.
69. Grande D, Leone M, Rizzo C et al. A multiparametric approach based on NT-proBNP, ST2, and Galectin3 for stratifying One year prognosis of chronic heart failure outpatients. *J Cardiovasc Dev Dis* 2017;4.
70. Iacoviello M, Aspromonte N, Leone M, et al. Galectin-3 serum levels Are independently associated with microalbuminuria in chronic heart failure outpatients. *Res Cardiovasc Med.* 2016;5:e28952.
71. Iacoviello M, Di Serio F, Rizzo C, et al. Association between high Gal-3 serum levels and worsening of renal function in chronic heart failure outpatients. *Biomark Med.* 2019;13:707–13.
72. Leone M, Iacoviello M. The predictive value of plasma biomarkers in discharged heart failure patients: role of galectin-3. *Minerva Cardioangiol.* 2016;64:181–94.
73. O’Seaghdha CM, Hwang SJ, Ho JE, Vasan RS, Levy D, Fox CS. Elevated galectin-3 precedes the development of CKD. *J Am Soc Nephrol.* 2013;24:1470–7.
74. Rebholz CM, Selvin E, Liang M, et al. Plasma galectin-3 levels are associated with the risk of incident chronic kidney disease. *Kidney Int.* 2018;93:252–9.
75. Suthahar N, Meijers WC, Silljé HHW, Ho JE, Liu FT, de Boer RA. Galectin-3 activation and inhibition in heart failure and cardiovascular disease: an update. *Theranostics.* 2018;8:593–609.
76. Knight EL, Verhave JC, Spiegelman D, et al. Factors influencing serum cystatin C levels other than renal function and the impact on renal function measurement. *Kidney Int.* 2004;65:1416–21.
77. Dupont M, Shrestha K, Singh D, Finucan M, Tang WH. Lack of concordance in defining worsening renal function by rise in creatinine vs rise in cystatin C. *Congest Heart Fail.* 2013;19:E17–21.
78. Arimoto T, Takeishi Y, Niizeki T, et al. Cystatin C, a novel measure of renal function, is an independent predictor of cardiac events in patients with heart failure. *J Card Fail.* 2005;11:595–601.
79. Shlipak MG, Katz R, Fried LF, et al. Cystatin-C and mortality in elderly persons with heart failure. *J Am Coll Cardiol.* 2005;45:268–71.
80. Lassus J, Harjola VP, Sund R, et al. Prognostic value of cystatin C in acute heart failure in relation to other markers of renal function and NT-proBNP. *Eur Heart J.* 2007;28:1841–7.
81. Masson S, Latini R, Milani V, et al. Prevalence and prognostic value of elevated urinary albumin excretion in patients with chronic heart failure: data from the GISSI-Heart Failure trial. *Circ Heart Fail.* 2010;3:65–72.
82. Jackson CE, Solomon SD, Gerstein HC, et al. Albuminuria in chronic heart failure: prevalence and prognostic importance. *Lancet.* 2009;374:543–50.
83. Brinkkoetter PT, Ising C, Benzinger T. The role of the podocyte in albumin filtration. *Nat Rev Nephrol.* 2013;9:328–36.
84. Comper WD, Hilliard LM, Nikolic-Paterson DJ, Russo LM. Disease-dependent mechanisms of albuminuria. *Am J Physiol Renal Physiol.* 2008;295:F1589–600.
85. Miller WG, Bruns DE, Hortin GL, et al. Current issues in measurement and reporting of urinary albumin excretion. *Clin Chem.* 2009;55:24–38.
86. Webster AC, Nagler EV, Morton RL, Masson P. Chronic kidney disease. *Lancet.* 2017;389:1238–52.
87. van de Wal RM, Asselbergs FW, Plokker HW, et al. High prevalence of microalbuminuria in chronic heart failure patients. *J Card Fail.* 2005;11:602–6.
88. Schmidt-Ott KM, Mori K, Li JY, et al. Dual action of neutrophil gelatinase-associated lipocalin. *J Am Soc Nephrol.* 2007;18:407–13.
89. Waring WS, Moonie A. Earlier recognition of nephrotoxicity using novel biomarkers of acute kidney injury. *Clin Toxicol (Phila).* 2011;49:720–8.
90. van Deursen VM, Damman K, Voors AA, et al. Prognostic value of plasma neutrophil gelatinase-associated lipocalin for mortality in patients with heart failure. *Circ Heart Fail.* 2014;7:35–42.
91. Imoto Y, Wakasaki A, Izumida K, et al. Analysis of the diagnostic capabilities of urinary neutrophil gelatinase-associated lipocalin and serum procalcitonin for acute kidney injury at the early stage of critical care intensive care unit admission. *J Clin Lab Anal.* 2021;35:e23852.
92. Collins SP, Hart KW, Lindsell CJ, et al. Elevated urinary neutrophil gelatinase-associated lipocalin after acute heart failure treatment is associated with worsening renal function and adverse events. *Eur J Heart Fail.* 2012;14:1020–9.
93. Haase M, Bellomo R, Devarajan P, Schlattmann P, Haase-Fielitz A. Accuracy of neutrophil gelatinase-associated lipocalin (NGAL) in diagnosis and prognosis in acute kidney injury: a systematic review and meta-analysis. *Am J Kidney Dis.* 2009;54:1012–24.
94. Bazzi C, Petrini C, Rizza V, et al. Urinary N-acetyl-beta-glucosaminidase excretion is a marker of tubular cell dysfunction and a predictor of outcome in primary glomerulonephritis. *Nephrol Dial Transplant.* 2002;17:1890–6.
95. Liangos O, Perianayagam MC, Vaidya VS, et al. Urinary N-acetyl-beta-(D)-glucosaminidase activity and kidney injury molecule-1 level are associated with adverse outcomes in acute renal failure. *J Am Soc Nephrol.* 2007;18:904–12.

96. Damman K, Masson S, Hillege HL, et al. Clinical outcome of renal tubular damage in chronic heart failure. *Eur Heart J*. 2011;32:2705–12.
97. Damman K, Van Veldhuisen DJ, Navis G, et al. Tubular damage in chronic systolic heart failure is associated with reduced survival independent of glomerular filtration rate. *Heart (British Cardiac Society)*. 2010;96:1297–302.
98. Jungbauer CG, Birner C, Jung B, et al. Kidney injury molecule-1 and N-acetyl- β -D-glucosaminidase in chronic heart failure: possible biomarkers of cardiorenal syndrome. *Eur J Heart Fail*. 2011;13:1104–10.
99. Han WK, Wagener G, Zhu Y, Wang S, Lee HT. Urinary biomarkers in the early detection of acute kidney injury after cardiac surgery. *Clin J Am Soc Nephrol*. 2009;4:873–82.
100. Damman K, Ng Kam Chuen MJ, MacFadyen RJ et al. Volume status and diuretic therapy in systolic heart failure and the detection of early abnormalities in renal and tubular function. *J Am Coll Cardiol* 2011;57:2233–41.
101. Thöni S, Mayer G. Urinary Dickkopf-3 and kidney injury in patients with chronic pulmonary disease. *Kidney Int*. 2021;100:983–5.
102. Sánchez-Álamo B, García-Iñigo FJ, Shabaka A, et al. Urinary Dickkopf-3: a new biomarker for CKD progression and mortality. *Nephrol Dial Transplant*. 2021;36:2199–207.
103. Zewinger S, Rauen T, Rudnicki M, et al. Dickkopf-3 (DKK3) in urine identifies patients with short-term risk of eGFR loss. *J Am Soc Nephrol*. 2018;29:2722–33.
104. Schunk SJ, Speer T, Petrakis I, Fliser D. Dickkopf 3-a novel biomarker of the “kidney injury continuum.” *Nephrol Dial Transplant*. 2021;36:761–7.
105. Fonarow GC, Corday E. Overview of acutely decompensated congestive heart failure (ADHF): a report from the ADHERE registry. *Heart Fail Rev*. 2004;9:179–85.
106. Felker GM, Lee KL, Bull DA, et al. Diuretic strategies in patients with acute decompensated heart failure. *N Engl J Med*. 2011;364:797–805.
107. Chitturi C, Novak JE. Diuretics in the Management of Cardiorenal Syndrome. *Adv Chronic Kidney Dis*. 2018;25:425–33.
108. Verma D, Firoz A, Garlapati SKP, et al. Emerging treatments of cardiorenal syndrome: an update on pathophysiology and management. *Cureus*. 2021;13:e17240.
109. Rubinstein J, Sanford D. Treatment of cardiorenal syndrome. *Cardiol Clin*. 2019;37:267–73.
110. Knauf H, Mutschler E. Functional state of the nephron and diuretic dose-response—rationale for low-dose combination therapy. *Cardiology*. 1994;84(Suppl 2):18–26.
111. Bart BA, Goldsmith SR, Lee KL, et al. Ultrafiltration in decompensated heart failure with cardiorenal syndrome. *N Engl J Med*. 2012;367:2296–304.
112. Mullens W, Verbrugge FH, Nijst P, et al. Rationale and design of the ADVOR (Acetazolamide in Decompensated Heart Failure with Volume Overload) trial. *Eur J Heart Fail*. 2018;20:1591–600.
113. Mullens W, Dauw J, Martens P et al. Acetazolamide in decompensated heart failure with volume overload trial (ADVOR): baseline characteristics. *Eur J Heart Fail* 2022.
114. Rao VS, Ahmad T, Brisco-Bacik MA, et al. Renal effects of intensive volume removal in heart failure patients with preexisting worsening renal function. *Circ Heart Fail*. 2019;12:e005552.
115. KDIGO. Clinical practice guideline for the management of glomerular diseases. *Kidney Int*. 2021;2021(100):S1–s276.
116. Testani JM, Chen J, McCauley BD, Kimmel SE, Shannon RP. Potential effects of aggressive decongestion during the treatment of decompensated heart failure on renal function and survival. *Circulation*. 2010;122:265–72.
117. Costanzo MR, Guglin ME, Saltzberg MT, et al. Ultrafiltration versus intravenous diuretics for patients hospitalized for acute decompensated heart failure. *J Am Coll Cardiol*. 2007;49:675–83.
118. Metra M, Davison B, Bettari L, et al. Is worsening renal function an ominous prognostic sign in patients with acute heart failure? The role of congestion and its interaction with renal function. *Circ Heart Fail*. 2012;5:54–62.
119. Valente MA, Voors AA, Damman K, et al. Diuretic response in acute heart failure: clinical characteristics and prognostic significance. *Eur Heart J*. 2014;35:1284–93.
120. Aronson D, Burger AJ. Diuretic response: clinical and hemodynamic predictors and relation to clinical outcome. *J Card Fail*. 2016;22:193–200.
121. Verbrugge FH, Dupont M, Bertrand PB, et al. Determinants and impact of the natriuretic response to diuretic therapy in heart failure with reduced ejection fraction and volume overload. *Acta Cardiol*. 2015;70:265–73.
122. Krämer BK, Schweda F, Riegger GA. Diuretic treatment and diuretic resistance in heart failure. *Am J Med*. 1999;106:90–6.
123. Mulder H, Schopman W Jr, van der Lely AJ, Schopman W Sr. Acute changes in plasma renin activity, plasma aldosterone concentration and plasma electrolyte concentrations following furosemide administration in patients with congestive heart failure—interrelationships and diuretic response. *Horm Metab Res*. 1987;19:80–3.
124. Kaissling B, Bachmann S, Kriz W. Structural adaptation of the distal convoluted tubule to prolonged furosemide treatment. *Am J Physiol*. 1985;248:F374–81.
125. Ellison DH. The physiologic basis of diuretic synergism: its role in treating diuretic resistance. *Ann Intern Med*. 1991;114:886–94.
126. Komlosi P, Fintha A, Bell PD. Current mechanisms of macula densa cell signalling. *Acta Physiol Scand*. 2004;181:463–9.
127. Brater DC. Update in diuretic therapy: clinical pharmacology. *Semin Nephrol*. 2011;31:483–94.
128. Rudy DW, Gehr TW, Matzke GR, Kramer WG, Sica DA, Brater DC. The pharmacodynamics of intravenous and oral torsemide in patients with chronic renal insufficiency. *Clin Pharmacol Ther*. 1994;56:39–47.
129. Gehr TW, Rudy DW, Matzke GR, Kramer WG, Sica DA, Brater DC. The pharmacokinetics of intravenous and oral torsemide in patients with chronic renal insufficiency. *Clin Pharmacol Ther*. 1994;56:31–8.
130. Aronson D. The complexity of diuretic resistance. *Eur J Heart Fail*. 2017;19:1023–6.
131. Trullàs JC, Morales-Rull JL, Casado J, Freitas Ramírez A, Manzano L, Formiga F. Rationale and design of the “safety and efficacy of the combination of loop with thiazide-type diuretics in patients with decompensated heart failure (CLOROTIC) trial:” a double-blind, randomized, placebo-controlled study to determine the effect of combined diuretic therapy (loop diuretics with thiazide-type diuretics) among patients with decompensated heart failure. *J Card Fail*. 2016;22:529–36.
132. Schulze PC, Bogoviku J, Westphal J et al. Effects of early empagliflozin initiation on diuresis and kidney function in patients with acute decompensated heart failure (EMPAG-HF). *Circulation* 2022;101161circulationaha122059038.
133. Damman K, Beusekamp JC, Boorsma EM, et al. Randomized, double-blind, placebo-controlled, multicentre pilot study on the effects of empagliflozin on clinical outcomes in patients with acute decompensated heart failure (EMPA-RESPONSE-AHF). *Eur J Heart Fail*. 2020;22:713–22.
134. Mullens W, Damman K, Testani JM, et al. Evaluation of kidney function throughout the heart failure trajectory - a position statement from the heart failure association of the European society of cardiology. *Eur J Heart Fail*. 2020;22:584–603.
135. Rossignol P, Lainscak M, Crespo-Leiro MG et al. Unravelling the interplay between hyperkalaemia, renin-angiotensin-aldosterone inhibitor use and clinical outcomes. Data from 9222 chronic heart

- failure patients of the ESC-HFA-EORP Heart Failure Long-Term Registry. *Eur J Heart Failure* 2020;22:1378–1389.
136. Voors AA, Mulder H, Reyes E et al. Renal function and the effects of vericiguat in patients with worsening heart failure with reduced ejection fraction: insights from the VICTORIA (Vericiguat Global Study in Subjects with HFrEF) trial. *Eur J Heart Failure* 2021;23:1313–1321. **This article demonstrates that renal function trajectories were similar between vericiguat- and placebo-treated patients and the beneficial effects of vericiguat on the primary outcome were consistent across the full range of eGFR and irrespective of worsening renal function.**
 137. Clark H, Krum H, Hopper I. Worsening renal function during renin-angiotensin-aldosterone system inhibitor initiation and long-term outcomes in patients with left ventricular systolic dysfunction. *Eur J Heart Fail.* 2014;16:41–8.
 138. Szymanski MK, Damman K, van Veldhuisen DJ, van Gilst WH, Hillege HL, de Boer RA. Prognostic value of renin and prorenin in heart failure patients with decreased kidney function. *Am Heart J.* 2011;162:487–93.
 139. Poletti R, Vergaro G, Zyw L, Prontera C, Passino C, Emdin M. Prognostic value of plasma renin activity in heart failure patients with chronic kidney disease. *Int J Cardiol.* 2013;167:711–5.
 140. Parikh CR, Coca SG. “Permissive AKI” with treatment of heart failure. *Kidney Int.* 2019;96:1066–8.
 141. Armstrong PW, Pieske B, Anstrom KJ, et al. Vericiguat in patients with heart failure and reduced ejection fraction. *N Engl J Med.* 2020;382:1883–93.
 142. Armstrong PW, Roessig L, Patel MJ, et al. A Multicenter, randomized, double-blind, placebo-controlled trial of the efficacy and safety of the oral soluble guanylate cyclase stimulator: the VICTORIA trial. *JACC Heart Fail.* 2018;6:96–104.
 143. Pieske B, Patel MJ, Westerhout CM, et al. Baseline features of the VICTORIA (vericiguat global study in subjects with heart failure with reduced ejection fraction) trial. *Eur J Heart Fail.* 2019;21:1596–604.
 144. Packer M, Anker SD, Butler J, et al. Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med.* 2020;383:1413–24.
 145. McMurray JJV, DeMets DL, Inzucchi SE, et al. The dapagliflozin and prevention of adverse-outcomes in heart failure (DAPA-HF) trial: baseline characteristics. *Eur J Heart Fail.* 2019;21:1402–11.
 146. Drazner MH. SGLT2 inhibition in heart failure with a preserved ejection fraction - a Win against a formidable Foe. *N Engl J Med.* 2021;385:1522–4.
 147. Nassif ME, Windsor SL, Borlaug BA, et al. The SGLT2 inhibitor dapagliflozin in heart failure with preserved ejection fraction: a multicenter randomized trial. *Nat Med.* 2021;27:1954–60.
 148. Kalra R, Duval S, Thenappan T, et al. Comparison of balloon pulmonary angioplasty and pulmonary vasodilators for inoperable chronic thromboembolic pulmonary hypertension: a systematic review and meta-analysis. *Sci Rep.* 2020;10:8870.
 149. Wilcox CS. Antihypertensive and Renal Mechanisms of SGLT2 (Sodium-Glucose Linked Transporter 2) Inhibitors. *Hypertension.* 2020;75:894–901.
 150. Ghezzi C, Yu AS, Hirayama BA, et al. Dapagliflozin binds specifically to sodium-glucose cotransporter 2 in the proximal renal tubule. *J Am Soc Nephrol.* 2017;28:802–10.
 151. Chu C, Lu YP, Yin L, Hoher B. The SGLT2 inhibitor empagliflozin might Be a New approach for the prevention of acute kidney injury. *Kidney Blood Press Res.* 2019;44:149–57.
 152. Zelniker TA, Braunwald E. Cardiac and Renal Effects of Sodium-Glucose Co-Transporter 2 Inhibitors in Diabetes: JACC State-of-the-Art Review. *J Am Coll Cardiol.* 2018;72:1845–55.
 153. Scheen AJ. An update on the safety of SGLT2 inhibitors. *Expert Opin Drug Saf.* 2019;18:295–311.
 154. Ata F, Yousaf Z, Khan AA, et al. SGLT-2 inhibitors associated euglycemic and hyperglycemic DKA in a multicentric cohort. *Sci Rep.* 2021;11:10293.
 155. Vinod P, Krishnappa V, Chauvin AM, Khare A, Raina R. Cardiorenal syndrome: role of arginine vasopressin and vaptans in heart failure. *Cardiol Res.* 2017;8:87–95.
 156. Costello-Boerrigter LC, Boerrigter G, Burnett JC Jr. V2 receptor antagonism with tolvaptan in heart failure. *Expert Opin Investig Drugs.* 2007;16:1639–47.
 157. Obi Y, Kim T, Kovesdy CP, Amin AN, Kalantar-Zadeh K. Current and potential therapeutic strategies for hemodynamic cardiorenal syndrome. *Cardiorenal Med.* 2016;6:83–98.
 158. Konstam MA, Gheorghiadu M, Burnett JC Jr, et al. Effects of oral tolvaptan in patients hospitalized for worsening heart failure: the EVEREST outcome trial. *JAMA.* 2007;297:1319–31.
 159. Aghel A, Tang WH. Tolvaptan: the evidence for its therapeutic value in acute heart failure syndrome. *Core Evid.* 2008;3:31–43.
 160. Bielecka-Dabrowa A, Godoy B, Schefold JC, Koziolok M, Banach M, von Haehling S. Decompensated Heart Failure and Renal Failure: What Is the Current Evidence? *Curr Heart Fail Rep.* 2018;15:224–38.
 161. Webb DJ, Vachiery JL, Hwang LJ, Maurey JO. Sildenafil improves renal function in patients with pulmonary arterial hypertension. *Br J Clin Pharmacol.* 2015;80:235–41.
 162. Harjola VP, Mebazaa A, Celutkienė J, et al. Contemporary management of acute right ventricular failure: a statement from the Heart Failure Association and the Working Group on Pulmonary Circulation and Right Ventricular Function of the European Society of Cardiology. *Eur J Heart Fail.* 2016;18:226–41.
 163. Kramer T, Dumitrescu D, Gerhardt F, et al. Therapeutic potential of phosphodiesterase type 5 inhibitors in heart failure with preserved ejection fraction and combined post- and pre-capillary pulmonary hypertension. *Int J Cardiol.* 2019;283:152–8.
 164. Vachiéry JL, Adir Y, Barberà JA, et al. Pulmonary hypertension due to left heart diseases. *J Am Coll Cardiol.* 2013;62:D100–8.
 165. Rosenkranz S, Gibbs JS, Wachter R, De Marco T, Vonk-Noordegraaf A, Vachiery JL. Left ventricular heart failure and pulmonary hypertension. *Eur Heart J.* 2016;37:942–54.
 166. Guazzi M, Naeije R. Pulmonary hypertension in heart failure: pathophysiology, pathobiology, and emerging clinical perspectives. *J Am Coll Cardiol.* 2017;69:1718–34.
 167. Guazzi M, Ghio S, Adir Y. Pulmonary hypertension in HFpEF and HFrEF: JACC review topic of the week. *J Am Coll Cardiol.* 2020;76:1102–11.
 168. Redfield MM, Chen HH, Borlaug BA, et al. Effect of phosphodiesterase-5 inhibition on exercise capacity and clinical status in heart failure with preserved ejection fraction: a randomized clinical trial. *JAMA.* 2013;309:1268–77.
 169. Hoendermis ES, Liu LC, Hummel YM, et al. Effects of sildenafil on invasive haemodynamics and exercise capacity in heart failure patients with preserved ejection fraction and pulmonary hypertension: a randomized controlled trial. *Eur Heart J.* 2015;36:2565–73.
 170. Opitz CF, Hoepfer MM, Gibbs JS, et al. Pre-capillary, combined, and post-capillary pulmonary hypertension: a pathophysiological continuum. *J Am Coll Cardiol.* 2016;68:368–78.
 171. Guazzi M, Vicenzi M, Arena R, Guazzi MD. Pulmonary hypertension in heart failure with preserved ejection fraction: a target of phosphodiesterase-5 inhibition in a 1-year study. *Circulation.* 2011;124:164–74.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.